ORGANIC CHEMISTRY







FRONTIERS

CORRECTION

View Article Online
View Journal | View Issue



Cite this: *Org. Chem. Front.*, 2020, **7**, 3842

Correction: A removable directing group-assisted Rh(III)-catalyzed direct C-H bond activation/annulation cascade to synthesize highly fused isoquinolines

Yilang Cheng, a,b,c Xu Han, a,b Junyou Li, a,b Yu Zhou*a,b and Hong Liu*a,b,c

DOI: 10.1039/d0qo90077j

rsc.li/frontiers-organic

Correction for 'A removable directing group-assisted Rh(III)-catalyzed direct C-H bond activation/annulation cascade to synthesize highly fused isoquinolines' by Yilang Cheng *et al.*, *Org. Chem. Front.*, 2020, **7**, 3186–3192, DOI: 10.1039/D0QO00786B.

The authors regret that some related work was not cited in the original manuscript.

The text "we serendipitously detected that the azomethine imine group could be used as a removable DG when treating aryl azomethine imines (1) with 4-diazoisochroman-3-imine (2) in the presence of a Rh(\mathfrak{m})-complex catalyst, in which the highly fused isoquinoline scaffolds could be effectively and facilely constructed via a C-H activation and annulation cascade (Scheme 1d)." should read "we serendipitously detected that the azomethine imine group could be used as a removable DG when treating aryl azomethine imines (1) with 4-diazoisochroman-3-imine (2), an intriguing substrate for the synthesis of isochroman polycyclic scaffolds, ^{1,2} in the presence of a Rh(\mathfrak{m})-complex catalyst, in which the highly fused isoquinoline scaffolds could be effectively and facilely constructed via a C-H activation and annulation cascade (Scheme 1d)."

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

References

- 1 A. Ren, P. Lu and Y. Wang, Convenient preparation of 4-diazoisochroman-3-imines and 3-substituted 3,5-dihydroisochromeno [3,4-d][1,2,3]triazoles, *Chem. Commun.*, 2017, 53, 3769–3772.
- 2 Z. Li, J. Xie, P. Lu and Y. Wang, Synthesis of 8-Alkoxy-5H-isochromeno[3,4-c]isoquinolines and 1-Alkoxy-4-arylisoquinolin-3-ols through Rh(III)-Catalyzed C–H Functionalization of Benzimidates with 4-Diazoisochroman-3-imines and 4-Diazoisoquinolin-3-ones, *J. Org. Chem.*, 2020, **85**, 5525–5535.

^aState Key Laboratory of Drug Research and CAS Key Laboratory of Receptor Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 555 Zu Chong Zhi Road, Shanghai, 201203, China. E-mail: zhouyu@simm.ac.cn, hliu@simm.ac.cn

^bUniversity of Chinese Academy of Sciences, No. 19A Yuquan Road, Beijing 100049, China

^cSchool of Life Science and Technology, ShanghaiTech University, 100 Haike Road, Shanghai 201210, China