



 Cite this: *RSC Adv.*, 2020, 10, 338

 Received 26th October 2019
 Accepted 15th December 2019

DOI: 10.1039/c9ra08792c

rsc.li/rsc-advances

Palladium mediated domino reaction: synthesis of isochromenes under aqueous medium†

 Lavisha Punia, Karu Ramesh and Gedu Satyanarayana *

Isochromenes have been synthesized using palladium-catalyzed C–C and C–O bond forming reactions starting from *ortho*-bromo tertiary benzylic alcohols and internal acetylenes. Notably, this domino process is feasible by using the green solvent, water. The protocol exhibited a broad substrate scope and afforded various isochromenes.

Introduction

Domino reactions are effective in the construction of two or more bonds and thereby facilitate obtaining products with reasonable structural complexity, under a particular set of reaction conditions, which may be devoid of additional reagents/catalysts.¹ Moreover, in such a reaction, there is no need to isolate the intermediate product(s). Mainly, domino reactions play an important role in achieving complex hetero- and carbo-cyclic products, which are of great importance in the realm of biologically active molecules. Amongst them, fused bicyclic ethers (isochromenes) consisting of an aromatic ring and a pyran moiety are widely available heterocyclic motifs comprising natural products of biological significance.² They are often identified in bacteria, fungi, plants, *etc.*³ and exhibit useful biological properties, such as antitumor,⁴ anticancer,⁵ antifungal,⁶ antibacterial, antibiotic,⁷ and antimicrobial activities.⁸ Some of the representative examples of natural products are as depicted in Fig. 1.^{9–12} Due to their interesting structural features and biological properties, isochromenes have captured the attention of chemists.¹³ As a consequence, a good number of synthetic routes were developed towards the synthesis of isochromenes.¹⁴ Especially, palladium-catalyzed annulation using relatively more reactive *ortho*-iodo tertiary benzylic alcohols with internal alkynes was also reported.¹⁵ These annulations were also driven by means of a stable palladacycle catalyst, as reported in ref. 16. Nevertheless, very few examples have been synthesized in these above two reports. In this regard, to the best of our knowledge, there are no reports of, particularly, using relatively less sensitive *ortho*-bromo tertiary benzylic alcohols catalyzed by a simple palladium-based catalyst and by employing water as the whole solvent medium, for the synthesis of isochromenes. In our efforts to establish metal-catalyzed

coupling reactions,¹⁷ herein, we describe the synthesis of isochromenes by utilizing the solvent water as the reaction medium.

Results and discussion

The optimization was begun with *ortho*-bromo tertiary benzylic alcohol **1a** and diphenylacetylene **2a**. In general, quaternary ammonium salts (QX; X = Cl, Br, I and HSO₄) can act as a phase transfer catalyst mostly when the solvent is water or also can act as an additive to accelerate the catalytic activity of metal-catalyst.¹⁸ With this viewpoint in our mind, firstly, the alcohol **1a** and diphenylacetylene **2a** were treated at different temperatures using Pd(OAc)₂ catalyst (5 mol%), tetrabutylammonium iodide (TBAI) (1 equiv.), base K₂CO₃ (4 equiv.) and with different ligands (entries 1 to 3, Table 1). Conversely, no progress of the reaction was noted. While treating **1a** and **2a** by using Xantphos ligand at 140 °C for 36 h, resulted the expected chromene **3aa**, although in poor yield (entry 4, Table 1). Gratifyingly, the combination of Pd(OAc)₂ and *L*-proline, TBAI, and base K₂CO₃ resulted in 76% yield of the desired chromene **3aa** (entry 5,




Fig. 1 Isochromene based natural products.

Department of Chemistry, Indian Institute of Technology (IIT) Hyderabad, Kandi – 502 285, Sangareddy District, Telangana, India. E-mail: gvsatya@iith.ac.in; Web: <https://sites.google.com/site/gsresearchgrouphomepage/home>; Fax: +91 40 2301 6032

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra08792c



Table 1 Optimizations of Pd-catalyzed domino reaction between substituted *ortho*-bromo tertiary benzylic alcohol **1a** and internal alkyne **2a**^{a,b,c,d,e,f,g}



Entry	Ligand (20 mol%)	Quaternary ammonium salt (equiv.) & base (equiv.)	Temp. (°C)	Time (h)	Yield of 3aa ^a (%)
1	BINAP	TBAI (1) + K ₂ CO ₃ (4)	140	36	— ^b
2	2,2'-Bipy	TBAI (1) + K ₂ CO ₃ (4)	120	38	— ^b
3	P(Cy) ₃	TBAI (1) + K ₂ CO ₃ (4)	100	42	— ^b
4	Xantphos	TBAI (1) + K ₂ CO ₃ (4)	140	36	30
5	l-Proline	TBAI (1) + K₂CO₃ (4)	140	36	76
6	l-Proline	TBAI (1) + Na ₂ CO ₃ (4)	140	38	20
7	l-Proline	TBAI (1) + Li ₂ CO ₃ (4)	140	36	15
8	l-Proline	BTEAC (1) + K ₂ CO ₃ (4)	140	36	15
9 ^c	l-Proline	TBAI (1) + K ₂ CO ₃ (4)	140	36	50
10 ^d	l-Proline	TBAI (1) + K ₂ CO ₃ (4)	140	36	— ^b
11 ^e	l-Proline	TBAI (1) + K ₂ CO ₃ (4)	140	36	— ^b
12	l-Proline	TBAI (1) + Na ₂ CO ₃ (2)	140	46	— ^f
13	l-Proline	TBAI (1) + K ₂ CO ₃ (2)	140	42	30
14	l-Proline	TBAI (1) + NEt ₃ (4)	140	36	— ^b

^a Conditions: **1a** (54 mg, 0.25 mmol), **2a** (111.2 mg, 0.625 mmol), Pd(OAc)₂ (2.8 mg, 0.012 mmol), base (1 mmol), ligand (0.05 mmol), quaternary ammonium salt (0.25 mmol), water (0.5 mL), 140 °C, 36 h. ^b Isolated yields of the product **3aa**. ^c Lead to the decomposition. ^d Used 0.2 mL of water. ^e The reaction was performed using 0.1 mL of water. ^f Used neat reaction conditions. ^g Starting materials were recovered. TBAI: tetrabutylammonium iodide. BTEAC: benzyltriethylammonium chloride.

Table 1). Whereas other attempts with the combination of tetrabutylammonium iodide (TBAI/Na₂CO₃, TBAI/Li₂CO₃) and benzyltriethylammonium chloride (BTEAC)/K₂CO₃ in the presence of l-proline as the ligand were not that much efficient (entries 6 to 8, Table 1). Further, decreasing the quantity of solvent (0.2 mL of water), gave **3aa** in 50% yield (entry 9, Table 1). While with 0.1 mL solvent water (entry 10, Table 1) or under neat reaction conditions (entry 11, Table 1), lead to the decomposition. On the other hand, when Na₂CO₃ (2 equiv.) was used as the base, there was no initiation in the reaction (entry 12, Table 1). While 30% yield of **3aa** was formed with 2 equiv. of K₂CO₃ (entry 13, Table 1). Treatment of **1a** and **2a** with 4 equiv. of Et₃N, lead to the decomposition (entry 14, Table 1).

With the above best-optimized conditions (entry 5, Table 1), to test the feasibility of the process, it was decided to test the protocol with other substrates. Thus, *ortho*-bromo tertiary benzylic alcohols **1a–1c** were reacted with different internal alkynes **2a–2k**, under standard conditions. Interestingly, this protocol was quite successful and afforded corresponding isochromenes **3aa–3cg** (Table 2). In addition, the protocol exhibited good compatibility with various substituents bearing aromatic rings of acetylenes. For example, the reaction was also smooth on internal alkyne flanked to a heteroaromatic ring (**3ad**, Table 2). Also, furnished the products **3ah** and **3ai** bearing electron deactivating F and CF₃ moieties. Besides, when *ortho*-bromo tertiary benzylic alcohol **1a** was subjected to the reaction

using unsymmetrical acetylenes **2j** and **2k**, as anticipated, furnished regioisomeric mixtures [(**3aj** : **3aj'** in 4 : 3 ratio) and (**3ak** : **3ak'** in 1 : 1 ratio) (Table 2)].


Furthermore, to establish the usefulness of the strategy, this time, the reaction was tested with the aromatic variant of *ortho*-bromo tertiary benzylic alcohols **1d–1g** with internal alkynes **2a–2f**. In general, the reaction exhibited a broad substrate scope and delivered the products **3da–3ga** (Table 3). The protocol was tolerable to different functionalities of either aromatic rings. Notably, the strategy was found to be smooth with electron-donating OMe groups of *ortho*-bromo tertiary benzylic alcohols **1d–1f** and also electron-withdrawing Cl group of *ortho*-bromo tertiary benzylic alcohol **1g**. On the other hand, amenable to simple phenyl, tolyl, anisyl, and thiophenyl rings of internal alkynes as well.

Furthermore, to show the effectiveness of the protocol, the annulation of *ortho*-bromo secondary benzylic alcohol **1h** was attempted with internal alkynes (**2a** & **2h**), using optimized conditions. Notably, as anticipated, afforded the cyclic ethers **3ha** and **3hh** in 41% and 44% yields, respectively (Table 4).

A plausible mechanistic path of this annulation process is as depicted in Scheme 1.

Initially, Pd(II) species **A** could be formed *via* oxidative incorporation of active Pd(0) catalyst onto the sigma C–Br bond of *ortho*-bromo tertiary benzylic alcohols **1**. The subsequent, *syn*-addition reaction of **A** upon C≡C bond of alkynes **2**, gives



Table 2 Scope to generate isochromenes **3aa–3cg** from *ortho*-bromo tertiary benzylic alcohols **1a–1c** and internal alkynes **2a–2k**^{a,b,c}

^a Conditions: **1a–1c** (0.25 mmol, 1 equiv.), **2a–2k** (0.625 mmol, 2.5 equiv.), Pd(OAc)₂ (2.8 mg, 0.012 mmol, 5 mol%), K₂CO₃ (138.6 mg, 1 mmol, 4 equiv.), L-proline (5.6 mg, 0.05 mol, 20 mol%), TBAI (tetrabutylammonium iodide) (92.3 mg, 0.25 mmol, 1 equiv.), 140 °C, 36 h. ^b Yields are isolated pure products **3aa–3cg**. ^c Determined by NMR analysis.

acyclic Pd(II)-species **B**, which on subsequent intramolecular coupling with the nucleophilic hydroxyl group, would furnish a seven-membered palladacycle **C** [Pd(II)]. Removal of HBr from **C** could yield seven-membered Pd(II)-cycle **D**. Ultimately, reductive elimination of **D**, gives isochromenes **3** along with the regeneration of Pd(0) active catalyst and hence, fulfils the overall catalytic path of the reaction.¹⁹

Conclusions


In summary, a palladium-catalyst promoted annulation between *ortho*-bromo benzylic alcohols and internal acetylenes is established, for achieving isochromenes. Significantly, this dual bond-forming process was feasible in water solvent and enabled the formation of various isochromenes.

Experimental section

General

IR spectra were recorded on a Bruker Tensor 37 (FTIR) spectrophotometer. ¹H NMR spectra were recorded on Bruker Avance 400 (400 MHz) spectrometer at 295 K in CDCl₃; chemical shifts (δ ppm) and coupling constants (Hz) are reported in standard fashion with reference to either internal standard tetramethylsilane (TMS) ($\delta_{\text{H}} = 0.00$ ppm) or CDCl₃ ($\delta_{\text{H}} = 7.25$ ppm). ¹³C{¹H} NMR spectra were recorded on Bruker Avance 400 (100 MHz) spectrometer at RT in CDCl₃; chemical shifts (δ ppm) are reported relative to CDCl₃ [$\delta_{\text{C}} = 77.00$ ppm (central line of triplet)]. In the ¹³C{¹H} NMR, the nature of carbons (C, CH, CH₂ and CH₃) was determined by recording the DEPT-135 spectra, and is given in parentheses and noted as s = singlet (for C), d = doublet (for CH), t = triplet (for CH₂) and q =



Table 3 Scope to yield isochromenes **3da–3ga** from *ortho*-bromo tertiary benzylic alcohols **1d–1g** and acetylenes **2a–2f**^{a,b}

^a Conditions: **1d–1g** (0.25 mmol, 1 equiv.), **2a–2f** (0.625 mmol, 2.5 equiv.), Pd(OAc)₂ (2.8 mg, 0.012 mmol, 5 mol%), K₂CO₃ (138.6 mg, 1 mmol, 4 equiv.), L-proline (5.6 mg, 0.05 mol, 20 mol%), TBAI (tetrabutylammonium iodide) (92.3 mg, 0.25 mmol, 1 equiv.), 140 °C, 36 h. ^b Yields are isolated pure products **3da–3ga**.


quartet (for CH₃). In the ¹H-NMR, the following abbreviations were used throughout: s = singlet, d = doublet, t = triplet, q = quartet, quint = quintet, sept = septet, dd = doublet of doublets, m = multiplet and br. s = broad singlet. The assignment of signals was confirmed by ¹H, ¹³C{¹H} CPD and DEPT spectra. High-resolution mass spectra (HR-MS) were recorded on an Agilent 6538 UHD Q-TOF electron spray ionization (ESI) mode and atmospheric pressure chemical ionization (APCI) modes. Solvents were distilled prior to use; petroleum ether with a boiling range of 60 to 80 °C was used. Palladium acetate, L-proline, TBAI (tetrabutylammonium iodide) and K₂CO₃ were

purchased from Sigma-Aldrich/local sources and used as received. Acme's silica gel (60–120 mesh) was used for column chromatography (approximately 20 g per one gram of crude material). It is worth noting that these sort of experimental procedures have already been published elsewhere.¹⁷

GP (general procedure for the synthesis of isochromenes)

An oven dried Screwcap vial was equipped with a magnetic stir bar and Pd(OAc)₂ (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.3 mL) were added *via* syringe. After the addition of



Table 4 Scope with secondary benzylic alcohol **1h**^{a,b}

^a Conditions: **1h** (57 mg, 0.25 mmol, 1 equiv.), **2a** or **2h** (0.625 mmol, 2.5 equiv.), Pd(OAc)₂ (2.8 mg, 0.012 mmol, 5 mol%), K₂CO₃ (138.6 mg, 1 mmol, 4 equiv.), L-proline (5.6 mg, 0.05 mol, 20 mol%), TBAI (92.3 mg, 0.25 mmol, 1 equiv.), 140 °C, 36 h. ^b Yields are isolated pure products **3ha** & **3hh**.

solvent, the solution was heated at 140 °C for 10 min for pre catalyst formation and brought to ambient temperature. A second oven dried vial was equipped with the starting materials, *ortho*-bromobenzyl alcohols **1** (54.0–84.8 mg, 0.25 mmol), internal alkynes **2** (111.2–196.2 mg, 0.625 mmol) and water (0.2 mL) was added at room temperature. This solution was then transferred to the first vial in which active catalyst was there *via* syringe. The reaction mixture was stirred at 140 °C for 36 h. Reaction mixture was then cooled to room temperature, quenched by the addition of aqueous NaHCO₃ solution and extracted by using ethyl acetate (3 × 20 mL). The organic layer was washed with saturated NH₄Cl solution, dried by Na₂SO₄ and then filtered. Evaporation of the solvent(s) under reduced


pressure and refinement of the crude mixture by silica gel column chromatography (petroleum ether/ethyl acetate) gave the isochromenes **3** (19–82%) as semi-solid/solid.

1,1-Dimethyl-3,4-bis(3-methylphenyl)-1H-isochromene (**3ab**)

GP was carried out with 2-(2-bromophenyl)propan-2-ol **1a** (53.4 mg, 0.25 mmol), 1-methyl-3-[(3-methylphenyl)ethynyl]benzene **2b** (128.8 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 100 : 0 to 99 : 1) to obtain isochromene **3ab** (69.6 mg, 82%) as a light yellow solid compound, melting point = 140 °C, [TLC control (petroleum ether/ethyl acetate 99 : 1), R_f (**2b**) = 0.8, R_f (**3ab**) = 0.7, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 3057, 3018, 1688, 1471, 1355, 1249, 1040, 956, 744, 598 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.26–7.17 (m, 3H, Ar-H), 7.16–7.08 (m, 3H, Ar-H), 7.07–6.99 (m, 4H, Ar-H), 6.96 (s, 1H, Ar-H), 6.88 (d, 1H, J = 7.8 Hz, Ar-H), 2.31 (s, 3H, CH₃), 2.20 (s, 3H, CH₃), 1.78 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 147.9 (s, Ar-C), 138.0 (s, Ar-C), 137.0 (s, Ar-C), 136.7 (s, Ar-C), 136.4 (s, Ar-C), 135.9 (s, Ar-C), 132.2 (d, 2 × Ar-CH), 129.3 (d, Ar-CH), 128.7 (d, Ar-CH), 128.5 (d, Ar-CH), 128.3 (d, Ar-CH), 127.6 (d, Ar-CH), 127.2 (d, Ar-CH), 127.2 (d, Ar-CH), 126.7 (d, Ar-CH), 126.0 (d, Ar-CH), 123.6 (d, Ar-CH), 122.1 (d, Ar-CH), 115.7 (s, Ar-CH), 77.5 (s, Ar-CH), 27.0 (q, 2 × CH₃), 21.4 (q, CH₃), 21.4 (q, CH₃) ppm. HR-MS (ESI+) *m/z* calculated for [C₂₅H₂₅O]⁺ = [M + H]⁺: 341.1900; found 341.1901.

1,1-Dimethyl-3,4-dithien-2-yl-1H-isochromene (**3ad**)

GP was carried out with 2-(2-bromophenyl)propan-2-ol **1a** (53.4 mg, 0.25 mmol), 2-(thien-2-ylethynyl)thiophene **2d** (118.6 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 100 : 0 to 99 : 1) to obtain isochromene **3ad** (32.3 mg, 40%) as a light yellow jelly compound, [TLC control (petroleum ether/ethyl acetate 99 : 1), R_f (**2d**) = 0.8, R_f (**3ad**) = 0.7, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 2981, 1714, 1589, 1445, 1238, 1158, 1093, 1017, 922, 750, 680 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.52 (dd, 1H, J = 5.4 and 1.0 Hz, Ar-H), 7.24–7.13 (m, 5H, Ar-H), 7.05 (dd, 1H, J = 5.4 and 1.0 Hz, Ar-H), 7.00–6.92 (m, 2H, Ar-H), 6.91–6.83 (m, 1H, Ar-H), 1.75 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 145.5 (s, Ar-C) 138.5 (s, Ar-C) 137.1 (s, Ar-C), 136.0 (s, Ar-C), 132.1 (s, Ar-C), 129.8 (d, Ar-CH), 127.8 (d, Ar-CH), 127.7 (d, Ar-CH), 127.6 (d, Ar-CH), 127.5 (d, Ar-CH) 127.5 (d, Ar-CH), 126.9 (d, Ar-CH), 126.8 (d, Ar-CH), 123.6 (d, Ar-CH), 122.0 (d, Ar-CH), 106.2 (s, Ar-CH), 78.5 (s, Ar-CH), 27.0 (q, 2 × CH₃) ppm. HR-MS (ESI+) *m/z* calculated for [C₁₉H₁₇OS₂]⁺ = [M + H]⁺: 325.0715; found 325.0705.

Scheme 1 Plausible mechanism for the formation of isochromenes **3**.

3,4-Bis(3-methoxyphenyl)-1,1-dimethyl-1H-isochromene (3ae)

GP was carried out with 2-(2-bromophenyl)propan-2-ol **1a** (53.4 mg, 0.25 mmol), 1-methoxy-3-[(3-methoxyphenyl)ethynyl]benzene **2e** (148.7 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 99 : 1 to 98 : 2) to obtain isochromene **3ae** (51 mg, 55%) as a light yellow solid compound, melting point = 82 °C, [TLC control (petroleum ether/ethyl acetate 98 : 2), R_f (**2e**) = 0.6, R_f (**3ae**) = 0.5, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 2839, 1596, 1511, 1458, 1249, 1140, 1027, 810, 761 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.26 (ddd, *J* = 7.8, 7.8 and 1.0 Hz, 1H, Ar-H), 7.23–7.18 (m, 2H, Ar-H), 7.14–7.10 (m, 1H, Ar-H), 7.06 (ddd, *J* = 7.8, 7.8 and 1.0 Hz, 1H, Ar-H), 6.93 (ddd, *J* = 7.8, 7.8 and 1.0 Hz, 2H, Ar-H), 6.89–6.77 (m, 4H, Ar-H), 6.76–6.69 (m, 1H, Ar-H), 3.72 (s, 3H, OCH₃), 3.55 (s, 3H, OCH₃), 1.77 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 159.8 (s, Ar-C), 158.6 (s, Ar-C), 147.6 (s, Ar-C), 138.5 (s, Ar-C), 137.2 (s, Ar-C), 136.3 (s, Ar-C), 131.8 (s, Ar-C), 129.6 (d, Ar-CH), 128.5 (d, Ar-CH), 127.2 (d, Ar-CH), 126.9 (d, Ar-CH), 124.1 (d, Ar-CH), 123.7 (d, Ar-CH), 122.1 (d, Ar-CH), 121.1 (d, Ar-CH), 116.9 (d, Ar-CH), 115.6 (s, Ar-CH), 114.2 (d, Ar-CH), 113.7 (d, Ar-CH), 112.6 (d, Ar-CH), 77.6 (s, Ar-C), 55.2 (q, OCH₃), 54.9 (q, OCH₃), 27.0 (q, 2 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₅H₂₅O₃]⁺ = [M + H]⁺: 373.1798; found 373.1784.

1,1-Dimethyl-3,4-bis[4-(trifluoromethyl)phenyl]-1H-isochromene (3ai)

GP was carried out with 2-(2-bromophenyl)propan-2-ol **1a** (53.4 mg, 0.25 mmol), 1-(trifluoromethyl)-4-[[4-(trifluoromethyl)phenyl]ethynyl]benzene **2i** (196.2 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 99 : 1 to 98 : 2) to obtain isochromene **3ai** (84.9 mg, 76%) as a light yellow jelly compound, [TLC control (petroleum ether/ethyl acetate 98 : 2), R_f (**2i**) = 0.7, R_f (**3ai**) = 0.6, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 2922, 2854, 1619, 1444, 1329, 1245, 1166, 1124, 1073, 914, 803, 753, 700 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.65–7.55 (m, 1H, Ar-H), 7.55–7.34 (m, 6H, Ar-H), 7.33–7.21 (m, 3H, Ar-H), 7.21–7.07 (m, 1H, Ar-H), 6.91–6.69 (m, 1H, Ar-H), 1.80 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 147.5 (s, Ar-C), 137.5 (s, 1 Ar-C), 136.3 (s, Ar-C), 136.2 (s, Ar-C), 135.0 (d, Ar-CH), 131.7 (d, Ar-CH), 131.3 (d, *J* = 32.3 Hz, Ar-C), 130.8 (d, Ar-CH), 130.1 (d, *J* = 5.1 Hz, Ar-C), 129.3 (d, Ar-CH), 128.3 (q, *J*_{c-f} = 3.7 Hz, Ar-C), 128.1 (d, Ar-CH), 127.7 (d, Ar-CH), 127.5 (d, Ar-CH), 125.6 (q, *J*_{c-f} = 3.7 Hz, Ar-C), 124.6 (q, *J*_{c-f} = 3.7 Hz, Ar-C), 124.2 (q, *J*_{c-f} = 3.7 Hz, Ar-C), 124.0 (q, *J*_{c-f} = 272.9 Hz, Ar-CF₃), 123.8 (q, *J*_{c-f} = 272.9 Hz, Ar-CF₃), 123.4 (d, Ar-CH), 122.6 (d, Ar-CH), 115.5 (s, Ar-C), 78.4 (s, Ar-C), 27.2 (q, 2 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₅H₁₉F₆O]⁺ = [M + H]⁺: 449.1335; found 449.1338.

3-(3-Methoxyphenyl)-1,1-dimethyl-4-phenyl-1H-isochromene (3aj) and 4-(3-methoxyphenyl)-1,1-dimethyl-3-phenyl-1H-isochromene (3aj')

GP was carried out with 2-(2-bromophenyl)propan-2-ol **1a** (53.4 mg, 0.25 mmol), 1-methoxy-3-(phenylethynyl)benzene **2j** (130 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 100 : 0 to 99 : 1) to obtain isochromene **3aj** + **3aj'** (29.8 mg, 35%) as a light yellow jelly compound, [TLC control (petroleum ether/ethyl acetate 100 : 0), R_f (**2j**) = 0.7, R_f (**3aj** + **3aj'**) = 0.6, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 2975, 2927, 1695, 1591, 1481, 1253, 1158, 1044, 761, 700 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.41–7.33 (m, 2H, Ar-H), 7.33–7.22 (m, 6H, Ar-H), 7.22–7.18 (m, 4H, Ar-H), 7.18–7.10 (m, 5H, Ar-H), 7.06 (ddd, *J* = 7.8, 7.8 and 1.0 Hz, 1H, Ar-H), 6.97–6.87 (m, 3H, Ar-H), 6.86–6.81 (m, 2H, Ar-H), 6.79–6.76 (m, 1H, Ar-H), 6.75–6.72 (m, 1H, Ar-H), 6.72–6.66 (m, 1H, Ar-H), 3.73 (s, 3H, OCH₃), 3.53 (s, 3H, OCH₃), 1.79 (s, 12H, 4 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 159.7 (s, Ar-C), 158.5 (s, Ar-C), 148.0 (s, Ar-C), 147.7 (s, Ar-C), 138.4 (s, Ar-C), 137.2 (s, Ar-C), 136.3 (s, Ar-C), 136.2 (s, Ar-C), 136.0 (s, Ar-C), 131.9 (s, Ar-C), 131.8 (s, Ar-C), 131.6 (d, 3 × Ar-CH), 129.5 (d, Ar-CH), 128.6 (d, 4 × Ar-CH), 128.4 (d, Ar-CH), 127.8 (d, Ar-CH), 127.5 (d, 3 × Ar-CH), 127.2 (d, Ar-CH), 127.2 (d, Ar-CH), 126.9 (d, Ar-CH), 126.9 (d, Ar-CH), 126.9 (d, Ar-CH), 124.2 (d, Ar-CH), 123.6 (d, Ar-CH), 123.6 (d, Ar-CH), 122.2 (d, 2 × Ar-CH), 121.2 (d, Ar-CH), 117.0 (d, Ar-CH), 115.8 (s, Ar-C), 115.5 (s, Ar-C), 114.2 (d, Ar-CH), 113.9 (d, Ar-CH), 112.6 (d, Ar-CH), 77.7 (s, Ar-C), 55.2 (q, OCH₃), 54.9 (q, OCH₃), 27.08 (q, 2 × CH₃), 27.05 (q, 2 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₄H₂₃O₂]⁺ = [M + H]⁺: 343.1693; found 343.1693.

3-(4-Fluorophenyl)-1,1-dimethyl-4-phenyl-1H-isochromene (3ak) and 4-(4-fluorophenyl)-1,1-dimethyl-3-phenyl-1H-isochromene (3ak')

GP was carried out with 2-(2-bromophenyl)propan-2-ol **1a** (53.4 mg, 0.25 mmol), 1-fluoro-4-(phenylethynyl)benzene **2k** (122.5 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 99 : 1 to 98 : 2) to obtain isochromene **3ak** + **3ak'** (30.4 mg, 37%) as a light yellow jelly compound, [TLC control (petroleum ether/ethyl acetate 98 : 2), R_f (**2k**) = 0.8, R_f (**3ak** + **3ak'**) = 0.7, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 3062, 2978, 2925, 1695, 1610, 1507, 1230, 1155, 1099, 962, 834, 760, 700 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.42–7.28 (m, 4H, Ar-H), 7.28–7.18 (m, 12H, Ar-H), 7.17–7.10 (m, 4H, Ar-H), 7.06–7.00 (m, 2H, Ar-H), 6.93–6.75 (m, 4H, Ar-H), 1.79 (s, 12H, 4 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 162.0 (d, *J* = 245.0 Hz, Ar-C), 160.8 (s, Ar-C), 147.1 (s, Ar-C), 136.9 (s, Ar-C), 136.2 (s, Ar-C), 135.9 (s, Ar-C), 133.3 (d, Ar-CH), 133.2 (d, Ar-CH), 131.8 (s, Ar-CH), 131.6 (d, 4 × Ar-CH), 130.7 (d, Ar-CH), 130.6 (d, Ar-CH), 128.7 (d, 3 × Ar-CH), 128.6 (d, 5 × Ar-CH),



127.9 (d, Ar-CH), 127.5 (d, 3 × Ar-CH), 127.3 (d, Ar-CH), 127.0 (d, 3 × Ar-CH), 127.0 (d, Ar-CH), 126.9 (d, Ar-CH), 123.5 (d, Ar-CH), 123.3 (d, Ar-CH), 122.3 (d, Ar-CH), 122.2 (d, Ar-CH), 115.7 (d, Ar-CH), 115.4 (d, Ar-CH), 114.5 (d, Ar-CH), 114.3 (d, Ar-CH), 77.8 (s, Ar-C), 27.1 (q, 4 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₃H₂₀FO]⁺ = [M + H]⁺: 331.1493; found 331.1494.

1-Methyl-3,4-diphenyl-1-propyl-1H-isochromene (3ba)

GP was carried out with 2-(2-bromophenyl)pentan-2-ol **1b** (60.5 mg), (phenylethynyl)benzene **2a** (111.2 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 100 : 0 to 99 : 1) to obtain isochromene **3ba** (36.5 mg, 43%) as a light yellow jelly compound, [TLC control (petroleum ether/ethyl acetate 100 : 0), *R_f* (**2a**) = 0.8, *R_f* (**3ba**) = 0.7, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 3055, 2947, 1702, 1605, 1477, 1445, 1348, 1278, 1231, 1026, 703 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.37–7.27 (m, 3H, Ar-H), 7.27–7.19 (m, 4H, Ar-H), 7.19–7.05 (m, 6H, Ar-H), 6.85 (dd, *J* = 7.3 and 1.0 Hz, 1H, Ar-H), 2.22–1.99 (m, 2H, CH₂), 1.74 (s, 3H, CH₃), 1.58–1.39 (m, 2H, CH₂), 0.93 (t, *J* = 7.3 Hz, 3H, CH₃) ppm. ¹³C {¹H} NMR (CDCl₃, 100 MHz): δ = 148.0 (s, Ar-C), 137.2 (s, Ar-C), 136.1 (s, Ar-C), 135.3 (s, Ar-C), 132.4 (s, Ar-C), 131.7 (d, 2 × Ar-CH), 128.7 (d, 3 × Ar-CH), 128.5 (d, Ar-CH), 127.6 (d, Ar-CH), 127.4 (d, 2 × Ar-CH), 127.0 (d, Ar-CH), 126.9 (d, Ar-CH), 126.5 (d, Ar-CH), 123.5 (d, Ar-CH), 122.9 (d, Ar-CH), 115.1 (s, Ar-C), 80.1 (s, Ar-C), 41.8 (t, CH₂), 25.7 (q, CH₃), 17.4 (t, CH₂), 14.6 (q, CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₅H₂₅O]⁺ = [M + H]⁺: 341.1900; found 341.1906.

1-Methyl-3,4-bis(4-methylphenyl)-1-propyl-1H-isochromene (3bc)

GP was carried out with 2-(2-bromophenyl)pentan-2-ol **1b** (60.5 mg, 0.25 mmol), 1-methyl-4-[(4-methylphenyl)ethynyl]benzene **2c** (128.8 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 100 : 0 to 99 : 1) to obtain isochromene **3bc** (18.4 mg, 20%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 99 : 1), *R_f* (**2c**) = 0.7, *R_f* (**3bc**) = 0.6, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 3025, 2920, 1670, 1507, 1446, 1261, 1181, 1108, 1033, 946, 819, 732 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.20–6.99 (m, 8H, Ar-H), 6.94 (d, *J* = 1.0 Hz, 1H, Ar-H), 6.93 (dd, *J* = 7.8 and 1.0 Hz, 1H, Ar-H), 6.84 (dd, *J* = 7.8 and 1.0 Hz, 1H, Ar-H), 6.64 (d, *J* = 2.0 Hz, 1H, Ar-H), 2.36 (s, 3H, CH₃), 2.24 (s, 3H, CH₃), 2.15–1.87 (m, 2H, CH₂), 1.54 (s, 3H, CH₃), 1.52–1.36 (m, 2H, CH₂), 0.91 (t, *J* = 7.3 Hz, 3H, CH₃) ppm. ¹³C {¹H} NMR (CDCl₃, 100 MHz): δ = 147.9 (s, Ar-C), 137.4 (s, Ar-C), 136.3 (s, Ar-C), 135.3 (s, Ar-C), 134.2 (s, Ar-C), 133.3 (s, Ar-C), 132.7 (s, Ar-C), 131.5 (d, 2 × Ar-CH), 131.3 (d, Ar-CH), 129.3 (d, 2 × Ar-CH), 128.6 (d, 2 × Ar-CH), 128.1 (d, 2 × Ar-CH), 127.0 (d, Ar-CH), 126.3 (d, Ar-CH), 123.5 (d, Ar-CH), 122.8 (s, Ar-C), 79.8 (s, Ar-C), 41.8 (t, CH₂), 25.6 (q, CH₃), 21.3 (q, CH₃), 21.2 (q, CH₃),

17.4 (t, CH₂), 14.6 (q, CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₇H₂₉O]⁺ = [M + H]⁺: 369.2213; found 369.2218.

3,4-Bis(3,4-dimethoxyphenyl)-1,1-diphenyl-1H-isochromene (3cg)

GP was carried out with (2-bromophenyl)(diphenyl)methanol **1c** (84.5 mg, 0.25 mmol), 4-[(3,4-dimethoxyphenyl)ethynyl]-1,2-dimethoxybenzene **2g** (186.3 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 73 : 27 to 70 : 30) to obtain isochromene **3cg** (34.7 mg, 25%) as a light yellow jelly compound, [TLC control (petroleum ether/ethyl acetate 70 : 30), *R_f* (**2g**) = 0.5, *R_f* (**3cg**) = 0.4, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 2925, 2850, 1511, 1459, 1254, 1151, 1028, 763, 702 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.37–7.26 (m, 10H, Ar-H), 7.23 (ddd, *J* = 7.8, 7.8 and 1.5 Hz, 1H, Ar-H), 7.11 (ddd, *J* = 7.8, 7.8 and 1.5 Hz, 1H, Ar-H), 7.08–6.99 (m, 2H, Ar-H), 6.81 (d, *J* = 8.3 Hz, 1H, Ar-H), 6.74–6.62 (m, 3H, Ar-H) 6.56 (dd, *J* = 8.3 and 2.0 Hz, 1H, Ar-H), 6.50 (d, *J* = 2.0 Hz, 1H, Ar-H), 3.85 (s, 3H, OCH₃), 3.80 (s, 3H, OCH₃), 3.68 (s, 3H, OCH₃), 3.48 (s, 3H, OCH₃) ppm. ¹³C {¹H} NMR (CDCl₃, 100 MHz): δ = 149.1 (s, Ar-C), 148.5 (s, Ar-C), 148.0 (s, Ar-C), 147.9 (s, Ar-C), 147.4 (s, Ar-C), 144.0 (d, Ar-CH), 134.2 (s, Ar-C), 133.5 (s, Ar-C), 131.5 (s, Ar-C), 130.0 (s, Ar-C), 129.8 (s, Ar-C), 128.8 (d, 3 × Ar-CH), 128.2 (s, Ar-CH), 127.9 (d, Ar-CH), 127.7 (d, 2 × Ar-CH), 127.5 (d, 4 × Ar-CH), 126.8 (s, Ar-C), 126.0 (d, Ar-CH), 123.6 (d, Ar-CH), 123.1 (d, Ar-CH), 121.3 (d, Ar-CH), 116.5 (s, Ar-C), 114.5 (d, Ar-CH), 111.9 (d, Ar-CH), 111.4 (d, Ar-CH), 110.0 (d, Ar-CH), 86.7 (s, Ar-C), 55.9 (q, OCH₃), 55.8 (q, OCH₃), 55.6 (q, OCH₃), 55.3 (q, OCH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₃₇H₃₃O₅]⁺ = [M + H]⁺: 557.2323; found 557.2331.

7-Methoxy-1,1-dimethyl-3,4-diphenyl-1H-isochromene (3da)

GP was carried out with 2-(2-bromo-5-methoxyphenyl)propan-2-ol **1d** (61 mg, 0.25 mmol), (phenylethynyl)benzene **2a** (111.2 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 100 : 0 to 99 : 1) to obtain isochromene **3da** (57.2 mg, 67%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 100 : 0), *R_f* (**2a**) = 0.8, *R_f* (**3da**) = 0.7, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 2925, 2364, 1743, 1694, 1511, 1301, 1069, 758, 699 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.39–7.28 (m, 2H, Ar-H), 7.28–7.19 (m, 4H, Ar-H), 7.17–7.08 (m, 3H, Ar-H), 6.89–6.75 (m, 3H, Ar-H), 6.67 (dd, *J* = 8.6 and 2.7 Hz, 1H, Ar-H), 3.81 (s, 3H, OCH₃), 1.77 (s, 6H, 2 × CH₃) ppm. ¹³C {¹H} NMR (CDCl₃, 100 MHz): δ = 158.8 (s, Ar-C), 146.0 (s, Ar-C), 138.3 (s, Ar-C), 137.3 (s, Ar-C), 136.1 (s, Ar-C), 131.6 (d, 2 × Ar-CH), 131.4 (d, Ar-CH), 128.6 (d, Ar-CH), 128.5 (d, 2 × Ar-CH), 127.4 (d, Ar-CH), 127.4 (d, Ar-CH), 126.9 (d, Ar-CH), 126.5 (d, Ar-CH), 125.3 (s, Ar-C), 125.1 (d, Ar-CH), 115.6 (s, Ar-C), 111.3 (d, Ar-CH), 109.0 (d, Ar-CH), 77.4 (s, Ar-C), 55.4 (q, OCH₃), 26.9 (q, 2 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₄H₂₃O₂]⁺ = [M + H]⁺: 343.1693; found 343.1686.



7-Methoxy-1,1-dimethyl-3,4-bis(3-methylphenyl)-1H-isochromene (3db)

GP was carried out with 2-(2-bromo-5-methoxyphenyl)propan-2-ol **1d** (61 mg, 0.25 mmol), 1-methyl-3-[(3-methylphenyl)ethynyl]benzene **2b** (128.8 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 99 : 1 to 98 : 2) to obtain isochromene **3db** (55.5 mg, 60%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 98 : 2), *R_f* (**2b**) = 0.7, *R_f* (**3db**) = 0.6, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{\max} = 2924, 2364, 1691, 1611, 1506, 1295, 1223, 1043, 785, 704 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.28–7.19 (m, 1H, Ar-H), 7.18–7.09 (m, 2H, Ar-H), 7.09–6.98 (m, 4H, Ar-H), 6.95 (m, 1H, Ar-H), 6.81 (dd, *J* = 8.3 and 2.5 Hz, 1H, Ar-H), 6.76 (d, *J* = 2.5 Hz, 1H, Ar-H), 6.67 (dd, *J* = 8.3 and 2.5 Hz, 1H, Ar-H), 3.81 (s, 3H, OCH₃), 2.32 (s, 3H, CH₃), 2.21 (s, 3H, CH₃), 1.77 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 158.7 (s, Ar-C), 145.8 (s, Ar-C), 138.4 (s, Ar-C), 138.0 (s, Ar-C), 137.2 (s, Ar-C), 136.8 (s, Ar-C), 136.0 (s, Ar-C), 132.0 (d, Ar-CH), 129.1 (d, Ar-CH), 128.6 (d, Ar-CH), 128.3 (d, Ar-CH), 128.2 (d, Ar-CH), 127.5 (d, Ar-CH), 127.2 (d, Ar-CH), 125.8 (d, Ar-CH), 125.5 (d, Ar-CH), 125.1 (d, Ar-CH), 115.6 (s, Ar-C), 111.3 (d, Ar-CH), 108.9 (d, Ar-CH), 77.2 (s, Ar-C), 55.3 (q, OCH₃), 26.9 (q, 2 × CH₃), 21.4 (q, 2 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₆H₂₇O₂]⁺ = [M + H]⁺: 371.2006; found 371.1992.

7-Methoxy-1,1-dimethyl-3,4-bis(4-methylphenyl)-1H-isochromene (3dc)

GP was carried out with 2-(2-bromo-5-methoxyphenyl)propan-2-ol **1d** (61 mg, 0.25 mmol), 1-methyl-4-[(4-methylphenyl)ethynyl]benzene **2c** (128.8 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.67 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 99 : 1 to 98 : 2) to obtain isochromene **3dc** (25.9 mg, 28%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 98 : 2), *R_f* (**2c**) = 0.7, *R_f* (**3dc**) = 0.6, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{\max} = 2923, 2856, 1701, 1614, 1498, 1295, 1222, 1044, 964, 816, 756 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.20–7.04 (m, 6H, Ar-H), 6.94 (dd, *J* = 7.8 and 1.0 Hz, 2H, Ar-H), 6.81 (d, *J* = 8.8 Hz, 1H, Ar-H), 6.76 (d, *J* = 2.9 Hz, 1H, Ar-H), 6.69–6.58 (m, 1H, Ar-H), 3.80 (s, 3H, OCH₃), 2.37 (s, 3H, CH₃), 2.25 (s, 3H, CH₃), 1.74 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 158.6 (s, Ar-C), 145.8 (s, Ar-C), 138.3 (s, Ar-C), 137.1 (s, Ar-C), 136.3 (s, Ar-C), 134.3 (s, Ar-C), 133.3 (s, Ar-C), 131.3 (d, 2 × Ar-CH), 129.3 (d, 2 × Ar-CH), 128.4 (d, 2 × Ar-CH), 128.1 (d, 2 × Ar-CH), 125.6 (s, Ar-C), 125.0 (d, Ar-CH), 115.0 (s, Ar-C), 111.3 (d, Ar-CH), 108.9 (d, Ar-CH), 77.2 (s, Ar-C), 55.3 (q, OCH₃), 26.9 (q, 2 × CH₃), 21.3 (q, CH₃), 21.2 (q, CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₆H₂₇O₂]⁺ = [M + H]⁺: 371.2006; found 371.1991.

7-Methoxy-1,1-dimethyl-3,4-dithien-2-yl-1H-isochromene (3dd)

GP was carried out with 2-(2-bromo-5-methoxyphenyl)propan-2-ol **1d** (61 mg, 0.25 mmol), 2-(thien-2-ylethynyl)thiophene **2d** (118.6 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 99 : 1 to 98 : 2) to obtain isochromene **3dd** (36.2 mg, 41%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 98 : 2), *R_f* (**2d**) = 0.6, *R_f* (**3dd**) = 0.5, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{\max} = 3041, 2923, 1712, 1595, 1482, 1446, 1264, 1209, 1167, 1081, 989, 907, 745, 695 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.51 (dd, *J* = 5.4 and 1.5 Hz, 1H, Ar-H), 7.23–7.15 (m, 2H, Ar-H), 7.04 (dd, *J* = 3.4 and 1.0 Hz, 1H, Ar-H), 6.96–6.85 (m, 3H, Ar-H), 6.75 (d, *J* = 2.9 Hz, 1H, Ar-H), 6.70 (dd, *J* = 8.8 and 3.0 Hz, 1H, Ar-H), 3.80 (s, 3H, OCH₃), 1.73 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 158.9 (s, Ar-C), 143.5 (s, Ar-C), 138.7 (s, Ar-C), 138.0 (s, Ar-C), 137.3 (s, Ar-C), 129.6 (d, Ar-CH), 127.7 (d, Ar-CH), 127.4 (d, Ar-CH), 127.1 (d, Ar-CH), 126.9 (d, Ar-CH), 126.7 (d, Ar-CH), 125.4 (s, Ar-C), 125.1 (d, Ar-CH), 111.6 (d, Ar-CH), 108.9 (d, Ar-CH), 106.3 (s, Ar-C), 78.2 (s, Ar-C), 55.4 (q, OCH₃), 26.8 (q, 2 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₀H₁₉O₂S₂]⁺ = [M + H]⁺: 355.0821; found 355.0818.

7-Methoxy-3,4-bis(3-methoxyphenyl)-1,1-dimethyl-1H-isochromene (3de)

GP was carried out with 2-(2-bromo-5-methoxyphenyl)propan-2-ol **1d** (61 mg, 0.25 mmol), 1-methoxy-3-[(3-methoxyphenyl)ethynyl]benzene **2e** (148.7 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 99 : 1 to 98 : 2) to obtain isochromene **3de** (40.2 mg, 40%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 98 : 2), *R_f* (**2e**) = 0.6, *R_f* (**3de**) = 0.5, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{\max} = 3087, 2974, 2838, 1608, 1487, 1425, 1297, 1216, 1043, 834, 701 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.27 (ddd, *J* = 7.8, 7.8 and 1.0 Hz, 1H, Ar-H), 7.06 (ddd, *J* = 7.8, 7.8 and 1.0 Hz, 1H, Ar-H), 6.92 (dt, *J* = 7.8 and 1.0 Hz, 1H, Ar-H), 6.88–6.82 (m, 3H, Ar-H), 6.82–6.72 (m, 3H, Ar-H), 6.67 (m, 2H, Ar-H), 3.80 (s, 3H, OCH₃), 3.73 (s, 3H, OCH₃), 3.55 (s, 3H, OCH₃), 1.75 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 159.8 (s, Ar-C), 158.9 (s, Ar-C), 158.6 (s, Ar-C), 145.5 (s, Ar-C), 138.8 (s, Ar-C), 138.3 (s, Ar-C), 137.3 (s, Ar-C), 129.6 (d, Ar-CH), 128.4 (d, Ar-CH), 125.1 (d, Ar-CH), 125.1 (s, Ar-C), 124.0 (d, Ar-CH), 120.9 (d, Ar-CH), 116.8 (d, Ar-CH), 115.6 (s, Ar-C), 113.9 (q, Ar-CH), 113.5 (d, Ar-CH), 112.7 (d, Ar-CH), 111.4 (d, Ar-CH), 109.0 (d, Ar-CH), 77.4 (s, Ar-C), 55.3 (q, OCH₃), 55.2 (q, OCH₃), 54.9 (q, OCH₃), 26.9 (q, 2 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₆H₂₇O₄]⁺ = [M + H]⁺: 403.1904; found 403.1901.

6,7-Dimethoxy-1,1-dimethyl-3,4-diphenyl-1H-isochromene (3ea)

GP was carried out with 2-(2-bromo-4,5-dimethoxyphenyl)propan-2-ol **1e** (68.5 mg, 0.25 mmol), (phenylethynyl)benzene



2a (111.2 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), *L*-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K_2CO_3 (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 85 : 15 to 80 : 20) to obtain isochromene **3ea** (37.2 mg, 40%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 80 : 20), R_f (**2a**) = 0.6, R_f (**3ea**) = 0.5, UV detection]. IR (MIR-ATR, 4000–600 cm^{-1}): ν_{max} = 2942, 2843, 1603, 1505, 1452, 1242, 1171, 1026, 825, 733 cm^{-1} . 1H NMR ($CDCl_3$, 400 MHz): δ = 7.42–7.17 (m, 7H, Ar-H), 7.17–7.00 (m, 3H, Ar-H), 6.73 (s, 1H, Ar-H), 6.45 (s, 1H, Ar-H), 3.91 (s, 3H, OCH₃), 3.64 (s, 3H, OCH₃), 1.76 (s, 6H, 2 × CH₃) ppm. $^{13}C\{^1H\}$ NMR ($CDCl_3$, 100 MHz): δ = 148.0 (s, Ar-C), 147.9 (s, Ar-C), 146.6 (s, Ar-C), 137.2 (s, Ar-C), 136.0 (s, Ar-C), 131.5 (d, 3 × Ar-CH), 129.0 (s, Ar-C), 128.6 (d, Ar-CH), 128.6 (d, 2 × Ar-CH), 127.5 (d, 2 × Ar-CH), 127.4 (d, Ar-CH), 126.9 (d, Ar-CH), 125.3 (s, Ar-C), 115.6 (s, Ar-C), 107.6 (d, Ar-CH), 106.2 (d, Ar-CH), 77.3 (s, Ar-C), 56.2 (q, OCH₃), 55.8 (q, OCH₃), 27.1 (q, 2 × CH₃) ppm. HR-MS (APCI+) m/z calculated for $[C_{25}H_{25}O_3]^+$ = $[M + H]^+$: 373.1798; found 373.1780.

6,7-Dimethoxy-1,1-dimethyl-3,4-bis(3-methylphenyl)-1H-isochromene (**3eb**)

GP was carried out with 2-(2-bromo-4,5-dimethoxyphenyl)propan-2-ol **1e** (68.5 mg, 0.25 mmol), 1-methyl-3-[(3-methylphenyl)ethynyl]benzene **2b** (128.8 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), *L*-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K_2CO_3 (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 93 : 7 to 90 : 10) to obtain isochromene **3eb** (52 mg, 52%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 90 : 10), R_f (**2b**) = 0.6, R_f (**3eb**) = 0.5, UV detection]. IR (MIR-ATR, 4000–600 cm^{-1}): ν_{max} = 3055, 2933, 1595, 1485, 1448, 1244, 1072, 1027, 747, 697 cm^{-1} . 1H NMR ($CDCl_3$, 400 MHz): δ = 7.22 (ddd, J = 7.8, 7.8 and 1.0 Hz, 1H, Ar-H), 7.16–7.07 (m, 2H, Ar-H), 7.07–7.01 (m, 2H, Ar-H), 7.01–6.89 (m, 3H, Ar-H), 6.72 (s, 1H, Ar-H), 6.46 (s, 1H, Ar-H), 3.90 (s, 3H, OCH₃), 3.65 (s, 3H, OCH₃), 2.30 (s, 3H, CH₃), 2.20 (s, 3H, CH₃), 1.76 (s, 6H, 2 × CH₃) ppm. $^{13}C\{^1H\}$ NMR ($CDCl_3$, 100 MHz): δ = 147.9 (s, Ar-C), 147.9 (s, Ar-C), 146.5 (s, Ar-C), 138.0 (s, Ar-C), 137.1 (s, Ar-C), 136.8 (s, Ar-C), 136.0 (s, Ar-C), 132.0 (d, Ar-CH), 129.1 (d, Ar-CH), 128.6 (d, Ar-CH), 128.3 (d, Ar-CH), 128.2 (d, Ar-CH), 127.6 (d, Ar-CH), 127.2 (d, Ar-CH), 125.8 (d, Ar-CH), 125.6 (s, Ar-C), 115.6 (s, Ar-C), 107.8 (d, Ar-CH), 106.2 (d, Ar-CH), 77.2 (s, Ar-C), 56.2 (q, OCH₃), 55.8 (q, OCH₃), 27.0 (q, 2 × CH₃), 21.4 (q, 2 × CH₃) ppm. HR-MS (APCI+) m/z calculated for $[C_{27}H_{29}O_3]^+$ = $[M + H]^+$: 401.2111; found 401.2101.

6,7-Dimethoxy-1,1-dimethyl-3,4-bis(4-methylphenyl)-1H-isochromene (**3ec**)

GP was carried out with 2-(2-bromo-4,5-dimethoxyphenyl)propan-2-ol **1e** (68.5 mg, 0.25 mmol), 1-methyl-4-[(4-methylphenyl)ethynyl]benzene **2c** (128.8 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), *L*-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K_2CO_3 (138.6 mg, 1

mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 93 : 7 to 90 : 10) to obtain isochromene **3ec** (44 mg, 44%) as a light yellow solid compound, melting point = 88 °C, [TLC control (petroleum ether/ethyl acetate 90 : 10), R_f (**2c**) = 0.6, R_f (**3ec**) = 0.5, UV detection]. IR (MIR-ATR, 4000–600 cm^{-1}): ν_{max} = 3094, 2975, 2936, 2841, 1703, 1609, 1490, 1419, 1290, 1219, 1044, 835, 703 cm^{-1} . 1H NMR ($CDCl_3$, 400 MHz): δ = 7.18–7.11 (m, 6H, Ar-H), 6.93 (dd, J = 7.8 and 1.0 Hz, 2H, Ar-H), 6.72 (s, 1H, Ar-H), 6.47 (s, 1H, Ar-H), 3.91 (s, 3H, OCH₃), 3.66 (s, 3H, OCH₃), 2.37 (s, 3H, CH₃), 2.25 (s, 3H, CH₃), 1.75 (s, 6H, 2 × CH₃) ppm. $^{13}C\{^1H\}$ NMR ($CDCl_3$, 100 MHz): δ = 147.9 (s, Ar-C), 147.8 (s, Ar-C), 146.5 (s, Ar-C), 137.2 (s, Ar-C), 136.3 (s, Ar-C), 134.2 (s, Ar-C), 133.3 (s, Ar-C), 131.3 (d, 2 × Ar-CH), 129.3 (d, 2 × Ar-CH), 129.1 (s, Ar-C), 128.5 (d, 2 × Ar-CH), 128.1 (d, 2 × Ar-CH), 125.7 (s, Ar-C), 114.9 (s, Ar-C), 107.6 (d, Ar-CH), 106.2 (d, Ar-CH), 77.1 (s, Ar-C), 56.2 (q, OCH₃), 55.8 (q, OCH₃), 27.0 (q, 2 × CH₃), 21.3 (q, CH₃), 21.2 (q, CH₃) ppm. HR-MS (APCI+) m/z calculated for $[C_{27}H_{29}O_3]^+$ = $[M + H]^+$: 401.2111; found 401.2107.

6,7-Dimethoxy-3,4-bis(3-methoxyphenyl)-1,1-dimethyl-1H-isochromene (**3ee**)

GP was carried out with 2-(2-bromo-4,5-dimethoxyphenyl)propan-2-ol **1e** (68.5 mg, 0.25 mmol), 1-methoxy-3-[(3-methoxyphenyl)ethynyl]benzene **2e** (148.7 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), *L*-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K_2CO_3 (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 73 : 27 to 70 : 30) to obtain isochromene **3ee** (44.2 mg, 41%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 70 : 30), R_f (**2e**) = 0.5, R_f (**3ee**) = 0.4, UV detection]. IR (MIR-ATR, 4000–600 cm^{-1}): ν_{max} = 2944, 2834, 1597, 1504, 1452, 1237, 1163, 1105, 1020, 825, 741 cm^{-1} . 1H NMR ($CDCl_3$, 400 MHz): δ = 7.28 (ddd, J = 7.8, 7.8 and 2.4 Hz, 1H, Ar-H), 7.06 (ddd, J = 7.8, 7.8 and 2.4 Hz, 1H, Ar-H), 6.94–6.88 (m, 2H, Ar-H), 6.87–6.82 (m, 1H, Ar-H), 6.82–6.76 (m, 2H, Ar-H), 6.72 (s, 1H, Ar-H), 6.71–6.66 (m, 1H, Ar-H), 6.49 (s, 1H, Ar-H), 3.91 (s, 3H, OCH₃), 3.73 (s, 3H, OCH₃), 3.66 (s, 3H, OCH₃), 3.56 (s, 3H, OCH₃), 1.75 (s, 6H, 2 × CH₃) ppm. $^{13}C\{^1H\}$ NMR ($CDCl_3$, 100 MHz): δ = 159.9 (s, Ar-C), 158.6 (s, Ar-C), 148.1 (s, Ar-C), 147.9 (s, Ar-C), 146.2 (s, Ar-C), 138.7 (s, Ar-C), 137.3 (s, Ar-C), 129.6 (d, Ar-CH), 129.1 (s, Ar-C), 128.4 (d, Ar-CH), 125.1 (s, Ar-C), 123.9 (d, Ar-CH), 121.0 (d, Ar-CH), 116.7 (d, Ar-CH), 115.5 (s, Ar-C), 114.0 (d, Ar-CH), 113.5 (d, Ar-CH), 112.8 (d, Ar-CH), 107.7 (d, Ar-CH), 106.3 (d, Ar-CH), 77.3 (s, Ar-C), 56.2 (q, OCH₃), 55.8 (q, OCH₃), 55.2 (q, OCH₃), 54.9 (q, OCH₃), 27.1 (q, 2 × CH₃) ppm. HR-MS (APCI+) m/z calculated for $[C_{27}H_{29}O_5]^+$ = $[M + H]^+$: 433.2010; found 433.2016.

6,7-Dimethoxy-3,4-bis(4-methoxyphenyl)-1,1-dimethyl-1H-isochromene (**3ef**)

GP was carried out with 2-(2-bromo-4,5-dimethoxyphenyl)propan-2-ol **1e** (68.5 mg, 0.25 mmol), 1-methoxy-4-[(4-methoxyphenyl)ethynyl]benzene **2f** (148.7 mg, 0.625 mmol),



palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 73 : 27 to 70 : 30) to obtain isochromene **3ef** (52.9 mg, 49%) as a light yellow jelly compound, [TLC control (petroleum ether/ethyl acetate 70 : 30), R_f (**3f**) = 0.5, R_f (**3ef**) = 0.4, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 3090, 1792, 1652, 1529, 1409, 1266, 1198, 834, 697 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.23–7.12 (m, 4H, Ar-H), 6.94–6.84 (m, 2H, Ar-H), 6.71 (s, 1H, Ar-H), 6.70–6.61 (m, 2H, Ar-H), 6.47 (s, 1H, Ar-H), 3.90 (s, 3H, OCH₃), 3.83 (s, 3H, OCH₃), 3.73 (s, 3H, OCH₃), 3.66 (s, 3H, OCH₃), 1.74 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 158.7 (s, Ar-C), 158.4 (s, Ar-C), 147.9 (s, Ar-C), 147.7 (s, Ar-C), 146.4 (s, Ar-C), 132.5 (d, 2 × Ar-CH), 129.9 (d, 2 × Ar-CH), 129.6 (s, Ar-C), 128.9 (s, Ar-C), 128.7 (s, Ar-C), 125.9 (s, Ar-C), 114.0 (d, 2 × Ar-CH), 113.9 (s, Ar-C), 112.8 (d, 2 × Ar-CH), 107.5 (d, Ar-CH), 106.3 (d, Ar-CH), 77.1 (s, Ar-C), 56.2 (q, OCH₃), 55.8 (q, OCH₃), 55.1 (q, OCH₃), 55.0 (q, OCH₃), 27.0 (q, 2 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₇H₂₉O₅]⁺ = [M + H]⁺: 433.2010; found 433.2018.

5,6,7-Trimethoxy-1,1-dimethyl-3,4-diphenyl-1*H*-isochromene (**3fa**)

GP was carried out with 2-(2-bromo-3,4,5-trimethoxyphenyl)propan-2-ol **1f** (76 mg, 0.25 mmol), (phenylethynyl)benzene **2a** (111.2 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 85 : 15 to 80 : 20) to obtain isochromene **3fa** (19 mg, 19%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 80 : 20), R_f (**2a**) = 0.6, R_f (**3fa**) = 0.5, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 2936, 2364, 1742, 1695, 1649, 1515, 1461, 1258, 1099, 1025, 758, 700 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.24–7.19 (m, 4H, Ar-H), 7.18–7.12 (m, 3H, Ar-H), 7.11–7.08 (m, 3H, Ar-H), 6.57 (s, 1H, Ar-H), 3.91 (s, 3H, OCH₃), 3.78 (s, 3H, OCH₃), 3.09 (s, 3H, OCH₃), 1.75 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 152.6 (s, Ar-C), 150.4 (s, Ar-C), 148.1 (s, Ar-C), 142.3 (s, Ar-C), 139.6 (s, Ar-C), 136.6 (s, Ar-C), 134.6 (s, Ar-C), 130.9 (d, 2 × Ar-CH), 128.6 (d, 2 × Ar-CH), 127.4 (d, 2 × Ar-CH), 127.3 (d, Ar-CH), 127.3 (d, 2 × Ar-CH), 125.9 (d, Ar-CH), 119.2 (s, Ar-C), 115.3 (s, Ar-C), 102.1 (d, Ar-CH), 77.6 (s, Ar-C), 60.6 (q, OCH₃), 60.1 (q, OCH₃), 56.2 (q, OCH₃), 26.6 (q, 2 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₆H₂₇O₄]⁺ = [M + H]⁺: 403.1904; found 403.1907.

5,6,7-Trimethoxy-1,1-dimethyl-3,4-bis(3-methylphenyl)-1*H*-isochromene (**3fb**)

GP was carried out with 2-(2-bromo-3,4,5-trimethoxyphenyl)propan-2-ol **1f** (76 mg, 0.25 mmol), 1-methyl-3-[(3-methylphenyl)ethynyl]benzene **2b** (128.8 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined

by silica gel column chromatography (petroleum ether/ethyl acetate, 93 : 7 to 90 : 10) to obtain isochromene **3fb** (21.5 mg, 20%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 90 : 10), R_f (**2b**) = 0.5, R_f (**3fb**) = 0.4, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 2926, 2856, 2365, 1696, 1650, 1514, 1463, 1097, 753 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.08 (ddd, *J* = 7.8, 7.8 and 1.0 Hz, 1H, Ar-H), 7.05–7.01 (m, 3H, Ar-H), 7.00–6.86 (m, 4H, Ar-H), 6.54 (s, 1H, Ar-H), 3.89 (s, 3H, OCH₃), 3.76 (s, 3H, OCH₃), 3.09 (s, 3H, OCH₃), 2.26 (s, 3H, CH₃), 2.19 (s, 3H, CH₃), 1.72 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 152.5 (s, Ar-C), 150.4 (s, Ar-C), 147.9 (s, Ar-C), 142.3 (s, Ar-C), 139.5 (s, Ar-C), 136.7 (s, Ar-C), 136.5 (s, Ar-C), 136.4 (s, Ar-C), 134.8 (s, Ar-C), 131.3 (d, Ar-CH), 129.2 (d, Ar-CH), 128.1 (d, Ar-CH), 128.0 (d, Ar-CH), 127.1 (d, 2 × Ar-CH), 126.5 (d, Ar-CH), 125.7 (d, Ar-CH), 119.5 (s, Ar-C), 115.4 (s, Ar-C), 102.0 (d, Ar-CH), 77.5 (s, Ar-C), 60.6 (s, Ar-C), 60.1 (q, OCH₃), 56.1 (q, OCH₃), 26.6 (q, 2 × CH₃), 21.4 (q, 2 × CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₈H₃₁O₄]⁺ = [M + H]⁺: 431.2217; found 431.2214.

5,6,7-Trimethoxy-1,1-dimethyl-3,4-bis(4-methylphenyl)-1*H*-isochromene (**3fc**)

GP was carried out with 2-(2-bromo-3,4,5-trimethoxyphenyl)propan-2-ol **1f** (76 mg, 0.25 mmol), 1-methyl-4-[(4-methylphenyl)ethynyl]benzene **2c** (128.8 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 90 : 10 to 85 : 15) to obtain isochromene **3fc** (21.5 mg, 20%) as a brown jelly compound, [TLC control (petroleum ether/ethyl acetate 85 : 15), R_f (**2c**) = 0.5, R_f (**3fc**) = 0.4, UV detection]. IR (MIR-ATR, 4000–600 cm⁻¹): ν_{max} = 2925, 2855, 2365, 1742, 1697, 1649, 1512, 1462, 1232, 1158, 1099, 828, 758, 700 cm⁻¹. ¹H NMR (CDCl₃, 400 MHz): δ = 7.15–7.06 (m, 4H, Ar-H), 7.01 (d, *J* = 7.3 Hz, 2H, Ar-H), 6.92 (d, *J* = 7.8 Hz, 2H, Ar-H), 6.53 (s, 1H, Ar-H), 3.88 (s, 3H, OCH₃), 3.75 (s, 3H, OCH₃), 3.07 (s, 3H, OCH₃), 2.30 (s, 3H, CH₃), 2.23 (s, 3H, CH₃), 1.70 (s, 6H, 2 × CH₃) ppm. ¹³C{¹H} NMR (CDCl₃, 100 MHz): δ = 152.4 (s, Ar-C), 150.4 (s, Ar-C), 147.9 (s, Ar-C), 142.3 (s, Ar-C), 137.0 (s, Ar-C), 136.5 (s, Ar-C), 135.2 (s, Ar-C), 134.7 (s, Ar-C), 133.8 (s, Ar-C), 130.7 (d, 2 × Ar-CH), 128.5 (d, 2 × Ar-CH), 128.1 (d, 2 × Ar-CH), 128.0 (d, 2 × Ar-CH), 119.6 (s, Ar-C), 114.9 (s, Ar-C), 102.0 (d, Ar-CH), 77.2 (s, Ar-C), 60.6 (q, OCH₃), 60.2 (q, OCH₃), 56.1 (q, OCH₃), 26.6 (q, 2 × CH₃), 21.3 (q, CH₃), 21.2 (q, CH₃) ppm. HR-MS (APCI+) *m/z* calculated for [C₂₈H₃₁O₄]⁺ = [M + H]⁺: 431.2217; found 431.2220.

7-Chloro-1,1-dimethyl-3,4-diphenyl-1*H*-isochromene (**3ga**)

GP was carried out with 2-(2-bromo-5-chlorophenyl)propan-2-ol **1g** (61.9 mg, 0.25 mmol), (phenylethynyl)benzene **2a** (111.2 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K₂CO₃ (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 100 : 1 to 98 : 2) to obtain isochromene **3ga**



(51.8 mg, 60%) as a light yellow jelly compound, [TLC control (petroleum ether/ethyl acetate 98 : 2), R_f (**2a**) = 0.7, R_f (**3ga**) = 0.6, UV detection]. IR (MIR-ATR, 4000–600 cm^{-1}): ν_{max} = 3405, 3042, 1692, 1602, 1453, 1320, 1260, 1088, 1025, 793, 744, 700 cm^{-1} . ^1H NMR (CDCl_3 , 400 MHz): δ = 7.36–7.28 (3H, m), 7.25–7.18 (4H, m), 7.18–7.11 (4H, m), 7.07 (1H, dd, J = 8.4, 2.1 Hz), 6.80 (1H, d, J = 8.4 Hz), 1.76 (6H, s) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3 , 100 MHz): δ = 148.3 (s, Ar-C), 137.9 (s, Ar-C), 136.6 (s, Ar-C), 135.6 (s, Ar-C), 132.1 (s, Ar-C), 131.5 (d, $2 \times$ Ar-CH), 128.8 (d, $2 \times$ Ar-CH), 128.7 (d, $2 \times$ Ar-CH), 127.9 (d, Ar-CH), 127.5 (d, $2 \times$ Ar-CH), 127.2 (d, Ar-CH), 127.1 (d, Ar-CH), 125.9 (s, Ar-C), 125.0 (d, Ar-CH), 122.5 (d, Ar-CH), 114.9 (s, Ar-C), 77.4 (s, Ar-C), 26.9 (q, $2 \times$ CH_3) ppm.

3,4-Diphenyl-1-propyl-1H-isochromene (3ha)

GP was carried out with 1-(2-bromophenyl)butan-1-ol **1h** (57 mg, 0.25 mmol), (phenylethynyl)benzene **2a** (111.2 mg, 0.62 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K_2CO_3 (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 99 : 1 to 98 : 2) to obtain isochromene **3ha** (33.4 mg, 41%) as a light yellow jelly compound, [TLC control (petroleum ether/ethyl acetate 98 : 2), R_f (**2a**) = 0.8, R_f (**3ha**) = 0.7, UV detection]. IR (MIR-ATR, 4000–600 cm^{-1}): ν_{max} = 3085, 2973, 2839, 1607, 1486, 1424, 1295, 1211, 1040, 831, 698 cm^{-1} . ^1H NMR (CDCl_3 , 400 MHz): δ = 7.40–7.28 (m, 3H, Ar-H), 7.28–7.20 (m, 4H, Ar-H), 7.20–7.03 (m, 6H, Ar-H), 6.86 (dd, J = 7.3 and 1.0 Hz, 1H, Ar-H), 5.28 (dd, J = 8.3 and 4.9 Hz, 1H, Ar-H), 2.29–2.11 (m, 1H, CH_2), 2.02–1.84 (m, 1H, CH_2), 1.77–1.61 (m, 1H, CH_2), 1.60–1.46 (m, 1H, CH_2), 1.01 (t, J = 7.34 Hz, 3H, CH_3) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3 , 100 MHz): δ = 148.9 (s, Ar-C), 137.0 (s, Ar-C), 135.7 (s, Ar-C), 132.8 (s, Ar-C), 131.6 (d, $2 \times$ Ar-CH), 128.7 (d, Ar-CH), 128.7 (s, Ar-C), 128.6 (d, $2 \times$ Ar-CH), 127.7 (s, Ar-C), 127.1 (d, Ar-CH), 127.5 (d, Ar-CH), 127.4 (d, $2 \times$ Ar-CH), 126.9 (d, Ar-CH), 126.5 (d, Ar-CH), 123.6 (d, Ar-CH), 123.2 (d, Ar-CH), 115.9 (s, Ar-C), 77.4 (s, Ar-C), 36.0 (t, CH_2), 18.7 (t, CH_2), 14.0 (q, CH_3) ppm. HR-MS (APCI+) m/z calculated for $[\text{C}_{24}\text{H}_{23}\text{O}]^+$ = $[\text{M} + \text{H}]^+$: 327.1743; found 327.1746.

3,4-Bis(4-fluorophenyl)-1-propyl-1H-isochromene (3hh)

GP was carried out with 1-(2-bromophenyl)butan-1-ol **1h** (57 mg, 0.25 mmol), 1-fluoro-4-[(4-fluorophenyl)ethynyl]benzene **2h** (133.7 mg, 0.625 mmol), palladium acetate (2.8 mg, 5 mol%), L-proline (5.6 mg, 20 mol%), TBAI (92.3 mg, 0.25 mmol), K_2CO_3 (138.6 mg, 1 mmol) and water (0.5 mL). The crude material was then refined by silica gel column chromatography (petroleum ether/ethyl acetate, 100 : 0 to 99 : 1) to obtain isochromene **3hh** (39.8 mg, 44%) as a light yellow jelly compound, [TLC control (petroleum ether/ethyl acetate 100 : 0), R_f (**2h**) = 0.7, R_f (**3hh**) = 0.6, UV detection]. IR (MIR-ATR, 4000–600 cm^{-1}): ν_{max} = 3096, 2974, 2930, 2842, 1610, 1491, 1423, 1297, 1220, 1044, 837, 701 cm^{-1} . ^1H NMR (CDCl_3 , 400 MHz): δ = 7.24–7.13 (m, 6H, Ar-H), 7.13–6.99 (m, 3H, Ar-H), 6.91–6.77 (m, 3H, Ar-H), 5.26 (dd, J = 8.6 and 5.1 Hz, 1H, Ar-H), 2.24–2.07 (m, 1H, CH_2), 1.99–1.83 (m, 1H, CH_2), 1.73–1.46 (m, 2H, CH_2), 1.01 (t, J = 7.3 Hz, 3H,

CH_3) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3 , 100 MHz): δ = 162.1 (d, J = 248.0 Hz, Ar-C), 161.9 (s, J = 246.5 Hz, Ar-C), 148.3 (s, Ar-C), 133.2 (d, Ar-CH), 133.1 (d, Ar-CH), 132.1 (d, J = 103.4 Hz, Ar-C), 131.9 (d, J = 103.4 Hz, Ar-C), 131.5 (d, J = 103.4 Hz, Ar-C), 130.6 (d, Ar-CH), 130.5 (d, Ar-CH), 127.6 (d, $2 \times$ Ar-CH), 126.7 (d, $2 \times$ Ar-CH), 123.7 (d, $2 \times$ Ar-CH), 123.0 (d, $2 \times$ Ar-CH), 115.7 (d, J = 21.3 Hz, Ar-CH), 114.6 (d, J = 21.3 Hz, Ar-CH), 77.5 (d, Ar-CH) 35.9 (t, CH_2) 18.70 (t, CH_2) 14.03 (q, CH_3) ppm. HR-MS (APCI+) m/z calculated for $[\text{C}_{24}\text{H}_{21}\text{F}_2\text{O}]^+$ = $[\text{M} + \text{H}]^+$: 363.1555; found 363.1563.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We are grateful to the Department of Science and Technology-Science and Engineering Research Board (DST-SERB) [No. EMR/2017/005312], New Delhi, for financial support. K. R. thanks MHRD, New Delhi, for the award of a research fellowship.

Notes and references

- (a) L. F. Tietze and N. Rackelmann, *Pure Appl. Chem.*, 2004, **76**, 1967–1983; (b) L. F. Tietze, *Chem. Rev.*, 1996, **96**, 115–136; (c) G. Satyanarayana, C. Maichle-Mössmer and M. E. Maier, *Chem. Commun.*, 2009, 1571–1573.
- (a) J. Lee and J. S. Panek, *Org. Lett.*, 2014, **16**, 3320–3323; (b) J. M. Gao, S. X. Yang and J. C. Qin, *Chem. Rev.*, 2013, **113**, 4755–4811.
- M. A. Brimble, L. J. Duncalf and M. R. Nairn, *Nat. Prod. Rep.*, 1999, **16**, 267–281.
- W. Wang, T. Li, R. Milburn, J. Yates, E. Hinnant, M. J. Luzzio, S. A. Noble and G. Attardo, *Bioorg. Med. Chem. Lett.*, 1998, **8**, 1579–1584.
- R. Mutter and M. Wills, *Bioorg. Med. Chem.*, 2000, **8**, 1841–1860.
- M. Masubuchi, K. Kawasaki, H. Ebiike, Y. Ikeda, S. Tsujii, S. Sogabe, T. Fujii, K. Sakata, Y. Shiratori, Y. Aoki, T. Ohtsuka and N. Shimma, *Bioorg. Med. Chem. Lett.*, 2001, **11**, 1833–1837.
- A. Shimbashi and S. Nishiyama, *Tetrahedron Lett.*, 2007, **48**, 1545–1548.
- (a) T. Oja, P. San Martin Galindo, T. Taguchi, S. Manner, M. Vuorela, K. Ichinose, M. Metsä-Ketelä and A. Fallarero, *Antimicrob. Agents Chemother.*, 2015, **59**, 6046–6052; (b) Y. S. Nanayakkara, R. M. Woodsb, Z. S. Breitbachb, S. Handac, L. M. Slaughterc and D. W. Armstrong, *J. Chromatogr. A*, 2013, **1305**, 94–101.
- H. S. Kang, E. M. Jun, S. H. Park, S. J. Heo, T. S. Lee, I. D. Yoo and J. P. Kim, *J. Nat. Prod.*, 2007, **70**, 1043–1045.
- H. He, H. Y. Yang, S. W. Luckman, D. M. Roll and G. T. Carter, *J. Antibiot.*, 2002, **55**, 1072–1075.



- 11 I. S. Chen, I. W. Tsai, C. M. Teng, J. J. Chen, Y. L. Chang, F. N. Ko, M. C. Lu and J. M. Pezzuto, *Phytochemistry*, 1997, **46**, 525–529.
- 12 H. Hussain, J. Hussain, S. Alib, A. Al-Harrasi, M. Saleem, G. A. Miana, M. Riaz, S. Anwar, S. Hussain and L. Alia, *J. Asian Nat. Prod. Res.*, 2012, **14**, 297–300.
- 13 (a) J. Kawai, P. K. Chikkade, Y. Shimizu and M. Kanai, *Angew. Chem., Int. Ed.*, 2013, **52**, 7177–7180; (b) C. Chan, Y. Chan and M. Chang, *Tetrahedron*, 2016, **72**, 1–8; (c) R. Mancuso, S. Mehta, B. Gabriele, G. Salerno, W. S. Jenks and R. C. Larock, *J. Org. Chem.*, 2010, **75**, 897–901.
- 14 (a) T. Uto, M. Shimizu, K. Ueura, H. Tsurugi, T. Satoh and M. Miura, *J. Org. Chem.*, 2008, **73**, 298–300; (b) K. Morimoto, K. Hirano, T. Satoh and M. Miura, *J. Org. Chem.*, 2011, **76**, 9548–9551; (c) M. Fukui, Y. Hoshino, T. Satoh, M. Miura and K. Tanakaa, *Adv. Synth. Catal.*, 2014, **356**, 1638–1644; (d) S. Nakanowatari and L. Ackermann, *Chem.–Eur. J.*, 2014, **20**, 5409–5413; (e) D. Yue, N. D. Ca and R. C. Larock, *J. Org. Chem.*, 2006, **71**, 3381–3388; (f) R. Mancuso, S. Mehta, B. Gabriele, G. Salerno, W. S. Jenks and R. C. Larock, *J. Org. Chem.*, 2010, **75**, 897–901; (g) A. Varela-Fernández, C. González-Rodríguez, J. A. Varela, L. Castedo and C. Saá, *Org. Lett.*, 2009, **11**, 5350–5353; (h) M. Dell'Acqua, B. Castano, C. Cecchini, T. Pedrazzini, V. Pirovano, E. Rossi, A. Caselli and G. Abbiati, *J. Org. Chem.*, 2014, **79**, 3494–3505.
- 15 E. K. Yum, M. J. Doty, K. K. C. Sham and R. C. Larock, *J. Org. Chem.*, 1995, **60**, 3270–3271.
- 16 W. J. Ang, C. Tai, L. Lo and Y. Lam, *RSC Adv.*, 2014, **4**, 4921–4929.
- 17 (a) A. Gopi Krishna Reddy and G. Satyanarayana, *J. Org. Chem.*, 2016, **81**, 12212–12222; (b) C. Sreenivasulu and G. Satyanarayana, *Eur. J. Org. Chem.*, 2018, 2846–2857; (c) K. Ramesh and G. Satyanarayana, *Eur. J. Org. Chem.*, 2018, 4135–4146; (d) K. Ramesh and G. Satyanarayana, *Green Chem.*, 2018, **20**, 369–374; (e) K. Ramesh, S. Basuli and G. Satyanarayana, *Eur. J. Org. Chem.*, 2018, 2171–2177; (f) L. Mahendar and G. Satyanarayana, *J. Org. Chem.*, 2014, **79**, 2059–2074; (g) J. Krishna, A. G. K. Reddy and G. Satyanarayana, *Synlett*, 2013, **24**, 967–972.
- 18 (a) N. Ohtani, T. Ohta, Y. Hosoda and T. Yamashita, *Langmuir*, 2004, **20**, 409–415; (b) A. W. Herriott and D. Picker, *J. Am. Chem. Soc.*, 1975, **97**, 2345–2349; (c) N. Ohtani and Y. Hosoda, *Bull. Chem. Soc. Jpn.*, 2000, **73**, 2263–2268; (d) C. M. Starks and M. Halpern, *Phase-Transfer Catalysis: Fundamentals, Applications, and Industrial Perspectives*, Springer Science & Business Media, 2012; (e) P. William, W. George and W. Gokel, *Phase Transfer Catalysis in Organic Synthesis*, Springer Science & Business Media, 2012; (f) A. Bharat Kumar and S. Krishna Nand, *Synthesis*, 2011, **7**, 1125; (g) A. de Meijere and F. E. Meyer, *Angew. Chem., Int. Ed.*, 1994, **33**, 2379.
- 19 (a) F. Bellina and R. Rossi, *Chem. Rev.*, 2010, **110**, 3850; (b) T. Konno, J. Chae, T. Ishihara and H. Yamanaka, *Tetrahedron*, 2004, **60**, 11695–11700; (c) A. Casnati, R. Maggi, G. Maestri, N. Della Ca and E. Motti, *J. Org. Chem.*, 2017, **82**, 8296–8303; (d) Q. Tian, A. A. Pletnev and R. C. Larock, *J. Org. Chem.*, 2003, **68**, 339–347; (e) F. Bellina and R. Rossi, *Chem. Rev.*, 2010, **110**, 3850; (f) R. C. Larock, M. J. Doty and X. Han, *J. Org. Chem.*, 1999, **64**, 8770–8779; (g) H. Miao and Z. Yang, *Org. Lett.*, 2000, **2**, 1765–1768.

