



Cite this: *RSC Adv.*, 2020, 10, 1829

Benzyl isothiocyanate fumigation inhibits growth, membrane integrity and mycotoxin production in *Alternaria alternata*†

Tiaolan Wang,  Yongcai Li,* Yang Bi, Miao Zhang, Tingting Zhang, Xiaoyuan Zheng, Yupeng Dong and Yi Huang

The antifungal activity of benzyl isothiocyanate (BITC) against pear pathotype-*Alternaria alternata*, the causal agent of pear black spot, and its possible mechanisms were studied. The results indicated that both the spore germination and mycelial growth of *A. alternata* were significantly inhibited by BITC in a dose-dependent manner. BITC concentrations at 1.25 mM completely suppressed mycelial growth of *A. alternata* and prevented $\geq 50\%$ of black spot development in wounded pears inoculated with *A. alternata*. Microscopic analyses and propidium iodide (PI) staining showed that spore morphology in *A. alternata* treated with BITC at 0.625 mM was severely damaged. Relative electrical conductivity and lysis ability assays further showed that BITC treatment destroyed the integrity of the plasma membrane. Additionally, mycotoxin production was inhibited by 0.312 mM BITC, and the inhibitory rates of alternariol monomethyl ether (AME), alternariol (AOH), altenuene (ALT) and tentoxin (TEN) were 89.36%, 84.57%, 91.41% and 67.78%, respectively. The above results suggest that BITC exerts antifungal activity through membrane-targeted mechanisms.

Received 7th November 2019
Accepted 30th December 2019

DOI: 10.1039/c9ra09225k

rsc.li/rsc-advances

1. Introduction

Alternaria sp. is a common genus of ascomycete fungi that are both saprophytic on organic materials and pathogenic to many plants.¹ Infection by *Alternaria* sp. leads to black spot rot of fresh fruits and vegetables, including pears, citrus, apricots, cherry tomatoes and blueberries. During the development and storage, black spots in pears caused by *Alternaria alternata* is one of the major postharvest diseases leading to fruit deterioration and rotting, which results in economic losses.^{2–5} *A. alternata* is a dematiaceous fungus, characterized by dark colonies ranging from grey to olive/brown.⁶ Microscopic characteristics of *A. alternata* conidia were ovoid or elliptical, with one to five transverse septa and none or three longitudinal septum. When the conidia adhere to the surface of the fruit, they germinate from the top and partially form the infectious hyphae which colonize the fruit cells and tissues, thereby capturing the nutrients and water of the host, and also producing harmful metabolites such as mycotoxins.⁷ The necrotrophic pathogen *A. alternata* in citrus exerts pathogenesis through the production of host-specific toxins.⁸ The most common host-specific toxins of *Alternaria* sp. in food items are alternariol monomethyl ether (AME), alternariol (AOH),

altenuene (ALT) and tentoxin (TEN),^{9–11} which accumulate in fruits and processed food, leading to food safety issues and a threat to human health.^{1,12} Currently, chemical fungicides have been extensively used as a primary means of combating black spot rot.¹³ In addition to its antifungal activity, inhibiting mycotoxin synthesis and reducing mycotoxin levels are major methods to control the postharvest decay of vegetables and fruits.¹⁴ However, pathogen resistance to various types of fungicides can develop after frequent usage. In addition, synthetic fungicides are not friendly for environment and human health issues.¹⁵ Accordingly, natural antifungal products require development to meet environmentally friendly health demands.

Isothiocyanates (ITCs) are naturally-occurring constituents that bear a wide biocidal spectrum, with potential as post-harvest fumigants.^{16–19} ITCs exhibit biocidal activity against fungi,^{20,21} bacteria,²² insects and nematodes.^{23–25} ITCs vary according to the R-side groups on the parent glucosinolate, which can be aliphatic, aromatic or heteroaromatic.²⁶ Aromatic ITCs are more antifungal than ITCs with aliphatic R groups,²⁷ and aromatic BITC show higher biocidal activity against *Sclerotinia sclerotiorum* and *Rhizoctonia solani* compared to ITCs derived from the aliphatic allyl-isothiocyanate and aromatic *p*-hydroxybenzyl.²⁸ BITC treatment of low-density polyethylene film (LDPE) bags is an active control strategy for *Alternaria* rot in tomato fruit.²⁹ BITC also shows potent inhibitory effects against *Fusarium culmorum* and *Staphylococcus aureus* among the several isothiocyanate flavors separately treated.^{30,31} However,

College of Food Science and Engineering, Gansu Agricultural University, Lanzhou 730070, China. E-mail: lyc@gsau.edu.cn

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra09225k



the mechanisms by which BITC inhibits fungal activity have not been defined.

The cell membrane is the major target of toxic lipophilic compounds in eukaryotic cells.³² BITC is lipophilic and therefore, it may react with enzymes present at the plasma membrane, causing fungal growth inhibition and cell death.²⁹ In addition, mycotoxin production during fungal growth might interact with ITCs after membrane destruction, reducing the levels of aflatoxins produced by *Aspergillus parasiticus* CECT 2681 in nuts (peanut, cashew, almond, hazelnut and pistachio).³³ I. Clemente *et al.* used active packaging containing BITC to reduce or eliminate ochratoxin production.³⁴ However, M. Mari *et al.* found that benzyl isothiocyanate showed little or no activity against the pathogens tested on *Botrytis cinerea* and *R. stolonifera*.³⁵ There are few studies focusing on the growth, membrane integrity and related mycotoxin production of *A. alternata* by BITC fumigated. In this study, we evaluated the antifungal activity of BITC against *A. alternata* through fumigation and *in vitro/in vivo* assessments. The effects of BITC on cell membrane integrity and mycotoxin production of *A. alternata* were also investigated.

2. Materials and methods

2.1 Chemicals

Standard benzyl isothiocyanate (CAS: 622-78-6) was purchased by the Tokyo Chemical Industry (Shanghai). BITC was reagent-grade with 99.9% purity and dissolved in 20% ethanol-sterile distilled water (v/v) and then filtrated through a 0.22 μm microporous membrane.

2.2 Fruit and pathogen

'Zaosu' pears (*Pyrus bretschneideri* Rehd) free of physical injury or infection were commercially harvested at the Tiaoshan Farm in Jingtai County, Gansu Province, China, and individually packed and stored at 4 °C cold storage at the laboratory. Pear fruits were used for *in vivo* assessments within a week.

A. alternata (JT03) was obtained from the post-harvest laboratory at Gansu Agricultural University, Gansu Province, China. The spores were harvested from 6 day-old PDA cultivates and suspended in 20 mL of sterile distilled water. Suspensions were followed by filtration through four layers of sterile gauze. The number of spores (1×10^5 or 1×10^6 spores per mL) were determined using hemocytometer for *in vivo* and *in vitro* testing.

2.3 Monitoring spore germination

The 20 μL *A. alternata* spore suspensions (1×10^5 spores per mL) were added to an agar plugs (diameter = 5 mm) (in triplicate) and placed onto a glass slides. Then the slides were placed in a glass Petri dish (diameter = 18 cm) with the same size of moist filter paper at the bottom to provide with 90–92% relative humidity. Sterile filter paper discs (diameter = 6 mm) containing BITC (0, 0.039, 0.078, 0.156, 0.312, 0.625, and 1.25 mM) were placed on the surface of the Petri dish and incubated at 28 °C for 2, 4, 6 and 8 hours. The cultures were taken at the corresponding time point, and stained with 20 μL of

lactophenol cotton blue for 2 minutes. And 100 spores were counted under a microscope to determine germination rates. Each BITC concentration was assessed in triplicate.

2.4 Mycelial growth assays

The effects of BITC on the mycelial growth of *A. alternata* were assessed on PDA plates according to the methods described by N. L. Tatsadjieu *et al.*³⁶ Autoclaved PDA medium which was cooled to about 60 °C was poured into Petri dishes (diameter = 90 mm). Sterile filter paper discs (diameter = 6 mm) containing various concentrations of BITC (0, 0.039, 0.078, 0.156, 0.312, 0.625, and 1.25 mM) were placed on the surface of the Petri dish. A 6 mm plug of mycelial agar was obtained from the edge of 6 day-old cultures of *A. alternata* and transferred into the center of each dish. Plates were sealed with parafilm and the radial growth of *A. alternata* was measured after incubation at 28 °C for 3, 5 and 7 day. Each treatment was applied to three replicates and the entire experiment was repeated three.

2.5 *In vivo* antifungal activity assays

The *in vivo* assays were performed as previously described with subtle modifications.^{37–39} Pears lacking physical injury or infections were selected based on size uniformity, surface-disinfected with 1% (v/v) sodium hypochlorite for 2 min, rinsed with tap water, and air-dried prior to use. Pears fruits were wounded (3 mm wide and 3 mm deep) with a sterile dissecting needle at the equator prior to inoculation. Each wound site was inoculated with 20 μL 1×10^5 spores per mL *A. alternata* and air dried at room temperature. Selected pears were placed in sealed boxes and fumigated with BITC (0, 0.312, 0.625, 1.25 mM) for 12 h (concentrations were selected based on preliminary *in vitro* experiments), and then stored at room temperature (25 ± 2 °C), 90–92% relative humidity for 9 days. The lesion diameter was measured every 2 days. There were three replicates with 10 fruits in each treatment, and the experiment was repeated three times.

2.6 Determination of membrane integrity

PI staining was performed as previously described with minor modifications.⁴⁰ *A. alternata* spores were cultured on PDA for 6 day and fumigated with BITC at 0, 0.156, 0.312 and 0.625 mM. Spores were harvested from PDA and suspended in 20 mL of 100 mM PBS (pH 7.4). After incubation of *A. alternata* spores for 4 hours, PI was added to a final concentration at 2.5 $\mu\text{g mL}^{-1}$ and maintained at 28 °C for 10 minutes in the dark. Mixtures were centrifuged at $8000 \times g$ for 2 min at 4 °C and precipitates were washed twice in PBS to remove residual dye. Samples were then resuspended in 500 μL of PBS and then 10 μL suspensions were dropped on slides. Unstained spore suspensions were used as controls. Samples were imaged under a fluorescence microscope (U-LH100HG 19v 100w, Tokyo 163-0914, Japan). The percentages of fluorescent spores in each population were calculated. Each experiment was performed in triplicate and the experiment was repeated three.



2.7 Cytoplasmic leakage assays

The leakage of cytoplasmic contents from *A. alternata* mycelia were determined according to the methods of J. Tian *et al.* and D. R. N. A. Cezar *et al.* with minor modifications.^{41,42} A total of 6–8 plugs of *A. alternata* mycelial agar (diameter = 6 mm) fumigated with BITC at 0, 0.312 and 0.625 mM were obtained from the edge of 6 day-old cultures on PDA and transferred into PDB medium at 28 °C on a rotary shaker (150 rpm). After 3 days of incubation, mycelia were harvested onto filter paper (pore diameter 10 µm) and washed with sterile distilled water. Mycelia was resuspended in 50 mL sterile distilled water and incubated on a rotary shaker at 26 °C for 0, 30, 60, 90 and 120 min. Cell membrane permeability was determined using an electric conductivity meter (DDS-307A, Shanghai, China). Nucleic acids levels were measured through the assessment of optical density at 260 nm (OD₂₆₀). All experiments were performed in triplicate.

2.8 Extraction of mycotoxins

2.8.1 Standards. Certified standards of *Alternaria* toxins, namely AME, AOH, ALT and TEN were purchased from Shanghai Yuanye Bio-Technology Co., Ltd (China). Solid portions of standards were dissolved in MeCN to prepare 100 µg mL⁻¹ stock solutions, and a portion of solutions were prepared from the stock standard solutions at 10 µg mL⁻¹ in MeCN, which were stored in the dark at –20 °C. The spiking solution (1 µg mL⁻¹) was freshly prepared from the mix solutions, and the working solutions with concentrations of 1–200 ng mL⁻¹ were freshly prepared just before using with the blank matrix.

2.8.2 Sample preparation. The method have been appropriately modified as described by P. López *et al.*⁹ The fungus were cultured in PDA for 4 day at 28 °C and fumigated with BITC at 0 and 0.312 mM according to previous experiment. *A. alternata* mycelium (0.5 g) was grounded in an ice bath and the hyphae were transferred to sterile centrifuge tubes containing 2.5 mL of acetonitrile/water (80 : 20) and 0.3% formic acid. Extracts were sonicated for 60 min and agitated in a shaker agitator for 30 min at 28 °C and 150 rpm. Next, 0.25 g of anhydrous MgSO₄ and 0.04 g NaCl were added while shaking for 1 min, and samples were centrifuged at 10 000 rpm for 10 min. Supernatant was passed through a 0.22 µm organic filter to

a volume of 1.2 mL for high performance liquid chromatography-tandem mass spectrometry (parameters described in Table 1; standard curve for standard samples in Fig. S1†).

2.9 Statistical analysis

Experiments were performed on three independent occasions and each replicate was performed in triplicate. Data was statistically analyzed using an analysis of variance (ANOVA) and expressed as the mean ± standard deviation (SD). Mean separations were analyzed using Duncan's multiple range tests and differences amongst the different treatments were determined at the 5% level (IBM SPSS Statistics 20, USA).

3. Results

3.1 BITC inhibited spore germination and mycelial growth of *A. alternata*

BITC treatment effectively inhibited spore germination and the mycelial growth of *A. alternata* in a dose-dependent manner (Fig. 1A and B). Spore germination rates of 0.625 mM BITC fumigated *A. alternata* decreased by 82.89% compared to that of the control ($p < 0.05$). When the concentration of BITC was increased to 1.25 mM, spores germination were entirely inhibited (Fig. 1A). Visible hyphae of 0.625 mM BITC treated *A. alternata* was observed after 2 days of cultivation (Fig. 1B), and the inhibition rate of mycelial growth was up to 41.05% ($p < 0.05$) 4 days after treatment. BITC at 1.25 mM completely inhibited mycelial growth. Colony morphologies of the treated groups showed a more pronounced and regular round shape (Fig. 1C).

3.2 Efficacy of BITC against *A. alternata* in pears

As shown in Fig. 2, the development of black spots in pears inoculated with *A. alternata* was effectively inhibited by BITC fumigation. The lesion diameters of the black spots in pears treated with 0.625 and 1.25 mM BITC were reduced by 34.42% and 52.83% ($p < 0.05$) compared to control pears after 7 days of inoculation, respectively. BITC concentrations of 1.25 mM showed the highest level of disease control (Fig. 2A). Dark brown infected areas were observed on 1.25 mM BITC fumigated pear fruits (Fig. 2B).

Table 1 Optimized LC-MS/MS instrumental parameters^a

Analyte	Ionization mode	Retention time (min)	MRM (m/z , positive)	Fragmentor	Collision energy
Alternariol monomethyl ether (AME)	ESI [−]	2.85	271.0 → 228.0 (Q) 271.0 → 256.0 (I)	32	20
Alternariol (AOH)	ESI [−]	2.37	257.0 → 147.2 (Q) 257.0 → 213.0 (I)	32	20
Allenuene (ALT)	ESI ⁺	3.33	293.1 → 239.1 (Q) 293.1 → 257.2 (I)	85	15
Tentoxin (Ten)	ESI ⁺	3.66	415.2 → 189.0 (Q) 415.2 → 312.3 (I)	110	30

^a MRM: (Q) transition used for quantification and (I) transition employed to confirm the identification.






Fig. 1 Effects of BITC on spore germination rates (A), the mycelial diameter (B) and colony morphology (C) of *A. alternata* after treatment with 0.312, 0.625 and 1.25 mM BITC. Vertical bars represent standard errors. Columns marked by different letters (such as a and b) indicate that there are significant differences in the treatment within the group, and identical letters (such as c and d) indicate no significant difference in treatment within the group according to Duncan's multiple range tests ($p < 0.05$).

3.3 Cell membrane integrity by PI staining

PI staining was performed for *A. alternata* cells in the presence and absence of BITC. As shown in Fig. 3A, BITC treatment led to

a loss of cell membrane integrity of *A. alternata* in a concentration dependent manner, evidenced by the increased intensity of PI staining. The percentage of spores with PI fluorescence which




Fig. 2 Harvested pears were treated with BITC at 0.312, 0.625 and 1.25 mM and inoculated with *A. alternata* and stored at 25 °C. Lesion diameter (A) was measured and imaged (B) every two days post inoculation. Vertical bars represent the standard error of treatment means. Columns marked by differed lowercase letters are significantly different according to Duncan's multiple range tests ($p < 0.05$).






Fig. 3 Effects of BITC treatment on the plasma membrane integrity of *A. alternata*. Spores incubated with different concentrations of BITC (0, 0.312 and 0.625 mM) were stained with propidium iodide (PI) and imaged under a fluorescence microscope. Spores with damaged plasma membranes showed fluorescence (A), and the percentage of spores with reduced plasma membrane integrity were analyzed (B). Three fields of view were randomly chosen for each treatment. Experiments were repeated on three occasions. Vertical bars represent standard errors. Columns marked by different lowercase letters significantly differed according to Duncan's multiple range tests ($p < 0.05$).

impaired membrane integrity was as high as 90% in the presence of 0.625 mM BITC (Fig. 3B).

3.4 Cellular leakage of *A. alternata* following BITC treatment

As shown in Fig. 4A, cell membrane conductivity values of non-treated *A. alternata* slowly increased slowly within 0–120 min of incubation, whilst the external conductivity of BITC treated *A. alternata* cells showed a rapid upwards trend. The conductivity values increased with the increasing of BITC concentrations. The conductivity of 0.625 mM BITC treated *A. alternata* increased and peaked at 60 min (Fig. 4A), which was significantly higher than observed in control samples (37.4 ± 0.063). In addition, significant differences in nucleic acid leakage were detected comparing to the control (Fig. 4B). The nucleic acid content characterized by OD_{260} in the filtrates of *A. alternata* treated with BITC significantly increased during the incubation

period. After 60 min, the OD_{260} values in 0.625 mM BITC-treated *A. alternata* filtrates were 7.2-fold higher than that in the controls (Fig. 4 B).

3.5 BITC treatment inhibited mycotoxin production in *A. alternata*

The retention time was determined by the standard curve and four mycotoxins were identified: AME, 2.845 min; AOH, 2.364 min; ALT, 2.333 min; TEN, 2.666 min (Table 1). The mass were 271.0 and 257.0 m/z under negative ionization mode for AME and AOH (Fig. S2A and C[†]). The quasi-molecular ions of ALT and TEN (m/z 293.1, $[M + H]^+$; m/z 415.2, $[M + H]^+$) suggest a molecular mass of 292 and 414 respectively (Fig. S2E and F[†]). Samples fumigated with 0.312 mM BITC showed significantly lower levels of mycotoxins (AME, AOH, ALT and TEN) in the *A. alternata* hyphae, and inhibitory effects varied with each type of




Fig. 4 Effects of BITC on electrical conductivity (A) and nucleic acid leakage (B) of *A. alternata*. Vertical bars represent standard errors. Asterisk (*) are significantly different according to the Duncan's multiple range test ($p < 0.05$).

mycotoxins. The inhibitory rates of BITC treatment on AME and AOH were 89.36% and 84.57% ($p < 0.05$), respectively (Fig. 5A and B). The TEN content decreased to $32.60 \pm 0.46 \mu\text{g g}^{-1}$ compared to the control ($141.70 \pm 1.73 \text{ ng L}^{-1}$) (Fig. 5D). The inhibitory rates of ALT were greater than 90% (Fig. 5C).

4. Discussion

ITCs is natural compounds with antimicrobial properties.¹⁷ Previous studies have shown that 0.1 mg L^{-1} ITCs reduced the incidence of *Botrytis cinerea* in strawberries by over 45% without detrimental effects on post-harvest quality.³⁸ Our results showed that the incidence of *A. alternata* infection in pears fumigated with BITC at 1.25 mM for 12 h was significantly lower after 7 d of storage, in which the fruit showed the smallest infection areas, consistent with the findings of L. Ugolini *et al.*³⁸ However, BITC concentrations of 1.25 mM led to a complete inhibition of mycelial growth in *A. alternata*. Thus inhibition of BITC on the growth was proved to be lower *in vivo* compared with the *in vitro* trial, these differences may be due to the diverse characteristics of stability through time and volatility in the compounds studied, as well as the diffusive effect of the active substance inside the wound.³⁵ Visible mycelial began to grow after 2 days of treatment with 0.625 mM BITC. However,

concentrations of 0.75 mg mL^{-1} BITC completely inhibited *Fusarium culmorum* growth *in vitro*.³⁰ Those differences may be due to variable compound purity or application methods. M. Mari *et al.* demonstrated the inhibition of conidial germination and mycelial growth of several postharvest fruit pathogens by BITC and other natural ITCs.^{39,43} Our results were consistent with these studies, as BITC efficiently inhibited spore germination and colony expansion of *A. alternata* on PDA medium. Spore germination was incompletely inhibited at concentrations of 0.039 – 1.25 mM for 8 h, with spore germination rates decreasing from 86.67% to 3%. S. G. Harvey reported that the spore germination of *Sclerotium rolfsii* inhibition of 100% was not recorded even at the highest concentrations of 2-propenyl-isothiocyanate (528.8 mM), indicating that the spores were more resistant than mycelia.⁴⁴ Our results suggested that BITC efficiently inhibits spore germination rates and colony expansion of *A. alternata* on PDA medium, and alleviates disease severity in pear fruits.

To further elucidate the mechanisms underlying the anti-fungal activity of BITC against *A. alternata*, a loss of membrane integrity and the leakage of cellular constituents of BITC-treated *A. alternata* were determined. BITC is lipophilic and may destroy membrane integrity.²⁹ PI is a DNA-stain that shows red fluorescence when inserted into double-stranded




Fig. 5 Effects of 0.312 mM BITC treatment on AME, AOH, ALT and TEN mycotoxins content in *A. alternata*. Vertical bars represent standard errors. Asterisk (*) indicates significant differences amongst sample groups ($P < 0.05$).



DNA in cells that have lost membrane integrity.^{41,45} A higher number of PI stained *A. alternata* spores were observed in response to BITC fumigation, with the membrane integrity of *A. alternata* almost completely destroyed at BITC concentration of 0.625 mM. These were similar to the values previously reported by F. Yan *et al.*⁴⁶

Lipophilic compound accumulates in the phospholipid bilayer and alters the permeability of the plasma membrane for K⁺ and H⁺, resulting in the disruption of ionic homeostasis.^{47,48} A loss of membrane integrity results in the leakage of small molecules ions and electrolytes from the cells.^{49,50} Membrane permeability parameters which are commonly used to indicate gross and irreversible damage to the cytoplasmic and plasma membranes included the loss of 260 nm absorbing materials (K⁺ and Na⁺ leakage).^{51,52} The release of electrolytes (Fig. 4A) and nucleic acids (Fig. 4B) in the fungal suspensions visibly increased with increasing BITC concentrations. The conductivity value of *A. alternata* suspensions treated with 0.312 and 0.625 mM of BITC increased over the initial 60 min, whilst the control group remained relatively stable within 0–120 min. The maximum release of cell constituents was observed in *A. alternata* suspensions treated with 0.625 mM BITC for 60 min, showing absorbances of 0.683. These findings suggest that the cytoplasmic membranes of *A. alternata* incurred irreversible damage after BITC fumigation. The results observed were in agreement with previous studies that suggested that the antibacterial activity of ITCs acts through the disturbance of membrane integrity.^{53,54}

I. Clemente *et al.* observed the effects of BITC on enzymes on the P450 family (Ochratoxins A-synthase).^{30,55,56} BITC can reduce the biosynthesis of secondary metabolites (Ochratoxins A and B) by affecting the relevant metabolic pathways of *Aspergillus ochraceus*.⁵⁷ *A. alternata* toxins are secondary metabolites produced by the *A. alternata* genera.⁵⁸ Some early reports have shown that *Alternaria* species do not produce the most known and frequent *Alternaria* toxins (e.g. AOH, AME, TEN),^{11,59} thus we conduct relevant exploratory experiments. The data presented in this study showed that high levels AME, AOH, ALT and TEN were detected in non-BITC treated *A. alternata*, similar to the report by T. T. T. Nguyen *et al.*⁶⁰ AME and AOH produced a high intensity (271.0 and 257.0 *m/z*) in negative mode (ESI[−]), and these results were consistent with the previous studies of A. A. Hildebrand *et al.* (2015).⁶¹ And significant reduction in mycotoxins contents was observed in BITC treated *A. alternata* (Fig. 5A and B), especially the inhibition rate of AME and ALT in response to BITC fumigation were 89.36% and 91.41%, respectively (*p* < 0.05). These results showed that BITC hold value for reducing mycotoxin contamination in *A. alternata*. Combination with the effect of BITC on the integrity and permeability of *A. alternata* plasma membrane, mycotoxin of *A. alternata* may leak out to extracellular medium and interact with BITC after the damage of cell membrane, leading to degradation of mycotoxins. However, its specific mechanism of action governing these effects needs further clarification.

5. Conclusion

BITC could inhibit *in vitro* spore germination and mycelial growth in *A. alternata* and reduce the occurrence of black spots in pears. BITC treatment destroyed the morphology of the spores and mycelia of the pathogens increased the leakage of intercellular electrolytes and nucleic acids of *A. alternata* leading to lethal effects. The mechanisms underlying the anti-fungal toxic activity of BITC were through membrane disruption and cell growth hindrance. BITC also significantly inhibited the production of *Alternaria* toxins (AME, AOH, ALT and TEN). BITC may thus offer an economical and environmentally-friendly strategy to control postharvest disease in fruit and vegetables.

Conflicts of interest

There is no conflict of interest.

Acknowledgements

This study was financed by the National Key R&D Plans of China (2018YFD0401302) and the National Natural Science Foundation of China (31860456, 31460534).

References

- 1 N. Estiarte, A. Crespo-Sempere, S. Marín, V. Sanchis and A. J. Ramos, Exploring polyamine metabolism of *Alternaria alternata* to target new substances to control the fungal infection, *Food Microbiol.*, 2017, **65**, 193–204.
- 2 C. Y. Diao, Y. Bi and Y. C. Li, The inhibition of chitosan on *Alternaria alternata* *in vitro*, *Chinese Agricultural Science Bulletin*, 2010, **26**, 91–94.
- 3 S. M. Sanzani, M. Reverberi and R. Geisen, Mycotoxins in harvested fruits and vegetables: insights in producing fungi, biological role, conducive conditions, and tools to manage postharvest contamination, *Postharvest Biol. Technol.*, 2016, **122**, 95–105.
- 4 C. Pane, F. Fratianni, M. Parisi, F. Nazzaro and M. Zaccardelli, Control of *Alternaria* post-harvest infections on cherry tomato fruits by wild pepper phenolic-rich extracts, *Crop Prot.*, 2016, **84**, 81–87.
- 5 M. Greco, A. Patriarca, L. Terminiello, V. F. Pinto and G. Pose, Toxigenic *Alternaria* species from Argentinean blueberries, *Int. J. Food Microbiol.*, 2012, **154**, 187–191.
- 6 I. Kustrzeba-Wójcicka, E. Siwak, G. Terlecki, A. Wolańczyk-Mędrala and W. Mędrala, *Alternaria alternata* and its allergens: a comprehensive review, *Clin. Rev. Allergy Immunol.*, 2014, **47**, 354–365.
- 7 Y. C. Li and Y. Bi, Occurrence and infection process of apple pear black spot disease, *J. Plant Prot.*, 2006, **33**, 131–135.
- 8 F. Kassel and C. Lillie, Coordinate control of oxidative stress tolerance, vegetative growth, and fungal pathogenicity via the AP1 pathway in the rough lemon pathotype of *Alternaria alternata*, *Physiol. Mol. Plant Pathol.*, 2010, **74**, 100–110.



- 9 P. López, D. Venema, H. Mol, M. Spanjer, J. De Stoppelaar and F. Pfeiffer, *Alternaria* toxins and conjugates in selected foods in the Netherlands, *Food Control*, 2016, **69**, 153–159.
- 10 V. Ostry, *Alternaria* mycotoxins: an overview of chemical characterization, producers, toxicity, analysis and occurrence in foodstuffs, *World Mycotoxin J.*, 2008, **1**, 175–188.
- 11 A. Patriarca, M. P. Azcarate, L. Terminiello and V. F. Pinto, Mycotoxin production by *Alternaria* strains isolated from Argentinean wheat, *Int. J. Food Microbiol.*, 2007, **119**, 219–222.
- 12 M. Wang, N. Jiang, Y. Wang, D. Jiang and X. Feng, Characterization of phenolic compounds from early and late ripening sweet cherries and their antioxidant and antifungal activities, *J. Agric. Food Chem.*, 2017, **65**, 5413–5420.
- 13 E. M. Soyulu and F. Kose, Antifungal activities of essential oils against citrus black rot disease agent *Alternaria alternata*, *J. Essent. Oil-Bear. Plants*, 2015, **18**, 894–903.
- 14 L. Xu, N. Tao and W. Yang, Cinnamaldehyde damaged the cell membrane of *Alternaria alternata* and induced the degradation of mycotoxins in vivo, *Ind. Crops Prod.*, 2018, **112**, 427–433.
- 15 J. D. S. Gabriel, M. J. Tiera and V. A. D. O. Tiera, Synthesis, characterization, and antifungal activities of amphiphilic derivatives of diethylaminoethyl chitosan against *Aspergillus flavus*, *J. Agric. Food Chem.*, 2015, **63**, 5725–5731.
- 16 S. K. Srivastava and S. V. Singh, Cell cycle arrest apoptosis induction and inhibition of nuclear factor kappa B activation in anti-proliferative activity of benzyl isothiocyanate against human pancreatic cancer cells, *Carcinogenesis*, 2004, **25**, 1701–1709.
- 17 E. A. S. Rosa and P. M. F. Rodrigues, Towards a more sustainable agriculture system: the effect of glucosinolates on the control of soil-borne diseases, *J. Hortic. Sci. Biotechnol.*, 1999, **74**, 667–674.
- 18 D. D. Archbold, T. R. Hamilton-Kemp and M. M. Barth, Identifying natural volatile compounds that control gray mold (*Botrytis cinerea*) during postharvest storage of strawberry, blackberry, and grape, *J. Agric. Food Chem.*, 1997, **45**, 4032–4037.
- 19 J. A. Kirkegaard and M. Sarwar, Biofumigation potential of brassicas—I. Variation in glucosinolate profiles of diverse field-grown brassicas, *Plant Soil*, 1998, **201**, 71–89.
- 20 U. Smolinska, M. J. Morra, G. R. Knudsen and R. L. James, Isothiocyanates produced by Brassicaceae species as inhibitors of *Fusarium oxysporum*, *Plant Dis.*, 2003, **87**, 407–412.
- 21 O. D. Dhingra, M. L. N. Costa and G. J. Silva, Potential of allyl isothiocyanate to control *Rhizoctonia solani* seedling damping off and seedling blight in transplant production, *J. Phytopathol.*, 2004, **152**, 352–357.
- 22 M. V. Galan, A. A. Kishan and A. L. Silverman, Oral broccoli sprouts for the treatment of *Helicobacter pylori* infection: a preliminary report, *Dig. Dis. Sci.*, 2004, **49**, 1088–1090.
- 23 R. Kermanshah, B. E. McCarry, J. Rosenfeld, P. S. Summers, E. A. Weretilnyk and G. J. Sorger, Benzyl isothiocyanate is the chief or sole anthelmintic in papaya seed extracts, *Phytochemistry*, 2001, **57**, 427–435.
- 24 I. A. Zasada and H. Ferris, Sensitivity of *Meloidogyne javanica* and *Tylenchulus semipenetrans* to isothiocyanates in laboratory assays, *Phytopathology*, 2003, **93**, 747–750.
- 25 R. F. Mithen, Glucosinolates and their degradation products, *Adv. Bot. Res.*, 2001, **35**, 213–262.
- 26 J. W. Fahey, A. T. Zalcmann and P. Talalay, The chemical diversity and distribution of glucosinolates and isothiocyanates among plants, *Phytochemistry*, 2001, **56**, 5–51.
- 27 R. Troncoso, C. Espinoza, A. Sánchez-Estrada, M. E. Tiznado and H. S. García, Analysis of the isothiocyanates present in cabbage leaves extract and their potential application to control *Alternaria* rot in bell peppers, *Food Res. Int.*, 2005, **38**, 701–708.
- 28 L. M. Manici, L. Lazzeri and S. Palmieri, *In vitro* fungitoxic activity of some glucosinolates and their enzyme-derived products toward plant pathogenic fungi, *J. Agric. Food Chem.*, 1997, **45**, 2768–2773.
- 29 R. Troncoso-Rojas, A. Sánchez-Estrada, C. Ruelas, H. S. García and M. E. Tiznado-Hernández, Effect of benzyl isothiocyanate on tomato fruit infection development by *Alternaria alternata*, *J. Sci. Food Agric.*, 2005, **85**, 1427–1434.
- 30 M. E. Tiznado-Hernández and R. Troncoso-Rojas, Control of fungal diseases with isothiocyanates, *Stewart Postharvest Review*, 2006, **2**(1), 1–14.
- 31 Y. Wang, Q. Q. Zhou and R. L. Wan, Inhibition of two gram-positive pathogenic bacteria by isothiocyanate spices, *Food Sci.*, 2017, **21**, 22–27.
- 32 J. Sikkema, J. A. de Bont and B. Poolman, Mechanisms of membrane toxicity of hydrocarbons, *Microbiol. Rev.*, 1995, **59**, 201–222.
- 33 C. Hontanaya, G. Meca, F. B. Luciano, J. Manes and G. Font, Inhibition of the Aflatoxins B1, B2, G1, and G2, production by *Aspergillus parasiticus* in nuts using yellow and oriental mustard flours, *Food Control*, 2015, **47**, 154–160.
- 34 I. Clemente, M. Aznar and C. Nerín, Effect of an active label based on benzyl isothiocyanate on the morphology and ochratoxins production of *Aspergillus ochraceus*, *Food Res. Int.*, 2017, **101**, 61–72.
- 35 M. Mari, R. Iori, O. Leoni and A. Marchi, Bioassay of glucosinolate-derived isothiocyanates against post-harvest pest pathogens, *Plant Pathol.*, 1996, **45**, 753–760.
- 36 N. L. Tatsadjieu, P. M. J. Dongmo, M. B. Ngassoum, F. X. Etoa and C. M. F. Mbofung, Investigations on the essential oil of *Lippia rugosa* from Cameroon for its potential use as antifungal agent against *Aspergillus flavus* Link ex. Fries, *Food Control*, 2009, **20**(2), 161–166.
- 37 H. Li, Z. Q. Zhang, C. He, G. Z. Qin and S. P. Tian, Comparative proteomics reveals the potential targets of *BcNoxR*, a putative regulatory subunit of NADPH oxidase of *Botrytis cinerea*, *Mol. Plant-Microbe Interact.*, 2016, **29**, 990–1003.
- 38 L. Ugolini, C. Martini, L. Lazzeri, L. D'Avino and M. Mari, Control of postharvest grey mould (*Botrytis cinerea* Per.: Fr.) on strawberries by glucosinolate-derived allyl-



- isothiocyanate treatments, *Postharvest Biol. Technol.*, 2014, **90**, 34–39.
- 39 M. Mari, O. Leoni, R. Bernardi, F. Neri and S. Palmieri, Control of brown rot on stonefruit by synthetic and glucosinolate-derived isothiocyanates, *Postharvest Biol. Technol.*, 2007, **47**, 61–67.
 - 40 F. Wang, F. Y. Wei, C. X. Song, Y. X. Huang and Y. X. Li, *Dodartia orientalis* L. essential oil exerts antibacterial activity by mechanisms of disrupting cell structure and resisting biofilm, *Ind. Crops Prod.*, 2017, **109**, 358–366.
 - 41 J. Tian, Y. Wang, H. Zeng, Z. Li, P. Zhang, A. Tessema and X. Peng, Efficacy and possible mechanisms of perillaldehyde in control of *Aspergillus niger* causing grape decay, *Int. J. Food Microbiol.*, 2015, **202**, 27–34.
 - 42 D. R. N. A. Cezar, M. Maraschin and R. M. Di Piero, Antifungal activity of salicylic acid against *Penicillium expansum* and its possible mechanisms of action, *Int. J. Food Microbiol.*, 2015, **215**, 64–70.
 - 43 H. Wu, X. Zhang, G. Zhang, S. Zeng and K. Kin, Antifungal vapour-phase activity of a combination of allyl isothiocyanate and ethyl isothiocyanate against *Botrytis cinerea* and *Penicillium expansum* infection on apples, *J. Phytopathol.*, 2011, **159**, 4050–4455.
 - 44 S. G. Harvey, H. N. Hannahan and C. E. Sams, Indian mustard and allyl-isothiocyanate inhibit *Sclerotium rolfsii*, *J. Am. Soc. Hortic. Sci.*, 2002, **127**(1), 27–31.
 - 45 D. J. Novo, N. G. Perlmutter, R. H. Hunt and H. M. Shapiro, Multiparameter flow cytometric analysis of antibiotic effects on membrane potential, membrane permeability, and bacterial counts of *staphylococcus aureus* and *micrococcus luteus*, *Antimicrob. Agents Chemother.*, 2000, **44**(4), 827–834.
 - 46 F. Yan, Q. Dang, C. Liu, J. Yan, T. Wang and B. Fan, 3,6-O-[N-(2-Aminoethyl)-acetamide-yl]-chitosan exerts antibacterial activity by a membrane damage mechanism, *Carbohydr. Polym.*, 2016, **149**, 102–111.
 - 47 L. da Cruz Cabral, V. F. Pinto and A. Patriarca, Application of plant derived compounds to control fungal spoilage and mycotoxin production in foods, *Int. J. Food Microbiol.*, 2013, **166**, 1–14.
 - 48 D. G. Yun and D. G. Lee, Silymarin exerts antifungal effects via membrane-targeted mode of action by increasing permeability and inducing oxidative stress, *Biochim. Biophys. Acta, Biomembr.*, 2017, **1859**, 467–474.
 - 49 V. K. Bajpai, A. Sharma and K. H. Baek, Antibacterial mode of action of *Cudrania tricuspidata* fruit essential oil, affecting membrane permeability and surface characteristics of food-borne pathogens, *Food Control*, 2013, **32**, 582–590.
 - 50 J. Liu, Y. Y. Zong, G. Z. Qin, B. Q. Li and S. P. Tian, Plasma membrane damage contributes to antifungal activity of silicon against *Penicillium digitatum*, *Curr. Microbiol.*, 2010, **61**, 274–279.
 - 51 S. Paul, R. C. Dubey, D. K. Maheswari and S. C. Kang, *Trachyspermum ammi* (L.) fruit essential oil influencing on membrane permeability and surface characteristics in inhibiting food-borne pathogens, *Food Control*, 2011, **22**, 725–731.
 - 52 M. Turgis, J. Han, S. Caillet and M. Lacroix, Antimicrobial activity of mustard essential oil against *Escherichia coli* O157: H7 and *Salmonella typhi*, *Food Control*, 2009, **20**, 1073–1079.
 - 53 C. M. Lin, J. F. Preston and C. I. Wei, Antibacterial mechanism of allyl isothiocyanate, *J. Food Prot.*, 2000, **63**, 727–734.
 - 54 A. C. Chan, D. Ager and I. P. Thompson, Resolving the mechanism of bacterial inhibition by plant secondary metabolites employing a combination of whole-cell biosensors, *J. Microbiol. Methods*, 2013, **93**, 209–217.
 - 55 I. Clemente, M. Aznar and C. Nerin, Raman imaging spectroscopy as a tool to investigate the cell damage on *Aspergillus ochraceus* caused by an antimicrobial packaging containing Benzyl isothiocyanate, *Anal. Chem.*, 2016, **88**, 4772–4779.
 - 56 T. C. Goosen, D. E. Mills and P. F. Hollenberg, Effects of benzyl isothiocyanate on rat and human cytochromes P450: Identification of metabolites formed by P4502B1, *J. Pharmacol. Exp. Ther.*, 2001, **296**, 198–206.
 - 57 A. Gallo, K. S. Bruno, M. Solfrizzo, G. Perrone, G. Mule, A. Visconti and S. E. Baker, New insight into the Ochratoxin A biosynthetic pathway through deletion of a nonribosomal peptide synthetase gene in *aspergillus carbonarius*, *Appl. Environ. Microbiol.*, 2012, **78**, 8208–8218.
 - 58 M. Wang, N. Jiang, H. Xian, D. Wei and X. Feng, A single-step solid phase extraction for the simultaneous determination of 8 mycotoxins in fruits by ultra-high performance liquid chromatography tandem mass spectrometry, *J. Chromatogr. A*, 2015, **1429**, 22–29.
 - 59 B. Andersen, E. Kröger and R. G. Roberts, Chemical and morphological segregation of *Alternaria arborescens*, *A. infectoria* and *A. tenuissima* species-groups, *Mycol. Res.*, 2002, **106**, 170–182.
 - 60 T. T. T. Nguyen, J. Kim, S. J. Jeon, C. W. Lee, N. Magan and H. B. Lee, Mycotoxin production of, *Alternaria*, strains isolated from Korean barley grains determined by LC-MS/MS, *Int. J. Food Microbiol.*, 2018, **268**, 44–52.
 - 61 A. A. Hildebrand, B. N. Kohn, E. Pfeiffer, D. Wefers and M. Bunzel, Conjugation of the mycotoxins alternariol and alternariol monomethyl ether in tobacco suspension cells, *J. Agric. Food Chem.*, 2015, **63**, 4728–4736.

