



 Cite this: *RSC Adv.*, 2020, 10, 6017

A unimolecular DNA fluorescent probe for determination of copper ions based on click chemistry†

 Yingfeng Qin,^a Ming Li,^a Yingying Yang,^a Zhiying Gao,^a Huaisheng Zhang^b
 and Jingjin Zhao *^{ab}

A homogenous fluorescence method was constructed for Cu²⁺ detection by employing DNA-templated click chemistry and exonuclease reaction. In this strategy, a dumbbell shaped DNA probe, which contained an alkyne group and an azide group at its ends, was designed as the template for the click chemistry reaction, and also the signal probe. In the absence of Cu²⁺, the DNA probe was digested into small oligonucleotide fragments by exonuclease, resulting in a low fluorescence background. However, this DNA probe can be sealed at its two ends by Cu²⁺-induced click chemistry ligation in the presence of Cu²⁺. This closed structure of DNA would remain stable after addition of exonuclease, and could then be stained by SYBR Green I. A strong fluorescence signal was observed, which was related to the concentration of Cu²⁺. This assay showed high selectivity and reached the detection limit of 39 nM. Moreover, the proposed strategy exhibited satisfactory detection results in real complex sample analysis, and has promising application in environmental monitoring and food safety.

Received 4th December 2019

Accepted 30th January 2020

DOI: 10.1039/c9ra10174h

rsc.li/rsc-advances

1. Introduction

Copper (Cu²⁺) is an essential trace element which plays a critical role in human health and various physiological fundamental life processes.¹ In addition, as an important transition metal, copper is involved in many biological processes as a structural component of certain proteins.² However, excessive amounts of copper lead to high toxicity in the human body. Some serious diseases have been found to be associated with the toxicity of copper, such as Alzheimer's disease, Wilson's disease, Menkes disease, and hereditary aceruloplasminemia.^{3–5} The accumulation of copper in the human body is mainly attributed to food and drinking water, as well as some copper come from the ecosystem that may be caused by environmental pollution. Therefore, determination of Cu²⁺ in the environment and humans for ecosystem balance and human health is of great significance.

To date, several approaches including atomic absorption spectroscopy (AAS),^{6,7} inductively coupled plasma atomic emission spectroscopy (ICP-AES),⁸ inductively coupled plasma mass spectrometry (ICP-MS),⁹ colorimetric methods,¹⁰ surface-

enhanced Raman scattering spectroscopy,¹¹ electrochemical assay,¹² fluorescence methods,^{13,14} have been used for Cu²⁺ detection in environmental samples and human bodies. Among these, fluorescence methods have gained an increasing interest due to its high sensitivity, low cost, ease of handling, and fast analysis speed.

Cu⁺-catalyzed azide-alkyne cycloaddition (CuAAC) is considered as a representative reaction of "Click Chemistry".¹⁵ It has gained an increasing interest in many fields due to its important merits such as high efficiency and selectivity, mild reaction conditions, and almost no byproducts.¹⁶ In addition, DNA-templated click chemistry reaction can be achieved by CuAAC as the linking tool to trigger azide and alkyne-modified DNA chemical ligation. Due to this merit, DNA-templated click chemistry reaction provides new signal conversion strategies^{17–20} for the detection of alkaline phosphatase activity, single nucleotide polymorphism, and protein, especially Cu²⁺. Nie's group developed a fluorescent method for Cu²⁺ detection based on DNA-templated click chemistry reaction and magnetic microparticle.²¹ Xiang's group reported a new glucometer-based sensing strategy using DNA-templated click chemistry reaction and multi-invertase loaded magnetic bead labels.²² Recently, our group established a novel fluorescence polarization assay based on DNA-templated click chemistry and silica nanoparticles-assisted fluorescence polarization enhancement.²³ Nevertheless, some methods for Cu²⁺ analysis normally required tedious separation process, complex sequence design or coupling of nanoparticles. Thus, developing new DNA-

^aKey Laboratory of Ecology of Rare and Endangered Species and Environmental Protection (Guangxi Normal University), Ministry of Education, Guilin 541004, PR China. E-mail: jzhao12@163.com; Fax: +86-773-583-2294; Tel: +86-773-5846141

^bDepartment of Chemistry and Biochemistry, Jackson State University, 1400 Lynch St., Jackson, MS, 39217, USA

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra10174h



templated click chemistry-based simple strategy for quantitative detection of Cu^{2+} is still needed.

Herein, we proposed a simple fluorescence assay for selectivity and sensitive detection of Cu^{2+} by DNA-templated click chemistry reaction and exonuclease digestion. A dumbbell shaped DNA (DS-DNA) is used as the DNA template and the signal probe, which bears an alkyne group and an azide group at its ends, respectively. In the presence of Cu^{2+} , the DS-DNA can be closed to form a sealed structure through DNA-templated click chemistry reaction. This stable structure is able to resist to the digestion of exonuclease I (Exo I) and exonuclease III (Exo III). The fluorescence signal can be assessed using SYBR Green I (SG I) as the label-free fluorescent dye. The change of fluorescence signal is related to the quantitation of Cu^{2+} . This method is not only simple, stable and specific, but also exhibits a good assay performance in complex samples.

2. Experimental

2.1 Reagents

The oligonucleotide (DS-DNA: 5'-CH \equiv CH-ATA TAT ATA TAT CGA CCC CCC TCG ATA TAT ATA TAT ATA TAT CGA CCC CCC TCG ATA TAT ATA TAT-N₃-3') was synthesized by Shanghai Sangon Biotech Co. Ltd (Shanghai, China) (<https://www.sangon.com>). Exonuclease III (Exo III), exonuclease I (Exo I) and 10 \times NEBuffer (10 mM Tris-HCl, 50 mM NaCl, 10 mM MgCl₂, 1 mM dithiothreitol, pH = 7.9) were gained from New England Biolabs (Beijing, China) (<https://www.neb-china.com>). Tris(hydroxypropyltriazolymethyl)amine (THPTA) was procured from Sigma-Aldrich (www.sigmaaldrich.com). SYBR Green I (SG I) (10 000 \times) and sodium ascorbate (NaVC) were purchased from Shanghai Sangon Biotech Co. Ltd (Shanghai, China) (<https://www.sangon.com>). GoldView II (5000 \times) was gained from solarbio (<http://www.solarbio.com>). All other reagents were of analytical grade. Ultrapure water (18.2 M Ω cm) was employed throughout the work.

2.2 Assay of Cu^{2+} analysis

The DS-DNA was denatured in 10 mM Tris-HCl buffer (50 mM NaCl, 10 mM MgCl₂) at 95 $^{\circ}\text{C}$ for 5 min and slowly cooled to room temperature to form dumbbell shaped structure. Then, 2 μL of 2 μM DS-DNA, different concentration of Cu^{2+} , 1 μL of 10 mM THPTA, 1 μL of 10 mM NaVC and 10 μL 2 \times click chemistry reaction buffer (0.2 M Tris-HCl, 1.0 M NaCl, pH = 7.4) were incubated at the room temperature for 30 min in dark. After that, Exo III (15 U), Exo I (2 U) and 10 mM Tris-HCl buffer (50 mM NaCl, 10 mM MgCl₂) were mixed and incubated at 37 $^{\circ}\text{C}$ for 60 min. Then 3.0 μL of SG I (50 \times) and ultrapure water were added with the final volume of 100 μL . The solutions were carried out at 37 $^{\circ}\text{C}$ for 30 min. Finally, the fluorescence spectra were measured by using an agilent G9800A fluorescence spectrometer (Agilent Technologies, USA) with the excitation wavelength at 480 nm. Fluorescence intensity at 525 nm was used to evaluate the assay performance.

2.3 Agarose gel electrophoresis

Three samples were prepared as follows: (1) DS-DNA (1 μM) was used as sample one; (2) the mixture of DS-DNA (1 μM), Exo III (90 U) and Exo I (12 U) were incubated at 37 $^{\circ}\text{C}$ for 60 min was used as sample two; (3) DS-DNA (1 μM) and Cu^{2+} (80 μM) were incubated at the room temperature for 30 min in dark, and then added Exo III (90 U) and Exo I (12 U) to the mixture for incubating 60 min at 37 $^{\circ}\text{C}$ was used as sample three. The samples were stained by GoldView II (5 \times) for incubating 20 min at 37 $^{\circ}\text{C}$. After adding loading buffer, the gel electrophoresis assay was carried out on 3% agarose gels and run in 0.5 \times TBE buffer (pH 8.3) at a 75 V constant voltage for 80 min. Then the gel was scanned and recorded by using the Tanon 1600 Gel imaging system.

2.4 Analysis of the river sample

The river water samples were obtained from Xiangsi River near Guangxi Normal University (Guangxi Province, China). The samples were filtered through a 0.22 μm membrane filter. Then, Cu^{2+} spiked river water samples were prepared by adding different concentration of Cu^{2+} into the filtered river water samples. The followed procedure was the same as the experiment mentioned above for assay of Cu^{2+} .

3. Results and discussion


3.1 Principle of fluorescence assay

A fluorescent assay was developed for Cu^{2+} detection based on click chemistry and exonuclease, which used only one DNA probe. The principle of this proposed method was illustrated in Scheme 1. As can be seen, a dumbbell shaped DNA probe (DS-DNA) was designed as the DNA-template of click chemistry reaction, which was labeled with an alkyne group (-CH \equiv CH) at the 5' end and an azide group (-N₃) at the 3' end, respectively. In the absence of Cu^{2+} , the click chemistry reaction was hard to carry out because of lacking of the catalyst. So, the ends of DS-DNA were still exposed, and then was recognized by Exo III and Exo I. Exo III was able to catalyze double-stranded region of DS-DNA from 3' \rightarrow 5' and Exo I could hydrolyze the single-stranded region of DS-DNA. Under the reaction of exonuclease, the DS-DNA was completely digested into small oligonucleotide fragments. Since SG I could not bind with the digestive products, a low fluorescence background signal was observed. On the contrary, the added Cu^{2+} was reduced to Cu^{+} by sodium ascorbate in the solution. Cu^{+} would catalyze the cycloaddition of click chemistry between 5'-CH \equiv CH and 3'-N₃ of DS-DNA. Thus, the DS-DNA probe was sealed to form a closed structure, which resisted to the degradation by Exo III and Exo I. As a result, SG I could easily bind with double-stranded region of DS-DNA, obtaining a high fluorescence signal.

3.2 Feasibility study of the method

The feasibility of the method was verified by investigating the fluorescence emission spectra with and without Cu^{2+} . As shown in Fig. 1A, a strong fluorescence signal was obtained when there was only one DS-DNA in the solution (curve a). However, DS-





Scheme 1 Schematic illustration of Cu^{2+} assay based on DNA-templated click chemistry reaction and exonuclease digestion.

DNA was hydrolyzed into small oligonucleotide fragments after the addition of Exo III and Exo I, causing a low fluorescence intensity (curve b). As expected, when Cu^{2+} was added to the solution before exonuclease reaction, the fluorescence intensity decreased slightly (curve c). Because DS-DNA was sealed by Cu^{2+} -catalyzed click ligation, which can stand up to exonuclease digestion. Curve b and curve c demonstrated that the fluorescence intensity change was attributed to Cu^{2+} . Thus, the proposed method can be used for the detection of Cu^{2+} .

To further verify the viability of the proposed method, gel electrophoresis was carried out. The results are shown in Fig. 1B. DS-DNA exhibited a bright band (lane 1). Upon adding Exo III and Exo I, the band of DS-DNA disappeared (lane 2), repeatedly implying that exonuclease digestion happened. After adding Cu^{2+} into the systems, one bright band was obtained (lane 3), which exhibited the same migration shift compared




Fig. 1 (A) Fluorescence spectra of different samples. (a) DS-DNA + SG I, (b) DS-DNA + Exo I + Exo III + SG I, (c) DS-DNA + Cu^{2+} + Exo I + Exo III + SG I. The concentration of Cu^{2+} was $10 \mu\text{M}$. (B) Agarose gel electrophoresis images of different samples. Lane 1: DS-DNA only; lane 2: DS-DNA + Exo I + Exo III; lane 3: DS-DNA + Cu^{2+} + Exo I + Exo III.

with lane 1. The gel electrophoresis experiments confirmed that Cu^{2+} -catalyzed click ligation could prevent exonuclease digestion.

3.3 Study of experimental conditions

In order to achieve the best assay performance, some important conditions such as the digestion capacity of exonuclease, the concentrations of Exo III and Exo I, incubation time of exonuclease, reaction time of click chemistry and the amount of SG I were investigated. First, the digestive capacity of exonuclease plays an important role in the system. In the absence of Cu^{2+} , the digestion capacity of exonuclease on DS-DNA was evaluated. As shown in Fig. 2, DS-DNA could be completely hydrolyzed only in the presence of both Exo I and Exo III.

The concentrations and incubation time of Exo III and Exo I have substantial impact on the exonuclease digestion, which directly affects the change of fluorescence intensity ΔF ($\Delta F = F - F_0$, where F and F_0 are the fluorescence intensity in the presence and absence of Cu^{2+} , respectively). As shown in Fig. S1,† with increasing concentrations or incubation time of Exo III and Exo I, ΔF increases significantly and reach a constant at 15 U Exo III (Fig. S1A†) and 2 U Exo I (Fig. S1B†) for 60 min (Fig. S1C†), and these results were chosen and used in the following experiments.

As one of the main parameters, the reaction time of click chemistry is closely related to chemical ligation efficiency. The amount of fluorescence dye SG I also can affect the fluorescence intensity of system. The effects of these two parameters were shown in Fig. S2.† The change of fluorescence intensity (ΔF) increases with the reaction time of click chemistry from 0 min to 30 min and reaches a plateau after 30 min (Fig. S2A†), indicating that the click chemistry reaction almost finished with 30 min, and 30 min was chosen as the optimal click chemistry reaction time. The amount of SG I of $50\times$ used was also investigated in Fig. S2B,† and ΔF increases markedly when SG I volume from $0 \mu\text{L}$ to $3 \mu\text{L}$. Consequently, the optimal amount of SG I used was selected to be $3 \mu\text{L}$ in this system.




Fig. 2 Study of the digestion capacity of exonuclease. The concentration of Exo III and Exo I are 15 U and 2 U, respectively. Error bars represent standard deviations of three repetitive measurements.






Fig. 3 (A) Fluorescence spectra of solutions after adding various concentrations of Cu^{2+} . Concentrations of Cu^{2+} from a to n were: 0, 0.1, 0.2, 0.4, 0.6, 1.0, 2.0, 5.0, 7.5, 10.0, 15.0, 50.0, 100.0 and 150.0 μM , respectively. (B) The relationship of the fluorescence intensity with Cu^{2+} concentration. Inset: linear correlation of the method. Error bars represent standard deviations of three repetitive measurements.

3.4 Assay performance

The assay performance of the proposed method was investigated under the optimal conditions by using a series of different concentrations of Cu^{2+} . Fig. 3A displays the fluorescence




Fig. 4 Selectivity test of the proposed method for Cu^{2+} . The concentration of Cu^{2+} was 10 μM and other metal ions was 100 μM . Each experiment was performed three times.

Table 1 Recovery experiments of Cu^{2+} in river samples

Samples	Add (μM)	Found (μM)	Recovery (%)	RSD (% , $n = 3$)
1	0.50	0.52	104.0	2.2
2	2.00	1.94	97.0	1.6
3	5.00	4.86	97.2	4.3
4	8.00	8.34	104.3	2.8

emission spectra obtained, and Fig. 4B shows the fluorescence intensity at 525 nm arises as increasing concentration of Cu^{2+} . A good linear correlation was achieved between the fluorescence intensity and Cu^{2+} concentration in the range from 0.1 μM to 10 μM (Fig. 3B, inset). The corresponding equation is $F = 44.169C + 100.864$ (F is the fluorescence intensity; C is the Cu^{2+} concentration, μM) with a correlation coefficient of 0.9980. The detection limit was estimated to be 39 nM ($S/N = 3$). The sensitivity of the proposed assay was comparable to or better than that of previously reported click chemistry reaction and magnetic microparticle-based fluorescence method,²¹ carbon dots-based fluorescence method,²⁴ DNzyme signal amplification-based lateral flow assay,²⁵ gold nanoparticles-based colorimetric assay,²⁶ and gold nanoclusters-based colorimetric assay.²⁷ Additionally, the proposed method is more convenience compared with these methods.

3.5 Analysis of selectivity and river samples

To evaluate the selectivity of the proposed method, other metal ions (Ag^+ , Co^{2+} , Mn^{2+} , Cr^{3+} , Ni^{2+} , Pb^{2+} , Al^{3+} , Zn^{2+} , Fe^{3+} , Hg^{2+}) were investigated as controlled experiments under the same experimental conditions. As shown in Fig. 4, only Cu^{2+} resulted in a remarkable increase in the fluorescence intensity, while other metal ions could not cause obvious fluorescence intensity change, even they are 10 times concentration than copper. These results indicated this assay exhibits an excellent selectivity for discriminating Cu^{2+} from other metal ions.

To further demonstrate the applicability of the developed method in complex matrixes, recovery experiments were carried out in river water samples. The obtained results are shown in Table 1, the recovery rates were in the range of 97.0–104.3% and the relative standard deviations (RSDs) were lower than 5%. In addition, the recoveries of standard addition were 101.4%, 98.5%, 99.6% and 102.3%, respectively for Cu^{2+} concentrations of 0.50, 2.00, 5.00 and 8.00 μM using ICP-MS method. The results obtained by the proposed method are comparable to ICP-MS, indicating the promising potential of this method for Cu^{2+} in real samples.

4. Conclusions

The proposed fluorescence assay for Cu^{2+} detection is based on click chemistry and exonuclease reaction. It designed only one dumbbell shaped DNA probe in the whole system, which served as both a template for click ligation and a signal probe for fluorescence dye. Compared with most of other reported



methods, this assay is convenient without troublesome synthesis of materials, separation procedures and fluorescence dye labeling. Moreover, the proposed strategy shows high specificity by click chemistry reaction to recognize Cu²⁺. In addition, the developed assay exhibits satisfactory detection results in real complex sample analysis. The above merits allow this method to hold the potential applications in environmental and biological fields.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the Project for Guangxi College Young and Middle-aged Teachers Basic Capability Improvement (No. 2019KY0082), Natural Science Foundation of Guangxi Province (No. 2017GXNSFFA198014) and BAGUI Scholar Program.

Notes and references

- 1 D. Denoyer, S. Masaldan, S. La Fontaine and M. A. Cater, *Metallomics*, 2015, **7**, 1459–1476.
- 2 B. E. Kim, T. Nevitt and D. J. Thiele, *Nat. Chem. Biol.*, 2008, **4**, 176–185.
- 3 P. G. Georgopoulos, A. Roy, M. J. Yonone-Lioy, R. E. Opiekun and P. J. Lioy, *J. Toxicol. Environ. Health, Part B*, 2001, **4**, 341–394.
- 4 D. R. Brown and H. Kozlowski, *Dalton Trans.*, 2004, **13**, 1907–1917.
- 5 G. J. Brewer, *Biofactors*, 2012, **38**, 107–113.
- 6 T. W. Lin and S. D. Huang, *Anal. Chem.*, 2001, **73**, 4319–4325.
- 7 D. Citak and M. Tuzen, *Food Chem. Toxicol.*, 2010, **48**, 1399–1404.
- 8 M. D. Ioannidou, G. A. Zachariadis, A. N. Anthemidis and J. A. Stratis, *Talanta*, 2005, **65**, 92–97.
- 9 N. G. Beck, R. P. Franks and K. W. Bruland, *Anal. Chim. Acta*, 2002, **455**, 11–22.
- 10 Y. Guo, Z. Wang, W. Qu, H. Shao and X. Jiang, *Biosens. Bioelectron.*, 2011, **26**, 4064–4069.
- 11 F. Li, J. Wang, Y. Lai, C. Wu, S. Sun, Y. He and H. Ma, *Biosens. Bioelectron.*, 2013, **39**, 82–87.
- 12 B. Kumar Jena and C. Retna Raj, *Anal. Chem.*, 2008, **80**, 4836–4844.
- 13 C. Lei, Z. Wang, Z. Nie, H. Deng, H. Hu, Y. Huang and S. Yao, *Anal. Chem.*, 2015, **87**, 1974–1980.
- 14 P. Yang, Y. Zhao, Y. Lu, Q. Xu, X. Xu, L. Dong and S. Yu, *ACS Nano*, 2011, **5**, 2147–2154.
- 15 V. V. Rostovtsev, L. G. Green, V. V. Fokin and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2002, **41**, 2596–2599.
- 16 Q. Wang, T. R. Chan, R. Hilgraf, V. V. Fokin, K. B. Sharpless and M. G. Finn, *J. Am. Chem. Soc.*, 2003, **125**, 3192–3193.
- 17 D. Yang, Z. Guo, Y. Tang and P. Miao, *ACS Appl. Nano Mater.*, 2018, **1**, 168–174.
- 18 Q. Y. Zhou, F. Yuan, X. H. Zhang, Y. L. Zhou and X. X. Zhang, *Chem. Sci.*, 2018, **9**, 3335–3340.
- 19 D. X. Wen, Q. R. Liu, Y. Cui, J. M. Kong, H. X. Yang and Q. Y. Liu, *Sens. Actuators, B*, 2018, **276**, 279–287.
- 20 L. Zhou, Q. Shen, P. Zhao, B. Xiang, Z. Nie, Y. Huang and S. Yao, *Methods*, 2013, **64**, 299–304.
- 21 Q. Shen, S. Tang, W. Li, Z. Nie, Z. Liu, Y. Huang and S. Yao, *Chem. Commun.*, 2012, **48**, 281–283.
- 22 J. Su, J. Xu, Y. Chen, R. Yuan and Y. Chai, *Biosens. Bioelectron.*, 2013, **45**, 219–222.
- 23 S. Liao, J. Zhao, Y. Qin and S. Zhao, *RSC Adv.*, 2017, **7**, 55668–55672.
- 24 J. Zong, X. Yang, A. Trinchì, S. Hardin, I. Cole, Y. Zhu and G. Wei, *Biosens. Bioelectron.*, 2014, **51**, 330–335.
- 25 Y. Wang, L. Wang, C. Zhang and F. Liu, *Microchim. Acta*, 2019, **186**, 82.
- 26 Z. Weng, H. Wang, J. Vongsivut, R. Li, A. M. Glushenkov, J. He, Y. Chen, C. J. Barrow and W. Yang, *Anal. Chim. Acta*, 2013, **803**, 128–134.
- 27 Y. Liu, D. Ding, Y. Zhen and R. Guo, *Biosens. Bioelectron.*, 2017, **92**, 140–146.

