


 Cite this: *RSC Adv.*, 2020, **10**, 8518

Unimolecular decomposition rates of a methyl-substituted Criegee intermediate *syn*-CH₃CHOO†

 Yu-Lin Li,^{ab} Mei-Tsan Kuo ^a and Jim Jr-Min Lin ^{*ab}

Criegee intermediates play important roles in atmospheric chemistry. Methyl Criegee intermediate, CH₃CHOO, has two conformers, *syn*- and *anti*-conformers. *Syn*-CH₃CHOO would undergo fast unimolecular decomposition to form OH radical via 1,4 H-atom transfer. In this work, unimolecular decomposition of *syn*-CH₃CHOO was probed in real time with UV absorption spectroscopy at 278–318 K and 100–700 torr. We used water vapor as the scavenger of *anti*-CH₃CHOO to distinguish the absorption signals of the two conformers. After removing the contributions from reactions with radical byproducts, reaction with water vapor and wall loss, we obtained the unimolecular reaction rate coefficient of *syn*-CH₃CHOO (at 300 torr), which increases from (67 ± 15) s⁻¹ at 278 K, (146 ± 31) s⁻¹ at 298 K, to (288 ± 81) s⁻¹ at 318 K with an Arrhenius activation energy of ca. 6.4 kcal mol⁻¹ and a weak pressure dependence for 100–700 torr. Compared to previous studies, this work provides temperature dependent unimolecular rates of *syn*-CH₃CHOO at higher pressures, which are more relevant to atmospheric conditions.

 Received 9th January 2020
 Accepted 18th February 2020

DOI: 10.1039/d0ra01406k

rsc.li/rsc-advances

Introduction

Criegee intermediates are very reactive carbonyl oxides formed in the reactions of ozone with alkenes (ozonolysis) and play important roles in atmospheric chemistry, including oxidation of water, SO₂, NO_x, etc. In particular, the decomposition of Criegee intermediates can be a significant source of OH radicals in the troposphere, especially during nighttime and winter.¹

After the breakthrough of the efficient preparation of Criegee intermediates in 2012,² the physical and chemical properties of simple Criegee intermediates have been widely investigated.^{3,4} Nowadays, it has been established that the reactivity of Criegee intermediates is highly structure-dependent.^{4–6} For example, CH₂OO and *anti*-CH₃CHOO react with water vapor very quickly but *syn*-CH₃CHOO and (CH₃)₂COO react with water vapor much slower. On the other hand, *syn*-CH₃CHOO and (CH₃)₂COO may undergo intramolecular H-atom transfer from the methyl group to the terminal oxygen atom, resulting in OH production.^{7,8}

However, there have been large discrepancies in the unimolecular (also bimolecular and termolecular) reaction rates of Criegee intermediates. If we focus on the unimolecular rate coefficients of *syn*-CH₃CHOO reported after 2012, the reported experimental value^{9–12} at 298 K ranges from 3 to 300 s⁻¹ whilst the reported theoretical value^{13–15} is from 24 to 330 s⁻¹.¹ Unimolecular rates may control the steady-state concentrations

and thus the impact of such a Criegee intermediate in the atmosphere. It is important to have reliable experimental data at atmospherically relevant pressure and temperature.

Lester and coworkers excited *syn*-CH₃CHOO, *syn*-C₂H₅CHOO and (CH₃)₂COO to a high vibrational energy via infrared laser and monitored the OH production by laser-induced fluorescence. The OH formation rates have been measured at specific vibrational energies. Master equation modeling based on these microcanonical rates and high-level quantum chemistry calculations has been used to estimate the thermal decay rate coefficients at 298 K to be 166/122 s⁻¹ for *syn*-CH₃CHOO,^{14,16} 279, and 276 s⁻¹ for *syn*-C₂H₅CHOO¹⁷ and (CH₃)₂COO,¹⁴ respectively.¹⁸ The fast rates indicate thermal decomposition is the main sink for these Criegee intermediates in the atmosphere.

Zhou *et al.* have monitored the OH products of *syn*-CH₃CHOO in a flow cell. The thermal decomposition rate at 298 K (25 to 100 torr) is reported to be 182 ± 66 s⁻¹.¹⁹ Their results also support that the thermal decomposition is the main atmospheric process for *syn*-CH₃CHOO.

In this work, we used water vapor to scavenge *anti*-CH₃CHOO²⁰ and directly monitored the kinetics of *syn*-CH₃CHOO via its strong UV absorption in real time.²¹ The unimolecular rates were determined at atmospherically relevant temperatures and pressures. The results are compared with previous experimental and theoretical works.

Experimental method

Following previous studies, we used CH₃CHI₂ (Aldrich, ≥98%) as the precursor and generated CH₃CHI radicals by applying photolysis laser (Excimer laser, KrF 248 nm, Coherent

^aInstitute of Atomic and Molecular Sciences, Academia Sinica, Taipei 10617, Taiwan. E-mail: jimlin@gate.sinica.edu.tw

^bDepartment of Chemistry, National Taiwan University, Taipei 10617, Taiwan

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0ra01406k



COMPexPro 205F). Under high $[O_2]$ (10 torr), the iodo-radicals would react with O_2 rapidly to form CH_3CHOO .¹² The photolysis laser beam was coupled into the reactor by a long-pass filter (275 nm, Eksma Optics, customized). After passing through the reactor, the photolysis beam was reflected out by another long-pass filter, and the laser pulse energy was monitored with an energy meter (Gentec QE25SP-H-MB-D0). Continuous UV probe beam originated from a broadband Xe lamp (Energetiq, EQ-99) went through the reactor six times. The probe beam overlapped collinearly with the photolysis beam, resulting in an effective absorption path length of *ca.* 426 cm.


The probe wavelength was selected by a band-pass filter (340 nm, 10 nm bandwidth, OD4, Edmund #65129). The time-dependent absorption change was detected with a balanced photodiode detector (Thorlab PDB450A) and recorded in real time with a high-definition oscilloscope (LeCroy HDO4034, 4096 vertical resolution). All time traces were averaged for *ca.* 120 laser shots to increase S/N ratio. We found a small absorption change even without adding the precursor, which was caused by the photolysis laser and the long-pass filters. This background could be subtracted by recording background traces before and after each experimental set. All reported time traces have undergone background subtraction.

CH_3CHI_2 vapor was carried by a stream of N_2 . All the gas flows were controlled by mass-flow controllers (Brooks, 5850E or 5800E). A small portion of CH_3CHI_2 vapor was introduced into a residual gas analyzer (SRS RGA200) to monitor the impurity. Most of CH_3CHI_2 vapor would mix with O_2 and then flowed into an absorption cell (90 cm in length) to measure the UV absorption spectrum with a deuterium lamp (Hamamatsu L10904) and a mini spectrometer (Ocean Optics USB-2000). The water vapor concentration was controlled by varying the ratio of dry and moisturized N_2 gases and monitored with a humidity sensor (Rotronic HC2-S, 0.8% RH accuracy at room temperature). The water vapor was pre-cooled or pre-heated by flowing through a copper tube (1/4" OD, *ca.* 75 cm long) which was immersed in a circulating water bath. Two streams of the gas mixtures (water vapor/ N_2 and precursor/ O_2) were mixed together right before entering the reactor. The laser repetition rate was set to *ca.* 1 Hz to fully refresh the gas between consecutive laser shots.

The temperature of the water-jacketed reactor was controlled (± 0.5 K) by using a circulating water bath (Yih Der BL-730), and detected by three resistance temperature detectors (RTD, Newport Omega F2020-1000-A). The pressures at the water reservoir, absorption cell and reactor were measured by diaphragm gauges (1000 torr, INFICON, $\pm 0.2\%$ of reading).

Results

We synthesized CH_3CHOO (a mixture of *anti* and *syn* conformers) *via* a well-known method.^{7,12}



Similar to the case of preparing CH_2OO , adduct formation (R3) is also expected and would reduce the yield of CH_3CHOO at high pressures.^{22,23}

Removing the contribution of *anti*- CH_3CHOO

Fig. 1 shows typical time traces of CH_3CHOO absorption signals recorded at 340 nm under various $[H_2O]$. We can see that the fast component decays faster at higher $[H_2O]$, while the effect of $[H_2O]$ on the slow component is rather weak. Because *anti*- CH_3CHOO reacts with water vapor much faster than *syn*- CH_3CHOO does,²⁰ we may separate their signals by fitting the time traces with two exponential functions (eqn (1)).

$$\begin{aligned} \Delta Abs(t) = & L\sigma_{anti}[anti-CH_3CHOO] + L\sigma_{syn}[syn-CH_3CHOO] \\ & + L\sigma_{prec}\Delta[CH_3CHI_2] = A_{anti} \exp(-t/\tau_{anti}) \\ & + A_{syn} \exp(-t/\tau_{syn}) + C_0 \end{aligned} \quad (1)$$

where L is the absorption path length; σ_{anti} , σ_{syn} and σ_{prec} are the absorption cross sections of *anti*- CH_3CHOO , *syn*- CH_3CHOO , and the CH_3CHI_2 precursor, respectively; A_{anti} and A_{syn} are the signal amplitudes for *anti*- CH_3CHOO and *syn*- CH_3CHOO , respectively; τ_{anti} and τ_{syn} are the corresponding lifetimes; C_0 accounts for the depletion of the precursor which has a weak absorption at 340 nm (1.07×10^{-18} cm²).²⁴ In further analysis, we have $k_{obs} = \tau_{syn}^{-1}$ and $[syn-CH_3CHOO]_0 = A_{syn}/(L\sigma_{syn})$.

The fast kinetics of *anti*- CH_3CHOO reaction with water vapor has been investigated in depth by a few groups.^{12,20,25} Therefore we do not repeat it here. Water vapor efficiently scavenges *anti*- CH_3CHOO ; Fig. 1 shows that at $[H_2O] > 1.9 \times 10^{17}$ cm⁻³, only *syn*- CH_3CHOO signal remains for $t > 1$ ms. The reaction of *syn*- CH_3CHOO with H_2O has been reported to be much slower than




Fig. 1 Typical time traces of CH_3CHOO absorption probed at 340 ± 5 nm under various $[H_2O]$ (308 K and 305 torr, Exp. W2-3: $[CH_3CHI_2] = 1.4 \times 10^{14}$ cm⁻³, laser fluence = 1.7 mJ cm⁻², see Table S1† for details). The photolysis laser pulse sets the time zero. The black lines are two-exponential fit to the data (see text). The fast decay signal is assigned to *anti*- CH_3CHOO , the slow one to *syn*- CH_3CHOO .



that of *anti*-CH₃CHOO.^{12,13,20,25} More kinetic investigation on the reaction of *syn*-CH₃CHOO with H₂O will be discussed later in this paper.

Effect of bimolecular reactions of *syn*-CH₃CHOO with radical byproducts

Fig. 2 shows the time traces of CH₃CHOO absorption signals recorded under various precursor concentrations [CH₃CHI₂]₀ and a fixed photolysis laser fluence. In this experiment, we added water ([H₂O] = 1.9 × 10¹⁷ cm⁻³) to scavenge *anti*-CH₃CHOO and the absorption signal of the remaining *syn*-CH₃CHOO predominates in the time traces for *t* > 1 ms. We can see that the lifetime of *syn*-CH₃CHOO becomes shorter at higher [CH₃CHI₂]₀. This is mainly due to the reactions of *syn*-CH₃CHOO with radical byproducts like iodine atoms, similar to previous results for CH₂OO,²³ *syn*-CH₃CHOO^{19,26} and (CH₃)₂COO.²⁷ The inset of Fig. 2 shows the observed decay rate coefficient *k*_{obs} of *syn*-CH₃CHOO plotted against its fitted peak height [*syn*-CH₃CHOO]₀, which was estimated by using the Beer–Lambert law with a reported cross section $\sigma_{syn} = 1.19 \times 10^{-17}$ cm² at 340 nm (ref. 21) (which is consistent with the results by Sheps *et al.*²⁵) and effective length *L* = 426 cm.

Under our experimental conditions, the concentrations of the radical byproducts (I, CH₃CHOO, CH₃CHIOO, *etc.*) are proportional to [*syn*-CH₃CHOO]₀. The related chemical reactions include (R1)–(3). Thus, we can use [X], [X] ∝ [*syn*-




Fig. 2 Typical time traces of CH₃CHOO probed at 340 ± 5 nm at [H₂O] = 1.9 × 10¹⁷ cm⁻³; different colors denote various precursor concentrations [CH₃CHI₂]₀ (4.8 × 10¹³ cm⁻³ (cyan), 7.7 × 10¹³ cm⁻³ (blue), 1.3 × 10¹⁴ cm⁻³ (green), and 2.5 × 10¹⁴ cm⁻³ (red)). These traces were acquired at 299 K and 301 torr with a laser fluence of 1.7 mJ cm⁻² (Exp. 2-1, see Table S2†). For each trace, the black line is the two-exponential fit (eqn (1)) to the signal of CH₃CHOO. The fast decay signal is assigned to *anti*-CH₃CHOO; the slow one is assigned to *syn*-CH₃CHOO. The slightly negative baseline is due to the depletion of the precursor CH₃CHI₂, which is a constant in the detection time window and does not affect the kinetics. We scaled the peak absorbance of *syn*-CH₃CHOO to the same value for easier visualization. Inset: the plot of *k*_{obs} versus [*syn*-CH₃CHOO]₀. The red line is the linear fit.

CH₃CHOO]₀, to represent the effective concentration of the total radical species. We used the following kinetic model to analyze the experimental data.

$$k_{\text{obs}} = k_{\text{uni}} + k_{\text{X}}[\text{X}] + k_{\text{w}}[\text{H}_2\text{O}] + k_{\text{wall}} \quad (2)$$

where *k*_{uni} is the unimolecular decomposition rate coefficient; *k*_X is the effective rate coefficient for the reaction of *syn*-CH₃-CHOO with total radicals X; *k*_w and *k*_{wall} represent the rate coefficients for the reaction of *syn*-CH₃CHOO with H₂O and the wall loss, respectively. The linear behavior of *k*_{obs} against [*syn*-CH₃CHOO]₀ is consistent with the above model.

Similar experiments have also been performed at temperatures from 278 K to 318 K. The results are shown in Fig. 3. Again, the positive slope indicates the observed decay rate coefficient becomes larger at higher radical concentrations. Remarkably, the intercept rate is substantially larger at higher temperature, suggesting that the unimolecular rate is faster at higher temperature.

Effect of water vapor

We observed that the decay rate of *syn*-CH₃CHOO becomes slightly faster at higher [H₂O], especially at higher temperatures (see Fig. 4). However, the slopes are not very significant, especially considering the large scatters of the data. The obtained slopes (*k*_w) at 298 K and 300 torr are on the order of 1 × 10⁻¹⁶ cm³ s⁻¹ (Exp. W3-1 and W4-1, see Table S1†), much larger than the theoretical results ranging from 1.9 × 10⁻¹⁹ to 2.4 × 10⁻¹⁸ cm³ s⁻¹.^{13,28} The source of this discrepancy is not fully clear. It is likely that minor impurity in the water vapor had some effect (different sets of gas tubing were used in Experiments W3 and W4†). It is difficult to determine the small rate coefficient of *syn*-CH₃CHOO reaction with water; as evidenced by previous investigations by Sheps *et al.*²⁵ and Taatjes *et al.*¹² which only reported upper limits of 2 × 10⁻¹⁶




Fig. 3 Plot of *k*_{obs} as a function of [*syn*-CH₃CHOO]₀ at different temperatures. The experiments were conducted at 300 torr and detailed experimental conditions are shown in Table S2†.






Fig. 4 Plot of k_{obs} as a function of $[\text{H}_2\text{O}]$ at three temperatures (color-coded) under 300 torr. Different symbols represent different sets of experiments at various $[\text{CH}_3\text{CH}_2]_0$, while each set of experiment was conducted at a constant laser power (see Table S1†). The lines are linear fit to the data. Larger values of the intercept rates are due to higher temperature or higher $[\text{CH}_3\text{CH}_2]_0$.

and $4 \times 10^{-15} \text{ cm}^3 \text{ s}^{-1}$, respectively, for the rate coefficient. A similar difficulty has also been reported for the case of $(\text{CH}_3)_2\text{COO}$ reaction with water ($k_w < 1.5 \times 10^{-16} \text{ cm}^3 \text{ s}^{-1}$).²⁹ Fig. S3† shows the Arrhenius plot of k_w . Because we cannot measure k_w at temperatures lower than 298 K, the values of k_w from the Arrhenius fit are used. Considering the difficulties in the experiments, here we report a conservative upper limit of $1 \times 10^{-16} \text{ cm}^3 \text{ s}^{-1}$ for k_w at 298 K.

Effect of wall loss

To estimate the wall loss in our system, we measured the decay rates of CH_2OO under similar experimental conditions. The thermal decomposition of CH_2OO has been calculated to be quite slow.⁶ In addition, Berndt *et al.* have performed time-resolved experiments of C_2H_4 ozonolysis, which yielded a rate coefficient of $(0.19 \pm 0.07) \text{ s}^{-1}$ for the unimolecular reaction of CH_2OO at 298 K and 1 bar.³⁰ Furthermore, Stone *et al.* directly measured the unimolecular reaction of CH_2OO at higher temperatures (450–650 K) and obtained a rate coefficient of $(1.1_{-1.1}^{+1.5} \times 10^{-3}) \text{ s}^{-1}$ at 298 K and 760 torr using a master-equation method.³¹ Thus, the observed CH_2OO decay (k_{obs} , see Fig. 5) in our system should mainly come from the wall loss (intercept) and bimolecular reactions (slope). As shown in Fig. S6,† the results at different temperatures and pressures are quite similar, with an average value of about 9 s^{-1} for k_{wall} . At our pressure range, diffusion is too slow to explain the observed wall loss rate; gas turbulence, which is expected to be more severe at higher pressures, should be the main cause of the observed wall loss (see ESI† for details). As gas turbulence would not be affected by the highly diluted precursor molecules and Criegee intermediates, we assume the wall loss rate of *syn*- CH_3CHOO is the same as that of CH_2OO .




Fig. 5 k_{obs} as a function of $[\text{CH}_2\text{OO}]_0$ at various temperatures (A) and pressures (B) (color-coded). Different symbols represent different experimental sets.

k_{uni} and its temperature dependence

Here we give one example of determining the unimolecular rate at 298 K.

$$\begin{aligned} k_{\text{uni}} &= k_{\text{obs}} - k_{\text{X}}[\text{X}] - k_{\text{w}}[\text{H}_2\text{O}] - k_{\text{wall}} \\ &= k_{\text{intercept}} - k_{\text{w}}[\text{H}_2\text{O}] - k_{\text{wall}} \end{aligned} \quad (2')$$

In Fig. 2 inset, we can see that the $k_{\text{X}}[\text{X}]$ term has a significant contribution to k_{obs} . This contribution can be removed by extrapolating k_{obs} to zero radical concentration (*i.e.*, using $k_{\text{intercept}}$). For Exp. 2-1 (299 K, 301 torr, see Table S2†), $k_{\text{intercept}} = 173 \text{ s}^{-1}$, $k_{\text{w}}[\text{H}_2\text{O}] = 17 \text{ s}^{-1}$, $k_{\text{wall}} = 9 \text{ s}^{-1}$ and we have $k_{\text{uni}} = 173 - 17 - 9 = 147 \text{ s}^{-1}$. See ESI† for details.

Fig. 6 shows the Arrhenius plot of k_{uni} , measured at 300 torr, as well as a few theoretical results. Our data are consistent with the theoretical results, especially those by Fang *et al.*¹⁴ and Yin *et al.*¹⁵ However, the theoretical slopes (activation energy) are all a bit higher than the experimental value. The reason for this difference is unclear. As mentioned above, the water reaction






Fig. 6 Arrhenius plot of the thermal decomposition rate coefficient of *syn*-CH₃CHOO at 300 torr (Table S2†). The dashed lines are theoretical data reported by various groups at the high pressure limit.^{13–15,32} The experimental data shown as triangles and circles represent the data with and without subtracting the water contribution, respectively.

rates $k_w[\text{H}_2\text{O}]$ are difficult to quantify. If we assume $k_w = 0$, the resulted activation energy would be a bit closer to the theoretical values (see Table 1).

Pressure dependence of k_{uni}

Fig. 7 shows the pressure dependence of k_{uni} at 298 K. Because we used water to scavenge *anti*-CH₃CHOO, we cannot measure k_{uni} at pressure lower than 100 torr. Our data from 100 to 700 torr suggest slightly higher unimolecular rate at higher pressure, albeit some scattering of the data points. The low-pressure results of Zhou *et al.* are also plotted in Fig. 7. As will be discussed below, Zhou *et al.*¹⁹ may have underestimated their diffusion loss and thus overestimated their k_{uni} values, especially for data at lower pressures. Overall, the observed pressure dependences are roughly consistent with the theoretical trend.

Table 1 Thermal decomposition rate coefficient and Arrhenius activation energy of *syn*-CH₃CHOO

	$k_{\text{uni}}/\text{s}^{-1}$	$E_a/\text{kcal mol}^{-1}$
This work	146 ± 31^a	6.4 ± 0.2^b
This work (assuming $k_w = 0$)	173 ± 31^a	7.1 ± 0.3^b
Fang <i>et al.</i> ¹⁴	166 ^c	8.5 ± 0.1^b
Long <i>et al.</i> ¹³	328 ^c	7.7 ± 0.2^b
Nguyen <i>et al.</i> ³²	124 ^c	8.2 ± 0.4^b
Yin <i>et al.</i> ¹⁵	182 ^c	8.6 ± 0.2^b

^a At 298 K and 300 torr. The error bar is one standard deviation (see ESI).

^b The activation energy (including theoretical ones) is obtained from the slope of the linear fitting of $\log(k)$ against $1/T$ within 278–318 K; the error bar is one standard deviation of the fitting. ^c Theoretical result at 298 K at the high pressure limit.




Fig. 7 Plot of k_{uni} of *syn*-CH₃CHOO against pressure from 100 to 700 torr at 298 K (Table S3†). The experimental data reported by Zhou *et al.* are shown as blue triangles,¹⁹ and the theoretical data reported by Nguyen *et al.* is shown as a red line.³² In order to compare with the experimental data, we also scaled the data of Nguyen *et al.* by a factor of 1.3 (grey line).

Discussion

Effect of second-order reactions

In previous works on the thermal decomposition reactions of (CH₃)₂COO by Smith *et al.*²⁷ and of *syn*-CH₃CHOO by Zhou *et al.*,¹⁹ the authors have considered the contributions of second-order reactions (including the reaction with iodine atoms and self-reaction of Criegee intermediates²⁶). Following the notation of Smith *et al.*, the corresponding kinetic equation can be expressed as below.²⁷

$$-\frac{d[\text{CI}]}{dt} = k_1[\text{CI}] + k_2[\text{CI}]^2 \quad (3)$$

where [CI] stands for the concentration of the Criegee intermediate of interest. However, Smith *et al.* have also demonstrated that after extrapolation to zero concentration, a simplified model of pseudo-first-order reaction would give essentially the same results for the intercept rates.²⁷ That is, we may use eqn (4), if [CI]₀ is low.

$$-\frac{d[\text{CI}]}{dt} \cong \left(k_1 + \frac{1}{2}k_2[\text{CI}]_0 \right) [\text{CI}] = k_{\text{obs}}[\text{CI}] \quad (4)$$

As mentioned by Smith *et al.*, the difference between the two models only shows up at high concentration data.²⁷ As shown in Fig. S5,† the simulated time traces of the solutions of eqn (3) and (4) are quite similar for [CI]₀ = $3 \times 10^{11} \text{ cm}^{-3}$, $k_1 = 150 \text{ s}^{-1}$, $k_2 = 1.6 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$. Therefore, we chose to use eqn (4) for more efficient data analysis.

Tunneling and potential energy surface

Lester, Klippenstein and coworkers have shown that H-atom tunneling is the main mechanism of the unimolecular decay



Table 2 Comparison of the effective first-order rate coefficients for *syn*-CH₃CHO at 298 K

Process	Assumed coreactant concentration/cm ⁻³	Bimolecular rate coefficient/cm ³ s ⁻¹	Effective first-order rate coefficient/s ⁻¹
Thermal decomposition	—	—	146 ± 31 ^a
<i>syn</i> -CH ₃ CHO + H ₂ O	3.8 × 10 ¹⁷ (50% RH)	<1 × 10 ⁻¹⁶ , ^b	<38
<i>syn</i> -CH ₃ CHO + SO ₂	9 × 10 ¹¹ (35 ppbv)	(2.4–2.9) × 10 ⁻¹¹ , ^c	22–26

^a This work. Note that the value of k_{uni} is quite sensitive to temperature. ^b Upper limit, this work. ^c The values are reported by Taatjes *et al.*¹² and Sheps *et al.*²⁵

of *syn*-CH₃CHO.^{14,16,33} Green *et al.* of the same group further investigated the tunneling dynamics by using *syn*-CD₃CHO sample.³⁴ Their theoretical calculation describes the energy-specific unimolecular rates quite well. Based on these successes, they can further estimate the unimolecular rates under thermalized conditions. The result is plotted in Fig. 6 (Fang *et al.*), which agrees with our experimental data. The subtle difference in the slopes of the Arrhenius plots may be due to minor differences in (i) potential energy surface, (ii) tunneling model, and (iii) experimental uncertainties.

Compare to the recent results of Zhou *et al.*

Recently Zhou *et al.* measured the unimolecular rates of *syn*-CH₃CHO by probing the OH products with laser-induced fluorescence.¹⁹ Because the barrier of OH production of *anti*-CH₃CHO is much higher than that for *syn*-CH₃CHO,^{13,15} Zhou *et al.* do not need to scavenge *anti*-CH₃CHO.

Their measurements were under 10 to 100 torr,¹⁹ limited by the lower sensitivity of the OH laser-induced fluorescence at higher pressure. Our work provides data at higher pressures up to 700 torr, much more relevant to the atmospheric conditions. In addition, perhaps due to sensitivity issue, Zhou *et al.* used higher concentrations of the Criegee intermediate in their experiments (judged by the faster observed decay rates), resulting in larger error bars for the unimolecular rates. Furthermore Zhou *et al.* did not investigate the temperature dependence.¹⁹

Diffusion would have a more significant effect at low pressures as the diffusion coefficient is inversely proportional to the pressure. However, Zhou *et al.* only argued that their experiments should be in the laminar flow condition (*i.e.*, no turbulence), without estimating the effect of diffusion.¹⁹ A simple but useful estimation of the diffusion effect can be found in textbooks³⁵ as

$$z_{\text{rms}} = (2Dt)^{0.5}$$

where z_{rms} is the root-mean-square value of the one-dimensional displacement of the molecule; D is the diffusion coefficient and t is the elapsed time. For CO₂-O₂ diffusion, $D = 0.14 \text{ cm}^2 \text{ s}^{-1}$ at 760 torr,³⁵ $10.6 \text{ cm}^2 \text{ s}^{-1}$ at 10 torr ($12.7 \text{ cm}^2 \text{ s}^{-1}$ for a more realistic estimation for their Criegee experiment at 10 torr,¹⁹ see ESI†). If $t = 10 \text{ ms}$, we would have $z_{\text{rms}} = 0.46 \text{ cm}$, which is already larger than the pump beam radius (0.3 cm) and the probe beam radius (0.1 cm) in the experiments of Zhou *et al.*¹⁹ This rough estimation indicates that at a low pressure of

ca. 10 torr, the diffusion length of the molecule is quite significant even for a short period of 10 ms.

Fig. S8† shows more-realistic simulation results for cases similar to the experimental conditions of Zhou *et al.* (with orthogonal pump and probe laser beams).¹⁹ Again, the diffusion loss has a significant effect of about 50 s^{-1} (10 torr) to $\approx < 10 \text{ s}^{-1}$ (100 torr) (see ESI†), indicating that the low-pressure results by Zhou *et al.* should be corrected by considering the diffusion loss.

Fate of *syn*-CH₃CHO in troposphere

Previous investigations⁶ have pointed out that unimolecular decay is one of the main processes of Criegee intermediates in the atmosphere. For *syn* conformers, the unimolecular decay would be the predominant process.⁴ For *syn*-CH₃CHO, we list the effective pseudo-first-order rates of some important processes in Table 2. It is clear that the unimolecular decomposition is the predominate decay pathway of *syn*-CH₃CHO in the troposphere. This fast unimolecular decay would control the steady-state concentration of *syn*-CH₃CHO. As a result, the probability for *syn*-CH₃CHO to oxidize SO₂ would be low due to its low concentration. For other Criegee intermediates of similar structures (alkyl substitution at the *syn* position), the situation is expected to be similar. As mentioned above, the thermal decomposition rate is quite sensitive to temperature. This work gives temperature dependent value of k_{uni} , which allows modelers to better estimate the unimolecular decay rates of *syn*-CH₃CHO at atmospherically relevant temperatures.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work is supported by Academia Sinica and Ministry of Science and Technology, Taiwan (MOST 106-2113-M-001-026-MY3). We thank Ms Yen-Hsiu Lin for help in the measurements of the wall loss rates.

References

- M. A. H. Khan, C. J. Percival, R. L. Caravan, C. A. Taatjes and D. E. Shallcross, *Environ. Sci.: Processes Impacts*, 2018, **20**, 437–453.



- 2 O. Welz, J. D. Savee, D. L. Osborn, S. S. Vasu, C. J. Percival, D. E. Shallcross and C. A. Taatjes, *Science*, 2012, **335**, 204–207.
- 3 C. A. Taatjes, *Annu. Rev. Phys. Chem.*, 2017, **68**, 183–207.
- 4 J. J. M. Lin and W. Chao, *Chem. Soc. Rev.*, 2017, **46**, 7483–7497.
- 5 L. Vereecken, D. R. Glowacki and M. J. Pilling, *Chem. Rev.*, 2015, **115**, 4063–4114.
- 6 L. Vereecken, A. Novelli and D. Taraborrelli, *Phys. Chem. Chem. Phys.*, 2017, **19**, 31599–31612.
- 7 F. Liu, J. M. Beames, A. S. Petit, A. B. McCoy and M. I. Lester, *Science*, 2014, **345**, 1596–1598.
- 8 F. Liu, J. M. Beames and M. I. Lester, *J. Chem. Phys.*, 2014, **141**, 234312.
- 9 T. Berndt, T. Jokinen, R. L. Mauldin, T. Petäjä, H. Herrmann, H. Junninen, P. Paasonen, D. R. Worsnop and M. Sipilä, *J. Phys. Chem. Lett.*, 2012, **3**, 2892–2896.
- 10 A. Novelli, L. Vereecken, J. Lelieveld and H. Harder, *Phys. Chem. Chem. Phys.*, 2014, **16**, 19941–19951.
- 11 M. J. Newland, A. R. Rickard, L. Vereecken, A. Muñoz, M. Ródenas and W. J. Bloss, *Atmos. Chem. Phys.*, 2015, **15**, 9521–9536.
- 12 C. A. Taatjes, O. Welz, A. J. Eskola, J. D. Savee, A. M. Scheer, D. E. Shallcross, B. Rotavera, E. P. F. Lee, J. M. Dyke, D. K. W. Mok, D. L. Osborn and C. J. Percival, *Science*, 2013, **340**, 177–180.
- 13 B. Long, J. L. Bao and D. G. Truhlar, *J. Am. Chem. Soc.*, 2016, **138**, 14409–14422.
- 14 Y. Fang, F. Liu, V. P. Barber, S. J. Klippenstein, A. B. McCoy and M. I. Lester, *J. Chem. Phys.*, 2016, **144**, 061102.
- 15 C. Yin and K. Takahashi, *Phys. Chem. Chem. Phys.*, 2017, **19**, 12075–12084.
- 16 Y. Fang, F. Liu, V. P. Barber, S. J. Klippenstein, A. B. McCoy and M. I. Lester, *J. Chem. Phys.*, 2016, **145**, 234308.
- 17 Y. Fang, F. Liu, S. J. Klippenstein and M. I. Lester, *J. Chem. Phys.*, 2016, **145**, 044312.
- 18 T. A. Stephenson and M. I. Lester, *Int. Rev. Phys. Chem.*, 2020, **39**, 1–33.
- 19 X. H. Zhou, Y. Q. Liu, W. R. Dong and X. M. Yang, *J. Phys. Chem. Lett.*, 2019, **10**, 4817–4821.
- 20 L. C. Lin, W. Chao, C. H. Chang, K. Takahashi and J. J. M. Lin, *Phys. Chem. Chem. Phys.*, 2016, **18**, 28189–28197.
- 21 M. C. Smith, W. L. Ting, C. H. Chang, K. Takahashi, K. A. Boering and J. J. M. Lin, *J. Chem. Phys.*, 2014, **141**, 074302.
- 22 Y.-H. Huang, L.-W. Chen and Y.-P. Lee, *J. Phys. Chem. Lett.*, 2015, **6**, 4610–4615.
- 23 W. L. Ting, C. H. Chang, Y. F. Lee, H. Matsui, Y. P. Lee and J. J. M. Lin, *J. Chem. Phys.*, 2014, **141**, 104308.
- 24 S. P. Sander, J. R. Barker, D. M. Golden, M. J. Kurylo, P. H. Wine, J. P. D. Abbatt, J. B. Burkholder, C. E. Kolb, G. K. Moortgat, R. E. Huie and V. L. Orkin, *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies Evaluation Number 17*, JPL Publication, 2011, pp. 10–6.
- 25 L. Sheps, A. M. Scully and K. Au, *Phys. Chem. Chem. Phys.*, 2014, **16**, 26701–26706.
- 26 P.-L. Luo, Y. Endo and Y.-P. Lee, *J. Phys. Chem. Lett.*, 2018, **9**, 4391–4395.
- 27 M. C. Smith, W. Chao, K. Takahashi, K. A. Boering and J. J. M. Lin, *J. Phys. Chem. A*, 2016, **120**, 4789–4798.
- 28 J. Anglada, J. Gonzalez and M. Torrent-Sucarrat, *Phys. Chem. Chem. Phys.*, 2011, **13**, 13034–13045.
- 29 H.-L. Huang, W. Chao and J. J.-M. Lin, *Proc. Natl. Acad. Sci. U. S. A.*, 2015, **112**, 10857–10862.
- 30 T. Berndt, R. Kaethner, J. Voigtländer, F. Stratmann, M. Pfeifle, P. Reichle, M. Sipilä, M. Kulmala and M. Olzmann, *Phys. Chem. Chem. Phys.*, 2015, **17**, 19862–19873.
- 31 D. Stone, K. Au, S. Sime, D. J. Medeiros, M. Blitz, P. W. Seakins, Z. Decker and L. Sheps, *Phys. Chem. Chem. Phys.*, 2018, **20**, 24940–24954.
- 32 T. L. Nguyen, L. McCaslin, M. C. McCarthy and J. F. Stanton, *J. Chem. Phys.*, 2016, **145**, 131102.
- 33 V. P. Barber, S. Pandit, V. J. Esposito, A. B. McCoy and M. I. Lester, *J. Phys. Chem. A*, 2019, **123**, 2559–2569.
- 34 A. M. Green, V. P. Barber, Y. Fang, S. J. Klippenstein and M. I. Lester, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, 12372–12377.
- 35 P. L. Houston, *Chemical Kinetics and Reaction Dynamics*, Dover Publications, Mineola, New York, 2006.

