RSC Advances



View Article Online

View Journal | View Issue

PAPER

Check for updates

Cite this: RSC Adv., 2020, 10, 21895

Construct indeno[1,2-b]oxepine or *cis*cyclopropylacrylate by sulfur ylides†

Qinfang Chen, Yihao Pan, Dongxin Zhao, Weiran Yang and Jing Zheng 吵 *

For the first time, the [4 + 3] or [2 + 1] annulation of crotonate-derived sulfur ylides with arylidenemalononitrile or arylidene-1*H*-indene-1,3(2*H*)-dione is reported using Na₂CO₃ as the base. This protocol is advantageous as it does not require prior preparation of arylidenemalononitrile or arylidene-1*H*-indene-1,3(2*H*)-dione substrates, due to the independent participation of the base in the two reactions. This mild, operationally multicomponent process can be employed for the transformation of a wide variety of commercially available aldehydes into the corresponding indeno[1,2-*b*]oxepine or cyclopropyl acrylate core in moderate to excellent yields under mild conditions.

1. Introduction

Received 18th May 2020

Accepted 29th May 2020

DOI: 10.1039/d0ra03919e

rsc.li/rsc-advances

Oxygen-containing heterocyclic compounds, especially unsaturated 7-membered oxacycle (oxepine) frameworks, exhibit a wide range of biological properties, such as ion-channel blocking, antiplasmodial, antiviral, antipsychotic, and antifungal activities.¹ Various natural and marine natural products containing the oxepine motif play a vital role in biological processes.² The cyclopropyl group is also a vital structural unit in several synthetic and naturally occurring compounds, exhibiting a wide spectrum of biological properties ranging from enzyme inhibition to herbicidal, antibiotic, antitumor, and antiviral activities.³ Fig. 1 shows some representative examples of these compounds.

Recently, crotonate-derived sulfur ylides have attracted attention in organic synthesis as sources of one-carbon or threecarbon synthons in the presence of an inorganic base.⁴ The ability of crotonate-derived sulfur ylides to undergo [m + n]cycloaddition reactions with various substrates is exploited to construct several cyclic compounds.5 Tang and co-workers have synthesized vinylaziridine by the diastereoselective annulation of crotonate-derived sulfur ylides with cyclic ketamine (Scheme 1a).6 Huang and co-workers have reported access to cyclic 2alkenyl aziridines by sequential annulation using crotonatederived sulfur ylides as the C3 synthon with α , β -unsaturated cyclic ketimines (Scheme 1b).5f,7 Huang has synthesized sevenmembered nitrogen-heterocycles with moderate-to-excellent yields by the development of a novel [4 + 3] annulation of azadienes with crotonate sulfonium salts (Scheme 1c).8 On the other hand, multicomponent reactions (MCRs), defined as

processes that combine at least three reactants in the same pot to generate a product containing most of the atoms of the starting material, have been extensively exploited to prepare small molecules with step economy, energy conservation, and



Fig. 1 Biologically active substances containing oxepine motif and cyclopropyl.



Scheme 1 Annulation with crotonate-derived sulfur-ylides.

School of Resources Environmental and Chemical Engineering, Nanchang University, 999 XuFu Road, Nangchang, 330031, China. E-mail: Zhengjing@ncu.edu.cn

[†] Electronic supplementary information (ESI) available: Copies of ¹H and ¹³C NMR spectra of isolated compounds. CCDC 1956080 and 1956081. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d0ra03919e

emission reduction.⁹ As highly reactive reagents, sulfur ylides can react with various electrophiles such as aldehydes, imines, and electron-deficient alkenes.¹⁰ Therefore, the design of MCRs involving sulfur ylides in the presence of aldehydes, 1,3-indanedione, and malononitrile might lead to selectivity problems.

Despite these challenges, it is interesting to develop sulfur ylide-mediated MCRs to construct core structures with medicinal interest. Based on the aforementioned motivation and our continued interest in organocatalysis, herein, the construction of indeno[1,2-b]oxepine by the three-component [4+3] or [1+2] annulation of crotonate-derived sulfur ylides, aldehydes, and 1,3-indanedione or malononitrile (Scheme 1d).

2. Results and discussion

Prior to investigation of the desired MCRs and sulfur ylides, studies were first initiated by investigating reactions between crotonate sulfonium salt **3a** and 2-benzylidene-1*H*-indene-1,3(2*H*)-dione, which was generated by the reaction of 1,3-indanedione (**1a**) and benzaldehyde (**2a**) (Table 1). The reaction was performed in DCM with NaHCO₃ as the base at 25 °C (Table 1, entry 1). Gratifyingly, the [4 + 3] annulation product was obtained in 76% yield. The structure of **4a** was confirmed by NMR and HRMS. Reaction conditions were optimized to improve the product yield. Optimization of base was first carried out. When the reaction was conducted in *t*-BuOK, the yield was improved to 85% (Table 1, entry 2). Moderate product yields were observed by using other bases, such as DABCO,

Table 1 Optimization of the MCRs reaction⁴

	$\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$ + $\begin{pmatrix} 0 \\ H \\ 0 \end{pmatrix}$ + $\begin{pmatrix} 0 \\ H \\ 2a \end{pmatrix}$	+ Br 3a	DEt Base Solvent	
Entry	Base	Solvent	Temp (°C)	Yield ^b (%)
1	NaHCO ₃	DCM	25	76
2	t-BuOK	DCM	25	85
3	DABCO	DCM	25	75
4	TMAF	DCM	25	76
5	Cs_2CO_3	DCM	25	56
6	NaOH	DCM	25	73
7	Et ₃ N	DCM	25	70
8	K_2CO_3	DCM	25	79
9	K_3PO_4	DCM	25	50
10	NaH	DCM	25	90
11	Na ₂ CO ₃	DCM	25	90
12	Na ₂ CO ₃	CH ₃ CN	25	79
13	Na_2CO_3	CH_3OH	25	48
14	Na_2CO_3	THF	25	55
15	Na_2CO_3	Toluene	25	67
16	Na ₂ CO ₃	DCE	25	70
17	Na_2CO_3	DCM	0	45
18	Na_2CO_3	DCM	60	88

^{*a*} Unless otherwise specified, the reactions were carried out with **1a** (0.2 mmol), **2a** (0.24 mmol) and **3a** (0.3 mmol) in the presence of base (0.4 mmol) in a solvent (2 mL) at 25 $^{\circ}$ C. ^{*b*} Isolated yield.

TMAF, Cs_2CO_3 , NaOH, Et_3N , K_2CO_3 and K_3PO_4 (Table 1, entries 3–9). To our surprise, the product yield was improved to 90% when the base was used instead of Na_2CO_3 . Although NaH was also found to afford yields similar to that obtained using Na_2CO_3 , the cost-effective base was used for [4 + 3] cyclization (Table 1, entries 10 and 11). After the optimization of the base, the scope of different solvents for the was investigated. Common solvents (CH₃CN, CH₃OH, THF, toluene, DCE) failed to further improve the product yields (Table 1, entries 12–16). In addition, the reaction temperature significantly affected the reaction. By decreasing the temperature from room temperature to 0 °C, the product yield decreased from 90% to 45% even for a longer reaction time of 6 h (Table 1, entry 17). Neither the decrease nor increase the temperature led to the improvement of the yield of **4a** (Table 1, entries 17 and 18).

Using optimized reaction conditions (Table 1, entry 11), the substrate scope of the substituted benzaldehydes was first investigated (Scheme 2). Different benzaldehydes with a variety of functional groups such as NO₂, CN, Cl, CH₃, N(CH₃)₂, and OCH₃ smoothly reacted with **1a** and **3a** to afford the corresponding indeno[1,2-*b*]oxepine (**4b**-**4o**) in moderate to excellent yields (42–90%), thus offering a broad range of opportunities for further derivatization. These results revealed that reactions of benzaldehydes bearing electron-donating and electron-



Scheme 2 Scope of the [4 + 3] reactions.^{*a,b a*} Unless otherwise specified, the reactions were carried out with **1** (0.2 mmol), **2a** (0.24 mmol) and **3** (0.3 mmol) in the presence of base (0.4 mmol) in a solvent (2 mL) at 25 °C. ^{*b*} Isolated yield. ^cThe reaction was carried out at 0 °C.

Paper

withdrawing groups at the para position do not clearly affect this transformation. In particular, aromatic aldehydes bearing a functional group at the para position exhibited higher reactivity than their ortho- or meta-substituted counterpart (4g vs. 4h and 4i). However, for polysubstituted aromatic aldehydes, yields for the desired products were less (4k, 4o) than those for the monosubstituted aromatic aldehydes. In addition, the structure of 40 was further confirmed by single-crystal X-ray analysis.¹¹ Moreover, reactions efficiently proceeded when sulfonium salt 3b was used, affording desired products in moderate to excellent yields (41-4n). As expected, heteroaryl aldehydes, such as 5methylfurfural, furfural, 2-thenaldehyde, and nicotinaldehyde, performed quite well in the MCRs (4p-4s). The use of indole-3carboxaldehyde afforded low yields of the desired product (4t). However, when indole-3-carboxaldehyde was protected using iodomethane, the product yield was increased to 70% (4u).

To demonstrate the generality of this transformation, malononitrile was investigated instead of 1,3-indanedione under the optimal conditions. As shown in Scheme 3, electron-withdrawing or electron-donating groups such as nitro, cyano and *N*,*N*-dimethyl, at the *para*-position \mathbb{R}^1 in the substrate 2 smoothly reacted with **5a** and **3a** to afford [2 + 1] annulation products **6a–6d** (69–89%) with an excellent *cis/trans* ratio. To our delight, the desired *cis*-cyclopropylacrylate **6a** was still exclusively obtained (*cis/trans* >99/1) in 80% yield using 1.0 mmol of substrate **5a**. While the methoxy group substituted at the *meta* position in **6e** afforded low product yield (75%), the structure of **6e** was further confirmed by single-crystal X-ray



Scheme 3 Scope of the [2 + 1] reactions.^{*a,b a*}Unless otherwise specified, the reactions were carried out with 5 (0.2 mmol), 2 (0.24 mmol) and 3 (0.3 mmol) in the presence of base (0.4 mmol) in a solvent (2 mL) at 25 °C. ^{*b*}Isolated yield.





analysis.¹² However, for polysubstituted aromatic aldehydes, the yield of the desired product **6f** was less than that of the monosubstituted aromatic aldehyde. As expected, heteroaryl aldehydes, such as 5-methylfurfural, furfural, 2-thenaldehyde, nicotinaldehyde, and 1-methyl-1*H*-indole-3-carbaldehyde performed quite well in the MCRs (**6g–6k**). In addition, aliphatic aldehydes proved to be suitable substrates to furnish the desired reaction in moderate yields (**6l**, 57%). Moreover, reactions proceeded efficiently proceeded when sulfonium salt **3b** was used, affording desired products in moderate to excellent yields (**6m–6o**, **6q**). Furthermore, cyanoacrylates also afforded the desired products (**6o–6q**). The relative configuration of **6o** was defined on the basis of the phase-sensitive NOESY spectrum. No correlation between hydrogen on cyclopropane and methyl by the NOESY spectrum was observed (see the ESI†).

Based on the present experimental data and the results described in the literature, ${}^{4f-h,5e,7,13}$ a plausible MCR reaction mechanism can be proposed as shown in Scheme 4. Crotonatederived ylide 2 was treated with base (Na₂CO₃) to afford allylic ylide **A**, which resonated to **B**. Next, the Michael addition of **B** to **C**, which was formed by **1a** and **2** in the presence of the base catalyst, afforded intermediate **D**. Intermediate **D** subsequently transformed into intermediate **G** *via* two proton-transfer processes, followed by an intramolecular SN₂ nucleophilic substitution to furnish [4 + 3] annulation product **4**. When we **5** was used instead of **1a**, the Michael addition of **A** to **I** in the presence of the base catalyst would lead to intermediate **J**. Then, the subsequent intramolecular nucleophilic addition and elimination of Me₂S further afforded the final product.

3. Conclusions

To the best of our knowledge, for the first time, a novel Na₂CO₃promoted MCR reaction using crotonate-derived sulfonium salts, aromatic aldehydes, and 1,3-indanedione or malononitrile was developed. Various indeno[1,2-*b*]oxepine and *cis*cyclopropylacrylate derivatives were obtained in good-toexcellent yields. The advantages of the current protocol included readily available starting materials, mild reaction conditions, good functional group tolerance, and broad substrate scope. Currently, further applications of this MCR reaction in organic synthesis, as well as investigations on the detailed mechanism, are underway in our laboratory.

4. Experimental section

4.1 General information

Infrared spectra were obtained on a FTIR spectrometer. ¹H NMR and ¹³C NMR spectra were recorded on Agilent DD2400 spectrometer. $CDCl_3$ was used as solvent. Chemical shifts were referenced relative to residual solvent. Melting points were measured with micro melting point apparatus. Infrared spectra were recorded on a FT-IR spectrometer. **1a**, **2** and **5** were prepared from Energy Chemical. **3** were prepared according to the literature procedure.¹⁴

4.2 Experimental procedures and characterization date

Ethyl-6-oxo-5-phenyl-5,6-dihydro-2H-indeno[1,2-b]oxepine-4carboxylate (4a). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-vlium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4-ethoxy-4oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, benzaldehyde 2a (25.4 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:10) as eluent to give 4a as a Yellow solid (62.3 mg, 90% yield), $R_f = 0.21$ (EtOAc/petroleum ether 1 : 10). Mp 115 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 7.36 (d, J = 6.8 Hz, 1H), 7.28–7.16 (m, 6H), 7.12–7.05 (m, 3H), 5.29 (s, 1H), 4.83 (dd, $J_1 = 12.4$ Hz, $J_2 = 7.6$ Hz, 1H), 4.54 $(dd, J_1 = 12.4 Hz, J_2 = 7.6 Hz, 1H), 4.28-4.16 (m, 2H), 1.25 (t, J =$ 7.2 Hz, 3H); $^{13}{\rm C}$ NMR (101 MHz, CDCl3), δ : 194.3, 173.6, 166.1, 144.5, 141.9, 139.7, 132.5, 132.1, 131.3, 130.0, 128.6, 127.1, 126.7, 121.3, 118.4, 109.6, 65.1, 61.8, 35.5, 14.3. IR(KBr) v: 2974, 2918, 1712, 1628, 1579, 1460, 1397, 1243, 1095, 1046. HRMS(EI) (m/z): calcd for C₂₂H₁₈O₄ (M + H)⁺: 347.1278; found: 347.1276.

Ethyl-5-(4-nitrophenyl)-6-oxo-5,6-dihydro-2*H*-indeno[1,2-*b*] oxepine-4-carboxylate (4b). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 4-nitrobenzaldehyde 2b (36.2 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:5) as eluent to give 4b as a yellow solid (32.8 mg, 42% yield), $R_{\rm f} = 0.12$ (EtOAc/petroleum ether 1 : 10). Mp 115 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 8.15 (t, 1H), 8.13 (t, 1H), 7.52–7.47 (m, 3H), 7.39 (t, J = 8.0 Hz, 1H), 7.33 (t, J = 8.0 Hz, 1H), 7.25–7.20 (m, 2H), 5.42 (s, 1H), 4.80 (dd, $J_1 =$

12.8 Hz, $J_2 = 7.2$ Hz, 1H), 4.74–4.69 (m, 1H), 4.39–4.26 (m, 2H), 1.35 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ : 194.2, 174.1, 165.8, 149.5, 146.8, 143.4, 139.4, 132.8, 132.8, 131.1, 130.5, 128.2, 123.9, 121.6, 118.9, 108.4, 65.2, 62.2, 35.9, 14.3. IR(KBr) ν : 2925, 2855, 1712, 1628, 1523, 1474, 1355, 1116, 990. HRMS(EI) (m/z): calcd for C₂₂H₁₇NO₆ (M + H)⁺: 392.1129; found: 392.1128.

Ethyl-5-(4-cyanophenyl)-6-oxo-5,6-dihydro-2*H*-indeno[1,2-*b*] oxepine-4-carboxylate (4c). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 4-formylbenzonitrile 2c (31.4 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:10) as eluent to give 4c as a yellow solid (45.3 mg, 61% yield), $R_{\rm f} = 0.15$ (EtOAc/petroleum ether 1 : 10). Mp 132 °C; ¹H NMR (CDCl₃, 400 MHz), δ: 7.57 (d, 2H), 7,46–7.44 (m, 3H), 7.37 (t, J = 7.2 Hz, 1H), 7.32 (d, J = 8.0 Hz, 1H), 7.21 (t, J = 8.0 Hz, 2H), 5.38 (s, 1H), 4.81-4.76 (m, 1H), 4.72-4.67 (m, 1H), 4.37–4.24 (m, 2H), 1.34 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 194.2, 174.1, 165.8, 147.4, 143.4, 139.4, 132.8, 132.7, 132.5, 131.0, 130.4, 128.1, 121.5, 118.9, 118.8, 110.7, 108.3, 65.2, 62.1, 35.9, 14.3. IR(KBr) v: 2981, 2918, 2231, 1705, 1621, 1515, 1452, 1396, 1242, 1179, 1102, 1046. HRMS(EI) (m/z): calcd for $C_{23}H_{17}NO_4 (M + H)^+$: 372.1230; found: 372.1231.

Ethyl-5-(4-chlorophenyl)-6-oxo-5,6-dihydro-2H-indeno[1,2-b] oxepine-4-carboxylate (4d). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 4chlorobenzaldehyde 2d (33.6 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:10) as eluent to give 4d as a yellow solid (63.1 mg, 83% yield), $R_{\rm f} = 0.21$ (EtOAc/petroleum ether 1 : 10). Mp 108 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 7.45 (d, J = 7.2 Hz, 1H), 7.35 (t, J = 8.4 Hz, 1H), 7.28 (t, J = 7.6 Hz, 1H), 7.25-7.22 (m, 4H), 7.19-7.15 (m, 2H), 5.31 (s, 1H), 4.89-4.84 (m, 1H), 4.69-4.64 $(m, 1H), 4.35-4.25 (m, 2H), 1.33 (t, J = 7.2 Hz, 3H); {}^{13}C NMR (101)$ MHz, CDCl₃), δ: 194.3, 173.8, 166.0, 144.1, 140.5, 139.6, 132.6, 132.6, 132.3, 131.2, 130.2, 128.7, 128.6, 121.4, 118.6, 109.1, 65.1, 62.0, 35.2, 14.3. IR(KBr) v: 2918, 1705, 1628, 1585, 1410, 1235, 1172, 1109, 1039, 955, 906, 765, 730. HRMS(EI) (m/z): calcd for $C_{22}H_{17}ClO_4 (M + H)^+$: 381.0888; found: 381.0890.

Ethyl-6-oxo-5-(p-tolyl)-5,6-dihydro-2H-indeno[1,2-b]oxepine-4-carboxylate (4e). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1Hinden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 4-methylbenzaldehyde 2e (28.8 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:10) as eluent to give 4e as a Yellow oil (59.8 mg, 83% yield), $R_{\rm f} = 0.18$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 7.45 (d, J = 6.8 Hz, 1H), 7.37–7.33 (m, 1H), 7.31– 7.27 (m, 1H), 7.22–7.13 (m, 4H), 7.08 (d, J = 8.0 Hz, 2H), 5.33 (s, 1H), 4.95 (dd, $J_1 = 12.4$ Hz, $J_2 = 7.2$ Hz, 1H), 4.63 (dd, $J_1 =$ 12.4 Hz, $J_2 = 8.0$ Hz, 1H), 4.36–4.24 (m, 2H), 2.29 (s, 3H), 1.34 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ : 194.4, 173.6, 166.2, 144.6, 139.8, 138.9, 136.3, 132.5, 132.0, 131.3, 130.0, 129.3, 127.1, 121.3, 118.4, 109.8, 65.2, 61.8, 35.3, 21.0, 14.3. IR(KBr) v: 2911, 2855, 1712, 1628, 1508, 1466, 1403, 1235, 1109, 1039. HRMS(EI) (m/z): calcd for C₂₃H₂₀O₄ $(M + H)^+$: 361.1434; found: 361.1436.

Ethyl-5-(4-(dimethylamino)phenyl)-6-oxo-5,6-dihydro-2H-

indeno[1,2-b]oxepine-4-carboxylate (4f). In a 5 mL vial, 1,3dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 4-(dimethylamino)benzaldehyde 2f (35.8 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:5) as eluent to give 4f as a yellow oil (68.5 mg, 88% yield), $R_{\rm f} = 0.15$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 7.44 (d, J = 7.2 Hz, 1H), 7.37–7.33 (m, 1H), 7.29 (d, J = 8.0 Hz, 1H), 7.21–7.09 (m, 4H), 6.65 (d, J = 8.8 Hz, 2H), 5.27 (s, 1H), 5.01 (dd, J₁ = 12.4 Hz, $J_2 = 7.3$ Hz, 1H), 4.63 (dd, $J_1 = 12.4$ Hz, $J_2 = 7.8$ Hz, 1H), 4.35-4.23 (m, 2H), 2.89 (s, 6H), 1.33 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 194.5, 173.4, 166.4, 149.5, 144.9, 140.0, 132.5, 131.8, 131.4, 129.9, 129.7, 127.9, 121.2, 118.3, 112.8, 110.3, 65.3, 61.8, 40.8, 34.9, 14.4. IR(KBr) v: 2911, 2841, 1705, 1628, 1515, 1473, 1403, 1340, 1228, 1109, 1046. HRMS(EI) (m/z): calcd for $C_{24}H_{23}NO_4 (M + H)^+$: 390.1700; found: 390.1699.

Ethyl-5-(4-methoxyphenyl)-6-oxo-5,6-dihydro-2*H*-indeno[1,2*b*]oxepine-4-carboxylate (4g). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1*H*-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (*E*)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 4methoxybenzaldehyde 2g (32.6 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1 : 10) as eluent to give **4g** as a Yellow oil (64.7 mg, 86% yield), $R_{\rm f} = 0.18$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 7.44 (d, J = 7.2 Hz, 1H), 7.35 (t, J = 7.6 Hz, 1H), 7.29 (t, J = 8.0 Hz, 1H), 7.24 (d, J = 9.2 Hz, 2H), 7.18–7.12 (m, 2H), 6.81 (d, J = 8.8 Hz, 2H), 5.30 (s, 1H), 4.95 (dd, $J_1 = 12.4$ Hz, $J_2 = 7.2$ Hz, 1H), 4.64 (dd, $J_1 = 12.4$ Hz, $J_2 = 8.0$ Hz, 1H), 4.36–4.24 (m, 2H), 3.76 (s, 3H), 1.34 (t, J = 7.1 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ : 194.4, 173.6, 166.2, 158.4, 144.7, 139.9, 134.0, 132.5, 132.0, 131.3, 130.1, 128.3, 121.3, 118.4, 114.0, 109.9, 65.2, 61.8, 55.4, 35.0, 14.3. IR(KBr) ν : 2925, 1712, 1628, 1515, 1466, 1403, 1249, 1179, 1102, 1039. HRMS(EI) (m/z): calcd for C₂₃H₂₀NO₅ (M + H)⁺: 377.1384; found: 377.1385.

Ethyl-5-(3-methoxyphenyl)-6-oxo-5,6-dihydro-2H-indeno[1,2b]oxepine-4-carboxylate (4h). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 3methoxybenzaldehyde 2h (32.6 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1: 10) as eluent to give 4h as a yellow oil (60.2 mg, 80% yield), $R_{\rm f} = 0.18$ (EtOAc/petroleum ether 1 : 10). ¹H NMR $(CDCl_3, 400 \text{ MHz}), \delta: 7.45 \text{ (d}, J = 6.8 \text{ Hz}, 1\text{H}), 7.35 \text{ (t}, J = 8.0 \text{ Hz},$ 1H), 7.29 (t, J = 8.0 Hz, 1H), 7.22–7.14 (m, 3H), 6.90 (d, J =6.8 Hz, 2H), 6.75–6.72 (m, 1H), 5.34 (s, 1H), 4.97 (dd, $J_1 =$ 12.4 Hz, $J_2 = 7.2$ Hz, 1H), 4.63 (dd, $J_1 = 12.4$ Hz, $J_2 = 8.0$ Hz, 1H), 4.37–4.24 (m, 2H), 3.76 (s, 3H), 1.34 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 194.3, 173.7, 166.1, 159.8, 144.4, 143.6, 139.8, 132.5, 132.2, 131.3, 130.1, 129.6, 121.3, 119.6, 118.5, 113.7, 111.5, 109.5, 65.1, 61.9, 55.3, 35.5, 14.4. IR(KBr) v: 2932, 2855, 1712, 1628, 1592, 1585, 1452, 1396, 1242, 1109, 1046. HRMS(EI) (m/z): calcd for C₂₃H₂₀O₅ $(M + H)^+$: 377.1384; found: 377.1387.

Ethyl-5-(3-chlorophenyl)-6-oxo-5,6-dihydro-2*H*-indeno[1,2-*b*] oxepine-4-carboxylate (4i). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 3chlorobenzaldehyde 2i (33.6 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1: 10) as eluent to give 4i as a yellow oil (60.8 mg, 80% yield), $R_{\rm f} = 0.22$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 7.46 (d, J = 7.2 Hz, 1H), 7.36 (t, J = 8.0 Hz, 1H), 7.30 (t, 2H), 7.22–7.16 (m, 5H), 5.33 (s, 1H), 4.87 (dd, J₁ =

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

12.4 Hz, $J_2 = 7.2$ Hz, 1H), 4.69–4.64 (m, 1H), 4.38–4.24 (m, 2H), 1.34 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ : 194.2, 173.9, 165.9, 144.0, 143.9, 139.6, 134.5, 132.6, 132.4, 131.2, 130.2, 129.9, 127.3, 127.0, 125.5, 121.4, 118.6, 108.9, 65.1, 62.0, 35.3, 14.3. IR(KBr) ν : 2925, 2855, 1712, 1628, 1585, 1466, 1403, 1235, 1179, 1102, 1032, 983. HRMS(EI) (m/z): calcd for C₂₂H₁₇ClO₄ (M + H)⁺: 381.0888; found: 381.0886.

Ethyl-5-(2-methoxyphenyl)-6-oxo-5,6-dihydro-2H-indeno[1,2b]oxepine-4-carboxylate (4j). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 2methoxybenzaldehyde 2j (32.6 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:10) as eluent to give 4j as a yellow solid (45.9 mg, 61% yield), $R_f = 0.19$ (EtOAc/petroleum ether 1 : 10). Mp 195 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 7.45 (d, J = 6.8 Hz, 1H), 7.36 (t, J = 8.0 Hz, 1H), 7.29 (t, J = 8.0 Hz, 1H), 7.26 (d, J = 7.2 Hz, 1H), 7.19 (t, J = 6.8 Hz, 2H), 6.88–6.81 (m, 2H), 6.70 (t, J = 8.0 Hz, 1H), 5.31 (s, 1H), 5.04 (dd, $J_1 = 12.4$ Hz, $J_2 = 8.0$ Hz, 1H), 4.53-4.48 (m, 1H), 4.36-4.27 (m, 2H), 3.73 (s, 3H), 1.36 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 193.7, 174.5, 166.9, 156.4, 146.5, 139.8, 132.4, 131.4, 130.3, 130.0, 128.7, 128.2, 128.0, 121.2, 120.2, 118.2, 110.8, 107.8, 64.5, 61.2, 55.0, 32.5, 14.4. IR(KBr) v: 2932, 2848, 1740, 1691, 1642, 1543, 1515, 1459, 1424, 1102, 1046, 997. HRMS(EI) (m/z): calcd for C₂₃H₂₀O₅ (M + H)⁺: 377.1384; found: 377.1386.

Ethyl-5-(2,3-dimethoxyphenyl)-6-oxo-5,6-dihydro-2H-indeno [1,2-b]oxepine-4-carboxylate (4k). In a 5 mL vial, 1,3-dioxo-2,3dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 2,3dimethoxybenzaldehyde 2k (39.8 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:5) as eluent to give 4k as a yellow solid (53.6 mg, 66% yield), $R_f = 0.15$ (EtOAc/petroleum ether 1 : 10). Mp 172 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 7.46 (d, J = 7.2 Hz, 1H), 7.37 (t, 1H), 7.30 (t, 1H), 7.19 (d, J = 7.2 Hz, 1H), 6.95 (t, 1H), 6.88 (d, J = 7.2 Hz, 1H), 6.82–6.76 (m, 2H), 5.36 (s, 1H), 4.51 (dd, $J_1 =$ $12.4 \text{ Hz}, J_2 = 8.0 \text{ Hz}, 1\text{H}, 4.51 \text{ (dd}, J_1 = 12.4 \text{ Hz}, J_2 = 7.6 \text{ Hz}, 1\text{H}),$ 4.30 (q, J = 7.2 Hz, 2H), 3.83 (s, 3H), 3.79 (s, 3H), 1.35 (t, J =7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 193.7, 174.3, 166.4, 153.2, 146.3, 146.2, 139.8, 136.2, 132.4, 131.5, 130.1, 128.0, 123.5, 121.3, 121.1, 118.3, 111.5, 108.3, 64.8, 61.5, 60.3, 55.9, 32.3, 14.4. IR(KBr) v: 2918, 2814, 1726, 1466, 1396, 1291, 1228,

1116, 1032, 983. HRMS(EI) (m/z): calcd for C₂₄H₂₂O₆ (M + H)⁺: 407.1489; found: 407.1488.

Methyl-5-(2,3-dimethoxyphenyl)-6-oxo-5,6-dihydro-2H-

indeno[1,2-b]oxepine-4-carboxylate (4l). In a 5 mL vial, 1,3dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 (E)-(4-methoxy-4-oxobut-2-en-1-yl)dimethylsulfonium equiv.), bromide **3b** (72.3 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 2,3-dimethoxybenzaldehyde 2l (39.8, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/ petroleum ether (v/v, 1: 10) as eluent to give 4l as a Yellow solid (52.5 mg, 67% yield), $R_{\rm f} = 0.2$ (EtOAc/petroleum ether 1 : 10). Mp 181 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 7.46 (d, J = 6.8 Hz, 1H), 7.37 (t, 1H), 7.30 (t, 1H), 7.19 (d, J = 7.2 Hz, 1H), 6.95 (t, 1H), 6.88 (d, J = 7.2 Hz, 1H), 6.79 (dd, $J_1 = 19.2$ Hz, $J_2 = 8.0$ Hz, 2H), 5.34 (s, 1H), 5.00 (dd, $J_1 = 12.4$ Hz, $J_2 = 8.0$ Hz, 1H), 4.54–4.49 (m, 1H), 3.84 (s, 3H), 3.83 (s, 3H), 3.79 (s, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 193.8, 174.4, 166.9, 153.2, 146.2, 145.9, 139.8, 136.1, 132.5, 131.5, 130.1, 128.3, 123.6, 121.3, 121.0, 118.4, 111.6, 108.1, 64.7, 60.3, 55.9, 52.5, 32.3. IR(KBr) v: 2915, 2808, 1733, 1465, 1387, 1283, 1216, 1109, 1022, 981. HRMS(EI) (m/z): calcd for $C_{23}H_{20}O_6 (M + H)^+$: 393.1333; found: 393.1335.

Methyl-6-oxo-5-phenyl-5,6-dihydro-2H-indeno[1,2-b]oxepine-4-carboxylate (4m). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1Hinden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4methoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3b (72.3 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, benzaldehyde 2a (25.4 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:10) as eluent to give 4m as a yellow oil (59.8 mg, 90% yield), $R_{\rm f} = 0.22$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 7.45 (d, J = 7.2 Hz, 1H), 7.37–7.27 (m, 6H), 7.21– 7.15 (m, 3H), 5.37 (s, 1H), 4.91 (dd, $J_1 = 12.4$ Hz, $J_2 = 7.6$ Hz, 1H), 4.63 (dd, $J_1 = 12.4$ Hz, $J_2 = 7.6$ Hz, 1H), 3.84 (s, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 194.3, 173.6, 166.6, 144.2, 141.7, 139.7, 132.5, 132.4, 131.2, 130.0, 128.6, 127.1, 126.7, 121.2, 118.4, 109.5, 65.1, 52.8, 35.5. IR(KBr) v: 2925, 2848, 1719, 1628, 1459, 1403, 1242, 1102, 1039, 976. HRMS(EI) (m/z): calcd for C₂₁H₁₆O₄ $(M + H)^+$: 333.1121; found: 323.1124.

Methyl-5-(4-methoxyphenyl)-6-oxo-5,6-dihydro-2*H*-indeno [1,2-*b*]oxepine-4-carboxylate (4n). In a 5 mL vial, 1,3-dioxo-2,3dihydro-1*H*-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (*E*)-(4-methoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3b (72.3 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and 1-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 4methoxybenzaldehyde 2g (32.6 mg, 1.2 equiv.) and DCM (2 mL)

Paper

were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:10) as eluent to give 4n as a yellow solid (60.1 mg, 83% yield), $R_f = 0.18$ (EtOAc/petroleum ether 1 : 10). Mp 112 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 7.44 (d, J = 7.0 Hz, 1H), 7.35 (t, 1H), 7.29 (t, 1H), 7.23 (d, 2H), 7.20-7.11 (m, 2H), 6.81 (d, J = 8.8 Hz, 1H), 5.30 (s, 1H); ¹³C NMR (101 MHz, CDCl₃), δ: 194.4, 173.5, 166.7, 158.4, 144.4, 139.8, 133.9, 132.5, 132.3, 131.3, 130.1, 128.2, 121.3, 118.5, 114.0, 109.9, 65.2, 55.4, 52.8, 35.0. IR(KBr) v: 2981, 2925, 1712, 1628, 1508, 1459, 1403, 1242, 1186, 1095, 1053. HRMS(EI) (m/z): calcd for C₂₂H₁₈O₅ $(M + H)^+$: 363.1227; found: 363.1225.

Ethyl-6-oxo-5-(2,3,4-trimethoxyphenyl)-5,6-dihydro-2H-

indeno[1,2-b]oxepine-4-carboxylate (40). In a 5 mL vial, 1,3dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium equiv.), bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 2,3,4-trimethoxybenzaldehyde 20 (47 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/ petroleum ether (v/v, 1:5) as eluent to give 40 as a Yellow solid (57.6 mg, 66% yield), $R_{\rm f} = 0.11$ (EtOAc/petroleum ether 1 : 10). Mp 155 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 7.44 (d, J = 7.2 Hz, 1H), 7.36 (t, 1H), 7.30 (t, 1H), 7.18 (d, J = 7.2 Hz, 1H), 6.90 (d, J =8.4 Hz, 1H), 6.79 (t, 1H), 6.52 (d, J = 8.4 Hz, 1H), 5.25 (s, 1H), 4.56-4.51 (m, 1H), 4.33-4.27 (m, 1H), 4.30 (q, 2H), 3.82 (s, 3H), 3.81 (s, 3H), 3.79 (s, 3H), 1.35 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 193.8, 174.3, 166.4, 153.1, 151.0, 146.3, 142.5, 139.8, 132.4, 131.4, 130.0, 128.2, 128.1, 123.1, 121.2, 118.3, 108.3, 106.1, 64.7, 61.4, 60.8, 60.4, 56.0, 32.0, 14.4. IR(KBr) v: 2925, 2855, 1726, 1628, 1466, 1396, 1249, 1109, 1039, 983. HRMS(EI) (m/z): calcd for C₂₅H₂₄O₇ (M + H)⁺: 437.1595; found: 437.1594.

Ethyl-5-(5-methylfuran-2-yl)-6-oxo-5,6-dihydro-2*H*-indeno

[1,2-*b*]oxepine-4-carboxylate (4p). In a 5 mL vial, 1,3-dioxo-2,3dihydro-1*H*-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (*E*)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 5methylfuran-2-carbaldehyde 2p (26.4 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/ petroleum ether (v/v, 1 : 10) as eluent to give 4p as a yellow oil (44.8 mg, 64% yield), $R_f = 0.23$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 7.43 (d, J = 7.2 Hz, 1H), 7.35 (t, J = 8.0 Hz, 1H), 7.28 (t, J = 8.0 Hz, 1H), 7.19 (t, 2H), 6.05 (d, J = 2.8 Hz, 1H), 5.84 (d, J = 2.0 Hz, 1H), 5.47 (dd, J_1 = 12.4 Hz, J_2 = 6.8 Hz, 1H), 5.28 (s, 1H), 4.75 (dd, J_1 = 12.4 Hz, J_2 = 8.0 Hz, 1H), 4.35-4.21 (m, 2H), 2.22 (s, 3H), 1.33 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ : 193.8, 173.5, 165.5, 151.8, 151.4, 141.6, 139.9, 133.1, 132.5, 131.3, 130.0, 121.3, 118.5, 107.6, 107.5, 106.4, 65.4, 61.7, 31.3, 14.3, 13.8. IR(KBr) ν : 2925, 2848, 1719, 1635, 1585, 1459, 1403, 1242, 1109. HRMS(EI) (m/z): calcd for C₂₁H₁₈O₅ (M + H)⁺: 351.1227; found: 351.1225.

Ethyl-5-(furan-2-yl)-6-oxo-5,6-dihydro-2*H*-indeno[1,2-*b*]

oxepine-4-carboxylate (4q). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, furan-2-carbaldehyde 2q (23.4 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:10) as eluent to give 4q as a yellow oil (40.3 mg, 60% yield), $R_{\rm f} = 0.21$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ: 7.43 (d, *J* = 7.2 Hz, 1H), 7.35 (t, 1H), 7.29 (d, 2H), 7.19 $(dd, J_1 = 8.0 Hz, J_2 = 12.0 Hz, 2H), 6.27 (dd, J_1 = 3.2 Hz, J_2 =$ 2.0 Hz, 1H), 6.18 (d, J = 3.2 Hz, 1H), 5.41–5.36 (m, 1H), 5.33 (s, 1H), 4.75 (dd, $J_1 = 12.4$ Hz, $J_2 = 8.0$ Hz, 1H), 4.34–4.23 (m, 2H), 1.32 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ : 193.8, 173.7, 165.4, 153.4, 142.1, 141.4, 139.8, 133.3, 132.5, 131.3, 130.1, 121.3, 118.6, 110.5, 107.3, 106.8, 65.3, 61.8, 31.3, 14.3. IR(KBr) v: 2918, 2848, 1712, 1635, 1543, 1508, 1459, 1403, 1242, 1109, 1046, 1004. HRMS(EI) (m/z): calcd for C₂₅H₂₁NO₄ $(M + H)^+$: 337.1071; found: 337.1076.

Ethyl-6-oxo-5-(thiophen-2-yl)-5,6-dihydro-2*H*-indeno[1,2-b] oxepine-4-carboxylate (4r). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, thiophene-2-carbaldehyde 2r (26.9 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:10) as eluent to give 4r as a yellow oil (50 mg, 71% yield), $R_{\rm f} = 0.24$ (EtOAc/petroleum ether 1 : 10). ¹H NMR $(CDCl_3, 400 \text{ MHz}), \delta: 7.44 \text{ (d}, J = 7.2 \text{ Hz}, 1\text{H}), 7.35 \text{ (t}, J = 8.0 \text{ Hz},$ 1H), 7.29 (t, 1H), 7.19-7.13 (m, 3H), 6.91-6.88 (m, 2H), 5.46 (s, 1H), 5.19–5.15 (m, 1H) 4.71 (dd, $J_1 = 12.8$ Hz, $J_2 = 8.0$ Hz, 1H), 4.37–4.24 (m, 2H), 1.34 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), *b*: 193.8, 173.5, 165.5, 145.5, 143.6, 139.6, 132.7, 132.6, 131.1, 130.2, 126.8, 124.7, 124.4, 121.4, 118.7, 109.6, 65.2, 61.9, 32.4, 14.3. IR(KBr) v: 2925, 2855, 1705, 1628, 1585, 1466, 1396,

1242, 1109, 1046. HRMS(EI) (m/z): calcd for C₂₀H₁₆O₄S (M + H)⁺: 353.0842; found: 353.0845.

Ethyl-6-oxo-5-(pyridin-3-yl)-5,6-dihydro-2*H*-indeno[1,2-*b*]

oxepine-4-carboxylate (4s). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, nicotinaldehyde 2s (25.7 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:10) as eluent to give 4s as a brown oil (39.6 mg, 57% yield), $R_{\rm f} = 0.18$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 8.56 (s, 1H), 8.44 (d, 1H), 7.67 (d, J = 8.0 Hz, 1H), 7.44 (d, J = 7.2 Hz, 1H), 7.36 (t, 1H), 7.29 (t, J = 7.2 Hz, 1H), 7.22–7.17 (m, 3H), 5.35 (s, 1H), 4.84 (dd, $J_1 = 12.4$ Hz, $J_2 = 7.2$ Hz, 1H), 4.73–4.68 (m, 1H), 4.34–4.25 (m, 2H), 1.33 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 194.2, 174.0, 165.8, 148.4, 148.1, 143.3, 139.5, 137.5, 135.2, 132.7, 132.7, 131.1, 130.3, 129.6, 123.5, 121.4, 118.7, 108.4, 65.2, 62.1, 33.9, 14.3. IR(KBr) v: 2974, 2925, 1705, 1621, 1585, 1550, 1459, 1410, 1256, 1116, 1046. HRMS(EI) (m/z): calcd for C₂₁H₁₇NO₄ $(M + H)^+$: 348.1230; found: 348.1227.

Ethyl-5-(1H-indol-3-yl)-6-oxo-5,6-dihydro-2H-indeno[1,2-b]

oxepine-4-carboxylate (4t). In a 5 mL vial, 1,3-dioxo-2,3-dihydro-1H-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 1Hindole-3-carbaldehyde 2t (34.8 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:5) as eluent to give 4t as a yellow solid (29.3 mg, 38% yield), $R_f = 0.15$ (EtOAc/petroleum ether 1 : 10). Mp 183 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 7.69 (d, J = 8.0 Hz, 1H), 7.46 (d, J= 8.0 Hz, 1H), 7.36 (t, 1H), 7.29 (d, J = 8.0 Hz, 2H), 4.84 (dd, $J_1 =$ 12.4 Hz, $J_2 = 7.2$ Hz, 1H), 7.12–7.05 (m, 2H), 6.98 (m, 1H), 5.54 (t, J = 1.2 Hz, 1H), 5.40–5.35 (m, 1H), 4.61 (dd, $J_1 = 12.4$ Hz, $J_2 =$ 8.0 Hz, 1H), 4.39–4.31 (m, 2H), 1.36 (t, J = 6.8 Hz, 3H); ¹³C NMR (101 MHz, $CDCl_3$), δ : 194.0, 173.6, 166.1, 144.1, 139.8, 137.0, 132.4, 132.1, 131.3, 129.9, 125.8, 122.9, 122.3, 121.2, 120.1, 119.7, 118.3, 117.4, 111.2, 109.8, 64.5, 61.8, 28.5, 14.3. IR(KBr) v: 2939, 1719, 1635, 1200, 1144. HRMS(EI) (m/z): calcd for $C_{25}H_{21}NO_4 (M + H)^+$: 86.1543; found: 386.1541.

Ethyl-5-(1-methyl-1H-indol-3-yl)-6-oxo-5,6-dihydro-2H-

indeno[1,2-*b*]oxepine-4-carboxylate (4u). In a 5 mL vial, 1,3dioxo-2,3-dihydro-1*H*-inden-2-ylium 1a (29.2 mg, 0.2 mmol, 1.0 equiv.), (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.) and L-proline (2.3 mg, 0.1 equiv.) was added. To this resultant mixture, 1-methyl-1H-indole-3-carbaldehyde 2u (38.1 mg, 1.2 equiv.) and DCM (2 mL) were added and stirred at room temperature overnight. The reaction was confirmed by TLC. Then the solution was diluted with ethyl acetate and transferred to a round bottom flask. Silica gel was added to the flask and volatiles were evaporated under vacuum. The purification was performed by flash column chromatography on silica gel using ethyl acetate/petroleum ether (v/v, 1:5) as eluent to give 4u as a yellow oil (55.9 mg, 70% yield), $R_{\rm f}$ = 0.12 (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ: 7.67 (d, J = 8.0 Hz, 1H), 7.46 (d, J = 7.2 Hz, 1H), 7.37 (t, J =8.0 Hz, 1H), 7.30 (t, J = 8.0 Hz, 1H), 7.24-7.18 (m, 3H), 7.12-7.05 (m, 2H), 6.87 (s, 1H), 5.53 (s, 1H), 5.44–5.39 (m, 1H), 4.65–4.60 (m, 1H), 4.39–4.33 (m, 2H), 3.70 (s, 3H), 1.37 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 194.1, 173.6, 166.2, 144.4, 139.9, 137.8, 132.5, 132.1, 131.4, 130.0, 127.7, 126.3, 122.0, 121.3, 120.4, 119.3, 118.3, 115.8, 110.1, 109.3, 64.7, 61.8, 32.9, 29.8, 28.6, 14.4. IR(KBr) v: 2925, 2855, 1719, 1628, 1585, 1473, 1403, 1235, 1109, 1039, 983. HRMS(EI) (m/z): calcd for C₂₅H₂₁NO₄ $(M + H)^+$: 400.1543; found: 400.1544.

Ethyl-(E)-3-(2,2-dicyano-3-phenylcyclopropyl)acrylate (6a). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (E)-(4-ethoxy-4oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and benzaldehyde 2a (25.4 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA : PE, 1:10) to afford compound 6a as a brown oil (44.2 mg, 83% yield), $R_{\rm f} = 0.17$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ: 7.43 (m, 3H), 7.32 (m, 2H), 6.43–6.34 (m, 2H), 4.23–4.18 (m, 2H), 3.57 (d, J =9.6 Hz, 1H), 3.15–3.10 (m, 1H), 1.28 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (101 MHz, $CDCl_3$), δ : 164.4, 136.1, 129.9, 129.6, 129.6, 129.1, 128.2, 114.8, 111.1, 61.2, 38.5, 36.4, 14.3, 12.3. IR(KBr) v: 2927.97, 2253.41, 1721.18, 1646.23, 1271.47, 1162.45, 1039.80, 971.66, 740.00, 692.30. HRMS(EI) (m/z): calcd for C₁₆H₁₄N₂O₂ (M $(+ H)^+$: 267.1128; found: 267.1123.

Ethyl-(E)-3-(2,2-dicyano-3-(4-nitrophenyl)cyclopropyl)acrylate (6b). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and 4-nitrobenzaldehyde 2b (36.2 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA: PE, 1:5) to afford compound **6b** as a yellow oil (46.7 mg, 75% yield), $R_{\rm f} = 0.23$ (EtOAc/petroleum ether 1 : 5).¹H NMR (CDCl₃, 400 MHz), δ : 8.29 $(d, 2H), 7.43 (d, 2H), 7.05 (s, 1H), 4.90 (s, 1H), 4.14 (dd, J_1 = 7.2 Hz)$ $J_2 = 4.0$ Hz, 2H), 3.48 (m, 2H), 1.17 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 161.7, 148.6, 141.2, 139.4, 136.6, 129.3, 124.6, 115.9, 113.1, 61.8, 59.6, 44.0, 39.8, 14.1. IR(KBr) v: 2918, 2245, 1719, 1649, 1494, 1466, 1256, 1165, 1032, 976, 751. HRMS(EI) (m/z): calcd for $C_{16}H_{13}N_{3}O_{4}$ (M + H)⁺: 312.0979; found: 312.0976.

Ethyl-(*E*)-3-(2,2-dicyano-3-(4-cyanophenyl)cyclopropyl)acrylate (6c). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (*E*)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline

Paper

(2.3 mg, 0.1 equiv.) and 4-formylbenzonitrile **2c** (31.4 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated *in vacuo*. The residue was purified by column chromatography (EA : PE, 1 : 3) to afford compound **6c** as a white solid (40.2 mg, 69% yield), $R_{\rm f}$ = 0.22 (EtOAc/petroleum ether 1 : 3). Mp 151 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 7.75 (d, 2H), 7.46 (d, 2H), 6.41 (d, J = 15.2 Hz, 1H), 6.27 (dd, J_1 = 15.6 Hz, J_2 = 10.0 Hz, 1H), 4.21 (dd, J_1 = 14.0 Hz, J_2 = 7.2 Hz, 2H), 3.57 (d, J = 10.0 Hz, 1H), 3.20 (t, J = 10.0 Hz, 1H), 1.28 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ : 164.1, 134.8, 133.3, 133.3, 130.8, 130.1, 117.9, 114.0, 113.9, 110.7, 61.4, 37.5, 35.9, 14.3, 12.4. IR(KBr) ν : 2925, 2231, 1733, 1642, 1438, 1284, 1228, 1046. HRMS(EI) (m/z): calcd for C₁₇H₁₃N₃O₂ (M + H)⁺: 292.1081; found: 292.1080.

Ethyl-(E)-3-(2,2-dicvano-3-(4-(dimethylamino)phenyl)cyclopropyl)acrylate (6d). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), Lproline (2.3 mg, 0.1 equiv.) and 4-(dimethylamino)benzaldehyde 2d (35.8 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA: PE, 1:5) to afford compound 6d as a brown oil (55 mg, 89% yield), $R_{\rm f} = 0.17$ (EtOAc/petroleum ether 1 : 5). ¹H NMR (CDCl₃, 400 MHz), δ: 7.13 (d, 2H), 6.74–6.67 (m, 3H), 6.30 (d, J = 16.0 Hz, 1H), 4.28–4.22 (m, 2H), 3.27 (d, J = 8.4 Hz, 1H), 3.06 (t, J = 8.8 Hz, 1H), 2.98 (s, 6H), 1.32 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 164.7, 151.2, 138.4, 129.1, 128.2, 115.9, 113.3, 112.7, 112.4, 61.2, 41.1, 40.3, 35.5, 14.9, 14.3. IR(KBr) v: 2925, 2245, 1712, 1614, 1536, 1354, 1277, 1165, 1095, 1039, 964, 822. HRMS(EI) (m/z): calcd for C₁₈H₁₉N₃O₂ (M + H)⁺: 310.1550; found: 310.1553.

Ethyl-(*E*)-3-(2,2-dicyano-3-(2-methoxyphenyl)cyclopropyl)

acrylate (6e). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and 2-methoxybenzaldehyde 2e (32.6 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA : PE, 1:5) to afford compound **6e** as a white solid (44.4 mg, 75% yield), $R_{\rm f}$ = 0.46 (EtOAc/petroleum ether 1:3). Mp 118 $^{\circ}$ C; ¹H NMR $(CDCl_3, 400 \text{ MHz}), \delta: 7.40 \text{ (t, } J = 8.0 \text{ Hz}, 1\text{H}), 7.12 \text{ (d, } J = 7.6 \text{ Hz},$ 1H), 7.01–6.94 (m, 2H), 6.45 (dd, $J_1 = 15.6$ Hz, $J_2 = 10.0$ Hz, 1H), 6.35 (d, J = 15.6 Hz, 1H), 4.24-4.19 (m, 2H), 3.93 (s, 3H), 3.29 (d, J = 9.6 Hz, 1H), 3.11 (t, J = 9.2 Hz, 1H), 1.28 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 164.6, 158.7, 137.1, 131.2, 130.3, 128.4, 120.8, 116.9, 115.1, 111.7, 111.3, 61.1, 55.8, 36.2, 35.0, 14.3, 13.1. IR(KBr) v: 2921, 2245, 1714, 1646, 1503, 1462, 1257, 1162, 1026, 978, 753. HRMS(EI) (m/z): calcd for C₁₇H₁₆N₂O₃ (M + H)⁺: 297.1234; found: 297.1229.

Ethyl-(E)-3-(2,2-dicyano-3-(2,3,4-trimethoxyphenyl)cyclo-

propyl)acrylate (6f). A mixture of malononitrile **5a** (13.2 mg, 1 equiv.), (*E*)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide **3a** (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and 2,3,4-trimethoxybenzaldehyde **2o** (47 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred

at 0 °C overnight. The reaction mixture was concentrated *in vacuo*. The residue was purified by column chromatography (EA : PE, 1 : 3) to afford compound **6f** as a white solid (49.8 mg, 70% yield), $R_{\rm f} = 0.22$ (EtOAc/petroleum ether 1 : 3). Mp 115 °C; ¹H NMR (CDCl₃, 400 MHz), δ : 6.81 (d, J = 8.4 Hz, 1H), 6.61 (d, J = 8.4 Hz, 1H), 6.53–6.46 (m, 1H), 6.34 (d, J = 15.6 Hz, 1H), 4.25–4.19 (m, 2H), 4.08 (s, 3H), 3.87 (s, 3H), 3.86 (s, 3H), 3.29 (d, J = 9.2 Hz, 1H), 3.08 (t, J = 10.0 Hz, 1H), 1.29 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ : 164.6, 155.4, 153.3, 141.9, 137.1, 128.5, 124.7, 115.0, 114.0, 111.8, 106.6, 61.4, 61.1, 61.1, 56.1, 36.2, 35.3, 14.3, 13.0. IR(KBr) ν : 2925, 2848, 2252, 1719, 1656, 1607, 1494, 1466, 1424, 1312, 1263, 1158, 1095, 1039, 976. HRMS(EI) (m/z): calcd for C₁₉H₂₀N₂O₅ (M + H)⁺: 357.1445; found: 357.1447.

Ethyl-(E)-3-(2,2-dicvano-3-(pyridin-3-yl)cyclopropyl)acrylate (6g). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and nicotinaldehyde 2s (25.7 mg, 1.2 equiv.) (25.4 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA: PE, 1:5) to afford compound 6g as a yellow oil (35.8 mg, 67% yield), $R_{\rm f} = 0.24$ (EtOAc/petroleum ether 1 : 5). ¹H NMR $(CDCl_3, 400 \text{ MHz}), \delta: 8.66 \text{ (d}, J = 3.6 \text{ Hz}, 1\text{H}), 8.52 \text{ (s, 1H)}, 7.56$ $(dt, J_1 = 8.0 Hz, J_2 = 2.0 Hz, 1H)$, 7.37 $(dd, J_1 = 7.6 Hz, J_2 =$ 4.8 Hz, 1H), 7.02 (dd, $J_1 = 4.0$ Hz, $J_2 = 2.4$ Hz, 1H), 4.81 (s, 1H), 4.19–4.08 (m, 2H), 3.51–3.39 (m, 2H), 1.16 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 161.8, 150.9, 149.2, 139.0, 136.6, 135.8, 130.1, 124.1, 116.1, 113.3, 61.7, 57.8, 43.8, 40.0, 14.1. IR(KBr) v: 1712, 1642, 1249, 1102, 1025, 709. HRMS(EI) (m/z): calcd for $C_{15}H_{13}N_3O_2 (M + H)^+$: 268.1081; found: 268.1085.

Ethyl-(E)-3-(2,2-dicyano-3-(5-methylfuran-2-yl)cyclopropyl) acrylate (6h). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and 5-methylfuran-2-carbaldehyde 2p (26.4 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA : PE, 1 : 10) to afford compound 6h as a brown oil (36.2 mg, 67% yield), $R_{\rm f} = 0.45$ (EtOAc/petroleum ether 1 : 5). ¹H NMR $(CDCl_3, 400 \text{ MHz}), \delta: 6.81 \text{ (dd}, J_1 = 15.6, J_2 = 10.0 \text{ Hz}, 1\text{H}), 6.33$ (d, 2H), 5.99 (s, 1H), 4.27–4.21 (m, 2H), 3.43 (d, *J* = 9.6 Hz, 1H), 3.05 (t, J = 10.0 Hz, 1H), 2.30 (s, 3H), 1.31 (t, J = 6.8 Hz, 3H);¹³C NMR (101 MHz, CDCl₃), δ: 164.5, 154.6, 140.5, 135.7, 129.5, 114.2, 113.6, 110.9, 107.2, 61.2, 35.4, 32.9, 14.3, 13.7, 12.7. IR(KBr) v: 2921, 2362, 1714, 1653, 1278, 1039, 978, 571. HRMS(EI) (m/z): calcd for C₁₅H₁₄N₂O₃ $(M + H)^+$: 271.1077; found: 277.1072.

Ethyl-(*E*)-3-(2,2-dicyano-3-(furan-2-yl)cyclopropyl)acrylate

(6i). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (*E*)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na_2CO_3 (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and furan-2-carbaldehyde 2q (23.4 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated *in vacuo*. The residue was purified by column chromatography (EA : PE, 1 : 10) to afford compound **6i** as a yellow oil (42 mg, 82% yield), $R_{\rm f} = 0.5$ (EtOAc/petroleum ether 1 : 5). ¹H NMR (CDCl₃, 400 MHz), δ : 7.46 (d, 1H), 6.69 (dd, $J_1 = 15.6$ Hz, $J_2 = 10.0$ Hz, 1H), 6.48–6.39 (m, 2H), 6.32 (t, J = 17.2 Hz, 1H), 4.26–4.21 (m, 2H), 2.78 (d, J = 10.0 Hz, 1H), 1.81 (s, 1H), 1.30 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ : 164.4, 146.5, 144.1, 143.7, 136.5, 128.9, 111.3, 111.2, 110.8, 61.1, 42.5, 37.6, 23.0, 14.2. IR(KBr) ν : 2918, 2245, 1712, 1656, 1452, 1270, 1039, 976, 744, 667. HRMS(EI) (m/z): calcd for C₁₄H₁₂N₂O₃ (M + H)⁺: 257.0921; found: 257.0923.

Ethyl-(E)-3-(2,2-dicyano-3-(thiophen-2-yl)cyclopropyl)acrylate (6j). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (E)-(4ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and thiophene-2-carbaldehyde 2s (26.9 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA: PE, 1:10) to afford compound 6j as a yellow oil (42.4 mg, 78% yield), $R_{\rm f} = 0.24$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 7.39 (d, J = 5.2 Hz, 1H), 7.16–7.14 (m, 1H), 7.07– 7.04 (m, 1H), 6.67–6.61 (m, 1H), 6.37 (d, J = 15.6 Hz, 1H), 4.23 (q, 2H), 3.59 (d, *J* = 9.6 Hz, 1H), 3.14 (t, 1H), 1.30 (t, *J* = 6.8 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 164.4, 135.3, 129.9, 129.8, 129.7, 128.1, 127.8, 114.2, 110.9, 61.3, 36.8, 34.2, 14.3, 13.9. IR(KBr) v: 2918, 2245, 1719, 1649, 1459, 1375, 1319, 1277, 1172, 1102, 1039, 969, 857, 702. HRMS(EI) (m/z): calcd for C₁₄H₁₂N₂O₂S (M + H)⁺: 273.0692; found: 273.0694.

Ethyl-(E)-3-(2,2-dicyano-3-(1-methyl-1H-indol-2-yl)cyclo-

propyl)acrylate (6k). A mixture of malononitrile 5a (13.2 mg, 1 (E)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium equiv.), bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), Lproline (2.3 mg, 0.1 equiv.) and 1-methyl-1H-indole-3-carbaldehyde 2u (38.1 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA: PE, 1:3) to afford compound 6k as a brown yellow (49.8 mg, 78% yield), $R_{\rm f} = 0.21$ (EtOAc/petroleum ether 1 : 3). Mp 113 °C; ¹H NMR $(CDCl_3, 400 \text{ MHz}), \delta$: 7.62 (d, J = 8.0 Hz, 1H), 7.37-7.31 (m, 2H),7.25–7.22 (m, 1H), 7.06 (s, 1H), 6.79 (dd, $J_1 = 15.6$ Hz, $J_2 = 8.8$ Hz, 1H), 6.35 (d, J = 16.0 Hz, 1H), 4.30–4.24 (m, 2H), 3.81 (s, 3H), 3.45 (d, J = 8.4 Hz, 1H), 3.02 (t, J = 8.8 Hz, 1H), 1.33 (t, J = 7.2 Hz, 3H);¹³C NMR (101 MHz, CDCl₃), δ: 164.7, 138.3, 137.2, 128.4, 127.5, 123.2, 120.7, 118.2, 113.2, 112.9, 110.2, 104.3, 61.3, 36.5, 33.9, 33.3, 14.9, 14.3. IR(KBr) v: 2918, 1719, 1656, 1466, 1375, 1270, 1214, 1039, 976, 737. HRMS(EI) (m/z): calcd for C₁₉H₁₇N₃O₂ (M + H)⁺: 320.1394; found: 320.1389.

Ethyl-(*E*)-3-(2,2-dicyano-3-phenethylcyclopropyl)acrylate (6l). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (*E*)-(4-ethoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and 3-phenylpropanal 2v (32.2 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated *in vacuo*. The residue was purified by column chromatography (EA : PE, 1 : 10) to afford compound 6l as a brown oil (33.5 mg, 57% yield), $R_f = 0.24$ (EtOAc/petroleum

ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 7.32 (t, 2H), 7.24– 7.18 (m, 3H), 6.48 (dd, $J_1 = 15.6$ Hz, $J_2 = 10.0$ Hz, 1H), 6.18 (d, J = 15.2 Hz, 1H), 4.26–4.20 (m, 2H), 2.91–2.78 (m, 2H), 2.73 (t, J = 10.0 Hz, 1H), 2.24–2.18 (m, 1H), 2.13–1.94 (m, 2H), 1.31 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ : 164.5, 139.2, 134.7, 129.5, 129.0, 128.7, 128.6, 127.0, 114.6, 111.4, 61.2, 35.6, 35.1, 34.1, 27.4, 14.3, 11.1. IR(KBr) ν : 2924, 1634.18, 1454.10, 1202.00, 1036.32, 748.19, 697. HRMS(EI) (m/z): calcd for C₁₈H₁₈N₂O₂ (M + H)⁺: 295.1441; found: 295.1445.

Methyl-(E)-3-(2,2-dicyano-3-phenylcyclopropyl)acrylate (6m). A mixture of malononitrile 5a (13.2 mg, 1 equiv.), (E)-(4-methoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3b (72.3 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and benzaldehyde 2a (25.4 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA: PE, 1:5) to afford compound 6m as a white solid (42.3 mg, 84% yield), $R_{\rm f} = 0.16$ (EtOAc/petroleum ether 1 : 10). Mp 122 °C; ¹H NMR (CDCl₃, 400 MHz), δ: 7.37 (s, 3H), 7.21–7.15 (m, 1H), 6.33–6.32 (m, 2H), 3.69 (d, J = 3.6 Hz, 3H), 3.52 (d, J = 9.6 Hz, 1H), 3.10–3.05 (m, 1H). ¹³C NMR (101 MHz, CDCl₃), δ: 164.4, 135.3, 130.0, 129.7, 129.7, 128.1, 127.8, 114.2, 110.9, 61.3, 36.8, 34.2, 14.3, 13.9. IR(KBr) v: 2918, 2848, 2252, 1726, 1656, 1438, 1270, 1039, 737, 695. HRMS(EI) (m/z): calcd for $C_{15}H_{12}N_2O_2 (M + H)^+$: 253.0972; found: 253.0966.

Methyl-(E)-3-(2,2-dicyano-3-(4-(dimethylamino)phenyl)cyclopropyl)acrylate (6n). A mixture of malononitrile 5a (13.2 mg, 1 (E)-(4-methoxy-4-oxobut-2-en-1-yl)dimethylsulfonium eauiv.). bromide 3b (72.3 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), Lproline (2.3 mg, 0.1 equiv.) and 4-(dimethylamino)benzaldehyde 2f (35.8 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated *in vacuo*. The residue was purified by column chromatography (EA: PE, 1:5) to afford compound 6n as a brown oil (52.5 mg, 89% yield), $R_{\rm f} = 0.34$ (EtOAc/petroleum ether 1 : 5). ¹H NMR $(CDCl_3, 400 \text{ MHz}), \delta: 7.13 \text{ (d}, J = 8.8 \text{ Hz}, 2\text{H}), 6.75-6.68 \text{ (m, 3H)},$ 6.31 (d, J = 15.2 Hz, 1H), 3.79 (s, 3H), 3.27 (d, J = 8.4 Hz, 1H), $3.06 (t, J = 8.8 \text{ Hz}, 1\text{H}), 2.98 (s, 6\text{H}); {}^{13}\text{C} \text{ NMR} (101 \text{ MHz}, \text{CDCl}_3),$ δ: 165.1, 151.2, 138.8, 129.1, 127.7, 115.9, 113.3, 112.6, 112.4, 52.3, 41.2, 40.3, 35.5, 14.9. IR(KBr) v: 2918, 1726, 1529, 1277, 1116, 1046, 983. HRMS(EI) (m/z): calcd for $C_{17}H_{17}N_3O_2 (M + H)^+$: 296.1394; found: 296.1396.

Methyl-1-cyano-2-((*E*)-3-methoxy-3-oxoprop-1-en-1-yl)-3phenylcyclopropane-1-carboxylate (6o). A mixture of methyl 2cyanoacetate 5b (19.8 mg, 1 equiv.), (*E*)-(4-methoxy-4-oxobut-2-en-1-yl)dimethylsulfonium bromide 3b (72.3 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and benzaldehyde 2a (25.4 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated *in vacuo*. The residue was purified by column chromatography (EA : PE, 1 : 5) to afford compound 6o as a white oil (42.2 mg, 74% yield), $R_{\rm f} = 0.15$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 7.42–7.36 (m, 3H), 7.29 (d, 2H), 6.95 (dd, $J_1 =$ 15.6 Hz, $J_2 = 9.6$ Hz, 1H), 6.25 (d, J = 15.6 Hz, 1H), 3.88 (s, 3H), 3.76 (s, 3H), 3.53 (d, J = 8.4 Hz, 1H), 3.08 (t, 1H); ¹³C NMR (101 MHz, CDCl₃), δ : 165.7, 165.1, 139.6, 132.0, 129.1, 129.0, 128.3, 126.6, 115.6, 54.1, 52.0, 39.6, 38.1, 29.8. IR(KBr) ν : 1726, 1649, 1438, 1277, 1249, 983, 702. HRMS(EI) (m/z): calcd for C₁₆H₁₅NO₄ (M + H)⁺: 286.1074; found: 286.1070.

Methyl-1-cyano-2-((E)-3-ethoxy-3-oxoprop-1-en-1-yl)-3-

phenylcyclopropane-1-carboxylate (6p). A mixture of 2-cyanoacetate 5b (19.8 mg, 1 equiv.), (E)-(4-ethoxy-4-oxobut-2-en-1-yl) dimethylsulfonium bromide 3a (76.5 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and benzaldehyde 2a (25.4 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA: PE, 1:5) to afford compound **6p** as a white oil (44.9 mg, 75% yield), $R_{\rm f}$ = 0.16 (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ : 7.42–7.36 (m, 3H), 7.30 (d, J = 6.8 Hz, 2H), 6.97–6.91 (m, 1H), 6.24 (d, 1H), 4.25-4.19 (m, 2H), 3.88 (s, 3H), 3.54 (d, J = 8.8 Hz, 1H), 3.07 (t, J = 9.2 Hz, 1H), 1.30 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 165.3, 165.1, 139.3, 132.0, 129.1, 129.0, 128.3, 127.1, 115.6, 60.9, 54.1, 39.6, 38.2, 29.7, 14.3. IR(KBr) v: 2245, 1712, 1649, 1438, 1277, 1242, 1179, 1123, 1032, 983, 695. HRMS(EI) (m/ z): calcd for $C_{17}H_{17}NO_4 (M + H)^+$: 300.1230; found: 300.1232.

Ethyl-1-cyano-2-((E)-3-methoxy-3-oxoprop-1-en-1-yl)-3phenylcyclopropane-1-carboxylate (6q). A mixture of ethyl 2cyanoacetate 5c (22.6 mg, 1 equiv.), (E)-(4-methoxy-4-oxobut-2en-1-yl)dimethylsulfonium bromide 3b (72.3 mg, 1.5 equiv.), Na₂CO₃ (42 mg, 2 equiv.), L-proline (2.3 mg, 0.1 equiv.) and benzaldehyde 2a (25.4 mg, 1.2 equiv.) in DCM. The reaction mixture was stirred at 0 °C overnight. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography (EA: PE, 1:5) to afford compound **6q** as a yellow oil (41.9 mg, 70% yield), $R_{\rm f} = 0.15$ (EtOAc/petroleum ether 1 : 10). ¹H NMR (CDCl₃, 400 MHz), δ: 7.43-7.36 (m, 3H), 7.31–7.29 (m, 2H), 6.96 (dd, $J_1 = 15.6$ Hz, $J_2 = 9.6$ Hz, 1H), 6.24 (d, J = 15.6 Hz, 1H), 4.36–4.30 (m, 2H), 3.76 (s, 3H), 3.53 (d, J = 8.4 Hz, 1H), 3.06 (t, J = 8.8 Hz, 1H), 1.36 (t, J = 7.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃), δ: 165.6, 164.5, 139.7, 132.0, 129.0, 128.9, 128.2, 126.4, 115.5, 63.5, 51.9, 39.3, 38.0, 29.9, 14.1. IR(KBr) v: 2925, 1733, 1649, 1508, 1284, 990, 702. HRMS(EI) (m/ z): calcd for $C_{17}H_{17}NO_4 (M + H)^+$: 300.1230; found: 300.1234.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We are grateful for the financial support from NSFC of China (21801114) and Jiangxi Province "1000 Talent Plan" High-Level Entrepreneurial and Innovative Program (1001-02102129).

Notes and references

 (a) V. K. Tandon, M. Kumar, A. K. Awasthi, H. O. Saxena and G. K. Goswamy, *Bioorg. Med. Chem. Lett.*, 2004, 14, 3177– 3180; (b) J. R. Vyvyan, J. M. Oaksmith, B. W. Parks and E. M. Peterson, *Tetrahedron Lett.*, 2005, 46, 2457–2460; (c) T. Sugaya, N. Kato, A. Sakaguchi and S. Tomioka, *Synthesis*, 1995, 10, 1257–1262; (d) L. S. Nicole, M. H. Heather and W. P. Mark, *Tetrahedron*, 2006, 62, 9301–9320; (e) C. R. Reddy, P. Ramesh, N. N. Rao and S. A. Ali, *Eur. J. Org. Chem.*, 2011, **11**, 2133–2141.

- 2 (a) J. W. Blunt, B. R. Copp, W.-P. Hu, M. H. G. Munro, P. T. Northcote and M. R. Prinsep, *Nat. Prod. Rep.*, 2008,
 25, 35–95; (b) T. F. Molinski, D. S. Dalisay, S. L. Lievens and J. P. Saludes, *Nat. Rev. Drug Discovery*, 2009, 8, 69–85;
 (c) V. J. Paul and R. Ritson-Williams, *Nat. Prod. Rep.*, 2008,
 25, 662–695; (d) D. Skropeta, *Nat. Prod. Rep.*, 2008, 25, 1131–1166.
- 3 (a) G. Boche and H. M. Walbirsky, Cyclopropane Derived Reactive Intermediates, John Wiley and Sons, New York, NY, 1990; (b) Z. Rappoport, The Chemistry of the Cyclopropyl Group, Wiley and Sons, New York, NY, 1996; (c) D. W. Graham, W. T. Ashton, L. Barash, J. E. Brown, R. D. Brown, L. F. Canning, A. Chen, J. P. Springer and E. F. Rogers, J. Med. Chem., 1987, 30, 1074-1090; (d) J. Salaun and M. S. Baird, Curr. Med. Chem., 1995, 2, 511-519; (e) Y. Baba, G. Saha, S. Nakao, C. Iwata, T. Tanaka, T. Ibuka, H. Ohishi and Y. Takemoto, J. Org. Chem., 2001, 66, 81-88; (f) D. L. Boger, T. V. Hughes and M. P. Hedrick, J. Org. Chem., 2001, 66, 2207-2216; (g) S. Yoshida, T. C. Rosen, O. G. J. Meyer, M. J. Sloan, S. Ye, G. Haufe and K. L. Kirk, Bioorg. Med. Chem., 2004, 12, 2645-2652; (h) K. Yamaguchi, Y. Kazuta, K. Hirano, S. Yamada, A. Matsuda and S. Shuto, Bioorg. Med. Chem., 2008, 16, 8875-8881; (i) W. A. Donaldson, Tetrahedron, 2001, 57, 8589-8627; (j) W. Brandt and T. Thiemann, Chem. Rev., 2003, 103, 1625-1647.
- 4 (a) W. P. Deng, A. H. Li, L. X. Dai and X.-L. Hou, *Tetrahedron*, 2000, 56, 2967–2974; (b) S. Ye, Z. Z. Huang, C. A. Xia, Y. Tang and L. X. Dai, *J. Am. Chem. Soc.*, 2002, 124, 2432–2433; (c) X. F. Yang, M. J. Zhang, X. L. Hou and L. X. Dai, *J. Org. Chem.*, 2002, 67, 8097–8103; (d) D. Morton, D. Pearson, R. A. Field and R. A. Stockman, *Org. Lett.*, 2004, 6, 2377–2380; (e) D. Morton, D. Pearson, R. A. Field and R. A. Stockman, *Org. Lett.*, 2004, 6, 2377–2380; (e) D. Morton, D. Pearson, R. A. Field and R. A. Stockman, *Chem. Commun.*, 2006, 17, 1833–1835; (f) X. M. Deng, P. Cai, S. Ye, X. L. Sun, W. W. Liao, K. Li, Y. Tang, Y. D. Wu and L. X. Dai, *J. Am. Chem. Soc.*, 2006, 128, 9730–9740; (g) Y. Q. Zhang, A. M. Yu, J. R. Jia, S. S. Ma, K. Li, Y. Wei and X. T. Meng, *Chem. Commun.*, 2017, 53, 10672–10675.
- 5 (a) A. H. Li, L. X. Dai and X. L. Hou, Chem. Commun., 1996, 4, 491-492; (b) Y.-G. Zhou, A.-H. Li, X. L. Hou and L. X. Dai, Chem. Commun., 1996, 11, 1353-1354; (c) A.-H. Li, L.-X. Dai, X.-L. Hou and M.-B. Chen, J. Org. Chem., 1996, 61, 4641-4648; (d) D. Morton, D. Pearson, R. A. Field and R. A. Stockman, Org. Lett., 2004, 6, 2377-2380; (e) B.-H. Zhu, J.-C. Zheng, C.-B. Yu, X.-L. Sun, Y.-G. Zhou, Q. Shen and Y. Tang, Org. Lett., 2010, 12, 504-507; (f) F. Gao and Y. Huang, Adv. Synth. Catal., 2014, 356, 2422-2428; (g) Z. Chen and J. Zhang, Chem.-Asian J., 2009, 4, 1527-1529.
- 6 (a) Q.-G. Wang, X.-M. Deng, B.-H. Zhu, L.-W. Ye, X.-L. Sun, C.-Y. Li, C.-Y. Zhu, Q. Shen and Y. Tang, J. Am. Chem. Soc., 2008, 130, 5408–5409; (b) B.-H. Zhu, R. Zhou, J.-C. Zheng, X.-M. Deng, X.-L. Sun, Q. Shen and Y. Tang, J. Org. Chem., 2010, 75, 3454–3457.

- 8 J. L. Chen, P. H. Jia and Y. Huang, *Org. Lett.*, 2018, **20**, 6715–6718.
- 9 (a) S. Brauch, S. S. VanBerkel and B. Westermann, *Chem. Soc. Rev.*, 2013, 42, 4948–4962; (b) R. C. Cioc, E. Ruijter and R. V. A. Orru, *Green Chem.*, 2014, 16, 2958–2975; (c) X. Guo and W. Hu, *Acc. Chem. Res.*, 2013, 46, 2427–2440; (d) P. Wu, M. Givskov and T. Nielsen, *Chem. Rev.*, 2019, 119, 11245–11290.
- 10 For selected reviews on sulfur ylides, see: (a) L.-Q. Lu, T.-R. Li, Q. Wang and W.-J. Xiao, *Chem. Soc. Rev.*, 2017, 46, 4135–4149; (b) J.-R. Chen, X.-Q. Hu, L.-Q. Lu and W.-J. Xiao, *Chem. Rev.*, 2015, 115, 5301–5365; (c) X.-L. Sun

and Y. Tang, Acc. Chem. Res., 2008, 41, 937–948; (d) A.-H. Li, L.-X. Dai and V. K. Aggarwal, Chem. Rev., 1997, 97, 2341–2372.

- 11 CCDC 1956080.†
- 12 CCDC 1956081.†
- 13 (a) W. H. Ding, Y. Q. Zhang, A. M. Yu, L. Zhang and X. T. Meng, *J. Org. Chem.*, 2018, 83, 13821–13833; (b) L. Satham and I. N. N. Namboothiri, *J. Org. Chem.*, 2018, 83, 9471–9477; (c) Y. H. Wang, J. R. Cao, D. I. Schuster and J. R. Wilson, *Tetrahedron Lett.*, 1995, 36, 6843–6846.
- 14 W. Yin, L. Fang, Z. Wang, F. Gao, Z. Li and Z. Wang, Org. Lett., 2019, 21, 7361-7364.