



Cite this: *RSC Adv.*, 2020, 10, 24830

# Copper(II)-catalyzed synthesis of multisubstituted indoles through sequential Chan–Lam and cross-dehydrogenative coupling reactions†

Xin Chen,<sup>ab</sup> Yunyun Bian,<sup>a</sup> Baichuan Mo,<sup>ab</sup> Peng Sun,<sup>ab</sup> Chunxia Chen<sup>\*ab</sup>  
and Jinsong Peng<sup>id</sup><sup>\*a</sup>

Starting from arylboronic acids and ester (Z)-3-aminoacrylates, one-pot syntheses of diverse indole-3-carboxylic esters have been described through copper(II)-catalyzed sequential Chan–Lam *N*-arylation and cross-dehydrogenative coupling (CDC) reactions. The initial Chan–Lam arylation can proceed in DMF at 100 °C for 24 h to give ester (Z)-3-(arylamino)acrylate intermediates in the presence of Cu(OAc)<sub>2</sub>/tri-*tert*-butylphosphine tetrafluoroborate, a catalytic amount of myristic acid as the additive, KMnO<sub>4</sub> and KHCO<sub>3</sub>. Sequentially, these *in situ* arylated intermediates can undergo an intramolecular oxidative cross-dehydrogenative coupling process in mixed solvents (DMF/DMSO = 2 : 1) at 130 °C to give C3-functionalized multi-substituted indole derivatives.

Received 24th May 2020  
Accepted 22nd June 2020

DOI: 10.1039/d0ra04592f


rsc.li/rsc-advances

## 1. Introduction

The indole unit is not only one of the most abundant structural motifs in natural products,<sup>1</sup> but is also ubiquitous among agrochemicals,<sup>2</sup> marketed medicines (such as dolasetron, tropisetron, indomethacin, proglumetacin, and ondansetron),<sup>3</sup> and progressive functional materials.<sup>4</sup> In particular, substituted indoles have been utilized as “privileged scaffolds” for drug discovery of anti-inflammatory, antihypertensive, anti-tumor, anti-HIV, and antimigraine agents.<sup>5</sup> Considering the importance of indole scaffolds for pharmaceutical research, the development of practical syntheses of indole derivatives is of immense interest to synthetic chemists.

Over the past century, a variety of approaches for indole preparation have been well established.<sup>6</sup> While numerous classical procedures based on condensation and cyclization have been developed,<sup>6b</sup> there is still a great demand to explore new strategies and methodologies for the modular synthesis of functionalized indoles from easily available starting materials. Over the past decades, transition metal-mediated inter- and intramolecular C–C and C–N bond forming reactions have emerged as one of the most powerful and popular tools for indole syntheses.<sup>7</sup> Following this tendency, an attractive C–H activation/cyclization strategy relying on the use of *N*-arylated

enamine or imine intermediates have emerged in recent years (Scheme 1). In 2008, Glorius group<sup>8a</sup> reported the first Pd-catalyzed oxidative cyclization of *N*-aryl enaminones/esters for indole syntheses with Cu(OAc)<sub>2</sub> as the oxidant. Shortly afterward, Cacchi<sup>9</sup> demonstrated a copper-catalyzed aerobic version for the synthesis of indoles from *N*-aryl enaminones in DMF. Subsequent to these original works, different catalytic system (Pd,<sup>10</sup> Cu,<sup>11</sup> Fe,<sup>12</sup> PIDA,<sup>13</sup> I<sub>2</sub>,<sup>14</sup> visible light,<sup>15</sup> photoredox/metal,<sup>16</sup> and electricity<sup>17</sup>) have been widely investigated to enlarge the substrate scope. Using simple and easily available substrates, domino one-pot processes combining the *in situ* formation of *N*-arylated enamines or imines with subsequent cyclization have also been developed for the synthesis of indoles. For instance, Jiao group<sup>18</sup> pioneered studies to construct an indole backbone through an efficient Pd-catalyzed aerobic oxidative C–H functionalization approach from simple anilines and activated




Scheme 1 Metal-catalyzed indole syntheses based on the *N*-arylated enamines and imines.

<sup>a</sup>College of Chemistry, Chemical Engineering and Resource Utilization, Northeast Forestry University, No. 26 Hexing Road, Harbin, 150040, P. R. China

<sup>b</sup>Material Science and Engineering College, Northeast Forestry University, No. 26 Hexing Road, Harbin, 150040, P. R. China. E-mail: ccx1759@163.com; jspeng1998@nefu.edu.cn

† Electronic supplementary information (ESI) available: Copies of <sup>1</sup>H, <sup>13</sup>C NMR, HRMS and IR spectra for new compounds. See DOI: 10.1039/d0ra04592f





Scheme 2 Prospected sequence for the synthesis of indoles.

alkynes. Recently, Zhang and Cao<sup>19</sup> reported a one-pot synthesis of 2-(perfluoroalkyl)indoles through sequential Michael-type addition and Pd(II)-catalyzed cross-dehydrogenative coupling

reaction of anilines and methyl perfluoroalk-2-ynoates with oxygen as the sole oxidant. Yoshikai group<sup>20</sup> developed a Pd-catalyzed method for the synthesis of indoles from *N*-aryl imines directly derived from simple and obtainable anilines and ketones. In addition, using simple anilines<sup>21</sup> or aniline derivatives with directing groups,<sup>22–27</sup> various transitional metal-catalyzed (such as Rh,<sup>22</sup> Ru,<sup>23</sup> Pd,<sup>24</sup> Au,<sup>25</sup> Ni,<sup>26</sup> and Co<sup>27</sup>) protocols have been well established through group-directed oxidative C–H annulation of alkynes.

We have been devoting our efforts to develop metal-catalyzed sequential one-pot processes containing a direct C–H functionalization step for the construction of heterocyclic frameworks.<sup>28</sup> Over the past decades, oxidative cross-dehydrogenative coupling (CDC) reactions have emerged as one of the most powerful routes for C–C bond formations in organic synthesis.<sup>29</sup>

Table 1 Condition optimization for the copper-catalyzed synthesis of indole 3aa<sup>a</sup>

Entry	Cu/ligand	Base	Solvent	Oxidant	Yield <sup>b</sup> [%]
1	CuI/L1	KHCO <sub>3</sub>	DMF	Air	10/6 <sup>c</sup> /7 <sup>d</sup>
2	CuBr/L1	KHCO <sub>3</sub>	DMF	Air	Trace
3	CuCl/L1	KHCO <sub>3</sub>	DMF	Air	Trace
4	Cu <sub>2</sub> O/L1	KHCO <sub>3</sub>	DMF	Air	Trace
5	Cu(OTf) <sub>2</sub> /L1	KHCO <sub>3</sub>	DMF	Air	5
6	CuCl <sub>2</sub> /L1	KHCO <sub>3</sub>	DMF	Air	8
7	CuBr <sub>2</sub> /L1	KHCO <sub>3</sub>	DMF	Air	7
8	CuSO <sub>4</sub> ·5H <sub>2</sub> O/L1	KHCO <sub>3</sub>	DMF	Air	10
9	Cu(OAc) <sub>2</sub> /L1	KHCO <sub>3</sub>	DMF	Air	20
10	Cu(acac) <sub>2</sub> /L1	KHCO <sub>3</sub>	DMF	Air	14
11	CuO/L1	KHCO <sub>3</sub>	DMF	Air	0
12 <sup>e</sup>	Cu(OAc) <sub>2</sub> /L2-L17	KHCO <sub>3</sub>	DMF	Air	0–25
13	Cu(OAc) <sub>2</sub>	KHCO <sub>3</sub>	DMF	Air	0
14 <sup>f</sup>	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMF	—	0
15	Cu(OAc) <sub>2</sub> /L12	K <sub>2</sub> CO <sub>3</sub>	DMF	Air	15
16	Cu(OAc) <sub>2</sub> /L12	NaOH	DMF	Air	0
17	Cu(OAc) <sub>2</sub> /L12	Li <sub>2</sub> CO <sub>3</sub>	DMF	Air	0
18	Cu(OAc) <sub>2</sub> /L12	K <sub>3</sub> PO <sub>4</sub>	DMF	Air	20
19	Cu(OAc) <sub>2</sub> /L12	NaHCO <sub>3</sub>	DMF	Air	17
20	Cu(OAc) <sub>2</sub> /L12	NaOAc	DMF	Air	18
21	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMSO	Air	10
22	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	THF	Air	15
23	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	1,4-Dioxane	Air	Trace
24	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMA	Air	19
25	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	<i>tert</i> -Pentyl alcohol	Air	20
26	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMF/DMSO	Air	33 (37) <sup>g</sup>
27 <sup>g</sup>	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMF/DMSO	KMnO <sub>4</sub>	48
28 <sup>g</sup>	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMF/DMSO	K <sub>2</sub> S <sub>2</sub> O <sub>8</sub>	21
29 <sup>g</sup>	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMF/DMSO	TBHP	20
30 <sup>g</sup>	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMF/DMSO	MnO <sub>2</sub>	30
31 <sup>g</sup>	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMF/DMSO	KMnO <sub>4</sub>	55 <sup>h</sup>
32 <sup>g</sup>	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMF/DMSO	KMnO <sub>4</sub>	47 <sup>i</sup>
33 <sup>g</sup>	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMF/DMSO	KMnO <sub>4</sub>	48 <sup>j</sup>
34 <sup>g</sup>	Cu(OAc) <sub>2</sub> /L12	KHCO <sub>3</sub>	DMF/DMSO	KMnO <sub>4</sub>	45 <sup>k</sup>

<sup>a</sup> Reaction conditions unless otherwise stated: **1a** (0.5 mmol), **2a** (0.2 mmol), base (0.6 mmol), catalyst (20 mol%), ligand (30 mol%), solvent (1.0 mL), air or oxidant, 70 °C, 24 h, air. <sup>b</sup> Isolated yield. <sup>c</sup> 1.5 equiv. of **1a**. <sup>d</sup> 2.0 equiv. of **1a**. <sup>e</sup> The effect of various ligands was investigated, see Scheme 1. <sup>f</sup> N<sub>2</sub>. <sup>g</sup> DMF (1.0 mL), 100 °C, 24 h; then DMSO (0.5 mL), 130 °C, 24 h. <sup>h</sup> 20 mol% of myristic acid was added. <sup>i</sup> 20 mol% of palmitic acid was added. <sup>j</sup> 20 mol% of stearic acid was added. <sup>k</sup> 20 mol% of trimethylacetic acid was added.









copper-catalyzed coupling reaction rate of arylboronic acids and amines through the coordination of aliphatic acid to the copper center to improve solubility of the copper catalyst in organic solvents.<sup>34</sup> Given the importance of aliphatic acids in achieving a homogeneous catalyst system, various carboxylic acids such as myristic acid, palmitic acid, stearic acid, and trimethylacetic acid were explored as the additive. It was found that the addition of 1 equiv of myristic acid relative to Cu(OAc)<sub>2</sub> improved the yield of **3aa** to 55% (entry 31), while other carboxylic acids had little effect on the reaction (entries 32–34). In general, in the presence of combinations of Cu(OAc)<sub>2</sub>, <sup>t</sup>Bu<sub>3</sub>P·HBF<sub>4</sub>, KHCO<sub>3</sub>, myristic acid and KMnO<sub>4</sub>, the synthesis of indoles **3** was conducted in a one-pot fashion using DMF/DMSO as mixed solvents.

Having established the feasibility of indole synthesis *via* copper(II)-catalyzed sequential oxidative Chan–Lam arylation/CDC process, we then explored the generality of arylboronic acids using methyl or ethyl (*Z*)-3-amino-3-phenylacrylate as the coupling partner (Table 2). As shown in Table 2, diverse substituents (such as Me, OMe, F, Cl, ester, and methylthio) on the aromatic moiety of boronic acids were applicable, and the corresponding indole products **3aa–3ja** can be obtained in 44–55% yields. Arylboronic acids containing electron-donating groups at the *para* (**1b**, **1c**, and **1g**), or *meta* (**1j**) position were generally more reactive than those bearing electron-withdrawing substituents (**1d–1f**) and provided higher yields (entries 2–7, and 10, Table 2). However, the incorporation of substituents in the *ortho* position of arylboronic acid seriously hampered copper-catalyzed oxidative annulation process, and the corresponding indoles cannot be obtained. In addition,  $\alpha$ - or  $\beta$ -naphthyl boronic acids **1h** and **1i** can also be smoothly transformed into the corresponding products **3ha** and **3ia** in 49% and 50% yields, respectively (entries 8 and 9). For  $\beta$ -naphthyl boronic acid **1i**, there are two possible C–H activation sites ( $\alpha$ - and  $\beta$ -position), the CDC process occurred only at the  $\alpha$ -site to give a single regioisomer **3ia** (entry 9). The specificity of site selectivity ( $\alpha/\beta$ ) implies that the aromatic C–H alkenylation is probably an outcome of electrophilic aromatic substitutions. However, regioselectivity issues surfaced for *meta*-substituted arylboronic acid (**1j**), and a mixture of two regioisomers (**3ja** and **3j'a**) was obtained in a nearly 2 : 1 ratio, indicating intramolecular CDC reaction occurred at the most sterically accessible site (entry 10).

The scope and limitation of ester (*Z*)-3-aminoacrylate substrates were finally investigated in this one-pot sequential oxidative process. As shown in Table 3, several functional groups (such as Me, OMe, OEt, <sup>t</sup>Bu, F, Cl, Br and CF<sub>3</sub>) are tolerated under the standard conditions. For ester (*Z*)-3-amino-3-arylpropiolates, the electronic nature of the aromatic motifs did not seem to affect the efficiency: both electron-donating (Me, MeO, EtO, and <sup>t</sup>Bu) and electron-withdrawing substituents (F, Cl, Br, and CF<sub>3</sub>) can be incorporated at the *para* (**2b–2j**, entries 1–9) and *meta* (**2k–2o**, entries 10–14) position, providing indole derivatives **3ab–3ao** in 42–53% yields. However, ester (*Z*)-3-amino-3-arylpropiolates (**2p–2r**, Fig. 1) with substituents in the *ortho* position of aromatic ring cannot proceed to give the corresponding products, probably because of the strong steric





Fig. 1 Unsuccessful enamine substrates.

repulsion of the *ortho* substituent. In addition, when the scope of substrates was extended to aliphatic and carbocyclic enamines [for example, ethyl (*Z*)-3-aminobut-2-enoate **2s** and 3-aminocyclohex-2-en-1-one **2u**, Fig. 1], the reaction did not happen to give the corresponding products. Enamine substrates with an amide (**2t**), aromatic ketone (**2v**) or nitro group (**2w**) were also examined under the standard conditions; however, no product can be obtained. These unsuccessful results indicated that the electron density of olefin moiety was highly essential for CDC process.

To gain insight into the mechanism of the reaction, some designed control experiments were conducted. When the reaction of **1a** and **2a** was conducted in DMF for 24 h, the Chan–Lam *N*-arylated intermediate **9** and **3aa** can be isolated in 68% and 10% yields, respectively. Then, the intermediate **9** was carried out under the optimized reaction conditions, 65% of the desired product **3aa** was obtained. Based on the above results and previous literature reports from our group<sup>30,31</sup> and others,<sup>35</sup> a reaction mechanism for the copper(II)-catalyzed oxidative annulation of enamines with arylboronic acids was proposed. The Scheme 4 showed a simplified sequence of events beginning with the Cu(OAc)<sub>2</sub>. First, a soluble active cupric tetradeanoate species was formed through the anion exchange reaction of Cu(OAc)<sub>2</sub> and myristic acid in the presence of base.<sup>34</sup> The coordination of (*Z*)-enamine **2** to the Cu(II) center to form **4**, which then reacted with a base to form a Cu–N bond and afforded an intermediate complex **5**. Engagement of arylboronic acid **1** led to transmetalation *via* 4-membered transition state<sup>35a</sup> and delivered aryl–Cu(II) species **6**. The intermediate **6** was then oxidized by Cu(II) to form an Cu(III) species **8**, and the subsequent reductive elimination gave *N*-arylated intermediate **9** and a Cu(I)OAc species **7**. Completion of the catalytic cycle was achieved *via* oxidation to Cu(II) in the presence of KMnO<sub>4</sub> (Chan–Lam C–N coupling process). The coordination of *N*-arylated intermediate **9** to Cu(II) gave a six-membered chelate ring complex **10**. A new C–Cu(II) complex **12** was then formed through sequential a base-promoted deprotonation of N–H, dissociation of acetate anion and complexation of the resulting cationic Cu(II) species at  $\alpha$ -carbon of **11**. A deprotonation/reprotonation process of alkyl–Cu(II) complex **12** afforded an alkenyl copper **13** under basic conditions. *Ortho*-cupration of





Scheme 4 Proposed catalytic pathway for the formation of indole 3.

phenyl ring of **13** provided a six-membered copper-cycle intermediate **14**,<sup>9</sup> which was then transformed into the product **3** and Cu(0) species **15** through reductive elimination process. Finally, Cu(0) was oxidized to the active Cu(II) catalyst by KMnO<sub>4</sub> for the next Chan-Lam arylation reaction.

### 3. Conclusions

In summary, we have developed a new one-pot approach to diverse multi-substituted indoles through copper-catalyzed oxidative annulations of enamines with readily accessible arylboronic acids. The accomplished reaction comprises an intermolecular Chan-Lam arylation followed by an intramolecular cross-dehydrogenative coupling reaction promoted by the same copper catalyst. The success of the reaction heavily relies on the careful selection of proper additive and oxidant. The combination of myristic acid and KMnO<sub>4</sub> was found to be essential for the formation of C3-functionalized multi-substituted indoles. Considering a broad substrate scope and considerable valance of the products for medicinal science, this novel synthetic method could be of utility for the discovery of drugs.

## 4. Experimental section

### 4.1. General information

Chemicals were all purchased from commercial supplies and used without further purification unless otherwise stated. Solvents were dried and purified according to the standard procedures before use. Reactions were monitored by analytical

thin-layer chromatography (TLC). All reactions were conducted in dried glassware. Purification of reaction products was done by flash chromatography with 230–400 mesh silica gel. Ester (Z)-3-aryl-3-aminoacrylate substrates were prepared according to the literature methods.<sup>36</sup> Melting points were determined on a melting point apparatus in open capillaries and are uncorrected. Infrared spectra of samples were recorded from 4000 to 500 cm<sup>-1</sup> in ATR (attenuated total reflectance) mode using an FT-IR instrument. <sup>1</sup>H NMR spectra were recorded on a 400 or 500 MHz spectrometer, and <sup>13</sup>C NMR spectra were recorded at 100, 125 or 150 MHz. Unless otherwise stated, deuteriochloroform (CDCl<sub>3</sub>) was used as a solvent. Chemical shifts (δ) are given in parts per million downfield relative to tetramethylsilane (TMS). Chemical shifts for carbon resonances are reported in parts per million and are referenced to the carbon resonance of the solvent CHCl<sub>3</sub> (δ = 77.16 ppm). The splitting patterns are reported as s (singlet), d (doublet), dd (double doublet), td (triplet of doublet), t (triplet), q (quartet), br (broad), and m (multiplet). Coupling constants are given in hertz. High-resolution mass spectra were recorded on a BIO TOF Q mass spectrometer equipped with an electrospray ion source (ESI), operated in the positive mode.

### 4.2. General procedure for synthesis of indole-3-carboxylate derivatives

A 10 mL Schlenk tube or standard vial equipped with a magnetic stirring bar was charged with ester (Z)-3-aryl-3-aminoacrylates (0.2 mmol, 1.0 equiv.), aryl boronic acid (0.5 mmol, 2.5 equiv.), KMnO<sub>4</sub> (0.1 mmol, 15.8 mg), and KHCO<sub>3</sub> (60 mg,



0.6 mmol, 3.0 equiv.), and then  $\text{Cu}(\text{OAc})_2$  (0.04 mmol, 8.0 mg),  $\text{tBu}_3\text{P} \cdot \text{HBF}_4$  (0.06 mmol, 17.4 mg) and myristic acid (0.04 mmol, 9.1 mg) were added. Finally, DMF (1.0 mL) was added to the mixture *via* syringe at room temperature under air. The vial was sealed and put into a preheated oil bath at 100 °C for 24 h. The mixture was cooled to room temperature, dimethyl sulfoxide (0.5 mL) was then added *via* syringe, and the reaction mixture was heated to 130 °C for another 24 h. Finally, the mixture was cooled to room temperature, quenched with water (3 mL), and diluted with ethyl acetate (5 mL). The layers were separated, and the aqueous layer was extracted with  $3 \times 5$  mL of ethyl acetate. The combined organic extracts were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. The crude product was then purified by a chromatography silica gel (H), eluting with ethyl acetate/petroleum ether (10–15%).

**Ethyl 2-phenyl-1H-indole-3-carboxylate (3aa).**<sup>37</sup> Yield, 55% (29.2 mg); white solid, mp 155–158 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3254, 1662, 1450, 1430, 1270, 1212, 1129, 1047, 744, 690;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.76 (br, 1H), 8.22–8.20 (d,  $J = 7.2$  Hz, 1H), 7.61–7.59 (m, 2H), 7.37–7.35 (m, 3H), 7.33–7.31 (m, 1H), 7.27–7.21 (m, 2H), 4.26 (q,  $J = 7.1$  Hz, 2H), 1.28 (t,  $J = 7.1$  Hz, 3H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  164.5, 143.6, 134.2, 131.0, 128.6, 128.1, 127.0, 126.5, 122.1, 121.0, 120.9, 110.1, 103.4, 58.7, 13.3.

**Methyl 5-chloro-2-(4-chlorophenyl)-1H-indole-3-carboxylate (3ab).** Yield, 47% (30.0 mg); light yellow solid, mp 163–165 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3249, 1677, 1666, 1485, 1444, 1296, 1210, 1135, 1091, 786, 800, 823;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.59 (s, 1H), 8.17–8.16 (d,  $J = 1.6$  Hz, 1H), 7.58–7.56 (d,  $J = 8.5$  Hz, 2H), 7.42–7.40 (d,  $J = 8.5$  Hz, 2H), 7.30–7.28 (m, 1H), 7.25–7.22 (m, 1H), 3.85 (s, 3H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  164.3, 143.3, 134.7, 132.5, 129.8, 128.7, 127.5, 127.4, 127.1, 122.9, 120.8, 111.1, 103.5, 50.2. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{16}\text{H}_{12}\text{Cl}_2\text{NO}_2^+$  ( $\text{M} + \text{H}$ )<sup>+</sup> 320.02396, found 320.02390.

**Ethyl 5-chloro-2-(4-fluorophenyl)-1H-indole-3-carboxylate (3ac).** Yield, 42% (26.7 mg); light yellow solid, mp 149–151 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3416, 1673, 1497, 1447, 1212, 1130, 840, 804, 786;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.58 (s, 1H), 8.19–8.18 (d,  $J = 1.2$  Hz, 1H), 7.64–7.60 (dd,  $J = 8.5, 5.4$  Hz, 2H), 7.30–7.27 (m, 1H), 7.24–7.21 (m, 1H), 7.13 (t,  $J = 8.6$  Hz, 2H), 4.32 (q,  $J = 7.1$  Hz, 2H), 1.33 (t,  $J = 7.1$  Hz, 3H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  164.9, 163.4 (164.2, 162.6, d,  $^1J_{\text{C-F}} = 249$  Hz), 144.5, 133.4, 131.5 (131.6, 131.5, d,  $^3J_{\text{C-F}} = 8$  Hz), 128.6, 128.0, 127.6 (127.6, 127.5, d,  $^4J_{\text{C-F}} = 3$  Hz), 123.8, 121.8, 115.3 (115.4, 115.3, d,  $^2J_{\text{C-F}} = 21$  Hz), 112.1, 104.6, 60.0, 14.4. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{17}\text{H}_{14}\text{ClFNO}_2^+$  ( $\text{M} + \text{H}$ )<sup>+</sup> 318.06916, found 318.06912.

**Methyl 2-(4-bromophenyl)-5-methyl-1H-indole-3-carboxylate (3ad).** Yield, 46% (31.7 mg); orange solid, mp 160–162 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3307, 1672, 1444, 1124, 830, 790;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.44 (s, 1H), 7.98 (s, 1H), 7.57–7.49 (q,  $J = 7.6$  Hz, 4H), 7.27 (d,  $J = 8.1$  Hz, 1H), 7.11 (d,  $J = 8.1$  Hz, 1H), 3.84 (s, 3H), 2.50 (s, 3H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  165.8, 143.1, 133.5, 131.9, 131.4, 131.1, 131.0, 127.7, 125.2, 123.6, 121.8, 110.7, 104.4, 51.0, 21.7. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{17}\text{H}_{15}\text{BrNO}_2^+$  ( $\text{M} + \text{H}$ )<sup>+</sup> 344.02807, found 344.02802.

**Ethyl 5-methyl-2-(*p*-tolyl)-1H-indole-3-carboxylate (3ae).** Yield, 49% (27.4 mg); white solid, mp 157–159 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3280, 1656, 1440, 1267, 1217, 1158, 1143, 1048, 800,

790;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.41 (s, 1H), 8.04 (s, 1H), 7.57 (d,  $J = 8.0$  Hz, 2H), 7.29–7.27 (m, 3H), 7.11 (d,  $J = 8.2$  Hz, 1H), 4.34 (q,  $J = 7.1$  Hz, 2H), 2.53 (s, 3H), 2.43 (s, 3H), 1.35 (t,  $J = 7.1$  Hz, 3H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  165.5, 144.7, 139.2, 133.4, 131.5, 129.5, 129.2, 128.8, 128.0, 124.6, 121.8, 110.6, 104.1, 59.6, 21.8, 21.4, 14.4. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{19}\text{H}_{20}\text{NO}_2^+$  ( $\text{M} + \text{H}$ )<sup>+</sup> 294.14886, found 294.14856.

**Methyl 2-(4-methoxyphenyl)-5-methyl-1H-indole-3-carboxylate (3af).** Yield, 53% (31.3 mg); white solid, mp 156–158 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3247, 1668, 1643, 1501, 1454, 1282, 1134, 832, 801, 792;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.47 (s, 1H), 7.97 (s, 1H), 7.56 (d,  $J = 8.6$  Hz, 2H), 7.23–7.21 (m, 1H), 7.06 (d,  $J = 8.8$  Hz, 1H), 6.92 (d,  $J = 8.6$  Hz, 2H), 3.83 (s, 3H), 3.81 (s, 3H), 2.49 (s, 3H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  165.1, 159.2, 143.8, 132.3, 130.4, 129.79, 126.8, 123.5, 123.2, 120.6, 112.6, 109.6, 102.4, 54.3, 49.8, 20.7. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{18}\text{H}_{18}\text{NO}_3^+$  ( $\text{M} + \text{H}$ )<sup>+</sup> 296.12812, found 296.12836.

**Methyl 5-methoxy-2-(4-methoxyphenyl)-1H-indole-3-carboxylate (3ag).** Yield, 51% (31.8 mg); yellow solid, mp 158–159 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3226, 1664, 1499, 1488, 1454, 1252, 1209, 1172, 1137, 842, 825, 811, 788;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.38 (s, 1H), 7.70 (d,  $J = 2$  Hz, 1H), 7.58 (d,  $J = 8.6$  Hz, 2H), 7.24 (s, 1H), 6.97 (d,  $J = 8.6$  Hz, 2H), 6.90 (dd,  $J = 8.6, 2.2$  Hz, 1H), 3.91 (s, 3H), 3.85 (s, 3H), 3.83 (s, 3H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  165.0, 159.3, 154.7, 144.0, 129.8, 129.0, 127.5, 123.3, 112.6, 112.1, 110.6, 102.7, 54.8, 54.3, 49.8. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{18}\text{H}_{18}\text{NO}_4^+$  ( $\text{M} + \text{H}$ )<sup>+</sup> 312.12303, found 312.12314.

**Methyl 2-[4-(*tert*-butyl) phenyl]-5-methyl-1H-indole-3-carboxylate (3ah).** Yield, 52% (33.4 mg); white solid, mp 189–190 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3267, 1674, 1443, 1201, 1284, 1158, 1119, 861, 789, 780;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.39 (s, 1H), 8.00 (s, 1H), 7.61 (d,  $J = 8.5$  Hz, 2H), 7.48 (d,  $J = 8.5$  Hz, 2H), 7.28 (s, 1H), 7.09 (d,  $J = 8.3$  Hz, 2H), 3.86 (s, 3H), 2.51 (s, 3H), 1.36 (s, 9H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  165.0, 151.3, 143.8, 132.3, 130.5, 128.1, 128.0, 126.8, 124.2, 123.6, 120.7, 109.6, 102.7, 49.9, 33.8, 30.2, 20.7. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{21}\text{H}_{24}\text{NO}_2^+$  ( $\text{M} + \text{H}$ )<sup>+</sup> 322.18016, found 322.18030.

**Methyl 2-(4-ethoxyphenyl)-5-methyl-1H-indole-3-carboxylate (3ai).** Yield, 46% (28.4 mg); white solid, mp 195–197 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3257, 1652, 1612, 1457, 1450, 1260, 1250, 1181, 1142, 1048, 848, 799, 790;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  8.40 (s, 1H), 7.98 (s, 1H), 7.58 (d,  $J = 8.8$  Hz, 2H), 7.24 (s, 1H), 7.07 (d,  $J = 9.0$  Hz, 2H), 6.94 (d,  $J = 8.8$  Hz, 2H), 4.06 (q,  $J = 7.0$  Hz, 2H), 3.85 (s, 3H), 2.50 (s, 3H), 1.44 (t,  $J = 7.0$  Hz, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  165.0, 158.7, 143.7, 132.3, 130.5, 129.8, 126.9, 123.5, 123.1, 120.7, 113.1, 109.5, 102.4, 62.5, 49.8, 20.7, 13.8. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{19}\text{H}_{20}\text{NO}_3^+$  ( $\text{M} + \text{H}$ )<sup>+</sup> 310.14377, found 310.14368.

**Methyl 5-methyl-2-(4-(trifluoromethyl) phenyl)-1H-indole-3-carboxylate (3aj).** Yield, 49% (32.6 mg); white solid, mp 165–167 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3288, 1665, 1446, 1325, 1221, 1166, 1134, 1068, 849, 802, 693, 622;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  8.66–8.56 (br, 1H), 8.00 (s, 1H), 7.77–7.69 (m, 4H), 7.31–7.29 (m, 1H), 7.13 (d,  $J = 8.2$  Hz, 1H), 3.85 (s, 3H), 2.51 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  164.7, 141.5, 134.6, 132.6, 131.0, 129.9 (130.2, 123.0, 129.7, 129.5, q,  $^2J_{\text{C-F}} = 33$  Hz), 128.8, 126.4, 125.2 (128.7, 126.5, 124.0, 121.8, q,  $^1J_{\text{C-F}} = 276$  Hz), 124.4, 124.0 (124.07, 124.04,



124.01, 123.98, q,  $^3J_{C-F} = 4$  Hz), 120.8, 109.8, 103.8, 50.0, 20.7. HRMS-ESI ( $m/z$ ) calcd for  $C_{18}H_{15}F_3NO_2^+$  ( $M + H$ ) $^+$  334.10494, found 334.10495.

**Methyl 2-(3-fluorophenyl)-5-methyl-1H-indole-3-carboxylate (3ak).** Yield, 47% (26.6 mg); orange solid, mp 121–122 °C; IR (KBr,  $cm^{-1}$ ): 3267, 1674, 1443, 1201, 1158, 1119, 861, 789, 680;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.55 (s, 1H), 7.98 (s, 1H), 7.38–7.33 (m, 3H), 7.24 (s, 1H), 7.09 (d,  $J = 7.7$  Hz, 2H), 3.83 (s, 3H), 2.49 (s, 3H).  $^{13}C$  NMR (151 MHz,  $CDCl_3$ )  $\delta$  165.8, 162.3 (163.1, 161.5, d,  $^1J_{C-F} = 246$  Hz), 142.8, 134.1 (134.1, 134.0, d,  $^3J_{C-F} = 8$  Hz), 133.5, 131.8, 129.7 (129.7, 129.6, d,  $^3J_{C-F} = 8$  Hz), 127.6, 125.2 (125.23, 125.21, d,  $^4J_{C-F} = 3$  Hz), 125.1, 121.8, 116.7 (116.8, 116.6, d,  $^2J_{C-F} = 23$  Hz), 116.1 (116.1, 116.0, d,  $^2J_{C-F} = 21$  Hz), 110.8, 104.4, 51.0, 21.7. HRMS-ESI ( $m/z$ ) calcd for  $C_{17}H_{15}FNO_2^+$  ( $M + H$ ) $^+$  284.10813, found 284.10825.

**Methyl 2-(3-chlorophenyl)-5-methyl-1H-indole-3-carboxylate (3al).** Yield, 46% (27.5 mg); light yellow solid, mp 152–153 °C; IR (KBr,  $cm^{-1}$ ): 3266, 1673, 1478, 1443, 1139, 782;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.53 (br, 1H), 7.98 (s, 1H), 7.60 (s, 1H), 7.51 (d,  $J = 7.2$  Hz, 1H), 7.37–7.31 (m, 2H), 7.24 (d,  $J = 5.4$  Hz, 1H), 7.09 (d,  $J = 7.1$  Hz, 1H), 3.83 (s, 3H), 2.49 (s, 3H).  $^{13}C$  NMR (151 MHz,  $CDCl_3$ )  $\delta$  164.7, 141.6, 133.0, 132.7, 132.5, 130.8, 128.4, 128.3, 128.1, 126.9, 126.6, 124.1, 120.8, 109.7, 103.5, 50.0, 20.7. HRMS-ESI ( $m/z$ ) calcd for  $C_{17}H_{15}ClNO_2^+$  ( $M + H$ ) $^+$  300.07858, found 300.07837.

**Methyl 5-methyl-2-(*m*-tolyl)-1H-indole-3-carboxylate (3am).** Yield, 49% (27.4 mg); light yellow solid, mp 136–138 °C; IR (KBr,  $cm^{-1}$ ): 3289, 1668, 1447, 1161, 1124, 1049, 786, 731, 698;  $^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$  8.52 (br, 1H), 7.92 (s, 1H), 7.36–7.34 (m, 2H), 7.24–7.21 (m, 1H), 7.17–7.13 (m, 2H), 7.01–6.99 (d,  $J = 7.9$  Hz, 1H), 3.74 (s, 3H), 2.42 (s, 3H), 2.30 (s, 3H).  $^{13}C$  NMR (126 MHz,  $CDCl_3$ )  $\delta$  164.8, 143.7, 136.8, 132.4, 131.0, 130.5, 128.9, 128.8, 127.0, 126.8, 125.8, 123.7, 120.7, 109.5, 102.9, 49.8, 20.7, 20.4. HRMS-ESI ( $m/z$ ) calcd for  $C_{18}H_{18}NO_2^+$  ( $M + H$ ) $^+$  280.13321, found 280.13309.

**Methyl 5-methoxy-2-(*m*-tolyl)-1H-indole-3-carboxylate (3an).** Yield, 49% (29.0 mg); light yellow solid, mp 138–140 °C; IR (KBr,  $cm^{-1}$ ): 3395, 1667, 1451, 1213, 1195, 1164, 1124, 1050, 1029, 796;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.71 (s, 1H), 7.68 (s, 1H), 7.39–7.36 (m, 2H), 7.26–7.24 (m, 1H), 7.20–7.16 (m, 2H), 6.87 (d,  $J = 9.0$  Hz, 1H), 3.87 (s, 3H), 3.77 (s, 3H), 2.32 (s, 3H).  $^{13}C$  NMR (151 MHz,  $CDCl_3$ )  $\delta$  165.1, 154.6, 144.2, 136.7, 130.9, 129.2, 128.8, 128.7, 127.4, 126.9, 125.7, 112.2, 110.9, 102.8, 102.5, 54.7, 49.8, 20.3. HRMS-ESI ( $m/z$ ) calcd for  $C_{18}H_{18}NO_3^+$  ( $M + H$ ) $^+$  296.12812, found 296.12802.

**Methyl 2-(3-methoxyphenyl)-5-methyl-1H-indole-3-carboxylate (3ao).** Yield, 48% (28.3 mg); yellow solid, mp 130–132 °C; IR (KBr,  $cm^{-1}$ ): 3354, 1686, 1466, 1440, 1286, 1123, 878, 804, 787;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.81 (br, 1H), 8.01 (s, 1H), 7.28–7.26 (m, 1H), 7.22–7.17 (m, 3H), 7.08 (d,  $J = 8.2$  Hz, 1H), 6.91 (d,  $J = 7.5$  Hz, 1H), 3.82 (s, 3H), 3.78 (s, 3H), 2.51 (s, 3H).  $^{13}C$  NMR (151 MHz,  $CDCl_3$ )  $\delta$  165.0, 158.0, 143.3, 132.4, 132.2, 130.4, 128.5, 128.0, 126.7, 123.7, 120.8, 120.5, 114.1, 113.6, 109.8, 102.7, 54.2, 49.8. HRMS-ESI ( $m/z$ ) calcd for  $C_{18}H_{18}NO_3^+$  ( $M + H$ ) $^+$  296.12812, found 296.12848.

**Methyl 2-phenyl-1H-indole-3-carboxylate (3ap).**<sup>8b,30</sup> Yield, 51% (25.6 mg); yellow solid, mp 137–139 °C; IR (KBr,  $cm^{-1}$ ):

3300, 1667, 1486, 1447, 1421, 1282, 1214, 1132, 792, 765, 740, 697;  $^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$  8.47 (s, 1H), 8.23–8.21 (m, 1H), 7.69–7.67 (m, 2H), 7.48–7.47 (m, 3H), 7.42–7.40 (m, 1H), 7.30–7.28 (m, 2H), 3.85 (s, 3H).  $^{13}C$  NMR (126 MHz,  $CDCl_3$ )  $\delta$  164.7, 143.5, 134.0, 130.9, 128.5, 128.3, 127.2, 126.5, 122.3, 121.2, 121.1, 109.9, 103.6, 49.9.

**Methyl 5-methyl-2-phenyl-1H-indole-3-carboxylate (3ba).**<sup>13,38</sup> Yield, 54% (28.6 mg); white solid, mp 154–156 °C; IR (KBr,  $cm^{-1}$ ): 3287, 2949, 1673, 1452, 1129, 800, 759, 696;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.60 (br, 1H), 7.99 (s, 1H), 7.62–7.60 (m, 2H), 7.40 (br, 3H), 7.23 (d,  $J = 8.1$  Hz, 1H), 7.07 (d,  $J = 8.1$  Hz, 1H), 3.81 (s, 3H), 2.49 (s, 3H).  $^{13}C$  NMR (151 MHz,  $CDCl_3$ )  $\delta$  164.9, 143.6, 132.4, 131.1, 130.5, 128.5, 128.5, 128.0, 127.1, 126.7, 123.7, 120.7, 109.7, 49.8, 20.7.

**Methyl 5-methoxy-2-phenyl-1H-indole-3-carboxylate (3ca).**<sup>38</sup> Yield, 49% (27.5 mg); white solid, mp 157–159 °C; IR (KBr,  $cm^{-1}$ ): 3294, 2950, 1678, 1485, 1462, 1268, 1206, 1167, 1129, 1047, 1035, 804, 696;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.70 (br, 1H), 7.71 (d,  $J = 3$  Hz, 1H), 7.61–7.59 (m, 2H), 7.41–7.39 (m, 3H), 7.23 (d,  $J = 8.8$  Hz, 1H), 6.90 (dd,  $J = 8.8, 2.5$  Hz, 1H), 3.90 (s, 3H), 3.79 (s, 3H).  $^{13}C$  NMR (151 MHz,  $CDCl_3$ )  $\delta$  165.0, 154.7, 143.9, 131.0, 129.2, 128.4, 128.0, 127.4, 127.1, 112.4, 110.9, 103.0, 102.6, 54.8, 49.8.

**Methyl 5-fluoro-2-phenyl-1H-indole-3-carboxylate (3da).** Yield, 47% (25.3 mg); white solid, mp 156–158 °C; IR (KBr,  $cm^{-1}$ ): 3282, 1667, 1487, 1464, 1454, 1270, 1213, 1138, 1047, 859, 701, 629;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.58 (br, 1H), 7.86 (dd,  $J = 10.0, 2.4$  Hz, 1H), 7.64–7.63 (m, 2H), 7.46–7.44 (m, 3H), 7.31–7.27 (m, 1H), 7.02 (td,  $J = 9, 2.4$  Hz, 1H), 3.83 (s, 3H).  $^{13}C$  NMR (151 MHz,  $CDCl_3$ )  $\delta$  165.5, 159.3 (160.0, 158.5, d,  $^1J_{C-F} = 236$  Hz), 146.1, 131.6 (131.59, 131.55, d,  $^3J_{C-F} = 6$  Hz), 129.5, 129.4, 128.4, 128.3, 111.8, 111.7, 111.6, 107.5 (107.6, 107.5, d,  $^2J_{C-F} = 26$  Hz), 104.7 (104.72, 104.69, d,  $^4J_{C-F} = 4$  Hz), 51.0. HRMS-ESI ( $m/z$ ) calcd for  $C_{16}H_{13}FNO_2^+$  ( $M + H$ ) $^+$  270.09248, found 270.09256.

**Ethyl 5-chloro-2-(*p*-tolyl)-1H-indole-3-carboxylate (3ea).** Yield, 46% (28.8 mg); white solid, mp 190–192 °C; IR (KBr,  $cm^{-1}$ ): 3429, 3270, 1671, 1431, 1210, 1128, 824, 786;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.60 (br, 1H), 8.17 (d,  $J = 1.6$  Hz, 1H), 7.50 (d,  $J = 8.0$  Hz, 2H), 7.35–7.05 (m, 5H), 4.30 (q,  $J = 7.1$  Hz, 2H), 2.37 (s, 3H), 1.33 (t,  $J = 7.1$  Hz, 3H).  $^{13}C$  NMR (151 MHz,  $CDCl_3$ )  $\delta$  165.1, 146.0, 139.6, 133.4, 129.4, 128.9, 128.8, 128.5, 127.7, 123.4, 121.7, 112.0, 104.1, 59.9, 21.4, 14.4. HRMS-ESI ( $m/z$ ) calcd for  $C_{18}H_{17}ClNO_2^+$  ( $M + H$ ) $^+$  314.09423, found 314.09406.

**Dimethyl 2-phenyl-1H-indole-3,5-dicarboxylate (3fa).** Yield, 44% (27.2 mg); white solid, mp 202–204 °C; IR (KBr,  $cm^{-1}$ ): 3328, 1693, 1674, 1452, 1437, 1288, 1105, 762, 696;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.93 (s, 1H), 8.68 (s, 1H), 8.00 (dd,  $J = 8.6, 1.3$  Hz, 1H), 7.69–7.67 (m, 2H), 7.48–7.41 (m, 4H), 3.96 (s, 3H), 3.88 (s, 3H).  $^{13}C$  NMR (151 MHz,  $CDCl_3$ )  $\delta$  166.9, 164.2, 144.8, 136.6, 130.3, 128.6, 128.5, 127.3, 126.1, 124.0, 123.7, 123.1, 109.8, 104.5, 51.0, 50.2. HRMS-ESI ( $m/z$ ) calcd for  $C_{18}H_{16}NO_4^+$  ( $M + H$ ) $^+$  310.10738, found 310.10748.

**Methyl 5-(methylthio)-2-phenyl-1H-indole-3-carboxylate (3ga).** Yield, 50% (29.7 mg); white solid, mp 131–133 °C; IR (KBr,  $cm^{-1}$ ): 3286, 1665, 1439, 1215, 1131, 1077, 766, 697;  $^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$  8.59 (s, 1H), 8.18 (s, 1H), 7.64–7.63 (m,





2H), 7.45–7.44 (m, 3H), 7.31–7.25 (m, 2H), 3.83 (s, 3H), 2.57 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  164.6, 144.0, 132.6, 130.7, 130.1, 128.4, 128.3, 127.3, 127.2, 123.4, 120.5, 110.5, 102.9, 50.0, 17.0. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{17}\text{H}_{16}\text{NO}_2\text{S}^+$  ( $\text{M} + \text{H}$ ) $^+$  298.08963, found 298.08957.

**Methyl 2-phenyl-1H-benzo[g]indole-3-carboxylate (3ha).** Yield, 49% (29.5 mg); white solid, mp 185–187 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3219, 1668, 1472, 1436, 1199, 1128, 811, 746, 684;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.16 (s, 1H), 8.31 (d,  $J = 8.8$  Hz, 1H), 8.02 (d,  $J = 8.1$  Hz, 1H), 7.97 (d,  $J = 8.1$  Hz, 1H), 7.73–7.70 (m, 3H), 7.56 (t,  $J = 7.6$  Hz, 1H), 7.51–7.45 (m, 4H), 3.88 (s, 3H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  164.8, 141.3, 131.1, 129.7, 129.0, 128.6, 128.0, 127.9, 127.2, 124.9, 123.6, 123.0, 121.8, 120.3, 120.0, 118.3, 105.2, 50.0. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{20}\text{H}_{16}\text{NO}_2^+$  ( $\text{M} + \text{H}$ ) $^+$  302.11756, found 302.11707.

**Methyl 2-phenyl-3H-benzo[e]indole-1-carboxylate (3ia).** Yield, 50% (30.0 mg); white solid, mp 192–193 °C; IR (KBr,  $\text{cm}^{-1}$ ): 3310, 1666, 1460, 1442, 1195, 1145, 806, 698;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.91 (d,  $J = 8.5$  Hz, 1H), 8.71 (s, 1H), 7.91 (d,  $J = 7.9$  Hz, 1H), 7.67 (d,  $J = 8.7$  Hz, 1H), 7.60–7.54 (m, 3H), 7.49–7.42 (m, 5H), 3.82 (s, 3H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  166.9, 138.7, 131.5, 131.4, 129.4, 127.8, 127.6, 127.5, 127.4, 127.0, 125.1, 124.3, 123.8, 123.0, 120.0, 111.1, 107.3, 50.6. HRMS-ESI ( $m/z$ ) calcd for  $\text{C}_{20}\text{H}_{16}\text{NO}_2^+$  ( $\text{M} + \text{H}$ ) $^+$  302.11756, found 302.11691.

**Mixture of methyl 6-methyl-2-phenyl-1H-indole-3-carboxylate (3ja) and methyl 4-methyl-2-phenyl-1H-indole-3-carboxylate (3ja').**<sup>30,39</sup> Yield, 52% (27.6 mg); yellow solid, the ratio (3ja : 3ja' = 2 : 1) is determined by NMR; IR (KBr,  $\text{cm}^{-1}$ ): 3297, 2949, 1682, 1452, 1214, 1127, 1048, 813, 768, 696;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.50 (br, 1.5H<sub>overlap</sub>), 8.07 (d,  $J = 8.2$  Hz, 1H<sub>3ja</sub>), 7.65–7.63 (m, 2H<sub>3ja</sub>), 7.55–7.51 (m, 1H<sub>3ja'</sub>), 7.45–7.40 (m, 4.50H<sub>overlap</sub>), 7.22 (d,  $J = 8.0$  Hz, 0.55H<sub>3ja'</sub>), 7.17–7.10 (m, 2.54H<sub>overlap</sub>), 7.01 (d,  $J = 7.1$  Hz, 0.53H<sub>3ja'</sub>), 3.83 (s, 3H<sub>3ja</sub>), 3.76 (s, 1.56H<sub>3ja'</sub>), 2.65 (s, 1.50H<sub>3ja'</sub>), 2.47 (s, 3H<sub>3ja</sub>).

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We are grateful for financial support from Special Project for Double First-Class-Cultivation of Innovative Talents (201815), the Fundamental Research Funds for the Central Universities (2572017DB07), and Natural Science Foundation of Heilongjiang Province (B2017002, LC2018003).

## Notes and references

- 1 A. J. Kochanowska-Karamyan and M. T. Hamann, *Chem. Rev.*, 2010, **110**, 4489.
- 2 C.-L. Do-Thanh, J. J. Vargas, J. W. Thomas, G. R. Armel and M. D. Best, *J. Agric. Food Chem.*, 2016, **64**, 3533.
- 3 (a) E. Vitaku, D. T. Smith and J. T. Njardarson, *J. Med. Chem.*, 2014, **57**, 10257; (b) N. Chadha and O. Silakari, *Eur. J. Med. Chem.*, 2017, **134**, 159.
- 4 (a) Y. Liu, K. Ai and L. Lu, *Chem. Rev.*, 2014, **114**, 5057; (b) M. Vlasselaer and W. Dehaen, *Molecules*, 2016, **21**, 785.
- 5 (a) F. R. de Sa Alves, E. J. Barreiro and C. A. M. Fraga, *Mini-Rev. Med. Chem.*, 2009, **9**, 782; (b) T. V. Sravanthi and S. L. Manju, *Eur. J. Pharm. Sci.*, 2016, **91**, 1; (c) N. K. Kaushik, N. Kaushik, P. Attri, N. Kumar, C. H. Kim, A. K. Verma and E. H. Choi, *Molecules*, 2013, **18**, 6620; (d) T. P. Singh and O. M. Singh, *Mini-Rev. Med. Chem.*, 2018, **18**, 9.
- 6 (a) G. R. Humphrey and J. T. Kuethe, *Chem. Rev.*, 2006, **106**, 2875; (b) F. T. Douglass and K. T. Pavan, *Tetrahedron*, 2011, **67**, 7195; (c) R. Vicente, *Org. Biomol. Chem.*, 2011, **9**, 6469; (d) M. Inman and C. J. Moody, *Chem. Sci.*, 2013, **4**, 29; (e) G. Bartoli, R. Dalpozzo and M. Nardi, *Chem. Soc. Rev.*, 2014, **43**, 4728; (f) M. M. Heravi, S. Rohani, V. Zadsirjan and N. Zahedi, *RSC Adv.*, 2017, **7**, 52852.
- 7 (a) G. Battistuzzi, S. Cacchi and G. Fabrizi, *Eur. J. Org. Chem.*, 2002, 2671; (b) K. Kruger, A. Tillack and M. Beller, *Adv. Synth. Catal.*, 2008, **350**, 2153; (c) S. Cacchi and G. Fabrizi, *Chem. Rev.*, 2011, **111**, PR215; (d) M. Platon, R. Amardeil, L. Djakovitch and J. C. Hierso, *Chem. Soc. Rev.*, 2012, **41**, 3929; (e) T. L. Guo, F. Huang, L. Yu and Z. Yu, *Tetrahedron Lett.*, 2015, **56**, 296; (f) S. W. Youn and T. Y. Ko, *Asian J. Org. Chem.*, 2018, **7**, 1467.
- 8 (a) S. Würtz, S. Rakshit, J. J. Neumann, T. Dröge and F. Glorius, *Angew. Chem., Int. Ed.*, 2008, **47**, 7230; (b) J. J. Neumann, S. Rakshit, T. Dröge, S. Würtz and F. Glorius, *Chem. - Eur. J.*, 2011, **17**, 7298.
- 9 R. Bernini, G. Fabrizi, A. Sferazza and S. Cacchi, *Angew. Chem., Int. Ed.*, 2009, **48**, 8078.
- 10 (a) X.-L. Lian, Z.-H. Ren, Y.-Y. Wang and Z.-H. Guan, *Org. Lett.*, 2014, **16**, 3360; (b) A. Mishra and I. Deb, *Adv. Synth. Catal.*, 2016, **358**, 2267.
- 11 (a) F. Huang, P. Wu, L. Wang, J. Chen, C. Sun and Z. Yu, *J. Org. Chem.*, 2014, **79**, 10553; (b) P. Drouhin and R. J. K. Taylor, *Eur. J. Org. Chem.*, 2015, 2333; (c) J. Liu, W. Wei, T. Zhao, X. Liu, J. Wu, W. Yu and J. Chang, *J. Org. Chem.*, 2016, **81**, 9326; (d) F.-Z. Hu, S.-H. Zhao, H. Chen, S.-W. Yu, X.-Y. Xu, W.-C. Yuan and X.-M. Zhang, *ChemistrySelect*, 2017, **2**, 1409.
- 12 Z.-H. Guan, Z.-Y. Yan, Z.-H. Ren, X.-Y. Liu and Y.-M. Liang, *Chem. Commun.*, 2010, 2823.
- 13 W. Yu, Y. Du and K. Zhao, *Org. Lett.*, 2009, **11**, 2417.
- 14 (a) Z. He, W. Liu and Z. Li, *Chem.-Asian J.*, 2011, **6**, 1340; (b) Z. Jia, T. Nagano, X. Li and A. S. C. Chan, *Eur. J. Org. Chem.*, 2013, 858.
- 15 (a) C.-J. Wu, Q.-Y. Meng, T. Lei, J.-J. Zhong, W.-Q. Liu, L.-M. Zhao, Z.-J. Li, B. Chen, C.-H. Tung and L.-Z. Wu, *ACS Catal.*, 2016, **6**, 4635; (b) W.-Q. Liu, T. Lei, Z.-Q. Song, X.-L. Yang, C.-J. Wu, X. Jiang, B. Chen, C.-H. Tung and L.-Z. Wu, *Org. Lett.*, 2017, **19**, 3251.
- 16 J. Zoller, D. C. Fabry, M. A. Ronge and M. Rueping, *Angew. Chem., Int. Ed.*, 2014, **53**, 13264.
- 17 S. Tang, X. Gao and A. Lei, *Chem. Commun.*, 2017, **53**, 3354.
- 18 (a) Z. Shi, C. Zhang, S. Li, D. Pan, S. Ding, Y. Cui and N. Jiao, *Angew. Chem., Int. Ed.*, 2009, **48**, 4572; (b) L. Ren, Z. Shi and N. Jiao, *Tetrahedron*, 2013, **69**, 4408.



- 19 D. Shen, J. Han, J. Chen, H. Deng, M. Shao, H. Zhang and W. Cao, *Org. Lett.*, 2015, **17**, 3283.
- 20 Y. Wei, I. Deb and N. Yoshikai, *J. Am. Chem. Soc.*, 2012, **134**, 9098.
- 21 (a) S. Kramer, K. Dooleweerd, A. T. Lindhardt, M. Rottländer and T. Skrydstrup, *Org. Lett.*, 2009, **11**, 4208; (b) G. Zhang, H. Yu, G. Qin and H.-M. Huang, *Chem. Commun.*, 2014, **50**, 4331; (c) D. Shen, J. Han, J. Chen, H. Deng, M. Shao, H. Zhang and W. Cao, *Org. Lett.*, 2015, **17**, 3283.
- 22 (a) D. R. Stuart, M. Bertrand-Laperle, K. M. N. Burgess and K. Fagnou, *J. Am. Chem. Soc.*, 2008, **130**, 16474; (b) J. Chen, G. Song, C.-L. Pan and X. Li, *Org. Lett.*, 2010, **12**, 5426; (c) D. R. Stuart, P. Alsabeh, M. Kuhn and K. Fagnou, *J. Am. Chem. Soc.*, 2010, **132**, 18326; (d) C. Wang, H. Sun, Y. Fang and Y. Huang, *Angew. Chem., Int. Ed.*, 2013, **52**, 5795; (e) C. Wang and Y. Huang, *Org. Lett.*, 2013, **15**, 5294; (f) D. Zhao, Z. Shi and F. Glorius, *Angew. Chem., Int. Ed.*, 2013, **52**, 12426; (g) B. Liu, C. Song, C. Sun, S. Zhou and J. Zhu, *J. Am. Chem. Soc.*, 2013, **135**, 16625; (h) T. Matsuda and Y. Tomaru, *Tetrahedron Lett.*, 2014, **55**, 3302; (i) L. Zheng and R. Hua, *Chem. - Eur. J.*, 2014, **20**, 2352; (j) K. Muralirajan and C. H. Cheng, *Adv. Synth. Catal.*, 2014, **356**, 1571; (k) Y. Hoshino, Y. Shibata and K. Tanaka, *Adv. Synth. Catal.*, 2014, **356**, 1577; (l) P.-Y. Tao and Y.-X. Jia, *Chem. Commun.*, 2014, **50**, 7367; (m) S. Kathiravan and I. A. Nicholls, *Chem. Commun.*, 2014, **50**, 14964; (n) B. Zhou, Y. Yang, H. Tang, J. Du, H. Feng and Y. Li, *Org. Lett.*, 2014, **16**, 3900; (o) Z.-L. Fan, S.-S. Song, W. Li and A. Zhang, *Org. Lett.*, 2015, **17**, 310; (p) L. Kong, F. Xie, S. Yu, Z. Qi and X. Li, *Chin. J. Catal.*, 2015, **36**, 925; (q) Z. Zhou, G.-X. Liu, Y. Chen and X.-Y. Lu, *Adv. Synth. Catal.*, 2015, **357**, 2944; (r) Y. Kim and S. Hong, *Chem. Commun.*, 2015, **51**, 11202; (s) H. Yan, H. Wang, X. Li, X. Xin, C. Wang and B. Wan, *Angew. Chem., Int. Ed.*, 2015, **54**, 10613; (t) H. J. Kim, D. C. Fabry, S. Madera and M. Rueping, *Org. Chem. Front.*, 2019, **6**, 2319.
- 23 (a) L. Ackermann and A. V. Lygin, *Org. Lett.*, 2012, **14**, 764; (b) Z. Zhang, H. Jiang and Y. Huang, *Org. Lett.*, 2014, **16**, 5976; (c) S. Allu and K. C. Kumara Swamy, *Adv. Synth. Catal.*, 2015, **357**, 2665.
- 24 (a) F. Zhou, X.-L. Han and X.-Y. Lu, *Tetrahedron Lett.*, 2011, **52**, 4681; (b) J.-L. Chen, Q. Y. Pang, Y. B. Sun and X. Li, *J. Org. Chem.*, 2011, **76**, 3523.
- 25 S. Kramer, K. Dooleweerd, A. T. Lindhardt, M. Rottländer and T. Skrydstrup, *Org. Lett.*, 2009, **11**, 4208.
- 26 W. Song and L. Ackermann, *Chem. Commun.*, 2013, **49**, 6638.
- 27 (a) Z.-Z. Zhang, B. Liu, J.-W. Xu, S.-Y. Yan and B.-F. Shi, *Org. Lett.*, 2016, **18**, 1776; (b) H. Wang, M. Moselage, M. J. González and L. Ackermann, *ACS Catal.*, 2016, **6**, 2705; (c) Q. Lu, S. Vázquez-Céspedes, T. Gensch and F. Glorius, *ACS Catal.*, 2016, **6**, 2352; (d) A. Lerchen, S. Vasquez-Céspedes and F. Glorius, *Angew. Chem., Int. Ed.*, 2016, **55**, 3208; (e) Y. J. Liang and N. Jiao, *Angew. Chem., Int. Ed.*, 2016, **55**, 4035.
- 28 (a) C. Chen, G. Shang, J. Zhou, Y. Yu, B. Li and J. Peng, *Org. Lett.*, 2014, **16**, 1872; (b) G. Zhao, C. Chen, Y. Yue, Y. Yu and J. Peng, *J. Org. Chem.*, 2015, **80**, 2827; (c) Y. Yue, J. Peng, D. Wang, Y. Bian, P. Sun and C. Chen, *J. Org. Chem.*, 2017, **82**, 5481; (d) X. Li, Y. Bian, X. Chen, H. Zhang, W. Wang, S. Ren, X. Yang, C. Lu, C. Chen and J. Peng, *Org. Biomol. Chem.*, 2019, **17**, 321; (e) X. Li, X. Chen, H. Wang, C. Chen, P. Sun, B. Mo and J. Peng, *Org. Biomol. Chem.*, 2019, **17**, 4014.
- 29 (a) C. Li, *Acc. Chem. Res.*, 2009, **42**, 335; (b) C. S. Yeung and V. M. Dong, *Chem. Rev.*, 2011, **111**, 1215; (c) C. Liu, J.-W. Yuan, M. Gao, S. Tang, W. Li, R.-Y. Shi and A.-W. Lei, *Chem. Rev.*, 2015, **115**, 12138.
- 30 Y. Li, J. Peng, X. Chen, B. Mo, X. Li, P. Sun and C. Chen, *J. Org. Chem.*, 2018, **83**, 5288.
- 31 P. Sun, J. Yang, Z. Song, Y. Cai, Y. Liu, C. Chen, X. Chen and J. Peng, *Synthesis*, 2020, **52**, 75.
- 32 Selected reviews, see: (a) J. X. Qiao and P. Y. S. Lam, *Synthesis*, 2011, 829; (b) K. S. Rao and T.-S. Wu, *Tetrahedron*, 2012, **68**, 7735; (c) X. Ma, F. Liu and D. Mo, *Chinese J. Org. Chem.*, 2017, **37**, 1069; (d) X. Duan, N. Liu, J. Wang and J. Ma, *Chinese J. Org. Chem.*, 2019, **39**, 661.
- 33 J. E. Steves and S. S. Stahl, *J. Am. Chem. Soc.*, 2013, **135**, 15742.
- 34 J. C. Antilla and S. L. Buchwald, *Org. Lett.*, 2001, **3**, 2077.
- 35 (a) J. C. Vantourout, H. N. Miras, A. Isidro-Llobet, S. Sproules and A. J. B. Watson, *J. Am. Chem. Soc.*, 2017, **139**, 4769; (b) A. E. King, T. C. Brunold and S. S. Stahl, *J. Am. Chem. Soc.*, 2009, **131**, 5044; (c) J. P. Collman and M. Zhong, *Org. Lett.*, 2000, **2**, 1233.
- 36 G. Hou, W. Li, M. Ma, X. Zhang and X. Zhang, *J. Am. Chem. Soc.*, 2010, **132**, 12844.
- 37 X. Guo, J. Han, Y. Liu, M. Qin, X. Zhang and B. Chen, *J. Org. Chem.*, 2017, **82**, 11505.
- 38 X. Zhang, D. Zhang-Negrerie, J. Deng, Y. Du and K. Zhao, *J. Org. Chem.*, 2013, **78**, 12750.
- 39 V. A. Bodunov, E. E. Galenko, A. V. Galenko, M. S. Novikov and A. F. Khlebnikov, *Synthesis*, 2018, **50**, 2784.

