






 Cite this: *RSC Adv.*, 2020, 10, 34729

 Received 23rd June 2020
 Accepted 3rd September 2020

DOI: 10.1039/d0ra05520d


rsc.li/rsc-advances

Kinetic investigation of the hydrolysis of uranium hexafluoride gas†

 Jason M. Richards, *^a Leigh R. Martin, ^a Glenn A. Fugate *^a
 and Meng-Dawn Cheng ^b

UF₆ is commonly employed in enrichment technologies and is known to react rapidly with water vapor to form radioactive particulates and hydrofluoric acid vapor. The kinetics of the UF₆ hydrolysis reaction have been observed directly for the first time. The rate appears to be half order and second order for UF₆ and water, respectively, with a rate constant of $1.19 \pm 0.22 \text{ Torr}^{-3/2} \text{ s}^{-1}$. The proposed mechanism involves formation of the [UF₆·2H₂O] adduct *via* two separate reactions.

Uranium hexafluoride (UF₆) is a key compound in the nuclear fuel cycle as the material employed in commercial enrichment, enabling production of the fuel for electricity production in nuclear power reactors. Studies evaluating the safety implications of small or large releases of UF₆ within a processing facility have been performed previously,^{1–3} as it is well known that this material undergoes a hydrolysis reaction in ambient air according to the simplified reaction:



producing both uranium particulates and hydrofluoric acid vapor. Recent work⁴ has explored the formation of the uranium particulate materials but the precise molecular mechanism of the initiating reaction has yet to be verified. Previous efforts to characterize this reaction have focused on spectroscopic measurements^{5–7} and computational studies.^{8–13} The experimental studies have focused on the spectroscopic detection of uranium species but lacked the temporal resolution to observe the reaction or identify any transient species. Unfortunately, there are major discrepancies between the experimentally observed reaction and computationally predicted mechanisms. For example, the computational studies suggest large thermodynamic energy barriers that would likely prevent the reaction of UF₆ and water vapor,^{9–13} yet the hydrolysis reaction is known to occur quickly and completely under ambient conditions.^{5–7} Additionally, one of the long-suspected intermediates of this reaction, UOF₄, has not been successfully identified under extreme conditions designed to optimize its production such as

reacting water with a 5000-fold excess of UF₆.¹⁴ Relevant spectroscopy measurements need to be performed to inform computational modelling because the mechanism may not be driven by a stepwise reaction of UF₆ and water molecules.

One prior study performed an indirect measurement of the rate of the UF₆ hydrolysis by monitoring the ingrowth of hydrogen fluoride with a laser-based analyzer.¹⁵ The reported rate constant of $4 \pm 4 \times 10^{-18} \text{ cm}^3 \text{ s}^{-1}$ remains questionable as limited experimental details were reported. Based on their findings, and assuming pseudo-first order kinetics under these reaction conditions in excess water vapor, the reaction half-life is calculated to be on the order of 1 ms or less. Standard infrared spectroscopy instrumentation cannot scan with sub-millisecond temporal resolution, requiring specialized instrumentation that has only recently become available.

This work focused on observing the disappearance of UF₆ under low pressure conditions by probing the most sensitive infrared (ν_3 antisymmetric stretching) vibrational band,¹⁶ located at 625 cm^{-1} , using a 5 m long-path length cell. The long path length combined with the sensitivity of the band allow the reaction rate to be investigated at sub-Torr partial pressures. The reaction is slow enough at these pressures to allow direct kinetic measurements of the hydrolysis to be made using a typical infrared spectrometer.

Experiments were performed in a 5 m long-path length gas cell fitted to an ABB MB3000 Fourier transform infrared spectrometer. The chamber was exposed to fluorine gas to passivate materials of construction and remove water from surfaces within the spectroscopy cell before each experiment. The UF₆ was then introduced into the evacuated cell at pressures between 10 and 30 mTorr and ambient temperature ($22.5 \pm 0.5 \text{ }^\circ\text{C}$). (**Warning:** UF₆ is a radioactive gas that forms highly toxic hydrogen fluoride in the presence of water.) After initial infrared spectroscopy measurements were made, 60 or 80 mTorr of water vapor was injected into the system and allowed to react. Typical infrared spectra collected during the hydrolysis reaction occurred during 0.5–2 min for these conditions, as shown in Fig. 1A. Note, the

^aIsotope and Fuel Cycle Technology Division, Oak Ridge National Laboratory, One Bethel Valley Road, Oak Ridge, TN, 37831, USA. E-mail: richardsjm@ornl.gov; fugatega@ornl.gov

^bEnvironmental Sciences Division, Oak Ridge National Laboratory, One Bethel Valley Road, Oak Ridge, TN, 37831, USA

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0ra05520d






Fig. 1 (A) Select infrared spectra showing the 625 cm^{-1} band of UF_6 just before introducing water vapor (red) and every 10 s after (blue) for 1 min. (B) Example of absorbance change of the 625 cm^{-1} band over time during a typical hydrolysis experiment. The rate of disappearance of the UF_6 increased about a factor of 10 when H_2O was introduced.

reactions that occurred when water vapor was introduced at these partial pressures were approximately an order of magnitude faster than any reactions of UF_6 with materials of construction. The observed rates (Fig. 1B) were significantly slower than those expected from diffusion-limited reactivity, suggesting that mixing occurred rapidly from the injection method.

Typical changes in the absorbance due to the reaction are shown in Fig. 2 after correction for reaction with materials of construction. Each figure is the average of three experiments and the data are corrected using a linear regression of the absorbance before time zero extrapolated through the points used for initial rate determination. The initial slope of the decay was fit using a linear regression on the first five data points for each reaction condition. The data were converted from absorbance units to mTorr, and the resulting conditional reaction rates in mTorr s^{-1} are shown in Table 1.

It should be noted that the conditions under which the hydrolysis reaction rate could be measured were limited by the scan rate of the spectroscopy equipment. The concentrations had to be maintained such that the reaction rate was significantly greater than the minimized rate of UF_6 consumption by the materials of construction but still had to be slow enough to have multiple measurements before the reaction was completed. Due to the high dependence of the hydrolysis reaction rate on water concentration, only two of the water concentrations measured gave reaction rates that were in the appropriate time frame bounding the measurable conditions.

The conditional reaction rates were plotted against the partial pressures of UF_6 and against the partial pressures of




Fig. 2 The normalized absorbance of the 625 cm^{-1} band in a 5 m path length cell for various partial pressures of UF_6 and water vapor. Water vapor was injected at time = 0 s.

Table 1 Conditional reaction rates of the hydrolysis of gaseous UF_6 in the presence of H_2O vapor

H_2O (mTorr)	UF_6 (mTorr)	Rate (mTorr s^{-1})
60 ± 2	8.2 ± 0.5	0.387 ± 0.008
60 ± 2	16.2 ± 1.0	0.556 ± 0.071
60 ± 2	30.2 ± 1.9	0.709 ± 0.321
80 ± 3	9.1 ± 0.6	0.702 ± 0.100
80 ± 3	18.6 ± 1.5	1.018 ± 0.066
80 ± 3	28.3 ± 1.6	1.312 ± 0.112

water on a log/log plot as shown in Fig. 3. The slope of the generated linear trends is the reaction order for UF_6 and water, respectively. This suggests a rate equation:

$$\text{Rate} = k[\text{UF}_6]^{0.5}[\text{H}_2\text{O}]^2,$$

where k is the rate constant. Applying the data in Table 1 to the suggested rate equation gives a rate constant of $1.19 \pm 0.22\text{ Torr}^{-3/2}\text{ s}^{-1}$.

Recent work⁴ explores the formation kinetics of uranyl fluoride particles from the reaction of gaseous UF_6 with water vapor. Some of the reactant concentrations in those studies are like those used in this work. Under humid condition (5% relative humidity and higher), the number concentration of uranyl fluoride particles produced in the hydrolysis reaction decrease during the observed time elapsed after start of the reaction (approximately 2–30 s). This is expected because the reaction is assumed to be complete in less than 2 s, and particles are undergoing various processes such as agglomeration and coagulation. Under very dry conditions (100 ppm and lower H_2O) the number concentration of uranyl fluoride particles increased over the same observed time (2–30 s). Applying the kinetic parameters determined in this work, after 30 s the reaction of 100 ppm (75 mTorr) UF_6 with 100 ppm (75 mTorr) of water is expected to only have consumed approximately 30% of the UF_6 and 60% of the water. The increase in number concentration of uranyl fluoride particles in very dry conditions observed




Fig. 3 Conditional reaction rates for the hydrolysis reaction as a function of the partial pressure of UF_6 (green, dashed) and water (blue, dotted) plotted on log-log scale. The dashed lines show the linear fit of the data for varied UF_6 pressures at 60 (■) and 80 (▲) mTorr H_2O . The dotted lines show the linear fit for varied H_2O pressures at 10 (●), 20 (■), and 30 (▲) mTorr UF_6 . The slope (m) of each line is displayed.




over the first 30 s in those studies agrees well with the reaction kinetics set forth in this work.

The kinetic model developed in this work can be applied to the recent work by Hu *et al.*¹³ that strongly suggests that the formation of two and three water molecule adduct species with UF₆ drastically reduced the calculated energy for the hydrolysis reaction. We propose the suggested reaction mechanism below based on the experimentally determined rate equation combined with those theoretical calculations.

Scheme 1 suggests a reaction mechanism that accounts for recent work suggesting high energy barriers for 1 : 1 UF₆ water adducts,^{8–11,13} and that multiple UF₆ centres, water molecules, or a combination of both may be involved,^{8,9,13} consistent with the reactant orders determined in this study. There is likely an initiation step where UF₆ reacts with water to form an adduct (eqn (1)). Note, theoretical studies suggest the formation of the single water adduct is slightly more energetically favourable than UF₆. Literature modelling suggests that further reactions toward the formation of uranyl fluoride are energetically disfavored.^{9–13} The increased energy stability coupled with the inability to react further suggests that this species may be formed in significant concentrations. The formation of an adduct containing two water molecules would confirm the second order behaviour in the rate equation. This may be formed by a direct reaction with water (eqn (2)) or potentially from reactions between two of the UF₆·H₂O species (eqn (3)). This may explain the fractional rate order, which is often explained by a complicated mechanism that have multiple pathways to reach the rate-limiting step. Both reactions may be viable because water and the UF₆ water adduct could be present in similar concentrations under the conditions of this work, and the formation of an anhydrous UF₆ molecule by eqn (3) would explain the fractional rate order associated with UF₆. This also suggests that the UF₆·2H₂O species is the reactant in the rate-determining step of the mechanism.

The conditions of this work were such that most of the water was consumed during the reaction and may have restricted the reaction to second order. If more water was available, Hu *et al.*¹³ suggest formation of a three water adduct species is more energetically favourable. This could be accommodated by the additional reactions shown in eqn (4) and (5), providing additional reaction pathways to react with UF₆.



Scheme 1 Proposed reaction scheme for initial UF₆ hydrolysis reaction.

In summary, direct observation of the hydrolysis reaction kinetics of gaseous UF₆ have been accomplished under low-pressure conditions. The results show the reaction was half order and second order with respect to UF₆ and water. The rate constant was determined to be $1.19 \pm 0.22 \text{ Torr}^{-3/2} \text{ s}^{-1}$. The results are consistent with recent experimental observations of the formation kinetics of uranyl fluoride particles from UF₆ hydrolysis under very dry conditions. The results also provide valuable data that support theoretical and computational studies.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors would also like to thank Dr Lee T. Trowbridge, Harold L. Jennings and Chris P. Boring for contributions to this work and the National Nuclear Security Administration's Defense Nuclear Nonproliferation Office of Research and Development for supporting this work.

Notes and references

- 1 S. P. Babenko, *Med. Tr. Prom. Ekol.*, 2005, 30–35.
- 2 M. Mohsendokht, *J. Loss Prev. Process Ind.*, 2017, **45**, 217–228.
- 3 Y.-f. Ma, Q. Li, R.-p. Hao, X.-y. Wu and X.-m. Xue, *Zhongguo Zhiye Yixue*, 2016, **43**, 467–470, 474.
- 4 M.-D. Cheng, J. M. Richards, M. A. Omana, J. A. Hubbard and G. A. Fugate, *React. Chem. Eng.*, 2020, **5**, 1708–1718.
- 5 S. A. Sherrow and R. D. Hunt, *J. Phys. Chem.*, 1992, **96**, 1095–1099.
- 6 D. Armstrong, W. Bostick and W. Fletcher, *Appl. Spectrosc.*, 1991, **45**, 1008–1016.
- 7 R. Reiner, *Moist air reaction with excess UF₆*, K/TCD-1122, Oak Ridge Gaseous Diffusion Plant, 1989.
- 8 T. Privalov, B. Schimmelpfennig, U. Wahlgren and I. Grenthe, *J. Phys. Chem. A*, 2002, **106**, 11277–11282.
- 9 S. L. Garrison and J. M. Becnel, *J. Phys. Chem. A*, 2008, **112**, 5453–5457.
- 10 S.-W. Hu, X.-Y. Wang, T.-W. Chu and X.-Q. Liu, *J. Phys. Chem. A*, 2008, **112**, 8877–8883.
- 11 S.-W. Hu, X.-Y. Wang, T.-W. Chu and X.-Q. Liu, *J. Phys. Chem. A*, 2009, **113**, 9243–9248.
- 12 M. C. Lind, S. L. Garrison and J. M. Becnel, *J. Phys. Chem. A*, 2010, **114**, 4641–4646.
- 13 S.-W. Hu, H. Lin, X.-Y. Wang and T.-W. Chu, *J. Mol. Struct.*, 2014, **1062**, 29–34.
- 14 M. Otey and R. LeDoux, *J. Inorg. Nucl. Chem.*, 1967, **29**, 2249–2256.
- 15 V. Klimov, Y. M. Kravetz and A. Besmelnitzin, *J. Fluorine Chem.*, 1992, **58**, 262.
- 16 B. Weinstock and G. L. Goodman, in *Advances in Chemical Physics*, ed. I. Prigogine, J Wiley, New York, 1965, vol. 9, pp. 169–319.

