# Chemical Science

# EDGE ARTICLE

Check for updates

Cite this: Chem. Sci., 2020, 11, 6479

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 27th February 2020 Accepted 16th March 2020

DOI: 10.1039/d0sc01171a

rsc.li/chemical-science

### Introduction

Photodynamic therapy (PDT) is an effective clinical treatment strategy for malignant tumors.<sup>1</sup> In the PDT process, a photosensitizer (PS) is activated under light irradiation, and the excited PS subsequently interacts with molecular oxygen to generate cytotoxic reactive oxygen species (ROS), which can oxidize biomolecules resulting in cancer cell death.<sup>2,3</sup> In particular, theranostics has recently been recognized as a promising medical technology that combines diagnostic and therapeutic capabilities in one dose to achieve the real-time and precise monitoring of the therapeutic effect of the drug.<sup>4</sup> Moreover, PDT combined with image-guided diagnosis has attracted considerable attention owing to its prominent advantages such as high spatiotemporal selectivity, noninvasiveness, and fewer side effects.<sup>5-7</sup> Therefore, the design of

# Fine-tuning the electronic structure of heavyatom-free BODIPY photosensitizers for fluorescence imaging and mitochondria-targeted photodynamic therapy<sup>†</sup>

Sujie Qi, 🝺 a Nahyun Kwon, a Yubin Yim, a Van-Nghia Nguyen 🕩 \*ab and Juyoung Yoon 🕩 \*a

Theranostics that combines both diagnosis and therapy into a single platform has recently emerged as a promising biomedical approach for cancer treatment; however, the development of efficient theranostic agents with excellent optical properties remains a challenge. Here, we report novel mitochondria-targeting BODIPY photosensitizers (R-BODs) that possess considerable singlet oxygen generation capabilities and good fluorescence properties for imaging-guided photodynamic therapy (PDT). The incorporation of sulfur atoms into the  $\pi$ -conjugated skeleton of BODIPY along with the introduction of different functional groups at the meso-position of the BODIPY core is essential for tuning the photophysical and photosensitizing properties. Notably, the MeOPh-substituted thiophenefused BODIPY (MeO-BOD, R = p-methoxyphenyl) displayed the highest singlet oxygen generation capability ( $\Phi_{\Delta} \approx 0.85$  in air-saturated acetonitrile) and a moderate fluorescence quantum yield ( $\Phi_{\rm f} =$ 17.11). Furthermore, MeO-BOD showed good biocompatibility, low dark toxicity and superior fluorescence imaging properties in living cells. More importantly, the PDT efficacy of mitochondriaspecific anchoring of MeO-BOD was remarkably amplified with an extremely low half-maximal inhibitory concentration (IC<sub>50</sub>) value of 95 nM. We believe that the incorporation of an electron-donating group at the meso-position of the thiophene-fused BODIPY platform may be an effective approach for developing theranostic agents for precision cancer therapy.

novel PSs that can effectively produce both fluorescence and ROS is in high demand.

To date, numerous organic dyes, such as porphyrins, phthalocyanines, cyanine, squaraine, diketopyrrolopyrrole (DPP), and boron dipyrromethane (BODIPY) derivatives, have been developed as theranostic agents.7-13 Among these, BODIPY dyes have attracted great interest as theranostic agents in photodynamic cancer therapy due to their excellent photochemical stability, good biocompatibility, high molar extinction coefficients, high quantum efficiencies of fluorescence and facile modification.<sup>11-17</sup> At present, for obtaining remarkable PDT efficiency of BODIPY dyes, the most popular approach is the introduction of heavy halogen atoms (Br and I) to promote spin-orbit coupling (SOC), which enhances intersystem crossing (ISC) and improves the singlet oxygen  $({}^{1}O_{2})$  generation capability.<sup>18-24</sup> However, the incorporation of heavy atoms increases the dark-toxicity and quench fluorescence.25 Thus, BODIPY PSs without heavy halogen atoms are preferred as theranostic agents. Recently, several approaches to enhance the ISC, such as the use of double excited states,<sup>26,27</sup> spin converters,<sup>28</sup> and photoinduced electron transfer (PET),<sup>29-31</sup> have been implemented in the development of PSs without



View Article Online

View Journal | View Issue

<sup>&</sup>lt;sup>a</sup>Department of Chemistry and Nanoscience, Ewha Womans University, Seoul 03760, Republic of Korea. E-mail: jyoon@ewha.ac.kr

<sup>&</sup>lt;sup>b</sup>Institute of Research and Development, Duy Tan University, Da Nang 550000, Vietnam. E-mail: nguyenvannghia8@duytan.edu.vn

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: 10.1039/d0sc01171a

heavy atoms; however, the search for novel PSs is still required for PDT to reach its full potential.

Intriguingly, one alternative strategy to strengthen the ISC with an efficient triplet population is to increase the SOC while decreasing the singlet-triplet energy gaps; to do this, a thiophene moiety was introduced into the  $\pi$ -conjugated system of **BODIPY**,<sup>32-35</sup> but the oncological applications of this approach have not been fully studied.<sup>35</sup> Furthermore, the introduction of electron-donating groups at the *meso*-site of the **BODIPY** platform is also an emerging strategy for achieving efficient PSs.<sup>36,37</sup> The abovementioned factors inspired us to consider designing PSs with different functional substituents incorporated at the *meso*-position of thiophene-fused **BODIPY** to achieve compounds suitable for clinical use. To the best of our knowledge, there have been no attempts to design efficient theranostic agents by taking advantage of the above two strategies.

Herein, we designed and synthesized a series of heavy-atomfree thiophene-**BODIPY** derivatives by incorporating different functional groups (pyridinyl (**PY**), phenyl (**PH**), *p*-methoxyphenyl (**MeO**) and *N*,*N*-dimethylaminophenyl (**DMA**)) at the *meso*-position of the thiophene-**BODIPY** platform (Scheme 1). The experimental results revealed that the <sup>1</sup>O<sub>2</sub> generation ability of these thiophene-**BODIPY** derivatives through a type II process gradually increased with increasing electron-donating ability of the substituent ( $\Phi_{\Delta}$ : **PY-BOD** < **PH-BOD** < **MeO-BOD**). Fortunately, **MeO-BOD** had prominent dual functions, showing both a high <sup>1</sup>O<sub>2</sub> quantum yield and moderate fluorescence intensity. Furthermore, the cellular experimental results showed that **MeO-BOD** could be utilized as a mitochondria-specific diagnostic agent to reinforce the PDT effect.

### **Results and discussion**

#### Synthesis and photophysical properties

The different thiophene-fused **BODIPY** derivatives were synthesized through the trifluoroacetic acid-catalyzed condensation of thiophene-fused pyrroles with different benzaldehydes, followed by oxidation and complexation (Scheme 1). Detailed experimental procedures are provided in the ESI.† The structures of all these thiophene-fused **BODIPY** derivatives were fully confirmed by <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy and high-resolution mass spectrometry (Fig. S1–S14†). UV-vis



Scheme 1 Synthetic route to the thiophene-fused BODIPY derivatives.

absorption and fluorescence spectra of the new thiophene-**BODIPY** derivatives were acquired (Fig. 1, S15 and S16<sup>†</sup>). These key photophysical property data are summarized in Table 1.

The absorption and fluorescence spectra of all the thiophene-fused BODIPY derivatives were similar in shape to those of H-BOD, indicating that the R substituents had little effect on the HOMO-LUMO energy gap of the thiophene-fused BODIPY chromophore because they are nearly orthogonal to the thiophene-fused BODIPY core (Fig. 1a).32 All the studied compounds retained high absorption molar extinction coefficients ( $\varepsilon \sim 100\ 000\ M^{-1}\ cm^{-1}$ ), and the maximum absorptions were shifted to longer wavelengths ( $\lambda_{abs} \sim 565 \text{ nm}$ ) compared with that of unmodified BODIPY. The absorption spectra of all the thiophene-BODIPY derivatives showed a slight blueshift with increasing solvent polarity. This phenomenon indicates that the dipole moments of the ground state of these thiophene-BODIPY derivatives might be larger than those of the first singlet excited states.32 On the other hand, PY-BOD, PH-BOD and MeO-BOD exhibited moderate fluorescence intensities, while DMA-BOD was poorly emissive due to a PET mechanism.<sup>38,39</sup> The fluorescence quantum yields of all the thiophene-BODIPY derivatives were evaluated in toluene (Table 1), and PY-BOD, PH-BOD and MeO-BOD were shown to be excellent candidates for fluorescence imaging.

#### <sup>1</sup>O<sub>2</sub> generation ability

Subsequently, the singlet oxygen generation capability of **PY-BOD**, **PH-BOD**, **MeO-BOD** and **DMA-BOD** was assessed in airsaturated acetonitrile (ACN) under 560 nm irradiation. A commercial <sup>1</sup>O<sub>2</sub> probe, 1,3-diphenylisobenzofuran (DPBF), was used as an indicator, and rose bengal (RB,  $\Phi_{\Delta} = 0.54$  in ACN) was used as the reference.<sup>40</sup> As shown in Fig. 2a and S17,† the absorbance at 410 nm of DPBF decreased gradually in the presence of the thiophene-**BODIPY** derivatives under continuous light irradiation. According to the linear relationship of the decay curves (Fig. 2b), the <sup>1</sup>O<sub>2</sub> quantum yields of **PY-BOD**, **PH-BOD**, **MeO-BOD** and **DMA-BOD** were calculated to be 0.52, 0.69, 0.85 and 0.04, respectively (Table 1). The strongest <sup>1</sup>O<sub>2</sub> generation ability of **MeO-BOD** among all the **R-BOD**s suggested



Fig. 1 (a) Design of thiophene-fused BODIPY derivatives, (b) UV-Vis absorption and (c) emission spectra of PY-BOD, PH-BOD, MeO-BOD and DMA-BOD in toluene at  $c = 5.0 \mu$ M and  $\lambda_{ex} = 540 nm$ .

Table 1 Photophysical properties of the synthesized compounds

Compounds	$\lambda_{abs}{}^{a} [nm]$	$\varepsilon  imes 10^{-3a} \left[ \mathrm{M}^{-1} \ \mathrm{cm}^{-1}  ight]$	$\lambda_{\rm em}{}^a [{\rm nm}]$	Stokes shift [nm]	$\Phi_{\mathrm{f}}^{\;b}\left(\% ight)$	$\Phi_{\Delta}{}^c$
BODIPY <sup>d</sup>	502	120.00	508	6	70.0	≈0
PY-BOD	568	107.09	585	17	9.82	0.52
PH-BOD	564	111.60	576	12	18.83	0.69
MeO-BOD	564	115.78	579	15	17.11	0.85
DMA-BOD	562	96.99	577	15	2.50	0.04

<sup>*a*</sup> In toluene ( $5.0 \times 10^{-6}$  M). <sup>*b*</sup> Fluorescence quantum yield estimated relative to rhodamine 101 as the standard ( $\Phi_{\rm f} = 1.0$  in methanol). <sup>*c*</sup> Singlet oxygen quantum yield was determined with respect to rose bengal ( $\Phi_{\Delta}$  (RB) = 0.54 in ACN). <sup>*d*</sup> Literature value.<sup>19</sup>

a<sub>1.2</sub> (A/A) 10 DPBF 0.8 0.8 vbsorbar 0.6 f 10 \$ 0.6 12 : 0.4 0.4 0 2 0.2 0 400 450 500 550 600 650 350 8 10 12 Time (s) Wavelength (nm) С d 0.15 0.12 0.09 0.06 0.03 0.00 500 550 600 650 700 450 Wavelength (nm)

Fig. 2 (a) Time-dependent photodegradation of DPBF with MeO-BOD; (b) the DPBF degradation rate curves with PY-BOD, PH-BOD, MeO-BOD, DMA-BOD and rose bengal in ACN; (c) a TEM image of MeO-BOD NPs (inset: high magnification TEM image); and (d) UV-Vis absorption spectrum of MeO-BOD (5  $\mu$ M) in deionized water.

the electron donating group (MeOPh-) in the thiophene-fused BODIPY derivatives played an important role in enhancing the  ${}^{1}O_{2}$  generation. Based on these results, we propose a plausible mechanism for MeO-BOD. The introduction of the donating group probably favors <sup>1</sup>O<sub>2</sub> generation first by increasing the formation of the charge transfer (CT) state via photoinduced charge transfer (PCT) (<sup>1</sup>BOD-Donor  $\rightarrow$  BOD<sup>[ $\delta$ -]</sup>-Donor<sup>[ $\delta$ +]</sup>). The charge recombination of the CT state further triggers the production of T1 of the thiophene-fused BODIPY derivatives  $(BOD^{[\delta-]}-Donor^{[\delta+]} \rightarrow {}^{3}BOD-Donor)$ . Finally,  ${}^{1}O_{2}$  is generated by energy transfer from  $T_1$  to molecular oxygen (<sup>3</sup>BOD-Donor +  $O_2$  $\rightarrow$  BOD-Donor + <sup>1</sup>O<sub>2</sub>) (Scheme S1<sup>†</sup>).<sup>29-31,37,41,42</sup> Unfortunately, **DMA-BOD** showed almost no  ${}^{1}O_{2}$  generating ability in ACN, which implied that the electron-donating group could also lead to a PET process, which triggered the quenching of fluorescence and forbidding of non-radiative transitions such as ISC, thereby prohibiting the <sup>1</sup>O<sub>2</sub> generation.<sup>43</sup> Thus, the <sup>1</sup>O<sub>2</sub> formation efficiency could be controlled by finely tuning the electronic properties of the meso-substituent on the thiophene-BODIPY platform, especially by the introduction of a suitable electrondonating group. Taken together, these results indicate that **MeO-BOD** has potential as a theranostic agent for cancer treatment.

#### Cellular fluorescence imaging and subcellular colocalization

Given the above inspiring results, we attempted to investigate the fluorescence (FL) imaging and PDT efficiency of MeO-BOD in living cells. To this end, MeO-BOD nanoparticles (NPs) were prepared by adding the DMSO stock solution of MeO-BOD into water.44 The transmission electron microscopy (TEM) image in Fig. 2c indicated that the MeO-BOD NPs had a regular spherical morphology with a diameter of approximately 72 nm. The size distribution of the nanoparticles was determined using dynamic light scattering (DLS), which showed that the average size was  $68 \pm 7$  nm (Fig. S18<sup>†</sup>). These particles are suitably sized for passive targeting through the enhanced permeability and retention (EPR) effect.45 In addition, the evident decrease in absorbance and the broad absorption spectrum and the redshift of the fluorescence spectrum of MeO-BOD in deionized water (DW) suggested that the formation of MeO-BOD NPs might be due to J-aggregation (Fig. 2d and S19b<sup>†</sup>).<sup>46</sup> The MeO-BOD NPs could be disassembled and the instinctive fluorescence peak of MeO-BOD could be restored in the presence of FBS (10%) in DW (Fig. S19<sup>†</sup>).

Then, we further explored the cellular uptake of MeO-BOD in HeLa cells by using confocal laser scanning microscopy. As illustrated in Fig. 3a, MeO-BOD could be rapidly internalized by living cells, and the FL images showed strong emission in the cell cytoplasm. Therefore, MeO-BOD could be employed as an imaging-guided PDT agent. Furthermore, to test the main organelle locations of MeO-BOD, we co-stained HeLa cells with MeO-BOD and commercial MitoTracker Green (MTG) or Lyso-Tracker Green (LTG). The colocalization experiments indicated that MeO-BOD were mainly localized in mitochondria, as indicated by the high Pearson's coefficient (0.98), instead of the lysosomes (Pearson's coefficient 0.62) (Fig. 3b-e). In addition, the time-dependent colocalization fluorescence imaging of MeO-BOD with MTG in HeLa cells was performed. As shown in Fig. S20,<sup>†</sup> MeO-BOD almost internalize into the mitochondria of HeLa cells after 1 h. Recent reports have demonstrated that the BODIPY dye itself could localize into the mitochondria of cells due to its low electron density character.47 The unmodified **BODIPY** platform has a low electron density  $(+\delta)$  character. Also, the incorporation of sulfur atoms into the  $\pi$ -conjugated skeleton of BODIPY perhaps led to more reduction of electron



Fig. 3 (a) Confocal laser scanning microscopy images of HeLa cells incubated with MeO-BOD (1.0  $\mu$ M) for different times (0, 15, 30 and 60 min); scale bar: 30  $\mu$ m. Confocal laser scanning microscopy colocalization fluorescence images of MeO-BOD (1.0  $\mu$ M) with (b) LTG and (c) MTG (500 nM) in HeLa cells, respectively. The fluorescence intensity correlation of MeO-BOD with (d) LTG and (e) MTG, respectively.  $R_p$  is Pearson's coefficient; scale bar: 20  $\mu$ m.

density of the **BODIPY** core,<sup>48</sup> which might facilitate the mitochondrial accumulation of **MeO-BOD**. Subcellular organelles are indispensable in maintaining cellular biological function.<sup>49</sup> In particular, the generation of <sup>1</sup>O<sub>2</sub> in mitochondria can induce direct dysfunction and trigger cell apoptosis. Thus, mitochondria-targeted theranostic agents could maximize cancer treatment efficiency.

#### In vitro PDT efficacy evaluation

To demonstrate the PDT efficacy of MeO-BOD in living cells, we first tested its cytotoxicity by the methyl thiazolyltetrazolium (MTT) assay in HeLa cells. As indicated in Fig. 4a, MeO-BOD had negligible cytotoxicity in the dark, revealing its excellent biocompatibility in vitro. Under 560 nm light irradiation (0.1 W  $cm^{-2}$ , 5 or 10 min), the viability of HeLa cells gradually decreased with the increasing concentration of MeO-BOD, and the growth inhibition ratio reached ~88% even at a very low concentration of 0.15 µM (Fig. 4b). The half-maximal inhibitory concentration  $(IC_{50})$  of MeO-BOD for HeLa cells was as low as 95 nM under 560 nm light irradiation ( $0.1 \,\mathrm{W}\,\mathrm{cm}^{-2}$ , 10 min). The extremely low IC<sub>50</sub> of MeO-BOD could be attributed to the high <sup>1</sup>O<sub>2</sub> quantum yield and efficient mitochondria-specific ROS generation upon light irradiation. To clarify the cytotoxicity in the PDT process, we tracked the morphological variations of HeLa cells in the presence of MeO-BOD under 559 nm laser irradiation using confocal laser scanning microscopy. As described in Fig. S21,† with increasing irradiation time (0-10 min), the morphology of the HeLa cells preincubated with MeO-BOD obviously changed; gradual thinning of the cell membrane and the formation of numerous blebs (red line) were observed.



Fig. 4 Concentration-dependent changes in the cell viability of HeLa cells treated with MeO-BOD using a typical MTT assay (a) in the dark and (b) under light irradiation. Cells were irradiated with 560 nm light (0.1 W cm<sup>-2</sup>, 5 and 10 min). (c) Fluorescence images of calcein AM/PI-stained HeLa cells after preincubation with MeO-BOD (0.5 and 1.0  $\mu$ M) for 1 h and irradiation with 560 nm light (0.1 W cm<sup>-2</sup>, 10 min). Scale bar: 40  $\mu$ m.

In contrast, the cells not exposed to **MeO-BOD** exhibited no appreciable morphological changes under the same laser irradiation.

Additionally, to intuitively establish the PDT efficacy of **MeO-BOD** (0.5 and 1.0  $\mu$ M), live/dead cell co-staining of HeLa cells with calcein AM (green, live cells) and PI (red, dead cells) was performed to visualize the cell viability (Fig. 4c). The control group showed strong green fluorescence for live cells, which also verified the low dark toxicity and good biocompatibility of **MeO-BOD**. In comparison, almost all the HeLa cells treated with **MeO-BOD** were killed, and intense red fluorescence was observed. All these results confirmed that **MeO-BOD** could be employed as a theranostic agent for cancer treatment *in vitro*.

#### Apoptosis mechanism of MeO-BOD-mediated PDT

Finally, to elucidate the potential therapeutic mechanism at the cellular level, we further evaluated the cellular  ${}^{1}O_{2}$  generation capability of **MeO-BOD** by using 2,7-dichlorodihydrofluorescein diacetate (DCFH-DA) as the  ${}^{1}O_{2}$  indicator in HeLa cells (Fig. 5a). The HeLa cells treated with only DCFH-DA and only **MeO-BOD** showed almost no fluorescence, whereas the group pretreated with **MeO-BOD** before light irradiation and then incubated with DCFH-DA showed apparent green fluorescence from oxidized 2,7-dichlorofluorescein (DCF). These experimental results validated the  ${}^{1}O_{2}$  generation ability of **MeO-BOD** in living cells. As mentioned above, the production of  ${}^{1}O_{2}$  in mitochondria causes mitochondrial destruction, resulting in cell apoptosis. The reduction of mitochondrial membrane potential (MMP) is



Fig. 5 (a) Fluorescence images of ROS generation in HeLa cells after incubation with MeO-BOD (1.0  $\mu$ M) for 1 h using DCFH-DA as an indicator (10  $\mu$ M). The green fluorescence indicates that DCFH-DA is oxidized to DCF. (b) Fluorescence images of the mitochondrial membrane potential of HeLa cells after incubation without or with MeO-BOD (1.0  $\mu$ M) using JC-1 as the indicator (2  $\mu$ M),  $\lambda_{ex} = 473$  nm, red channel for aggregates (healthy cells): 575–675 nm, and green channel for monomers (apoptotic cells): 490–540 nm. Samples were irradiated with 560 nm light (0.1 W cm<sup>-2</sup>, 10 min). Scale bar: 30  $\mu$ m.

a crucial signal of mitochondrial damage. Hence, the MMP changes were monitored by using confocal fluorescence images using JC-1 dye, as its fluorescence color changed between its aggregates (red, high MMP) and monomers (green, low MMP). The untreated control group under 560 nm light irradiation  $(0.1 \text{ W cm}^{-2} \text{ for 10 min})$  displayed strong red emission and weak green emission, indicating that the cells were healthy with a high MMP. By comparison, the cells treated with **MeO-BOD** suffered from depolarization of the mitochondrial membrane, as demonstrated by the increase of green fluorescence intensity (Fig. 5b). These observations further suggested that the cell death was induced by a mitochondria-associated pathway.

### Conclusions

We synthesized thiophene-fused **BODIPY** analogues with different functional groups at the *meso*-position of the **BODIPY** core as mitochondria-targeted theranostic agents for imaging and PDT. The varying R-substituents had little effect on the absorption and emission maxima. **PY-BOD**, **PH-BOD** and **MeO-BOD** exhibited moderate fluorescence quantum yields, while

that of DMA-BOD was much lower due to a PET process. Among the analogues, MeO-BOD exhibited excellent dual functionality, both considerable <sup>1</sup>O<sub>2</sub> generation ability and high brightness. The cell experiments manifested that MeO-BOD offered many advantages: good biocompatibility, mitochondria-specific fluorescence imaging, and a very low IC<sub>50</sub> value ( $\approx$  95 nM). Therefore, MeO-BOD could be employed as an imaging-guided PDT agent for cancer treatment. In addition, the apoptosis mechanism of light-induced PDT might be a result of ROS-induced damage to mitochondria, which was demonstrated by detecting the changes of the mitochondrial membrane potential. Briefly, finely tuning the electronic structure of the substituent at the meso-site of a heavy-atom-free thiophene-fused BODIPY core is a promising strategy for developing highly efficient theranostic agents with minimal side effects to accomplish the integration of diagnosis and therapy.

### Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This study was supported by grants from the National Research Foundation of Korea (NRF) funded by the Korean government (MSIP) (No. 2012R1A3A2048814 for J. Y.). Mass spectral data were obtained from the Korea Basic Science Institute (Daegu) on a Jeol JMS 700 high resolution mass spectrometer.

## Notes and references

- 1 X. Li, S. Lee and J. Yoon, *Chem. Soc. Rev.*, 2018, 47, 1174–1188.
- P. Agostinis, K. Berg, K. A. Cengel, T. H. Foster, A. W. Girotti,
   S. O. Gollnick, S. M. Hahn, M. R. Hamblin, A. Juzeniene,
   D. Kessel, M. Korbelik, J. Moan, P. Mroz, D. Nowis,
   J. Piette, B. C. Wilson and J. Golab, *Ca-Cancer J. Clin.*, 2011,
   61, 250–281.
- 3 D. E. J. G. J. Dolmans, D. Fukumura and R. K. Jain, *Nat. Rev. Cancer*, 2003, **3**, 380–387.
- 4 J. Zhang, L. Ning, J. Huang, C. Zhang and K. Pu, *Chem. Sci.*, 2020, **11**, 618–630.
- 5 J. Dai, Y. Li, Z. Long, R. Jiang, Z. Zhuang, Z. Wang, Z. Zhao, X. Lou, F. Xia and B. Z. Tang, ACS Nano, 2020, 14, 854–866.
- 6 Y. Yuan, C.-J. Zhang, M. Gao, R. Zhang, B. Z. Tang and B. Liu, *Angew. Chem., Int. Ed.*, 2015, **54**, 1780–1786.
- 7 Y. Cai, W. Si, W. Huang, P. Chen, J. Shao and X. Dong, *Small*, 2018, **14**, 1704247.
- 8 R. Kumar, W. S. Shin, K. Sunwoo, W. Y. Kim, S. Koo, S. Bhuniya and J. S. Kim, *Chem. Soc. Rev.*, 2015, 44, 6670– 6683.
- 9 X. Yi, F. Wang, W. Qin, X. Yang and J. Yuan, *Int. J. Nanomed.*, 2014, **9**, 1347–1365.
- 10 X. Li, C. y. Kim, S. Lee, D. Lee, H.-M. Chung, G. Kim, S.-H. Heo, C. Kim, K.-S. Hong and J. Yoon, *J. Am. Chem. Soc.*, 2017, **139**, 10880–10886.

- 11 M. Li, S. Long, Y. Kang, L. Guo, J. Wang, J. Fan, J. Du and X. Peng, J. Am. Chem. Soc., 2018, 140, 15820–15826.
- 12 R. Wang, K. Dong, G. Xu, B. Shi, T. Zhu, P. Shi, Z. Guo, W.-H. Zhu and C. Zhao, *Chem. Sci.*, 2019, **10**, 2785–2790.
- 13 H. Wang, W. Zhao, X. Liu, S. Wang and Y. Wang, *ACS Appl. Bio Mater.*, 2020, **3**, 593–601.
- 14 J. Zhao, W. Wu, J. Sun and S. Guo, *Chem. Soc. Rev.*, 2013, 42, 5323–5351.
- 15 S. G. Awuah and Y. You, RSC Adv., 2012, 2, 11169–11183.
- 16 A. Turksoy, D. Yildiz and E. U. Akkaya, *Coord. Chem. Rev.*, 2019, **379**, 47–64.
- 17 W. Sun, X. Zhao, J. Fan, J. Du and X. Peng, *Small*, 2019, **15**, 1804927.
- 18 A. Gorman, J. Killoran, C. O'Shea, T. Kenna, W. M. Gallagher and D. F. O'Shea, *J. Am. Chem. Soc.*, 2004, **126**, 10619–10631.
- 19 T. Yogo, Y. Urano, Y. Ishitsuka, F. Maniwa and T. Nagano, *J. Am. Chem. Soc.*, 2005, **127**, 12162–12163.
- 20 J. Tian, J. Zhou, Z. Shen, L. Ding, J.-S. Yu and H. Ju, *Chem. Sci.*, 2015, **6**, 5969–5977.
- 21 Q.-J. Hu, Y.-C. Lu, C.-X. Yang and X.-P. Yan, *Chem. Commun.*, 2016, **52**, 5470–5473.
- 22 L. Huang, Z. Li, Y. Zhao, Y. Zhang, S. Wu, J. Zhao and G. Han, J. Am. Chem. Soc., 2016, **138**, 14586–14591.
- 23 H. S. Jung, J. Han, H. Shi, S. Koo, H. Singh, H.-J. Kim, J. L. Sessler, J. Y. Lee, J.-H. Kim and J. S. Kim, *J. Am. Chem. Soc.*, 2017, **139**, 7595–7602.
- 24 X. Miao, W. Hu, T. He, H. Tao, Q. Wang, R. Chen, L. Jin, H. Zhao, X. Lu, Q. Fan and W. Huang, *Chem. Sci.*, 2019, 10, 3096–3102.
- 25 S. Xu, W. Wu, X. Cai, C.-J. Zhang, Y. Yuan, J. Liang, G. Feng,
   P. Manghnani and B. Liu, *Chem. Commun.*, 2017, 53, 8727– 8730.
- 26 Y. Cakmak, S. Kolemen, S. Duman, Y. Dede, Y. Dolen, B. Kilic, Z. Kostereli, L. T. Yildirim, A. L. Dogan, D. Guc and E. U. Akkaya, *Angew. Chem., Int. Ed.*, 2011, 50, 11937– 11941.
- 27 S. Kolemen, M. Işık, G. M. Kim, D. Kim, H. Geng, M. Buyuktemiz, T. Karatas, X.-F. Zhang, Y. Dede, J. Yoon and E. U. Akkaya, *Angew. Chem., Int. Ed.*, 2015, 54, 5340– 5344.
- 28 M. Üçüncü, E. Karakuş, E. Kurulgan Demirci, M. Sayar, S. Dartar and M. Emrullahoğlu, Org. Lett., 2017, 19, 2522– 2525.
- 29 J. T. Buck, A. M. Boudreau, A. DeCarmine, R. W. Wilson, J. Hampsey and T. Mani, *Chem*, 2019, 5, 138–155.

- 30 M. A. Filatov, S. Karuthedath, P. M. Polestshuk, H. Savoie,
  K. J. Flanagan, C. Sy, E. Sitte, M. Telitchko, F. Laquai,
  R. W. Boyle and M. O. Senge, *J. Am. Chem. Soc.*, 2017, 139, 6282–6285.
- 31 X.-F. Zhang and N. Feng, *Chem.-Asian J.*, 2017, **12**, 2447–2456.
- 32 S. Ji, J. Ge, D. Escudero, Z. Wang, J. Zhao and D. Jacquemin, *J. Org. Chem.*, 2015, **80**, 5958–5963.
- 33 S. G. Awuah, S. K. Das, F. D'Souza and Y. You, *Chem.-Asian J.*, 2013, 8, 3123–3132.
- 34 K. Tanaka, H. Yamane, R. Yoshii and Y. Chujo, *Bioorg. Med. Chem.*, 2013, **21**, 2715–2719.
- 35 R. L. Watley, S. G. Awuah, M. Bio, R. Cantu, H. B. Gobeze,
  V. N. Nesterov, S. K. Das, F. D'Souza and Y. You, *Chem.-Asian J.*, 2015, **10**, 1335–1343.
- 36 X. Xia and Y. Qian, Analyst, 2018, 143, 5218-5224.
- 37 C. Wang and Y. Qian, Biomater. Sci., 2020, 8, 830-836.
- 38 W. Lin, W. Zhang, S. Liu, Z. Li, X. Hu, Z. Xie, C. Duan and G. Han, ACS Appl. Mater. Interfaces, 2019, 11, 43928–43935.
- 39 X. Liu, Q. Qiao, W. Tian, W. Liu, J. Chen, M. J. Lang and Z. Xu, J. Am. Chem. Soc., 2016, 138, 6960–6963.
- 40 V.-N. Nguyen, S. Qi, S. Kim, N. Kwon, G. Kim, Y. Yim, S. Park and J. Yoon, *J. Am. Chem. Soc.*, 2019, **141**, 16243–16248.
- 41 V. N. Nguyen, Y. Yim, S. Kim, B. Ryu, K. M. K. Swamy, G. Kim, N. Kwon, C. Y. Kim, S. Park and J. Yoon, *Angew. Chem., Int. Ed.*, 2020, DOI: 10.1002/anie.202002843.
- 42 W. Hu, Y. Lin, X.-F. Zhang, M. Feng, S. Zhao and J. Zhang, *Dyes Pigm.*, 2019, **164**, 139–147.
- 43 Y. Liu, C. Xu, L. Teng, H.-W. Liu, T.-B. Ren, S. Xu, X. Lou,
  H. Guo, L. Yuan and X.-B. Zhang, *Chem. Commun.*, 2020, 56, 1956–1959.
- 44 X. Li, D. Lee, J.-D. Huang and J. Yoon, *Angew. Chem., Int. Ed.*, 2018, **57**, 9885–9890.
- 45 N. Kamaly, Z. Xiao, P. M. Valencia, A. F. Radovic-Moreno and
   O. C. Farokhzad, *Chem. Soc. Rev.*, 2012, 41, 2971–3010.
- 46 C. Duan, Y. Zhou, G.-G. Shan, Y. Chen, W. Zhao, D. Yuan,
  L. Zeng, X. Huang and G. Niu, *J. Mater. Chem. C*, 2019, 7, 3471–3478.
- 47 T. Gayathri, S. Karnewar, S. Kotamraju and S. P. Singh, ACS Med. Chem. Lett., 2018, 9, 618–622.
- 48 K. Tanaka, H. Yamane, R. Yoshii and Y. Chujo, *Bioorg. Med. Chem.*, 2013, **21**, 2715–2719.
- 49 P. Gao, W. Pan, N. Li and B. Tang, *ACS Appl. Mater. Interfaces*, 2019, **11**, 26529–26558.