

EDGE ARTICLE

[View Article Online](#)
[View Journal](#) | [View Issue](#)Cite this: *Chem. Sci.*, 2020, **11**, 8610

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 4th June 2020
Accepted 27th July 2020DOI: 10.1039/d0sc03129a
rsc.li/chemical-science

Palladium catalyzed carbonylative generation of potent, pyridine-based acylating electrophiles for the functionalization of arenes to ketones†

Yi Liu, Angela M. Kaiser and Bruce A. Arndtsen*

We describe here the design of a palladium catalyzed route to generate aryl ketones *via* the carbonylative coupling of (hetero)arenes and aryl- or vinyl-triflates. In this, the use of the large bite angle Xantphos ligand on palladium provides a unique avenue to balance the activation of the relatively strong C(sp²)-OTf bond with the ultimate elimination of a new class of potent Friedel–Crafts acylating agent: *N*-acyl pyridinium salts. The latter can be exploited to modulate reactivity and selectivity in carbonylative arene functionalization chemistry, and allow the efficient synthesis of ketones with a diverse array of (hetero)arenes.

Introduction


The design of catalytic methods for the efficient and selective functionalization of aromatic C–H bonds represents an important current thrust in synthetic chemistry.¹ Despite many significant recent advances in this area, adapting these transformations to carbonylative ketone synthesis has seen less success.² Aryl ketones are instead often generated from arenes with traditional Friedel–Crafts acylations using potent acyl electrophiles (Fig. 1a).³ Friedel–Crafts reactions have several attractive features, including their versatility, ability to proceed in an intermolecular fashion, predictable selectivity, and reactivity without directing groups. Nevertheless, a limitation of this chemistry is the need to generate the electrophiles themselves. The latter can require multiple steps and use high energy reagents (*e.g.* thionyl chloride, oxalyl chloride, or carbodiimides).⁴ Moreover, stoichiometric Lewis acids are usually employed to mediate Friedel–Crafts couplings, which leads to compatibility problems with basic functionalities or even decomposition, and generates significant metal-containing waste.⁵ These limitations are of relevance, since aryl ketones are useful motifs in a range of areas, including pharmaceuticals, photosensitizers and polymers.⁶

Recent studies by a number of labs including our own have described how carbonylations can offer an efficient alternative approach to the generation of potent acylating electrophiles.^{7–10} The reactivity of these products has allowed the application of carbonylation reactions to a range of challenging substrates, including the direct functionalization of arenes with acyl triflates.^{8c} However, as with classical Friedel–Crafts chemistry, the


reaction here also requires the use of stoichiometric metal additives: in this case silver triflate salts, in concert with reactive aryl iodides. In addition, the acyl triflates generated are potent electrophiles, and offer little opportunity to control their reactivity without changing the substrate itself.

From an efficiency standpoint, a potential solution to these issues would be to directly carbonylate aryl triflates to aroyl electrophiles. Such a system would obviate the need for metal triflate salts, and also allow the use of broadly available aryl- or


a. Friedel–Crafts Acylation



b. Catalyst Design for Carbonylative Acyl Electrophile Formation



c. This Work:



Department of Chemistry, McGill University, 801 Sherbrooke Street West, Montreal, QC, H3A 0B8, Canada. E-mail: bruce.arndtsen@mcgill.ca

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0sc03129a

Fig. 1 Friedel–Crafts acylations and this work: carbonylative generation of *N*-acyl pyridinium salts for C–H functionalization.

Chem. Sci., 2020, 11, 8610–8616 | 8611




Fig. 3 Mechanistic experiments on the carbonylative formation and reactivity of *N*-acyl pyridinium salts. ^aReactions with acid chlorides were performed in the presence of 1 equivalent of 2,4,6-collidine. ^b1 h.

presumably favor reductive elimination relative to the potential build-up of a labile acyl triflate. However, this is counter-balanced by the ability of these systems to react with arenes. The latter can be seen as well in control experiments (Fig. 3b). The 4-trifluoromethyl substituted pyridinium salt reacts within minutes at ambient temperature with *N*-benzyl pyrrole to form ketone. Replacing the pyrrole with thiophene also leads to ketone (Fig. S4[†]), although the latter requires more pressing conditions than acyl triflates themselves. In contrast, the more electron rich 4-methoxy substituted pyridinium salt does not react with pyrrole at ambient temperature, and requires heating to 80 °C to form product. For comparison, acid chlorides show low reactivity, 14%, under these conditions, and the electron rich 4-dimethylamino substituted pyridinium salt requires even more pressing conditions (130 °C) to form ketone.

Based upon this data, we postulate that this carbonylative functionalization reaction proceeds as shown in Fig. 3c. The

reactivity observed with the Xantphos ligand is presumably related in part to its ability to associate in a bidentate fashion to palladium and better activate the relatively strong C(sp²)-OTf bond towards carbonylation, in concert with its large bite angle, which can favor the reductive elimination.^{8,19} The combination of these features offers an avenue to build-up kinetically reactive acylating agents from C(sp²)-triflates.²⁰ The importance of pyridine, and the specific pyridine structure, in catalysis is tied to its ability to stabilize the reductive elimination product relative to a highly reactive acid triflate. The adducts **1** represent a new class of Friedel-Crafts acylating reagent, and incorporate several attractive features, including the ability to react with (hetero)arenes without added Lewis acids, and tune Friedel-Crafts acylation reactivity with the pyridine employed (*vide infra*).

The *in situ* generation of *N*-acyl pyridinium salts offers a versatile approach to the overall carbonylative conversion of arenes to ketones. As shown in Fig. 4, various pyrroles can be employed in this reaction in concert with 4-methoxypyridine, such as those with *N*-benzyl, -methyl and -aryl substituents. These each lead to the favored formation of the 2-substituted ketone product (**2a–2c**), which is consistent with a Friedel-Crafts mechanism of reaction.^{4a,21} The more sterically encumbered *N*-*tert*-butyl pyrrole is functionalized with high selectivity at the 3-position (**2d–i**). Disubstituted pyrroles such as 2,5-dimethyl-*N*-phenyl pyrrole also undergo efficient reaction to form ketone **2j**. This chemistry can be applied to other nitrogen heterocycles such as substituted indoles, (**2k–n**), and *N*-substituted benzimidazole (**2o**). In addition to the variation of the heteroarenes, a number of aryl triflates can be employed in the reaction, including 4-methoxy, -phenyl (**2g**) and -alkyl substituted (**2h**) reagents. Meta-substituted aryl triflates are tolerated (**2i**, **2m**, and **2n**), albeit with diminished yields. Catalysis also proceeds with less electron rich substrates, such as simple phenyl (**2d**) or naphthyl triflate (**2f**). However, the use of aryl triflates with electron-withdrawing substituents such as 4-cyano and 4-trifluoromethyl substituted aryl triflates leads instead to significant amounts of the phenyl-substituted ketone (**2d**) (44% and 43%, respectively, see Fig. S5[†]). The latter presumably arises from the ability of the palladium-aryl intermediate to scramble substituents with the Xantphos ligand.^{8h–i,22} Similarly, less polar aryl mesylates are not reactive under these conditions (Fig. S6[†]).

We next explored the compatibility of this system with vinyl triflates as a route to access α,β -unsaturated ketones. The transformation here is more efficient than reactions with aryl triflates, and coupling with pyrroles and indoles can proceed at ambient temperature with some substrates (*e.g.* **3f**). As with the aryl triflates, the reaction is compatible with *N*-substituted (**3a**) or 2,5-disubstituted pyrroles (**3b**), and similarly functionalized indoles using cyclic (**3c–e**) or acyclic (**3b** and **3f**) vinyl triflate reagents. These allow access to various α - or β,β -substituted acyclic enones.

In addition, the pyridine unit can be used to modulate electrophilic reactivity. Thus, while less activated (hetero)arenes (*e.g.* thiophenes or substituted arenes) do not lead to product formation using 4-methoxypyridine, the use of the






Fig. 4 Diversity of products available from the Pd catalyzed C–H functionalization of (hetero)arenes with organotriflates. Performed with 2 equiv. aryl or vinyl triflates for pyrroles and indoles, 1 equiv. with other (hetero)arenes, 4 atm CO, 130 °C; 10% (COD)Pd(CH₂TMS)₂, 48 h for aryl triflates; using 5% Pd₂dba₃·CHCl₃, for vinyl triflates. ^aDifficult isolation due to the presence of phenyl ketone **2d**; ^b10% Pd(Xantphos)₂; ^c1.25% Pd₂dba₃·CHCl₃; ^d150 °C; ^e100 °C; ^f80 °C; ^g25 °C.

weakly nucleophilic 4-trifluoromethylpyridine can allow the carbonylative C–H functionalization of various other classes of aromatic reagents. The more reactive vinyl triflates are employed here to allow the build-up of these less stable *N*-acyl pyridinium salts. Examples of hetero(arenes) that can now be functionalized include substituted thiophenes (**3g–l**), furans (**3m**, **3n**), and even substituted arenes, such as 1,3,5-trimethoxybenzene (**3o–s**, **3w**) and anisole (**3t–v**), each of which undergo C–H functionalization to afford the corresponding α,β -unsaturated ketone. This reactivity is compatible with a range of cyclic vinyl triflates (**3g**, **3l**, **3p**). Moreover, acyclic vinyl triflates bearing α - or/and β - (**3b**, **3f**), alkyl- (**3f**, **3k**) or aryl- (**3b**) substituents can also proceed to generate ketone

products. Interestingly, by using 4-trifluoromethylpyridine, the vinyl triflate bond can be selectively activated in the presence of an aryl triflate (**3w**). Together, this system provides access to an array of ketones *via* the palladium catalyzed carbonylative functionalization of arenes with accessible organic triflates.

Finally, we have preliminarily probed the ability of the pyridine to fine-tune product selectivity. One simple effect is through concentration, where lowering the amount of pyridine employed favours the formation of 3-substituted ketone **2e** for pyrroles bearing bulky *N*-substituents (Fig. 5a). The latter may affect the rate of deprotonation of the Wheland intermediate, and thereby allow migration of the aroyl




Fig. 5 Selectivity control in carbonylative C–H functionalization with pyridine employed. ^a1.1 equiv. 4-methoxypyridine, 100 °C, 2.5 h, 17% 3o also formed; ^b5% Pd₂dba₃·CHCl₃, 150 °C, 12 min; ^c5% Pd₂dba₃·CHCl₃, 20 min; ^d10% Pd(Xantphos)₂, 4-trifluoromethylpyridine, 16 h; ^e20 h; ^f4 h; ^g48 h.

fragment to the favoured 3-position under less basic conditions.²³ Alternatively, the pyridine can be used to influence chemoselectivity. An illustration is in the reaction of 1-cyclohexenyltriflate with both *N*-methyl indole and 1,3,5-trimethoxybenzene (Fig. 5b). While the use of 4-trifluoromethylpyridine leads to a mixture of products, 4-methoxypyridine moderates the electrophilicity of the intermediate formed, and allows functionalization to occur exclusively on the more activated indole substrate. The pyridine base can also affect the reactivity of the products (Fig. 5c). As previously noted, the catalytic formation of these products can be performed using 4-methoxypyridine. In contrast, employing the less nucleophilic 4-trifluoromethylpyridine can divert this reaction to instead generate cyclized products of indoles, pyrroles and thiophenes (4a–c). The generation of 4a–c presumably arises from the ability of these systems to undergo acid catalyzed Nazarov cyclization,²⁴ and offers a method to functionalize two C–H bonds at once to build up polycyclic products from heterocycles, CO and vinyl triflates.

Conclusions

In summary, we have described a new approach to the palladium catalyzed carbonylative generation of potent electrophiles with aryl and vinyl triflates for the overall functionalization of arenes. The reaction offers a route to access ketones from combinations of available substrates (organotriflates, (hetero) arenes, CO), and without the typical need for Lewis acids or stoichiometric metal-containing reagents. Mechanistic analysis shows that the Pd/Xantphos catalyst system, in concert with pyridine additives, can provide an unusual avenue to balance the activation of relatively strong C(sp²)–OTf bonds with the ultimate elimination of reactive acylating agents. In this, the pyridine additive serves not only as a base, but more crucially as a building block to drive the formation of a new class of Friedel–Crafts reagent: *N*-acyl pyridinium salts. The latter can be exploited to modulate reactivity and product selectivity in carbonylative functionalization chemistry with a diverse array of (hetero)arenes.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank the Natural Sciences and Engineering Research Council of Canada (NSERC), the Canadian Foundation for Innovation (CFI), McGill University (James McGill Research Fund), and the Fonds de recherche du Québec – Nature et Technologies (FQRNT) supported Centre for Green Chemistry and Catalysis for funding this research.

Notes and references

- (a) T. W. Lyons and M. S. Sanford, *Chem. Rev.*, 2010, **110**, 1147–1169; (b) J. F. Hartwig, *Acc. Chem. Res.*, 2012, **45**, 864–873; (c) X. Chen, K. M. Engle, D. H. Wang and J. Q. Yu, *Angew. Chem., Int. Ed.*, 2009, **48**, 5094–5115; (d) D. A. Colby, R. G. Bergman and J. A. Ellman, *Chem. Rev.*, 2010, **110**, 624–655; (e) V. N. Charushin and O. N. Chupakhin, *Metal Free C–H Functionalization of Aromatics: Nucleophilic Displacement of Hydrogen*, Springer, Berlin, 2014; (f) C. Liu, J. Yuan, M. Gao, S. Tang, W. Li, R. Shi and A. Lei, *Chem. Rev.*, 2015, **115**, 12138–12204; (g) T. Cernak, K. D. Dykstra, S. Tyagarajan, P. Vachal and S. W. Krska, *Chem. Soc. Rev.*, 2016, **45**, 546–576; (h) T. Gensch, M. N. Hopkinson, F. Glorius and J. Wencel-Delord, *Chem. Soc. Rev.*, 2016, **45**, 2900–2936.
- For examples: (a) M. A. Campo and R. C. Larock, *J. Org. Chem.*, 2002, **67**, 5616–5620; (b) X.-F. Wu, P. Anbarasan, H. Neumann and M. Beller, *Angew. Chem., Int. Ed.*, 2010, **49**, 7316–7319; (c) H. Zhang, R. Shi, P. Gan, C. Liu, A. Ding, Q. Wang and A. Lei, *Angew. Chem., Int. Ed.*, 2012, **51**, 5204–5207; (d) F. Zeng and H. Alper, *Org. Lett.*, 2013, **15**, 2034–2037; (e) Z. Lian, S. D. Friis and T. Skrydstrup, *Chem.*



- Commun.*, 2015, **51**, 1870–1873; (f) J. Liu, Z. Wei, H. Jiao, R. Jackstell and M. Beller, *ACS Cent. Sci.*, 2018, **4**, 30–38.
- 3 (a) G. A. Olah, *Friedel–Crafts and Related Reactions*, Interscience Publishers, New York, 1963; (b) G. Sartori and R. Maggi, *Advances in Friedel–Crafts Acylation Reactions: Catalytic and Green Processes*, CRC Press, Boca Raton, 2009.
- 4 (a) D. J. C. Constable, P. J. Dunn, J. D. Hayler, G. R. Humphrey, J. J. L. Leazer, R. J. Linderman, K. Lorenz, J. Manley, B. A. Pearlman, A. Wells, A. Zaks and T. Y. Zhang, *Green Chem.*, 2007, **9**, 411–420; (b) E. Valeur and M. Bradley, *Chem. Soc. Rev.*, 2009, **38**, 606–631; (c) V. R. Pattabiraman and J. W. Bode, *Nature*, 2011, **480**, 471–479.
- 5 J. A. Joule and K. Mills, *Heterocyclic chemistry*, Blackwell Science, Oxford, 5th edn, 2010.
- 6 (a) S.-B. Wu, C. Long and E. J. Kennelly, *Nat. Prod. Rep.*, 2014, **31**, 1158–1174; (b) G. Dorman and G. D. Prestwich, *Biochemistry*, 1994, **33**, 5661–5673; (c) F. Boscá and M. A. Miranda, *J. Photochem. Photobiol., B*, 1998, **43**, 1–26; (d) S. Hirata, *Adv. Optical Mater.*, 2017, **5**, 1700116; (e) W. Zhao, Z. He, J. W. Y. Lam, Q. Peng, H. Ma, Z. Shuai, G. Bai, J. Hao and B. Z. Tang, *Chem*, 2016, **1**, 592–602; (f) H. X. Nguyen and H. Ishida, *Polym. Compos.*, 1987, **8**, 57–73.
- 7 (a) C. F. J. Barnard, *Organometallics*, 2008, **27**, 5402–5422; (b) A. Brennfürer, H. Neumann and M. Beller, *Angew. Chem., Int. Ed.*, 2009, **48**, 4114–4133; (c) R. Grigg and S. P. Mutton, *Tetrahedron*, 2010, **66**, 5515–5548; (d) M. Beller and X.-F. Wu, *Transition Metal Catalyzed Carbonylation Reactions*, Springer, Berlin, 2013; (e) Y. Li, Y. Hu and X.-F. Wu, *Chem. Soc. Rev.*, 2018, **47**, 172–194; (f) D. U. Nielsen, K. T. Neumann, A. T. Lindhardt and T. Skrydstrup, *J. Labelled Compd. Radiopharm.*, 2018, **61**, 949–987; (g) S. Zhang, H. Neumann and M. Beller, *Chem. Soc. Rev.*, 2020, **49**, 3187–3210.
- 8 For recent approaches to acid chlorides/triflates: (a) J. S. Quesnel and B. A. Arndtsen, *J. Am. Chem. Soc.*, 2013, **135**, 16841–16844; (b) J. S. Quesnel, L. V. Kayser, A. Fabrikant and B. A. Arndtsen, *Chem.–Eur. J.*, 2015, **21**, 9550–9555; (c) J. S. Quesnel, S. Moncho, K. E. O. Ylijoki, G. M. Torres, E. N. Brothers, A. A. Bengali and B. A. Arndtsen, *Chem.–Eur. J.*, 2016, **22**, 15107–15118; (d) J. Tjutrins and B. A. Arndtsen, *J. Am. Chem. Soc.*, 2015, **137**, 12050–12054; (e) R. G. Kinney, J. Tjutrins, G. M. Torres, N. J. Liu, O. Kulkarni and B. A. Arndtsen, *Nat. Chem.*, 2018, **10**, 193–199; (f) T. A. Cernak and T. H. Lambert, *J. Am. Chem. Soc.*, 2009, **131**, 3124–3125; (g) X. Fang, B. Cachérat and B. Morandi, *Nat. Chem.*, 2017, **9**, 1105–1109; (h) Y. H. Lee and B. Morandi, *Nat. Chem.*, 2018, **10**, 1016–1022; (i) M. De La Higuera Macias and B. A. Arndtsen, *J. Am. Chem. Soc.*, 2018, **140**, 10140–10144; (j) D. R. Gauthier Jr, N. R. Rivera, H. Yang, D. M. Schultz and C. S. Shultz, *J. Am. Chem. Soc.*, 2018, **140**, 15596–15600.
- 9 For alternative electrophiles, such as acyl fluorides: (a) T. Sakakura, M. Chaisupakitsin, T. Hayashi and M. Tanaka, *J. Organomet. Chem.*, 1987, **334**, 205–211; (b) T. Okano, N. Harada and J. Kiji, *Bull. Chem. Soc. Jpn.*, 1992, **65**, 1741–1743; (c) T. Ueda, H. Konishi and K. Manabe, *Org. Lett.*, 2013, **15**, 5370–5373; Weinreb amides: (d) J. R. Martinelli, D. M. M. Freckmann and S. L. Buchwald, *Org. Lett.*, 2006, **8**, 4843–4846; (e) A. Takács, A. Petz and L. Kollár, *Tetrahedron*, 2010, **66**, 4479–4483; (f) A. Więckowska, R. Fransson, L. R. Odell and M. Larhed, *J. Org. Chem.*, 2011, **76**, 978–981; activated esters: (g) J. R. Martinelli, T. P. Clark, D. A. Watson, R. H. Munday and S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2007, **46**, 8460–8463; (h) T. Ueda, H. Konishi and K. Manabe, *Org. Lett.*, 2012, **14**, 5370–5373; (i) Y. Wang and V. Gevorgyan, *Angew. Chem., Int. Ed.*, 2017, **56**, 3191–3195; thioesters: (j) M. N. Burhardt, R. H. Taaning and T. Skrydstrup, *Org. Lett.*, 2013, **15**, 948–951; acyl azides: (k) F. M. Miloserdov and V. V. Grushin, *Angew. Chem., Int. Ed.*, 2012, **51**, 3668–3672; (l) F. M. Miloserdov, C. L. McMullin, M. M. N. Belmonte, J. Benet-Buchholz, V. I. Bakhmutov, S. A. Macgregor and V. V. Grushin, *Organometallics*, 2014, **33**, 736–752; acyl isocyanate: (m) Z. Li, S. Xu, B. Huang, C. Yuan, W. Chang, B. Fu, L. Jiao, P. Wang and Z. Zhang, *J. Org. Chem.*, 2019, **84**, 9497–9508; acyl-DMAP salts: (n) P.-L. Lagueux-Tremblay, A. Fabrikant and B. A. Arndtsen, *ACS Catal.*, 2018, **8**, 5350–5354.
- 10 Multicomponent applications: (a) G. M. Torres, J. S. Quesnel, D. Bijou and B. A. Arndtsen, *J. Am. Chem. Soc.*, 2016, **138**, 7315–7324; (b) G. M. Torres, M. De La Higuera Macias, J. S. Quesnel, O. P. Williams, V. Yempally, A. A. Bengali and B. A. Arndtsen, *J. Org. Chem.*, 2016, **81**, 12106–12115; (c) J. Tjutrins and B. A. Arndtsen, *Chem. Sci.*, 2017, **8**, 1002–1007; (d) S. Bontemps, J. S. Quesnel, K. Worrall and B. A. Arndtsen, *Angew. Chem., Int. Ed.*, 2011, **50**, 8948–8951.
- 11 (a) L. J. Goossen, N. Rodríguez and C. Linder, *J. Am. Chem. Soc.*, 2008, **130**, 15248–15249; (b) G. J. Lovinger, M. D. Aparece and J. P. Morken, *J. Am. Chem. Soc.*, 2017, **139**, 3153–3160.
- 12 (a) P. J. Stang and W. Treptow, *Synthesis*, 1980, 283–284; (b) K. Ritter, *Synthesis*, 1993, 735–762; (c) H. H. Patel and M. S. Sigman, *J. Am. Chem. Soc.*, 2016, **138**, 14226–14229.
- 13 Y.-R. Luo, *Handbook of bond dissociation energies in organic compounds*, CRC Press, Boca Raton, Fla, 2003.
- 14 (a) S. Cacchi, P. G. Ciattini, E. Morera and G. Ortá, *Tetrahedron Lett.*, 1986, **27**, 3931–3934; (b) A. M. Echavarren and J. Stille, *J. Am. Chem. Soc.*, 1988, **110**, 1557–1565; (c) F. Garrido, S. Raepel, A. Mann and M. Lautens, *Tetrahedron Lett.*, 2001, **42**, 265–266; (d) R. Lou, M. VanAlstine, X. Sun and M. P. Wentland, *Tetrahedron Lett.*, 2003, **44**, 2477–2480; (e) A. Brennfürer, H. Neumann and M. Beller, *Synlett*, 2007, 2537–2540; (f) J. Lindh, A. Fardost, M. Almeida and P. Nilsson, *Tetrahedron Lett.*, 2010, **51**, 2470–2472; (g) X.-F. Wu, B. Sundararaju, H. Neumann, P. H. Dixneuf and M. Beller, *Chem.–Eur. J.*, 2011, **17**, 106–110; (h) P. B. Wakchaure, S. R. Borhade, A. Sandström and P. I. Arvidsson, *Eur. J. Org. Chem.*, 2015, 213–219; (i) J. T. Joseph, A. M. Sajith, R. C. Ningegowda and S. Shashikanth, *Adv. Synth. Catal.*, 2017, **359**, 419–425; (j) C. Shen, Z. Wei, H. Jiao and X.-F. Wu, *Chem.–Eur. J.*, 2017, **23**, 13369–13378; (k) Y. Yuan and X.-F. Wu, *Org. Lett.*, 2019, **21**, 5310–5314.



- 15 P. W. N. M. van Leeuwen and P. C. J. Kamer, *Catal. Sci. Technol.*, 2018, **8**, 26–113.
- 16 The low volatility 4-methoxypyridine was the most convenient to employ and used for subsequent studies.
- 17 (a) G. A. Olah and P. J. Szilagyi, *J. Am. Chem. Soc.*, 1969, **91**, 2949–2955; (b) C. Lohse, S. Hollenstein and T. Laube, *Angew. Chem., Int. Ed.*, 1991, **30**, 1656–1658.
- 18 For selected carbonylations with vinyl triflates: (a) G. Crisp, W. J. Scott and J. K. Stille, *J. Am. Chem. Soc.*, 1984, **106**, 7500–7506; (b) R. Anacardio, A. Arcadi, G. D'Anniballe and F. Marinelli, *Synthesis*, 1995, 831–836; (c) G. T. Crisp and A. G. Meyer, *Tetrahedron*, 1995, **51**, 5585–5596; (d) T. Luker, H. Hiemstra and W. N. Speckamp, *J. Org. Chem.*, 1997, **62**, 8131–8140; (e) P. Larini, A. Guarna and E. G. Occhiato, *Org. Lett.*, 2006, **8**, 781–784; (f) S. Zhang, H. Neumann and M. Beller, *Org. Lett.*, 2019, **21**, 3528–3532.
- 19 For reviews in large bite angle diphosphines: (a) P. W. N. M. van Leeuwen, P. C. J. Kamer, J. N. H. Reek and P. Dierkes, *Chem. Rev.*, 2000, **100**, 2741–2770; (b) P. C. J. Kamer, P. W. N. M. van Leeuwen and J. N. H. Reek, *Acc. Chem. Res.*, 2001, **34**, 895–904; (c) P. W. N. M. van Leeuwen and Z. Freixa, in *Modern Carbonylation Methods*, ed. L. Kollár, Wiley-VCH, Weinheim, 2008, pp. 1–25; (d) G. M. Adams and A. S. Weller, *Coord. Chem. Rev.*, 2018, **355**, 150–172.
- 20 Other mechanistic pathways cannot be ruled-out for reactions with aryl triflates, including the direct reaction of heterocycles with the palladium-acyl intermediate, since we do not observe the *in situ* build-up of acyl-pyridinium salts with these reagents.
- 21 (a) A. R. Katritzky, K. Suzuki, S. K. Singh and H.-Y. He, *J. Org. Chem.*, 2003, **68**, 5720–5723; (b) F. He, H. Wu, J. Chen and W. Su, *Synth. Commun.*, 2008, **38**, 255–264; (c) J. E. Taylor, M. D. Jones, J. M. J. Williams and S. D. Bull, *Org. Lett.*, 2010, **12**, 5740–5743.
- 22 (a) B. Segelstein, T. Butler and B. Chenard, *J. Org. Chem.*, 1995, **60**, 12–13; (b) F. E. Goodson, T. I. Wallow and B. M. Novak, *J. Am. Chem. Soc.*, 1997, **119**, 12441–12453; (c) Y. H. Lee and B. Morandi, *Coord. Chem. Rev.*, 2019, **386**, 96–118.
- 23 J. R. Carson and N. M. Davis, *J. Org. Chem.*, 1981, **46**, 839–843.
- 24 (a) W. Yin, Y. Ma, J. Xu and Y. Zhao, *J. Org. Chem.*, 2006, **71**, 4312–4315; (b) T. Vaidya, R. Eisenberg and A. J. Frontier, *ChemCatChem*, 2011, **3**, 1531–1548.

