Chemical Science

EDGE ARTICLE

Check for updates

Cite this: Chem. Sci., 2020, 11, 12124

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 22nd July 2020 Accepted 28th September 2020

DOI: 10.1039/d0sc04007j

rsc.li/chemical-science

Rh(III)-catalyzed tandem annulative redox-neutral arylation/amidation of aromatic tethered alkenes[†]

Chao Chen,^{‡ab} Chen Shi,^{‡ab} Yaxi Yang^{abc} and Bing Zhou ^(b)*^{abc}

Transition-metal-catalyzed directed C–H functionalization has emerged as a powerful and straightforward tool to construct C–C bonds and C–N bonds. Among these processes, the intramolecular annulative alkene hydroarylation reaction has received much attention because this intramolecular annulation can produce more complex and high value-added structural motifs found in numerous natural products and bioactive molecules. Despite remarkable progress, these annulative protocols developed to date remain limited to hydroarylation and functionalization of one side of alkenes, thus largely limiting the structural diversity and complexity. Herein, we developed a rhodium(III)-catalyzed tandem annulative arylation/amidation reaction of aromatic tethered alkenes to deliver a variety of 2,3-dihydro-3-benzofuranmethanamine derivatives bearing an all-carbon quaternary stereo center by employing 3-substituted 1,4,2-dioxazol-5-ones as an amidating reagent to capture the transient C(sp³)–Rh intermediate. Notably, by simply changing the directing group, a second, unsymmetrical *ortho* C–H amidation/annulation can be achieved to provide tricyclic dihydrofuro[3,2-f]quinazolinones in good yields.

Transition-metal-catalyzed C-H functionalization for the direct conversion of C-H bonds to C-C bonds and C-N bonds has evolved into a widespread and effective strategy for fine chemical production.¹ Among these processes, the hydroarylation of C-C double bonds via a C-H addition has been well-established and become an effective strategy to access synthetically useful structural motifs.1 Recently, this alkene hydroarylation reaction has received much attention in an intramolecular fashion² (Scheme 1a) because this intramolecular annulation can produce more complex and high value-added structural motifs found in numerous natural products and bioactive molecules (Fig. 1).³ Despite remarkable progress in this area, most of the annulative protocols developed to date remain limited to onecomponent intramolecular alkene hydroarylation and functionalization of one side of alkenes. More challenging C-H arylation of intramolecular alkenes followed by a tandem coupling with a different coupling partner have unfortunately proven elusive thus far, thus largely limiting the structural diversity and complexity.



On the other hand, nitrogen-containing molecules have gained great attention due to their widespread presence in

ROYAL SOCIETY

OF CHEMISTRY

View Article Online

View Journal | View Issue

In this context, we wondered if a catalytic annulative C-H arylation of a O-bearing olefin-tethered arenes might be



Scheme 1 Transition-metal-catalyzed C–H functionalization to construct the C–C and C–N bonds.

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

Dpen Access Article. Published on 16 October 2020. Downloaded on 7/24/2025 7:07:48 PM.

[&]quot;State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 555 Zu Chong Zhi Road, Shanghai 201203, China. E-mail: zhoubing@simm.ac.cn; zhoubing2012@hotmail.com

^bUniversity of Chinese Academy of Sciences, 19 Yuquan Road, Beijing 100049, China ^cSchool of Pharmaceutical Science and Technology, Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou 310024, China

[†] Electronic supplementary information (ESI) available. CCDC 2015893 and 2014245. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d0sc04007j

[‡] These authors contributed equally.



Fig. 1 Representative bioactive 2,3-dihydrobenzofurans.

possible, thus leading to a C(sp³)–M intermediate, which upon capture with a amidation reagent to construct a new C–N bond and provide bioactive 2,3-dihydro-3-benzofuranmethanamine derivatives. Inherently, the tandem annulative 1,2-arylation/ amidation of alkenes has several challenges. First, the resulting C(alkyl)–M intermediate is liable to undergo protonation to provide the alkene hydroarylation products.^{1,2} Moreover, a potential competing β -H elimination of the resulting C(alkyl)– M intermediate also required to be suppressed. In addition, compared with the C(sp²)–M species, the resulting C(alkyl)–M species is relatively unstable and also has a low reactivity.

To address these challenges and with our continuing interest in the Rh(m)-catalyzed C–H functionalization,⁸ we introduced a Weinreb amide as a directing group and 3-substituted 1,4,2dioxazol-5-ones as the amide sources⁹ to trigger a new tandem annulative 1,2-arylation/amidation of alkenes *via* a Rh(m)-catalyzed C–H activation,¹⁰ providing a variety of synthetically challenging 2,3-dihydro-3-benzofuranmethanamine derivatives bearing an all-carbon quaternary stereo center (Scheme 1b). More importantly, through simply changing the directing group, a second, unsymmetrical *ortho* C–H amidation/ annulation could be realized to provide tricyclic dihydrofuro [3,2-*f*]quinazolinone derivatives. This protocol provides a good complement to previously reported carboamination reactions.¹¹

Table 1 Optimization of reaction conditions^a

To begin our studies, Weinreb amide 1a was reacted with methyl dioxazolone 3a in the presence of various catalyst and AgSbF₆ at 70 °C in DCE (Table 1, entries 1-4). The use of [Cp*RhCl₂]₂ as the catalyst was found to be crucial to give the desired tandem annulative product 4a, with other catalysts, such as $[Ru(p-cymene)Cl_2]_2$, $[Cp*IrCl_2]_2$, and $Cp*Co(CO)I_2$, resulting in no desired product. Attempt to increase or lower the reaction temperature led to a slightly low yield (entries 5 and 6). Interestingly, when employing a NH-OMe amide 2a as the substrate and using 3 equivalent of 3a, a second, unsymmetrical ortho C-H amidation/annulation was achieved to provide the tricyclic dihydrofuro[3,2-f]quinazolinone 5a in 50% yield (entry 7). A screen of additives (entries 8-10) identified LiOAc as the optimal additive, affording the desired product 5a in 93% yield (entry 10). The Rh(III) catalyst was found to be crucial for this tandem annulative arylation/amidation reaction, with no reactivity in its absence (entries 11 and 12).

Having determined the optimal reaction conditions, we sought to evaluate the substrate scope (Scheme 2). First, the amidation reagents were explored and 1,4,2-dioxazol-5-ones substituted with primary alkyl (4a and 4e), secondary alky (4b, 4c and 4f), tertiary alkyl (4d) and aryl group (4g-i) all coupled smoothly with 1a, providing the 2,3-dihydro-3benzofuranmethanamines 4a-4j in good yields. The structure of 4f was unambiguously confirmed by an X-ray crystallographic analysis (CCDC 2015893). The scope with regards to the arene moiety was then examined. The substrates 1 containing either electron-donating or electron-withdrawing substituents at different positions on the arene ring were well tolerated and provide the desired products 4k-t in good yields. We were pleased that 2-naphthalenecarboxamide effectively underwent this tandem annulative 1,2-arylation/amidation reaction,

H = H = M = M = M = M = M = M = M = M =							
Entry	х	Catalyst (5 mol%)	Additive (20 mol%)	Solvent	Т (°С)	Yield of 4a (%)	Yield of 5 a (%)
1	Ме	[Cp*RhCl ₂] ₂	_	DCE	70	89	0
2	Me	$[Ru(p-cymene)Cl_2]_2$	_	DCE	70	0	0
3	Ме	[Cp*IrCl ₂] ₂	_	DCE	70	0	0
4	Ме	$Cp*Co(CO)I_2$	_	DCE	70	0	0
5	Ме	[Cp*RhCl ₂] ₂		DCE	90	69	0
6	Ме	[Cp*RhCl ₂] ₂	_	DCE	50	70	0
7 ^b	Н	[Cp*RhCl ₂] ₂		DCE	70	0	50
8 ^b	Н	[Cp*RhCl ₂] ₂	$Cu(OAc)_2$	DCE	70	0	87
9^b	Н	[Cp*RhCl ₂] ₂	KOAc	DCE	70	0	86
10^b	Н	[Cp*RhCl ₂] ₂	LiOAc	DCE	70	0	93
11^b	Н		LiOAc	DCE	70	0	0
12	Me	_	_	DCE	70	0	0

^{*a*} Conditions: **1a** (0.1 mmol), **3a** (0.12 mmol), catalyst (5 mol%), AgSbF₆ (20 mol%) and additive (20 mol%) in DCE (1 mL) for 12 h. Yield isolated by column chromatography. ^{*b*} Conditions: **2a** (0.1 mmol), **3a** (0.3 mmol), catalyst (5 mol%), AgSbF₆ (20 mol%), additive (20 mol%) in DCE (1 mL) for 12 h. Yield isolated by column chromatography.



Scheme 2 Substrate scope of tandem annulative arylation/amidation reaction of aromatic tethered alkenes. Conditions: 1 (0.1 mmol), 3 (0.12 mmol), [Cp*RhCl₂]₂ (5 mol%), AgSbF₆ (20 mol%) in DCE (1 mL) at 70 °C for 12 h. Yield isolated by column chromatography.

affording the desired product **4r** in good yield. Notably, the various substituted allyl groups such as ethyl, cyclopentyl, phenyl, and phenoxymethyl groups were found to be compatible with the reaction conditions (**4u**-**x**). In addition, 3-N-tethered and 3-S-tethered substrates failed to give the desired tandem annulative products **4z** and **4za**.

Next, we proceeded to explore the scope of this unsymmetrical twofold C-H functionalization reaction (Scheme 3). Under the optimal reaction conditions, amidating reagents bearing alkyl or aryl groups are fully tolerated, affording the tricyclic dihydrofuro[3,2-*f*]quinazolinones **5a**-**5h** in good yields. The structure of **5e** was unambiguously confirmed by an X-ray crystallographic analysis (CCDC 2014245). Electronic and steric modification of the aryl group was also tolerated. Both electron-deficient (**5j**, **5m**-**o**) and electron-rich (**5i**, **5k**, **5l**, **5q** and **5r**) substrates gave the corresponding tricyclic systems in good yields. *Meta* and *para* substitutions of a methyl group were also tolerated and delivered the products **5i** and **5k**, indicating a high tolerance for steric hindrance. Interestingly, when 2-naphthalenecarboxamide was used, a third C-H amidation of



Scheme 3 Substrate scope of unsymmetrical twofold C–H functionalization reaction. Conditions: 2 (0.1 mmol), 3 (0.30 mmol), $[Cp*RhCl_2]_2$ (5 mol%), AgSbF₆ (20 mol%), LiOAc (20 mol%) and DCE (1 mL) at 70 °C for 12 h. Yield isolated by column chromatography.

naphthalene ring took place, affording the product **5s** in 65% yield. Notably, the current method effectively resulted in the ethyl-, cyclopentyl, phenyl, and phenoxymethyl-substituted products **5t-w** bearing an all-carbon quaternary stereo center in good yield, respectively.

To check the practicability of this protocol, this two procedures could be readily scaled up with comparable efficiency in the presence of 2.5 mol% of Rh(m) catalyst on a 2.0 mmol scale (eqn (1) and (2)). The product **5a** could be readily converted into potential useful intermediates, such as amines **6a** and free amino quinazolinone analog **6b**, respectively (eqn (3)).



To gain insight into the reaction mechanism, hydrogen/ deuterium (H/D) exchange were carried out. A H/D exchange at the *ortho*-position of the amide group in the re-isolated 1a and 2a was observed in the absence or presence of 3a, indicative of the reversibility of the *ortho* C–H activation (eqn (4)–(6)). Treatment of 2a with 1 equivalent of 3d at room temperature for 1 h delivered the 6d as the sole product, indicating that the intramolecular tandem annulative 1,2-arylation/amidation of alkenes is faster than *ortho* C–H amidation/annulation (eqn (7)). In addition, the use of 3 equivalent of 3d at 70 °C for 2 h provided 6e as the main product and subsequent treatment of 6e under the standard conditions gave 5d in 60% yield (eqn (8)), indicating that the second *ortho* C–H amidation to give the desired quinazolinone product. Finally, treatment of substrate 7 with 3a under the standard reaction conditions did not give any product 8, ruling out the possibility of the insertion of a nitrene to double bond (eqn (9)).



Based on above-mentioned experimental results, a plausible reaction pathway is proposed in Scheme 4. $[Cp*RhCl_2]_2$ precursor reacts with AgSbF₆ to form an active cationic Rh(m)



Scheme 4 Proposed reaction mechanism.

species, which undergoes a C–H bond activation to form cyclometalated complex Int-A. Coordination of the tethered olefin and a subsequent migratory insertion affords the intermediate Int-B, which undergoes an oxidative addition into the N–O bond of **3a**, followed by a CO₂ extrusion, to provide the Rh(v) nitrenoid species Int-C. Reductive elimination occurs to deliver the intermediate Int-D which then is protonated to release product **4a** or Int-E and regenerate the catalyst. Int-E can undergo a second *ortho* C–H activation to give Int-F, which can be oxidized by **3a** again to afford the Rh(v) nitrenoid species Int-G, with a CO₂ extrusion. Subsequent reductive elimination and protonation give the Int-I which undergoes an intramolecular dehydration to deliver the product **5a**.

Conclusions

In summary, we have developed an unprecedented rhodiumcatalyzed tandem annulative arylation/amidation reaction of aromatic tethered alkenes by using 3-substituted 1,4,2-dioxazol-5-ones as an amidating reagent. This robust transformation proceeds with a broad functional group tolerance under relatively mild and redox-neutral reaction conditions, releasing CO_2 as the single byproduct. A wide variety of 2,3-dihydro-3benzofuranmethanamine derivatives bearing an all-carbon quaternary stereo center can be accessed with high yields. Notably, by simply changing the directing group, a second, unsymmetrical *ortho* C–H amidation/annulation can be achieved to provide tricyclic dihydrofuro[3,2-*f*]quinazolinone derivatives in good yields.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work is financially supported by National Natural Science Foundation of China (No. 21702218, 81973166, 91753207, 81773568), the National major science and technology project "major new drug creation" (2018ZX09711002-006-001, 2018ZX09711002-008-005), Youth Innovation Promotion Association (2017333), Science and Technology Commission of Shanghai Municipality (18431907100) and K. C. Wong Education Foundation.

References

 Selected reviews on transition metal-catalyzed C-H bond funtionalization: (a) P. Gandeepan, T. Mgller, D. Zell, G. Cera, S. Warratz and L. Ackermann, *Chem. Rev.*, 2019, **119**, 2192-2452; (b) R. Hummel, J. A. Boerth and J. A. Ellman, *Chem. Rev.*, 2017, **117**, 9163-9227; (c) Y. Park, Y. Kim and S. Chang, *Chem. Rev.*, 2017, **117**, 9247-9301; (d) Z. Dong, Z. Ren, S. J. Thompson, Y. Xu and G. Dong, *Chem. Rev.*, 2017, **117**, 9333-9403; (e) J. He, M. Wasa, K. S. L. Chan, Q. Shao and J.-Q. Yu, *Chem. Rev.*, 2017, **117**, 8754-8786; (f) D.-S. Kim, W.-J. Park and C.-Ho. Jun, *Chem.* Rev., 2017, 117, 8977–9015; (g) T. Gensch, M. N. Hopkinson,
F. Glorius and J. Wencel-Delord, *Chem. Soc. Rev.*, 2016, 45, 2900–2936; (h) L. C. M. Castro and N. Chatani, *Chem. Lett.*, 2015, 44, 410–421; (i) J. Wencel-Delord and F. Glorius, *Nat. Chem.*, 2013, 5, 369–375; (j) J. Yamaguchi, A. D. Yamaguchi and K. Itami, *Angew. Chem., Int. Ed.*, 2012, 51, 8960–9009; (k) T. W. Lyons and M. S. Sanford, *Chem. Rev.*, 2010, 110, 1147–1169.

- 2 Selected examples: (a) R. K. Thalji, K. A. Ahrendt, R. G. Bergman and J. A. Ellman, J. Am. Chem. Soc., 2001, 123, 9692-9693; (b) T. A. Davis, T. K. Hyster and T. Rovis, Angew. Chem., Int. Ed., 2013, 52, 14181-14185; (c) T. A. Davis, C. Wang and T. Rovis, Synlett, 2015, 26, 1520-1524; (d) B. Ye, P. A. Donets and N. Cramer, Angew. Chem., Int. Ed., 2014, 53, 507-511; (e) Z. Shi, M. Boultadakis-Arapinis, D. C. Koester and F. Glorius, Chem. Commun., 2014, 50, 2650-2652; (f) K. Ghosh, R. K. Rit, E. Ramesh and A. K. Sahoo, Angew. Chem., Int. Ed., 2016, 55, 7821-7825; (g) Z. Ding and N. Yoshikai, Angew. Chem., Int. Ed., 2013, 52, 8574-8578; (h) D. F. Fernández, M. Gulías, J. L. Mascareñas and F. López, Angew. Chem., Int. Ed., 2017, 56, 9541-9545; (i) Z. Guan, S. Chen, Y. Huang and H. Yao, Org. Lett., 2019, 21, 3959-3962; (j) K. Ghosh, M. Shankar, R. K. Rit, G. Dubey, P. V. Bharatam and A. K. Sahoo, J. Org. Chem., 2018, 83, 9667-9681; (k) K. Mukherjee, E. Ramesh, K. Ghosh and A. K. Sahoo, Asian J. Org. Chem., 2018, 7, 1380-1384; (l) K. Ghosh, A. Ghosh, K. Mukherjee, R. K. Rit and A. K. Sahoo, J. Org. Chem., 2020, 85, 8618-8626.
- 3 (a) M. N. Attala and P. Diaz, PCT, WO /012221 A1, 2009; (b)
 A. Stoit, W. I. Iwema Barker, H. K. A. C. Coolen,
 M. J. P. Van Dongen and N. J.-L. D. Leflemme, PCT, WO /004378 A1, 2012; (c)
 S. Chowdhury, J. Fu, R. Kamboj,
 S. Liu, Q. Jia, V. Raina and J. Sun, PCT, WO/046087 A2, 2008.
- 4 (*a*) E. Vitaku, D. T. Smith and J. T. Njardarson, *J. Med. Chem.*, 2014, **57**, 10257–10274; (*b*) R. Hili and A. K. Yudin, *Nat. Chem. Biol.*, 2006, **2**, 284–287; (*c*) B. M. Dancy and P. A. Cole, *Chem. Rev.*, 2015, **115**, 2419–2452.
- 5 Selected examples: (a) J. Y. Kim, S. H. Park, J. Ryu, S. H. Cho, S. H. Kim and S. Chang, J. Am. Chem. Soc., 2012, 134, 9110-9113; (b) J. Ryu, K. Shin, S. H. Park, J. Y. Kim and S. Chang, Angew. Chem., Int. Ed., 2012, 51, 9904-9908; (c) S. Yu, B. Wan and X. Li, Org. Lett., 2013, 15, 3706-3709; (d) C. Grohmann, H. Wang and F. Glorius, Org. Lett., 2013, 15, 3014-3017; (e) D.-G. Yu, M. Suri and F. Glorius, J. Am. Chem. Soc., 2013, 135, 8802-8805; (f) H. Zhao, Y. Shang and W. Su, Org. Lett., 2013, 15, 5106-5109; (g) K.-H. Ng, Z. Zhou and W.-Y. Yu, Chem. Commun., 2013, 49, 7031-7033; (h) B. Zhou, J. Du, Y. Yang, H. Feng and Y. Li, Org. Lett., 2013, 15, 6302-6305; (i) J. Du, Y. Yang, H. Feng, Y. Li and B. Zhou, Chem.-Eur. J., 2014, 20, 5727-5731; (j) B. Zhou, J. Du, Y. Yang, H. Feng and Y. Li, Org. Lett., 2014, 16, 592-595; (k) S. H. Park, J. Kwak, K. Shin, J. Ryu, Y. Park and S. Chang, J. Am. Chem. Soc., 2014, 136, 2492–2502; (l) H. Kim, K. Shin and S. Chang, J. Am. Chem. Soc., 2014, 136, 5904-5907; (m) C. Suzuki, K. Hirano, T. Satoh and M. Miura, Org. Lett., 2015, 17, 1597-1600; (n) G. B. Boursalian, M.-Y. Ngai, K. N. Hojczyk and T. Ritter, J. Am. Chem. Soc., 2013, 135,

13278–13281; (*o*) D. Zhu, G. Yang, J. He, L. Chu, G. Chen, W. Gong, K. Chen, M. D. Eastgate and J.-Q. Yu, *Angew. Chem.*, *Int. Ed.*, 2015, **54**, 2497–2500.

- 6 For selected reviews: (a) H. M. L. Davies and J. R. Manning, Nature, 2008, 451, 417–424; (b) J. L. Roizen, M. E. Harvey and J. Du Bois, Acc. Chem. Res., 2012, 45, 911–922; (c) T. A. Ramirez, B. Zhao and Y. Shi, Chem. Soc. Rev., 2012, 41, 931–942.
- 7 For examples:(a) H.-Y. Thu, W.-Y. Yu and C.-M. Che, J. Am. Chem. Soc., 2006, 128, 9048-9049; (b) E. T. Nadres and O. Daugulis, J. Am. Chem. Soc., 2012, 134, 7-10; (c) T. Kang, Y. Kim, D. Lee, Z. Wang and S. Chang, J. Am. Chem. Soc., 2014, 136, 4141–4144; (d) X. Huang, Y. Wang, J. Lan and J. You, Angew. Chem., Int. Ed., 2015, 54, 9404-9408; (e) M. Yang, B. Su, Y. Wang, K. Chen, X. Jiang, Y.-F. Zhang, X.-S. Zhang, G. Chen, Y. Cheng, Z. Cao, Q.-Y. Guo, L. Wang and Z.-J. Shi, Nat. Commun., 2014, 5, 4707-4713; (f) Q. Zhang, K. Chen, W. Rao, Y. Zhang, F.-J. Chen and B.-F. Shi, Angew. Chem., Int. Ed., 2013, 52, 13588-13592; (g) Z. Wang, J. Ni, Y. Kuninobu and M. Kanai, Angew. Chem., Int. Ed., 2014, 53, 3496-3499; (h) X. Wu, Y. Zhao, G. Zhang and H. Ge, Angew. Chem., Int. Ed., 2014, 53, 3706-3710; (i) S. M. Paradine, J. R. Griffin, J. Zhao, A. L. Petronico, S. M. Miller and M. C. White, Nat. Chem., 2015, 7, 987-994; (j) J. He, T. Shigenari and J.-Q. Yu, Angew. Chem., Int. Ed., 2015, 54, 6545-6549.
- 8 (a) Y. Wu, Z. Chen, Y. Yang, W. Zhu and B. Zhou, J. Am. Chem. Soc., 2018, 140, 42–45; (b) B. Zhou, Z. Chen, Y. Yang, W. Ai, H. Tang, Y. Wu, W. Zhu and Y. Li, Angew. Chem., Int. Ed., 2015, 54, 12121–12126; (c) Y. Yang, X. Wang, Y. Li and B. Zhou, Angew. Chem., Int. Ed., 2015, 54, 15400–15404.
- 9 (a) Y. Park, K. T. Park, J. G. Kim and S. Chang, J. Am. Chem. Soc., 2015, 137, 4534-4542; (b) H. Wang, G. Tang and X. Li, Angew. Chem., Int. Ed., 2015, 54, 13049-13052; (c)
 S. Y. Hong, Y. Park, Y. Hwang, Y. B. Kim, M.-H. Baik and
 S. Chang, Science, 2018, 359, 1016-1021; (d) H. Lei and
 T. Rovis, J. Am. Chem. Soc., 2019, 141, 2268-2273; (e)
 T. Knecht, S. Mondal, J.-H. Ye, M. Das and F. Glorius, Angew. Chem., Int. Ed., 2019, 58, 7117-7121; (f)
 J. S. Burman, R. J. Harris, C. M. B. Farr, J. Bacsa and
 S. B. Blakey, ACS Catal., 2019, 9, 5474-5479.
- 10 For reviews on Rh(III)-catalyzed C-H activations, see: (a)
 D. A. Colby, R. G. Bergman and J. A. Ellman, Chem. Rev., 2010, 110, 624-655; (b) T. Satoh and M. Miura, Chem.-Eur. J., 2010, 16, 11212-11222; (c) G. Song, F. Wang and X. Li, Chem. Soc. Rev., 2012, 41, 3651-3678; (d) D. A. Colby, A. S. Tsai, R. G. Bergman and J. A. Ellman, Acc. Chem. Res., 2012, 45, 814-825; (e) F. W. Patureau, J. Wencel-Delord and F. Glorius, Aldrichimica Acta, 2012, 45, 31-41; (f) N. Kuhl, N. Schröder and F. Glorius, Adv. Synth. Catal., 2014, 356, 1443-1460; (g) G. Song and X. Li, Acc. Chem. Res., 2015, 48, 1308-1318; (i) S. Vasquez-Cespedes, X. Wang and F. Glorius, ACS Catal., 2018, 8, 242-257; (j) T. Piou and T. Rovis, Acc. Chem. Res., 2018, 51, 170-180.
- 11 For selected examples of non-annulative carboamination reactions, see:(*a*) T. Piou and T. Rovis, *Nature*, 2015, **527**,

86-90; (b) A. Lerchen, T. Knecht, C. G. Daniliuc and F. Glorius, Angew. Chem., Int. Ed., 2016, 55, 15166-15170;
(c) J. E. Ney and J. P. Wolfe, Angew. Chem., Int. Ed., 2004, 43, 3605-3608; (d) R. Lira and J. P. Wolfe, J. Am. Chem. Soc., 2004, 126, 13906-13907; (e) D. N. Mai and J. P. Wolfe, J. Am. Chem. Soc., 2010, 132, 12157-12159; (f) D. R. White, J. T. Hutt and J. P. Wolfe, J. Am. Chem. Soc., 2015, 137, 11246-11249; (g) W. Zeng and S. R. Chemler, J. Am. Chem. Soc., 2007, 129, 12948-12949; (h) C. F. Rosewall, P. A. Sibbald, D. V. Liskin and F. E. Michael, J. Am. Chem. Soc., 2009, 131, 9488-9489; (i) G. Zhang, L. Cui, Y. Wang

and L. Zhang, J. Am. Chem. Soc., 2010, 132, 1474–1475; (j) T. W. Liwosz and S. R. Chemler, J. Am. Chem. Soc., 2012, 134, 2020–2023; (k) A. Faulkner, J. S. Scott and J. F. Bower, J. Am. Chem. Soc., 2015, 137, 7224–7230; (l) T. Pinkert, T. Wegner, S. Mondal and F. Glorius, Angew. Chem., Int. Ed., 2019, 58, 15041–15045; (m) S. Maity, T. J. Potter and J. A. Ellman, Nat. Catal., 2019, 2, 756–762; (n) D. Zhao, S. Vasquez-Cespedes and F. Glorius, Angew. Chem., Int. Ed., 2015, 54, 1657–1661; (o) T. K. Hyster, L. Knorr, T. R. Ward and T. Rovis, Science, 2012, 338, 500–503.