


Showcasing research from Professor Nimse's laboratory, Department of Chemistry, Hallym University, Chuncheon, South Korea.

The detection of Al^{3+} and Cu^{2+} ions using isonicotinohydrazide-based chemosensors and their application to live-cell imaging

A novel receptor (3) synthesized from isonicotinic hydrazide and 2,4-dihydroxybenzaldehyde in ethanol demonstrated excellent selectivity and sensitivity towards Cu^{2+} ($1.3 \times 10^4 \text{ M}^{-1}$) and Al^{3+} (K_a of $4.8 \times 10^4 \text{ M}^{-1}$). Receptor 3 showed the LOD of 3.0 nM for Al^{3+} and LOD of 1.9 μM for Cu^{2+} . The quick response, easy-synthesis, and high sensitivity make receptor 3 an ideal sensor for detecting Al^{3+} ions in a semi-aqueous medium and living cells.

As featured in:



See Satish Balasaheb Nimse, Anil Kuwar *et al.*, *Mater. Adv.*, 2021, 2, 6306.

Cite this: *Mater. Adv.*, 2021,
2, 6306

The detection of Al³⁺ and Cu²⁺ ions using isonicotinohydrazide-based chemosensors and their application to live-cell imaging†

In-ho Song,^{‡,a} Pritam Torawane,^{§,b} Jung-Seop Lee,^a Shrikant Dashrath Warkad,^c Amulrao Borase,^b Suban K. Sahoo,^{§,d} Satish Balasaheb Nimse^{§,a} and Anil Kuwar^{§,b}

A new Schiff base receptor (**3**) was synthesized by an equimolar reaction between isonicotinic hydrazide and 2,4-dihydroxybenzaldehyde in ethanol. Receptor **3** demonstrated excellent selectivity and sensitivity towards Cu²⁺ ions and Al³⁺ ions by UV-vis absorption spectroscopy and fluorescence spectroscopy, respectively. Receptor **3** showed a detectable color change from colorless to yellow with a red-shift ($\Delta\lambda \approx 70$ nm) in the absorption spectra in the presence of Cu²⁺. In the emission study of **3**, Al³⁺ showed significant fluorescent enhancement ($\lambda_{em} = 473$ nm) over a wide range of tested metal ions. The quantum yield of receptor **3** ($\Phi = 0.0021$) increases ~ 230 folds in the presence of Al³⁺ ions to form receptor **3**·Al³⁺ complex ($\Phi = 0.484$). Receptor **3** showed high selectivity for Al³⁺ with a K_a of 4.8×10^4 M⁻¹ and LOD of 3.0 nM. In comparison, K_a for Cu²⁺ was 1.3×10^4 M⁻¹ and LOD of 1.9 μ M. Receptor **3** is an excellent chemosensor for detecting Al³⁺ ions as indicated by its nanomolar range LOD. The quick response, easy-synthesis, and high sensitivity make receptor **3** an ideal sensor for detecting Al³⁺ ions in a semi-aqueous medium and living cells.

Received 29th June 2021,
Accepted 15th August 2021

DOI: 10.1039/d1ma00564b

rsc.li/materials-advances

1. Introduction

Development of highly specific and sensitive chemosensors has attracted significant interest in detecting bioactive metal ions and toxic metal ions due to their importance in chemical, biological, medical, material, and environmental sciences.^{1–4} Development of colorimetric and fluorescence-based detection methods that target precise recognition of bioactive metals such as copper (Cu²⁺) and aluminum (Al³⁺) is a topic of high interest among researchers.^{5–11}

Cu²⁺ is the third most abundant heavy metal after Fe³⁺ and Zn²⁺ in the human body. Several biological processes require Cu²⁺ in optimum amounts.^{12–15} Enzymes such as tyrosinase, lysyl oxidase, cytochrome *c* oxidase, and superoxide dismutase

require Cu²⁺ for redox reactions.^{16,17} Even though Cu²⁺ is potentially toxic, it is an essential element.¹⁸ The deficiency of Cu²⁺ leads to Menkes disease.^{19,20} Whereas, the accumulation of Cu²⁺ is correlated to the Wilson disease,^{21,22} Amyotrophic Lateral Sclerosis,^{23,24} and Alzheimer's disease.^{25,26} The crucial physiological relevance of Cu²⁺ and its associated biomedical insinuations has resulted in substantial attention for the scheming of highly selective and sensitive copper chemosensors.²⁷

Al³⁺ is one of the most abundant biosphere elements at approximately 8% of the total mineral components. The neurotoxicity of Al³⁺ has been known to humans for over one hundred years.^{28,29} Al³⁺ can cause many health issues, including Alzheimer's disease³⁰ and osteomalacia,³¹ and increased risk of breast cancer.³² The World Health Organization (WHO) recommends 3–10 mg of Al³⁺ as an average daily intake. At the same time, the weekly tolerable dietary intake is about 7 mg kg⁻¹ body weight. Thus, recognizing Al³⁺ in life and environmentally significant samples is critical to address.^{33,34} The detection of Al³⁺ has been challenging compared to other metal ions because of poor spectroscopic characteristics, meager coordination ability, and easy hydration.^{35–37} The development of highly sensitive and selective chemosensor for Al³⁺ detection is in great demand. Therefore, it is of substantial importance to build receptors for the selective detection of Al³⁺.

According to recent literature, noncyclic receptors containing multiple coordination sites have considerably improved in

^a Institute of Applied Chemistry and Department of Chemistry, Hallym University, Chuncheon 200702, Republic of Korea. E-mail: satish_nimse@hallym.ac.kr

^b School of Chemical Sciences, KBC-North Maharashtra University, Jalgaon 425001, India. E-mail: kuwaras@gmail.com

^c Biometrix Technology, Inc., 2-2 Bio Venture Plaza 56, Chuncheon 24232, Korea

^d Department of Applied Chemistry, SV National Institute of Technology, Surat 395007, India

† Electronic supplementary information (ESI) available: Tables S1–S5 and Fig. S1–S9 are available. CCDC 2051947. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1ma00564b

‡ These authors contributed equally. Hence, both should be considered as first authors.






Fig. 1 (a) Scheme for the synthesis of receptor **3**, (b) Single X-ray crystal structure of receptor **3** (50% probability ellipsoids).

the chemosensor design because of their ability to recognize different ionic and neutral molecules.^{38–40} Such noncyclic receptors are well known to show interesting coordination properties due to heteroatoms in chelating sites. With this in mind, a Schiff base (receptor **3**, Fig. 1a) was prepared from isonicotinohydrazide and 2,4-dihydroxybenzaldehyde and evaluated as the target receptor for the metal ions. It was hypothesized that N atoms of the $-\text{CH}=\text{N}-$ bond and the O atom of *ortho* $-\text{OH}$ in receptor **3** could coordinate with the metal ion to form a stable complex. Herein, we report receptor **3** as a selective chemosensor for Cu^{2+} as it showed the distinctive red shift in the absorption maxima but without any fluorescence signal for the complex. Receptor **3** was found to be an excellent fluorescence sensor for Al^{3+} , as evident from the chelation-enhanced fluorescence (CHEF) effect. The application of receptor **3** in fluorescence imaging was evaluated by confocal fluorescence microscopy in A549 cells, proving receptor **3** as a convenient tool for tracking Al^{3+} *in vivo*.

2. Material and methods

2.1 Materials and instrumentation

All reagents were purchased from Sigma-Aldrich and were used as received unless stated otherwise. The reaction was carried out under an inert atmosphere (Argon gas) and monitoring the reaction using thin-layer chromatography (TLC) to confirm the formation of the product. The developed plates were visualized under UV light (254 nm). The synthesized compound was characterized by ^1H and ^{13}C NMR on a Jeol FT-NMR spectrometer (400 MHz; JEOL, Japan) in $\text{DMSO}-d_6$. The chemical shifts (δ ppm) and the coupling constants (J Hz) are reported. UV-visible spectra were recorded on a Shimadzu UV-24500 (Shimadzu, Japan) in the range of 200–600 nm at room temperature using a quartz cuvette of 1 cm optical path length. Fluorescence emission spectra were measured on an Agilent Cary Eclipse fluorescence spectrophotometer (Agilent Technologies, USA). FT-IR spectra were recorded on a Nicolet iS5 FT-IR (ThermoScientific) spectrometer in the range of 400–4000 cm^{-1} using KBr pellets. The JMS-700 MStation Mass Spectrometer (JEOL, Japan) was used for recording the high-resolution mass spectra of receptor **3**. The mass spectra of $3\cdot\text{Al}^{3+}$ complex were obtained by matrix-assisted laser deposition/ionization (time-of-flight), MALDI-TOF mass spectrometry on a Bruker Autoflex speed TOF/TOF spectrometer (Bruker Daltonics, Bremen, Germany). X-ray analysis was performed using a Bruker AXS D8 Quest CMOS diffractometer (Bruker, USA). The microplate reader was Spectramax Plus 384 (Molecular Devices, USA),

and the fluorescent microscope Zeiss-ScopeA1 (Germany) was used in this study.

2.2 Synthesis of receptor 3

Isonicotinohydrazide (0.198 g, 1.0 mmol) and 2,4-dihydroxybenzaldehyde (0.200 g, 1.0 mmol) were reacted in ethanol (25 mL) at room temperature until the completion of the reaction (5 h). The yellow color solid obtained was filtered and dried. Then, the obtained yellow colored crud product was recrystallization from ethanol (yield = 90%). ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 6.36 (d, 2H, Ar-H), 7.36 (s, 1H, Ar-H), 7.82 (d, 2H, Ar-H), 8.54 (s, 1H, CH=N), 8.77 (d, 2H, Ar-H), 10.08 (s, 1H, Ar-OH), 11.29 (s, 1H, NH), 12.14 (s, 1H, Ar-OH); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 102.61, 107.84, 110.40, 121.43, 131.20, 140.07, 150.29, 159.49, 161; mass: expt. m/z = 258.0870 ($\text{C}_{13}\text{H}_{11}\text{N}_3\text{O}_3$ [M + H]), calc. m/z = 257.245.

2.3 UV-vis and fluorescence spectral measurements

All stock and working solutions were prepared using double distilled water and spectroscopic grade DMSO. A stock solution of receptor **3** (1×10^{-3} M) was prepared in DMSO, and the corresponding working solutions (10×10^{-6} M) were prepared simply by diluting with DMSO. Similarly, stock solutions of cations (1×10^{-2} M) were prepared in double-distilled water, and the corresponding working solutions (1×10^{-3} M) were prepared by diluting with water. The UV-visible absorption and emission spectra of the receptor **3** (10×10^{-6} M) dissolved in DMSO were recorded by adding the aqueous solution of various metal ions (Na^+ , K^+ , Ag^+ , Cs^+ , Sr^{2+} , Ca^{2+} , Co^{2+} , Cu^{2+} , Pd^{2+} , Mn^{2+} , Mg^{2+} , Ba^{2+} , Ni^{2+} , Zn^{2+} , Cd^{2+} , Pb^{2+} , Al^{3+} , Cr^{3+} , Fe^{3+} , Fe^{2+}) to examine the selectivity at room temperatures (298 K). For the sensitivity study, UV-visible absorption and emission titration experiments were performed through a stepwise addition of four equivalents of metals (1×10^{-3} M) to a solution of receptor **3** (10×10^{-6} M) in DMSO. The absorbance intensity and emission intensity were recorded in the range of 200–600 nm and 360–600 nm, respectively, alongside a reagent blank. Receptor **3** showed selectivity for detecting Cu^{2+} in the absorption titrations and detecting Al^{3+} in the fluorescence titrations. The binding stoichiometry of receptor **3** with Cu^{2+} ion (UV-visible absorption spectroscopy) and Al^{3+} ion (fluorescence spectroscopy) were investigated by Job's plot method. A receptor **3** was titrated with successive addition of Cu^{2+} or Al^{3+} (1 μL , 1.0×10^{-3} M) in water to a receptor **3** (1.0 mL) solution in DMSO. The collected data were processed using the Benesi-Hildebrand equation⁴¹ to determine the association constant (K_a) of analyte Cu^{2+} and Al^{3+} ion with receptor **3**. The absorbance changes at 413 nm were used alongside a reagent blank for the



detection of Cu²⁺ ions. The fluorescence intensity was recorded at $\lambda_{\text{ex}}/\lambda_{\text{em}} = 343/473$ nm alongside a reagent blank with the excitation and emission slits set to 5.0 nm. The limit of detection (LOD) was estimated by applying the IUPAC recommended equation, $\text{LOD} = 3\sigma/\text{slope}$.⁴² Where σ is the standard deviation of ($n = 10$) blank samples and the slope is the slope for calibration curves.

2.4 Crystal growth for single X-ray crystallography

The single crystals of receptor **3** were obtained by slow diffusion of ethanol in DMSO. However, several attempts to obtain the single crystals of receptor **3** and Al³⁺ complex (**3**·Al³⁺), receptor **3** and Cu²⁺ complex (**3**·Cu²⁺) were unsuccessful. A suitable single crystal was carefully mounted for X-ray crystallography with the help of a trace of Fomblin oil on a Mitegen micromesh mount. Then it was transferred to the goniometer head with a fixed chi angle, a molybdenum K_α wavelength fine focus sealed X-ray tube ($\lambda = 0.71073$), a single crystal curved graphite incident beam monochromator, a Photon100 CMOS area detector, and an Oxford Cryosystems low-temperature device. X-ray diffraction data were collected at 150 K using ω and ϕ scans to a maximum resolution of $\Theta = 33.221^\circ$. Data reduction, scaling, and absorption corrections were performed using SAINT (Bruker, V8.38A). The final completeness is 89.00% out of 33.221° in Θ . Multi-scan absorption correction was performed using SADABS 2016/2.⁴³ The absorption coefficient μ of this material was 0.107 mm^{-1} at this wavelength ($\lambda = 0.71073 \text{ \AA}$). The space group was determined based on systematic absences using XPREP⁴⁴ as *Pna2*₁. The structure was solved using direct methods with ShelXS-97 and refined by full-matrix least-squares on F^2 using ShelXL-2018/3 and the graphical interface ShelXLE (Rev937).⁴⁵ All non-hydrogen atoms were refined anisotropically. Hydrogen atom positions were calculated geometrically and improved using a riding model. Mercury, PyMol, and POVray were utilized for molecular measurements and molecular visualization.⁴⁶

2.5 Effect of pH on the detection of Al³⁺ and reversibility of receptor **3**

The effect of pH (pH = 2–12) on receptor **3** was examined by fluorescence spectroscopy both in the absence and presence of Al³⁺ ions. The pH was adjusted by adding perchloric acid and tetrabutylammonium hydroxide in the HEPES buffered system. Reversibility is a critical aspect of the fluorescent recognition process. Hence, we examined the reversibility of receptor **3** in the presence of ethylenediaminetetraacetic acid disodium salt (EDTANa₂). For the reversibility study, EDTANa₂ (4 equiv.) was added to the solution containing receptor **3**·Al³⁺ complex obtained by adding Al³⁺ at a 1 : 4 mole ratio. The reversibility was recorded by alternate additions of Al³⁺ (4 equiv.) and EDTANa₂ (4 equiv.).

2.6 Determination of quantum yield of receptor **3** and receptor **3**·Al³⁺ complex

Quantum yields (Φ) of receptor **3** and its complexes with Al³⁺ were measured using the following formula.

$$\Phi_{\text{sample}} = \left\{ \frac{(\text{OD}_{\text{standard}} \times A_{\text{sample}} \times \eta_{\text{sample}}^2)}{(\text{OD}_{\text{sample}} \times A_{\text{standard}} \times \eta_{\text{standard}}^2)} \right\} \times \Phi_{\text{standard}}$$

where A is the area under the emission spectral curve, OD is the compound's optical density at the excitation wavelength, and η is the refractive index of the solvent. The quantum yield of receptor **3** and its complexes with Al³⁺ was determined using β -carboline ($\Phi = 0.570$) as the standard.⁴⁷

2.7 Cell culture studies

The cytotoxicity assay (MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay) and cell imaging study of receptor **3**, Al³⁺, and Al³⁺ combined with receptor **3** was conducted using A549 cells (colorectal carcinoma cell line). The A549 cells were procured from the Korea cell line bank, Seoul, South Korea. Dulbecco Modified Eagle Medium (DMEM), fetal bovine serum (FBS), trypsin, 3-(4,5-dimethyl thiazol-2-yl)-2,5-diphenyl tetrazolium bromide was procured from Thermo Fisher Scientific, USA. DMSO was procured from Biosesang, Korea. Cell culture plates and glass coverslips were procured from SPL Life Sciences, Korea. A549 cells were grown in an incubator at 37 °C and 5% CO₂ using DMEM media containing 2 mM glutamine and 10% FBS. Cells were trypsinized for seeding at 70–90% of cell confluency.

2.8 Cytotoxicity assay and cell imaging

About 2000 A549 cells per well were seeded in 96-well plates. After 24 h, the media containing receptor **3**, Al³⁺ ion, and receptor **3** with Al³⁺ (1, 10, 25, and 50 μM) were added to the wells and incubated for another 24 h. Control wells were treated with equivalent volumes of dimethyl sulfoxide (DMSO). 200 μL of fresh media containing MTT solution and incubated for four hours at 37 °C. The absorbance was recorded at 570 nm to evaluate the cell viability. Each experiment was executed three times. The data analysis was performed using the Origin software.

For cell imaging, A549 cells were seeded separately on poly-L-lysine coated 14 mm coverslips in 6 well plates and allowed to grow for 24 h. For cell imaging control experiments, 10 μM of Al³⁺ ion and 10 μM of receptor **3** were incubated separately for 30 min at 37 °C and 5% CO₂ in the dark. For cell imaging of receptor **3** and Al³⁺ complex, 10 μM of Al³⁺ ion was incubated separately for 30 min. The media was then replaced with 10 μM of receptor **3** and incubated for an additional 30 min. The cells were washed with PBS buffer (pH = 7.4), followed by fixing with 2% paraformaldehyde for 30 min after removing the media. Coverslip was mounted on a glass slide, and imaging was performed under a fluorescence microscope (Zeiss-ScopeA1, Germany). Images were taken through a green channel.

3. Results and discussion

3.1 Synthesis of receptor **3**

The Schiff base (*E*)-*N'*-(2,4-dihydroxybenzylidene)isonicotinohydrazide (receptor **3**) was synthesized with a slight modification of the reported⁴⁸ method through a direct reaction between isonicotinohydrazide and 2,4-dihydroxybenzaldehyde (Fig. 1a) in ethanol with stirring and refluxing for 5 h. The molecular structure of



receptor **3** was characterized using various spectral techniques (FT-IR, ^1H NMR, ^{13}C NMR, and Mass), data (ESI,† Fig. S1–S3) and finally confirmed by single-crystal X-ray crystallography (Fig. 1b). The crystallographic data, selected bond parameters, and hydrogen-bond parameters are presented in Tables S1, S2 and S3, respectively. The CIF file for receptor **3** was placed in the Cambridge Structure Database (CCDC 2051947†). The orange-colored crystal ($0.55 \times 0.42 \times 0.05 \text{ mm}^3$) of receptor **3** demonstrated an orthorhombic system having a $Pna2_1$ space group within the unit cell. The ORTEP diagram with numbering and packing diagram is shown in Fig. 1b. The receptor **3** in its free form displays the molecular association *via* intramolecular hydrogen bonding between the phenolic hydroxyl to imine nitrogen. Receptor **3** shows an intramolecular hydrogen bond (O(1)–H...N(1)) with a distance of 1.92 (3) Å and a bond angle of 150 (2) Å, which is in the expected range of such hydrogen bonds. Receptor **3** undergoes a solvent-assisted keto tautomerization suitable for the intramolecular charge transfer (ICT) process.⁴⁹ The fluorescence intensity enhancement of receptor **3** in the presence of Al^{3+} is attributed to the ICT process.

3.2 Determination of selectivity of receptor **3** as a chemosensor for metal ions

The selectivity of receptor **3** for cation detection was investigated using the UV-visible absorption and fluorescence spectroscopy. The UV-vis absorption spectra of receptor **3** ($10 \times 10^{-6} \text{ M}$, in DMSO) were recorded in the absence and presence of 4 equivalents of various metal ions, such as Na^+ , K^+ , Ag^+ , Cs^+ , Sr^{2+} , Ca^{2+} , Co^{2+} , Cu^{2+} , Pd^{2+} , Mn^{2+} , Mg^{2+} , Ba^{2+} , Ni^{2+} , Zn^{2+} , Cd^{2+} , Pb^{2+} , Al^{3+} , Cr^{3+} , Fe^{3+} , Fe^{2+} ($1 \times 10^{-3} \text{ M}$, in H_2O).

Receptor **3** showed an absorption band at 343 nm, most likely due to the π to π^* transition (Fig. 2a). Upon addition of Cu^{2+} ions, the absorption band at 343 nm was red-shifted to 413 nm ($\Delta\lambda \approx 70 \text{ nm}$), indicating that receptor **3** has a higher binding affinity towards Cu^{2+} ions than other surveyed metal ions. Receptor **3** showed two additional shoulder peaks at 428 nm and 475 nm to the major peak at 413 nm. In the presence of other metal ions, receptor **3** showed either no change or moderate decrease in the absorption intensity

relative to the receptor. These results indicated the intramolecular charge transfer (ICT) character of the synthesized receptor **3** by recognizing Cu^{2+} ions through imine-N, amide carbonyl, and hydroxyl groups.⁵⁰ The push-pull character of the ICT state due to multiple coordination resulted in a red-shift ($\Delta\lambda \approx 70 \text{ nm}$). These results indicated that receptor **3** shows selectivity for Cu^{2+} ions.

The fluorescence emission spectra of receptor **3** ($10 \times 10^{-6} \text{ M}$, in DMSO) were recorded in the absence and presence of 4 equivalents of various metal ions, such as Na^+ , K^+ , Ag^+ , Cs^+ , Sr^{2+} , Ca^{2+} , Co^{2+} , Cu^{2+} , Pd^{2+} , Mn^{2+} , Mg^{2+} , Ba^{2+} , Ni^{2+} , Zn^{2+} , Cd^{2+} , Pb^{2+} , Al^{3+} , Cr^{3+} , Fe^{3+} , Fe^{2+} ($1 \times 10^{-3} \text{ M}$, in H_2O). Receptor **3** showed weak fluorescence emission at 473 nm upon excitation at 343 nm. Fascinatingly, fluorescence was remarkably enhanced (~ 430 -folds) in the presence of Al^{3+} ions (Fig. 2b). Interestingly, there was no change in the emission performance of receptor **3** in the presence of other cations, including Cu^{2+} . The increase in fluorescence emission intensity was due to the azomethine group of receptor **3**. Receptor **3** is reducibly fluorescent due to the $\text{C}=\text{N}$ double bond's isomerization at the excited state and the excited-state proton transfer (ESPT). The ESPT involves the phenolic proton of the substituted dihydroxyl moieties of salicylaldehyde in receptor **3**. Upon stable chelation with Al^{3+} , the $\text{C}=\text{N}$ isomerization is inhibited. The coordination of receptor **3** with the Al^{3+} prohibits the ESPT process, as indicated by the fluorescence enhancement.^{51–53}

3.3 Binding mechanism and association constant

Job's plot was used to determine the binding stoichiometry of receptor **3** with Cu^{2+} ions (UV-visible absorption spectroscopy) and Al^{3+} ions (fluorescence spectroscopy). The molar ratio of metal ions was changed from 0.1 to 1.0 by keeping the total concentration of receptor **3** and Cu^{2+} ions at $10 \times 10^{-6} \text{ M}$. The absorption maxima ($\lambda = 413 \text{ nm}$) was observed when the molar ratio of the receptor **3** to Cu^{2+} was 0.33, indicating the formation of a 2 : 1 **3**- Cu^{2+} complex (Fig. S4a, ESI†). Similarly, the change in fluorescence intensity ($\lambda_{\text{ex}} = 343 \text{ nm}$, $\lambda_{\text{em}} = 473 \text{ nm}$) was used to determine the binding stoichiometry of receptor **3**




Fig. 2 (a) Changes in UV-vis absorption spectra and (b) fluorescence intensity ($\lambda_{\text{ex}} = 343 \text{ nm}$, $\lambda_{\text{em}} = 473 \text{ nm}$) of receptor **3** ($10 \times 10^{-6} \text{ M}$) upon the addition of 4 equivalents of different metal ions ($1 \times 10^{-3} \text{ M}$, in H_2O). Inset shows the color change of the solutions from colorless to pale green and fluorescence 'turn-on' effect.






Fig. 3 FT-IR spectra of (a) **3**-Cu²⁺ complex and (b) **3**-Al³⁺ complex recorded using the reflectance technique (4000–400 cm⁻¹).

with Al³⁺. The molar ratio of Al³⁺ was changed from 0.1 to 1.0 by keeping the total concentration of receptor **3**, and Al³⁺ ions at 10×10^{-6} M. The emission maxima ($\lambda = 473$ nm) was observed when the molar ratio of the receptor **3** to Al³⁺ was 0.33, indicating the formation of a 2:1 **3**-Al³⁺ complex (Fig. S4b, ESI[†]). The complexes **3**-Cu²⁺ and **3**-Al³⁺ were obtained by refluxing four equivalents of Cu²⁺ and Al³⁺ with receptor **3** in an ethanolic solution. To ascertain the formation of receptor **3**-Cu²⁺ complex and **3**-Al³⁺ complex, we compared FT-IR spectra for receptor **3** with that of respective complexes, as shown in Fig. 3.

The FT-IR spectrum of receptor **3** (Fig. 3a) demonstrated signals at 3436.05, 3224.39 (amide), 3086.99 (intramolecular bonded O-H), and 1647.39 (imine) cm⁻¹. For the receptor **3**-Cu²⁺ complex, the FT-IR spectrum demonstrated a broad signal at 3422.06 (amide), no frequency for intramolecular bonded O-H, and 1621.84 (imine) cm⁻¹. These results indicate that receptor **3** forms a stable complex with Cu²⁺. Similarly, the FT-IR spectrum of receptor **3**-Al³⁺ complex demonstrated a broad signal at 3430.74 (amide), no frequency for intramolecular bonded O-H, and 1623.77 (imine) cm⁻¹. The FT-IR spectrum shifts in the **3**-Cu²⁺ and **3**-Al³⁺ complexes compared to the receptor **3**, confirming the involvement of imine and amide groups in the complexation process.

As shown in Fig. 4, the ¹H NMR titration experiment was conducted using the mixture of 0.5% D₂O-d₂ in DMSO-d₆. The ¹H NMR of receptor **3** demonstrated sharp peaks at δ_{H} 12.12 (amide N-H), δ_{H} 11.25, and δ_{H} 10.01 (phenolic -OH). The signals for one N-H proton and two -OH protons gradually disappeared upon increasing the amount of Al³⁺ ions (0–1 equiv.). However, there was no significant change in the peak corresponding to imine C-H (δ_{H} 8.52). These results indicate that the amide group, imine group, and phenolic moiety of receptor **3** take part in the complexation with Al³⁺ ions. Further, the MALDI-TOF mass spectrum of receptor **3**-Al³⁺ complex showed a signal at an m/z value of 544.129 and 561.133 (Fig. S5, ESI[†]). These results rationalize the formation of the 2:1 complexation pattern for the **3**-Al³⁺ complex. The paramagnetic property of Cu²⁺ results in the peak broadening in proton NMR spectra. Thus, the binding process of receptor **3** with Cu²⁺ could not be monitored by NMR studies.⁵⁴




Fig. 4 ¹H NMR spectral changes of receptor **3** in the mixture of 0.5% D₂O-d₂ in DMSO-d₆, (a) **3** + Al³⁺ (0 equiv.), (b) **3** + Al³⁺ (0.25 equiv.), (c) **3** + Al³⁺ (0.5 equiv.), (d) **3** + Al³⁺ (1 equiv.).

The association constants (K_a) of Cu²⁺ (UV-visible absorption spectroscopy) and Al³⁺ (fluorescence spectroscopy) complexes with receptor **3** were determined by Benesi-Hildebrand equations (eqn (S1) and (S2), ESI[†]). As shown in Fig. 5a, the receptor **3** (1.0 mL, 10×10^{-6} M) in DMSO was titrated with successive addition of Cu²⁺ (0–20 μ L in H₂O, $c = 10 \times 10^{-4}$ M) to measure the association constant (K_a). The absorbance values at absorption maxima ($\lambda = 413$ nm) were processed using the Benesi-Hildebrand equation (eqn (S1), ESI[†]) to obtain the binding curve (Fig. S6a, ESI[†]). The K_a value for the complexation of Cu²⁺ with receptor **3** was 1.3×10^4 M⁻¹.

As shown in Fig. 5b, receptor **3** (1.0 mL, 10×10^{-6} M) in DMSO was titrated with successive addition of Al³⁺ (0–20 μ L in H₂O, $c = 10 \times 10^{-4}$ M). The changes in fluorescence intensity ($\lambda_{\text{ex}} = 343$ nm, $\lambda_{\text{em}} = 473$ nm) were used to determine the K_a value by plotting the binding curve (Fig. S6b, ESI[†]) according to the Benesi-Hildebrand equation (eqn (S2), ESI[†]). The K_a value for the complexation of Al³⁺ with receptor **3** was 4.8×10^4 M⁻¹. The binding affinity of Al³⁺ for receptor **3** is 4-fold higher than that of Cu²⁺ ions.






Fig. 5 Changes in (a) UV-vis absorption spectra and (b) fluorescence spectra ($\lambda_{\text{ex}} = 343$ nm, $\lambda_{\text{em}} = 473$ nm) of receptor **3** (in 1 mL DMSO, 10×10^{-6} M) up on successive addition of Cu^{2+} (0 to 20 μL in H_2O , $c = 10 \times 10^{-4}$ M), and Al^{3+} (0 to 20 μL in H_2O , $c = 10 \times 10^{-4}$ M), respectively.

Computational studies were conducted by applying the density functional theory (DFT) using Gaussian 09. The molecular geometries of the singlet ground state of receptor **3**, **3**· Cu^{2+} complex, and **3**· Al^{3+} complex were optimized using hybrid B3LYP functions with a 6-31G++(d,p) (C, H, N, O) and LANL2DZ (Cu, Al) basis sets.⁵⁵ The HOMO, LUMO results, information for bond length, and bond angles were obtained using Avogadro 1.2.0.⁵⁶ The 3D structures of the **3**· Cu^{2+} and **3**· Al^{3+} complexes were calculated by the DFT method using the 2:1 binding stoichiometry between receptor **3** and respective ions. The DFT computed structure of receptor **3** and its complexes are presented in Fig. S7 (ESI[†]). The LUMO–HOMO bandgap (ΔE (eV) = $E_{\text{LUMO}} - E_{\text{HOMO}}$) for receptor **3** was found to be 0.293.

In contrast, the LUMO–HOMO bandgap for **3**· Cu^{2+} and **3**· Al^{3+} complexes were 0.279 and 0.078, respectively. These results provided the basis for ascertaining the ICT between receptor **3** and Cu^{2+} and Al^{3+} ions. Comparing the electron densities of the HOMO and LUMO of receptor **3** with the **3**· Cu^{2+} complex supported the charge transfer occurring between the receptor and metal ions that further lowered the bandgap. The lowering of the bandgap upon complexation supported the red-shift of the absorbance of receptor **3** upon the addition of Cu^{2+} . On the contrary, the significant decrease in the LUMO–HOMO bandgap ($\Delta E = 0.078$) indicated the formation of a relatively stable **3**· Al^{3+} complex as compared to the **3**· Cu^{2+} complex. The increase in fluorescence enhancement was complemented by a sharp decrease in energy HOMO–LUMO bandgap of the receptor **3**.

3.4 Receptor **3** as a chemosensor for Cu^{2+} and Al^{3+} ions

The detection of the target analyte in the presence of possibly competing analytes is a crucial aspect for any compound to be an excellent chemosensor. Therefore, we evaluated the specificity of receptor **3** for Cu^{2+} in a competition experiment by recording the absorption ($\lambda = 413$ nm) receptor **3** in the presence of Cu^{2+} (1 equiv.) ion mixed with other cations (4 equiv.). The results of the competition experiments are presented in Fig. S8a (ESI[†]). Similarly, we determined the efficiency of receptor **3** for detecting Al^{3+} ions in a competition experiment by recording the fluorescence intensity

($\lambda_{\text{ex}} = 343$ nm, $\lambda_{\text{em}} = 473$ nm). As shown in Fig. S8b (ESI[†]), the fluorescence intensity for receptor **3** was measured in the presence of Al^{3+} (1 equiv.) ion mixed with other cations (4 equiv.). The coefficient of variation in the change of absorbance and fluorescence intensity for Cu^{2+} detection and Al^{3+} detection, were below $\pm 10\%$, respectively. These results indicate that receptor **3** is a highly valuable chemosensor for detecting Cu^{2+} by UV-vis absorption spectroscopy. Moreover, receptor **3** demonstrated its excellence in detecting Al^{3+} ions with high specificity by fluorescence spectroscopy. Therefore, the high selectivity and specificity of receptor **3** for Cu^{2+} and Al^{3+} ions make it an excellent chemosensor for analytical applications. The absorbance ($\lambda = 413$ nm) and fluorescence intensity ($\lambda_{\text{ex}} = 343$ nm, $\lambda_{\text{em}} = 473$ nm) were plotted at various concentrations of Cu^{2+} ions and Al^{3+} ions, respectively, to obtain the calibration plots (Fig. S9, ESI[†]). The LOD for detecting Cu^{2+} ions by UV-vis absorption spectroscopy was 1.86 μM . Whereas, the LOD for the detection of Al^{3+} ions by Fluorescence spectroscopy was 3.08 nM. The approximately 600-fold lower sensitivity of receptor **3** for Al^{3+} ions compared to Cu^{2+} ions was attributed to the ~ 4 -fold higher binding constant for **3**· Al^{3+} complex. It is important to notice that receptor **3** demonstrated relatively lower detection limits for Cu^{2+} and Al^{3+} than some of the reported methods presented in Tables S4 and S5 (ESI[†]).

3.5 Effect of pH on the detection of Al^{3+} and reversibility of receptor **3**

The effect of pH on receptor **3** for detecting Al^{3+} ions was tested in the pH range of 2.0–12.0, as shown in Fig. 6a. The emission intensity ($\lambda_{\text{em}} = 473$ nm) of receptor **3** did not change significantly with the pH change. However, equimolar Al^{3+} ions changed the fluorescence intensity of receptor **3**· Al^{3+} complex considerably with the pH change. The receptor **3**· Al^{3+} complex's emission intensity was significantly high at pH 2.0–8.0 than pH 9.0–12.0. These results indicate that interactions of Al^{3+} ions with receptor **3** are pH dependant. The decrease in emission intensity at higher pH values (9.0–12.0) specifies that the Al^{3+} ions are freed from the complex, possibly due to more robust interactions with increased $-\text{OH}$ levels. Nonetheless, receptor **3**






Fig. 6 (a) Changes in the fluorescence intensity of receptor **3** and receptor $\text{3}\cdot\text{Al}^{3+}$ complex in a binary mixture of DMSO : HEPES buffer (at various pH values) (9 : 1; v/v), (b) fluorescence intensities ($\lambda_{\text{ex}} = 343 \text{ nm}$, $\lambda_{\text{em}} = 473 \text{ nm}$) of $\text{3}\cdot\text{Al}^{3+}$ complex (1 : 1) in the presence of EDTA for many cycles in DMSO.

demonstrated significant fluorescence intensity at pH 5.0–8.0 in the presence of Al^{3+} ions, indicating its applicability for the detection of intracellular Al^{3+} ions.

It is of a great advantage if a sensor can be reversible and reusable for sensing cations with high selectivity. A literature survey revealed that most of the reported Al^{3+} ion sensors are based on chemodosimetry that is typically irreversible.⁵⁷ Reversibility test experiments were conducted by alternate additions of Al^{3+} ion and EDTA (Fig. 6b) to the solution of receptor **3**. As shown in Fig. 6b, the emission spectra of receptor **3** in the presence of Al^{3+} (4 equiv.) showed high emission intensity, which was quenched by EDTA (6 equiv.) on the solution. However, further addition of Al^{3+} (4 equiv.) demonstrated fluorescence signal, but this time slightly lower than the previous cycle. Adding Al^{3+} ions (4 equiv.) causes emission enhancement, which can be quenched by adding another portion of EDTA (6 equiv.). The reversibility of receptor **3** was repetitive, with a slight loss in fluorescence efficiency due to Al^{3+} /EDTA additions. The observed decrease in fluorescence intensity for each cycle results from an excess of EDTA (6 equiv.) compared to Al^{3+} (4 equiv.). This experiment suggests that receptor **3** can act as a likely environmental receptor for Al^{3+} detection.

3.6 Quantum yield of receptor **3** and receptor $\text{3}\cdot\text{Al}^{3+}$ complex

Quantum yields of receptor **3** and $\text{3}\cdot\text{Al}^{3+}$ complex were determined using norharmene as a standard. The quantum yield of receptor **3** ($\Phi = 0.0021$) increases ~ 230 folds in the presence of Al^{3+} ions to form receptor $\text{3}\cdot\text{Al}^{3+}$ complex ($\Phi = 0.484$).

3.7 Application of receptor **3** in cell imaging application for detection of intracellular Al^{3+} ions

The complexation-induced fluorescence “turn on” or “turn off” effect is crucial for bio-imaging applications of small molecular probes designed to detect the cationic analyte. Receptor **3** did not show any change in fluorescence intensity upon binding with Cu^{2+} ions. However, the fluorescence signal of receptor **3** was increased by a few hundred folds in the presence of Al^{3+} ions. Therefore, receptor **3** was used to detect Al^{3+} ions in the living A549 cell lines to explore their biological applications. The MTT assay allowed us to estimate the cytotoxicity of receptor **3**, Al^{3+} , and receptor $\text{3}\cdot\text{Al}^{3+}$ complex after exposure of cells to concentrations of 1, 10, 25, and 50 μM for 24 h with DMSO as a control. As depicted in Fig. 7a, the results are shown as the percent cell growth for each group compared to the control. There was no significant cell death even after 24 h of




Fig. 7 (a) Cytotoxicity's of receptor **3**, Al^{3+} , and receptor $\text{3}\cdot\text{Al}^{3+}$ on the A549 cells after 24 h, (b) bright-field images (top row) and green channel (bottom row), control, receptor **3**, Al^{3+} , and receptor $\text{3}\cdot\text{Al}^{3+}$.



treatment at 1–25 μM of the receptor **3**, Al^{3+} ions, and the receptor **3**· Al^{3+} complex. However, upon treatment at 50 μM of receptor **3** and receptor **3**· Al^{3+} complex, about a 20% decrease in cell growth was observed. Hence, 10 μM of receptor **3** was used for cell imaging applications. As shown in Fig. 7b, the cells did not show significant fluorescence upon incubation with Al^{3+} ions (10 μM) alone. However, very weak fluorescence was observed upon incubation of cells with receptor **3** (10 μM). Interestingly, the fluorescence intensity was increased upon incubation of cells with receptor **3** (10 μM) in the presence of Al^{3+} ions (10 μM). These results indicated that receptor **3** has a high potential in biological applications to detect Al^{3+} in an *in vitro* assay.

4. Conclusions

In conclusion, we have developed a new optical receptor **3** based on Schiff base chemistry that demonstrated excellent selectivity and sensitivity towards Cu^{2+} ions in UV-vis absorption spectroscopy. Whereas the developed receptor **3** showed excellent selectivity and sensitivity for the detection of Al^{3+} ion by using fluorescence spectroscopy. In either case, the receptor did not show any interference from other tested metal ions. The 2:1 binding stoichiometry of receptor **3** and Al^{3+} ions was confirmed by FT-IR, NMR, and mass spectroscopy. The reversibility of receptor **3** for Al^{3+} ions in the presence of EDTA ensures its ability as an excellent probe for detecting Al^{3+} in various samples, including living cells. Receptor **3** showed high selectivity for Al^{3+} with a K_a of $4.8 \times 10^4 \text{ M}^{-1}$ and LOD of 3.0 nM. In comparison, the K_a for Cu^{2+} was $1.3 \times 10^4 \text{ M}^{-1}$ and LOD of 1.9 μM . Receptor **3** is an excellent chemosensor for detecting Al^{3+} ions indicated by its nanomolar range LOD. The quick response, easy synthesis, and high sensitivity make receptor **3** an ideal sensor for detecting Cu^{2+} and Al^{3+} ions. Further, the synthesized receptor showed a highly sensitive and highly specific fluorescent ‘turn-on’ effect ($\lambda_{\text{em}} = 473 \text{ nm}$) for the 2:1 binding with Al^{3+} ions in a semi-aqueous medium and living cells.

Author contributions

I. S. and P. T. contributed equally. Hence, both should be considered as the first authors. S. B. N. and A. K. designed the study; I. S., P. T., J. L., A. B., and S. D. W. performed the experiments; S. B. N., S. K. S., and A. K. analyzed and interpreted the data. S. K. S. completed the theoretical calculation. S. B. N. and A. K. supervised the research. S. B. N. and A. K. wrote the paper. All authors analyzed the data and commented on the paper.

Conflicts of interest

The authors declare no competing interests.

Acknowledgements

Hallym University Research Fund (HRF-202012-009) supported this research.

References

- H. S. Jung, P. S. Kwon, J. W. Kwon, J. I. Kim, C. S. Hong, J. W. Kim, S. Yan, J. Y. Lee, J. H. Lee, T. Joo and J. S. Kim, *J. Am. Chem. Soc.*, 2009, **131**(5), 2008–2012, DOI: 10.1021/ja808611d.
- Z. Q. Guo, W. Q. Chen and X. M. Duan, *Org. Lett.*, 2010, **12**(10), 2202–2205, DOI: 10.1021/ol100381g.
- M. Royzen, A. Durandin, V. G. Young, N. E. Geacintov and J. W. Canary, *J. Am. Chem. Soc.*, 2006, **128**, 3854–3855, DOI: 10.1021/ja056631g.
- C. Gou, S.-H. Qin, H.-Q. Wu, Y. Wang, J. Luo and X.-Y. Liu, *Inorg. Chem. Commun.*, 2011, **14**(10), 1622–1625, DOI: 10.1016/j.inoche.2011.06.024.
- J. Ting, F. Mei, Z. Mengyao, Q. Jianwen, Z. Hu and G. Yong, *Spectrochim. Acta, Part A*, 2020, **227**, 117530, DOI: 10.1016/j.saa.2019.117530.
- J. J. Xiong, P. C. Huang, C. Y. Zhang and F. Y. Wu, *Sens. Actuators, B*, 2016, **226**, 30–36, DOI: 10.1016/j.snb.2015.11.113.
- G. Zhang, H. Zhang, J. Zhang, W. Ding, J. Xu and Y. Wen, *Sens. Actuators, B*, 2017, **253**, 224–230, DOI: 10.1016/j.snb.2017.06.144.
- K. P. Carter, A. M. Young and A. E. Palmer, *Chem. Rev.*, 2014, **114**, 4564–4601, DOI: 10.1021/cr400546e.
- N. Dey and S. Bhattacharya, *Dalton Trans.*, 2018, **47**, 2352–2359, DOI: 10.1039/C7DT03401F.
- H. Rao, W. Liu, K. He, S. Zhao, Z. Lu, S. Zhang, M. Sun, P. Zou, X. Wang, Q. Zhao, Y. Wang and T. Liu, *ACS Sustainable Chem. Eng.*, 2020, **8**, 8857–8867, DOI: 10.1021/acssuschemeng.0c03354.
- S. L. Gui, Y. Y. Huang, F. Hu, Y. L. Jin, G. X. Zhang, L. S. Yan, D. Q. Zhang and R. Zhao, *Anal. Chem.*, 2015, **87**, 1470–1474, DOI: 10.1021/ac504153c.
- M. Kumar, N. Kumar, V. Bhalla, P. R. Sharma and T. Kaur, *Org. Lett.*, 2012, **14**, 406–409, DOI: 10.1021/ol3034059.
- D. G. Barceloux, *J. Toxicol., Clin. Toxicol.*, 1999, **37**, 217–230, DOI: 10.1081/CLT-100102421.
- T. G. Thomas, K. Sreenath and K. R. Gopidas, *Analyst*, 2012, **137**, 5358–5362, DOI: 10.1039/C2AN35904A.
- A. K. Manna, K. Rout, S. Chowdhury and G. K. Patra, *Photochem. Photobiol. Sci.*, 2019, **18**(6), 1512–1525, DOI: 10.1039/c9pp00114j.
- S. N. Gacheru, P. C. Trackman, M. A. Shah, C. Y. O’Gara, P. Spacciapoli, F. T. Greenaway and H. M. Kagan, *J. Biol. Chem.*, 1990, **265**(31), 19022–19027, DOI: 10.1016/S0021-9258(17)30618-X.
- H. Beinert, *Eur. J. Biochem.*, 1997, **245**(3), 521–532, DOI: 10.1111/j.1432-1033.1997.t01-1-00521.x.
- J. F. Mercer, *Trends. Mol. Med.*, 2001, **7**(2), 64–69, DOI: 10.1016/s1471-4914(01)01920-7.
- H. Tapiero, D. M. Townsend and K. D. Tew, *Biomed. Pharmacother.*, 2003, **57**(9), 386–398, DOI: 10.1016/s0753-3322(03)00012-x.



- 20 O. Bandmann, K. H. Weiss and S. G. Kaler, *Lancet Neurol.*, 2015, **14**(1), 103–113, DOI: 10.1016/S1474-4422(14)70190-5.
- 21 R. Hait-Darshan, T. Babushkin and Z. Malik, *J. Environ. Pathol., Toxicol. Oncol.*, 2009, **28**(3), 209–221, DOI: 10.1615/jenvironpatholtoxicoloncol.v28.i3.20.
- 22 E. Koutsouraki, D. Michmizos, O. Patsi, J. Tzartos, M. Spilioti, M. Arnaoutoglou and M. Tsolaki, *Virol. J.*, 2020, **17**(1), 35, DOI: 10.1186/s12985-020-01309-x.
- 23 E. Tokuda and Y. Furukawa, *Int. J. Mol. Sci.*, 2016, **17**(5), 636, DOI: 10.3390/ijms17050636.
- 24 K. A. Trumbull and J. S. Beckman, *Antioxid. Redox Signaling*, 2009, **11**(7), 1627–1639, DOI: 10.1089/ars.2009.2574.
- 25 R. Squitti, D. Lupoi, P. Pasqualetti, G. Dal Forno, F. Vernieri, P. Chioventa, L. Rossi, M. Cortesi, E. Cassetta and P. M. Rossini, *Neurology*, 2002, **59**(8), 1153–1161, DOI: 10.1212/wnl.59.8.1153.
- 26 P. G. Georgopoulos, A. Roy, M. J. Yonone-Lioy, R. E. Opiekun and P. J. Lioy, *J. Toxicol. Environ. Health, Part B*, 2001, **4**(4), 341–394, DOI: 10.1080/109374001753146207.
- 27 U. A. Fegade, S. K. Sahoo, A. Singh, N. Singh, S. B. Attarde and A. S. Kuwar, *Anal. Chim. Acta*, 2015, **872**, 63–69, DOI: 10.1016/j.aca.2015.02.051.
- 28 A. K. Manna, S. Chowdhury and G. K. Patra, *New J. Chem.*, 2020, **44**, 10819, DOI: 10.1039/D0NJ01954B.
- 29 M. L. D. R. Crapper, W. J. Lukiw and T. P. Kruck, *Environ. Geochem. Health*, 1990, **12**, 103–114, DOI: 10.1007/BF01734059.
- 30 C. N. Martyn, D. J. Barker, C. Osmond, E. C. Harris, J. A. Edwardson and R. F. Lacey, *Lancet*, 1989, **333**, 61–62.
- 31 G. C. Woodson, *Bone*, 1998, **22**, 695–698, DOI: 10.1016/S8756-3282(98)00060-x.
- 32 P. D. Darbre, *J. Inorg. Biochem.*, 2005, **99**, 1912–1919, DOI: 10.1016/j.jinorgbio.2005.06.001.
- 33 S. Kim, J. Y. Noh, K. Y. Kim, J. H. Kim, H. K. Kang, S. W. Nam, S. H. Kim, S. Park, C. Kim and J. Kim, *Inorg. Chem.*, 2012, **51**(6), 3597–3602, DOI: 10.1021/ic2024583.
- 34 C. Gao, P. Zang, W. Liu and Y. Tang, *J. Fluoresc.*, 2016, **26**, 2015, DOI: 10.1007/s10895-016-1895-z.
- 35 X. Sun, Y. W. Wang and Y. Peng, *Org. Lett.*, 2012, **14**, 3420–3423, DOI: 10.1021/ol301390g.
- 36 K. K. Upadhyay and A. Kumar, *Org. Biomol. Chem.*, 2010, **8**, 4892–4897, DOI: 10.1039/C0OB00171F.
- 37 K. Soroka, R. Vithanage, D. A. Philips, B. Walker and P. K. Dasgupta, *Anal. Chem.*, 1987, **59**, 629–636, DOI: 10.1021/ac00131a019.
- 38 P. Torawane, K. Keshav, M. K. Kumawat, R. Srivastava, T. Anand, S. Sahoo, A. Borse and A. Kuwar, *Photochem. Photobiol. Sci.*, 2017, **16**, 1464–1470, DOI: 10.1039/C7PP00182G.
- 39 U. A. Fegade, S. K. Sahoo, A. Singh, N. Singh, S. B. Attarde and A. S. Kuwar, *Anal. Chim. Acta*, 2015, **872**, 63–69, DOI: 10.1016/j.aca.2015.02.051.
- 40 A. Kuwar, R. Patil, A. Singh, R. Bendre and N. Singh, *ChemPhysChem*, 2014, **15**, 3933–3937, DOI: 10.1002/cphc.201402534.
- 41 H. A. Benesi and J. H. Hildebrand, *J. Am. Chem. Soc.*, 1949, **71**(8), 2703–2707, DOI: 10.1021/ja01176a030.
- 42 G. L. Long and J. D. Winefordner, *Anal. Chem.*, 1983, **55**, 712A–724A, DOI: 10.1021/ac00258a724.
- 43 L. Krause, R. Herbst-Irmer, G. M. Sheldrick and D. Stalke, *J. Appl. Crystallogr.*, 2015, **48**, 3–10, DOI: 10.1107/S1600576714022985.
- 44 G. M. Sheldrick, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 2008, **64**, 112–122, DOI: 10.1107/S2053229614024218.
- 45 G. M. Sheldrick, *Acta Crystallogr., Sect. C: Struct. Chem.*, 2015, **71**, 3–8, DOI: 10.1107/S2053229614024218.
- 46 C. F. Macrae, P. R. Edgington, P. McCabe, E. Pidcock, G. P. Shields, R. Taylor, M. Towler and J. van de Streek, *J. Appl. Crystallogr.*, 2006, **39**, 453–457, DOI: 10.1107/S002188980600731X.
- 47 A. Pardo, D. Reyman, J. M. L. Poyato and F. Medina, *J. Lumin.*, 1992, **51**, 269–274, DOI: 10.1016/0022-2313(92)90077-M.
- 48 K. Tayade, S. K. Sahoo, B. Bondhopadhyay, V. K. Bhardwaj, N. Singh, A. Basu, R. Bendre and A. Kuwar, *Biosens. Bioelectron.*, 2014, **61**, 429–433, DOI: 10.1016/j.bios.2014.05.053.
- 49 S. Patil, U. Fegade, S. K. Sahoo, A. Singh, J. Marek, N. Singh, R. Bendre and A. Kuwar, *ChemPhysChem*, 2014, **15**, 2230–2235, DOI: 10.1002/cphc.201402076.
- 50 R. Patil, A. Moirangthem, R. Butcher, N. Singh, A. Basu, K. Tayade, U. Fegade, D. Hundiwale and A. Kuwar, *Dalton Trans.*, 2014, **43**, 2895–2899, DOI: 10.1039/c3dt52770k.
- 51 D. S. Huerta-Joséa, J. G. Hernández-Hernándezb, C. A. Huerta-Aguilarc and P. Thangarasua, *Sens. Actuators, B*, 2019, **293**, 357–365, DOI: 10.1016/j.snb.2019.04.011; C. A. Huerta-Aguilar, T. Pandiyan, N. Singh and N. Jayanthi, *Spectrochim. Acta, Part A*, 2015, **146**, 142–150, DOI: 10.1016/j.saa.2015.03.082; C. A. Huerta-Aguilar, B. Ramírez-Guzmán, P. Thangarasu, J. Narayanan and N. Singh, *Photochem. Photobiol. Sci.*, 2019, **18**, 1761–1772, DOI: 10.1039/C9PP00060G.
- 52 C. A. Huerta-Aguilar, T. Pandiyan, P. Raj, N. Singh and R. Zanella, *Sens. Actuators, B*, 2016, **223**, 59–67, DOI: 10.1016/j.snb.2015.09.064; C. A. Huerta-Aguilar, P. Raj, P. Thangarasu and N. Singh, *RSC Adv.*, 2016, **6**, 37944–37952, DOI: 10.1039/C6RA01231K; C. A. Huerta-Aguilar, T. Pandiyan, N. Singh and N. Jayanthi, *Spectrochim. Acta, Part A*, 2015, **146**, 142–150, DOI: 10.1016/j.saa.2015.03.082; J. A. O. Granados, P. Thangarasu, N. Singh and J. M. Vázquez-Ramos, *Food Chem.*, 2019, **278**, 523–532, DOI: 10.1016/j.foodchem.2018.11.086.
- 53 A. K. Manna, S. Chowdhury and G. K. Patra, *New J. Chem.*, 2020, **44**, 10819–10832, DOI: 10.1039/D0NJ01954B; A. K. Manna, K. Rout, S. Chowdhury and G. K. Patra, *Photochem. Photobiol. Sci.*, 2019, **18**, 1512–1525, DOI: 10.1039/C9PP00114J.
- 54 P. Kaur, D. Sareen and K. Singh, *Talanta*, 2011, **83**(5), 1695–1700, DOI: 10.1016/j.talanta.2010.11.072.
- 55 H. J. Xu, Z. D. Liu, L. Q. Sheng, M. M. Chen, D. Q. Huang, H. Zhang, C. F. Song and S. S. Chen, *New J. Chem.*, 2013, **37**, 274–277, DOI: 10.1039/C2NJ40767A.
- 56 J. S. Lee, S. D. Warkad, P. B. Shinde, A. Kuwar and S. B. Nimse, *Arabian J. Chem.*, 2020, **13**(12), 8697–8707, DOI: 10.1016/j.arabjc.2020.09.061.
- 57 S. Liu, L. Zhang, W. Zan, X. Yao, Y. Yang and X. Liu, *Sens. Actuators, B*, 2014, **192**, 386–392, DOI: 10.1016/j.snb.2013.10.134.

