



 Cite this: *RSC Adv.*, 2021, 11, 2315

# The synthesis and characterization of the octahedral Co<sup>III</sup> complex of a pyrrolopyrrolizine derivative formed with dicyanovinylene-bis-(*meso*-aryl)dipyrri<sup>†</sup>

 Ji-Young Shin \*

A low spin state and diamagnetic Co<sup>III</sup> complex **1** possessing pyrrolopyrrolizine ligands formed with dicyanovinylene-bis-(*meso*-aryl)dipyrri<sup>†</sup> was synthesized *via* the thermally activated metalation with CoCl<sub>2</sub> and isolated *via* column chromatography. The nuclear magnetic resonance of complex **1** revealed diamagnetism, thereby confirming the structure of the octahedral Co<sup>III</sup>-complex of strong-field ligands. The resulting molecular structure of **1** was elucidated by the X-ray diffraction analysis. An arrangement of two pyrrolizine-ligands for the metal chelation was found in the AB–BA order, which was distinct from the case observed during the formation of bis-Ni<sup>II</sup>-expanded porphyrinoids.

 Received 6th November 2020  
 Accepted 15th December 2020

DOI: 10.1039/d0ra09452h

[rsc.li/rsc-advances](https://rsc.li/rsc-advances)

## Introduction

Dipyrri<sup>†</sup> molecules have garnered considerable attention from synthetic and applied chemists due to their proficiency in various wide-ranged applications from medicinal/medical sciences to photo-electronic devices.<sup>1</sup> Especially, dipyrri<sup>†</sup> derivative-generated neutral metal complexes, whose separation, purification, and further handling are significantly beneficial.<sup>2</sup> Thus, dipyrri<sup>†</sup> segments, have been used in manufacturing either singly expanded molecules or largely assembled arrays/polymers that act as proficient building units.<sup>3</sup>

2,3-Dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) has been a suitable oxidant to completely obtain delocalized  $\pi$ -electron conjugations.<sup>4</sup> It has been established that DDQ-adducts were formed during the oxidation of dipyrromethane possessing electron-withdrawing *meso*-aryl. The subsequent transformations of DDQ-adducts were also investigated by treating with Lewis acid in various alcohols and with Lewis base, respectively. Dichlorofuranone-derivatives were formed in a concerted reaction involving the nucleophilic attack of alcohol, the partial elimination of the molecule, followed by the rearrangement of the  $\pi$ -conjugation pathway when treated with Lewis acid (**B** in Fig. 1).<sup>5</sup> Dicyanovinylene-derivatives were formed in a concerted reaction involving the initiation by the

nucleophilic attack of the Lewis base, the partial elimination of the DDQ-segment, followed by the regeneration of a  $\pi$ -conjugation network when treated with the Lewis base (**A** in Fig. 1).<sup>5a</sup>

Furthermore, the nickel metalation of pentafluoro-phenyl-substituted dicyanovinylene compound **A-C<sub>6</sub>F<sub>5</sub>** afforded




Fig. 1 The formation of DDQ adduct and further transformation towards compounds **A** and **B** in the presence of Lewis base and Lewis acid.




Fig. 2 The formation of bis-Ni<sup>II</sup>-expanded porphyrinoids during the metalation of **A-C<sub>6</sub>F<sub>5</sub>** with Ni(OAc)<sub>2</sub>.

Department of Molecular and Macromolecular Chemistry, Graduate School of Engineering, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8603, Japan. E-mail: jyshin@chembio.nagoya-u.ac.jp; Fax: +81-52-747-6771

<sup>†</sup> Electronic supplementary information (ESI) available: NMR, Mass, and absorption spectra as well as X-ray packing diagrams. CCDC 1870416. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d0ra09452h







Fig. 3 The formation of pyrrolo-diazepino-pyrrolizines in the presence of a non-transition metal Lewis acid, either  $\text{BF}_3\cdot\text{OEt}_2$  or  $\text{InBr}_3$ .



Scheme 1 The synthesis of  $\text{Co}^{\text{III}}$  complex 1.

enantiomeric pairs of regioisomeric pairs of bis-nickel complexes of expanded porphyrinoids (Fig. 2) when treated with a transition-metal Lewis acid such as  $\text{Ni}(\text{OAc})_2$ .<sup>6</sup> It was




Fig. 4 The crystal structure of 1: (a) a complex unit, (b) and (c) the first ligand moiety positioning horizontally, and (d) and (e) the second ligand moiety positioning vertically. Figures (c) and (e) show the dihedral angles between the terminal pyrrolones and the coordinating parts in the horizontal and vertical ligand moieties, respectively.

comprehended that the bicyclic pyrrolizine moiety was functionalized during this transformation. It was reported that the use of a semi-metallic Lewis-acid,  $\text{BF}_3\cdot\text{OEt}_2$  or a basic-metal Lewis-acid,  $\text{InBr}_3$ , resulted in the formation of pyrrolo-diazepino-pyrrolizine derivatives (Fig. 3) due to a setback in creating sizeable macrocyclic structures.<sup>7</sup>

Other transition-metal sources were then stimulated to foster the investigation of different metal-complexes of cyclic/chain ligands. It is noteworthy to mention that the use of  $\text{CoCl}_2$  afforded a different alignment (AB-BA order) of ligands as compared with the sequence obtained in the presence of  $\text{Ni}(\text{OAc})_2$  (AB-AB order). The resulting cobalt complex indicated a low spin, diamagnetic, and octahedral  $\text{Co}^{\text{III}}$  complex formed due to facing two strong-field ligands in the same direction. The formation of bis-nickel(II)-expanded porphyrins, involving cross-condensations and further oxidations were successful within the AB-AB alignment of ligands. Herein, the synthetic and structural details are reported.

## Results and discussion

Dicyanovinylene-bis(*meso*-aryl)dipyrin (DCVAD) was prepared following a previously method and used in the formation of cobalt complexes with  $\text{CoCl}_2$  (Scheme 1).

A toluene solution of DCVAD with  $\text{CoCl}_2$  was heated to 100 °C overnight, from which  $\text{Co}^{\text{III}}$  complex 1 was isolated as a yellow fraction *via* silica gel column chromatography. The low solubility of complex 1 made it difficult to isolate on the silica gel, and the isolation yield was about 10%. A by-product exhibited a larger molecular mass value ( $m/z = 2130.82$ ) than that of complex 1, whose structure was predicted to have two cobalt cations that coordinated with three ligand units. The mass per charge ratio ( $m/z$ ) in the MALDI-TOF mass spectrometry of 1 was 1491.89, which was consistent with  $[\text{M} + \text{H}]^+$  for the molecular formula  $\text{C}_{68}\text{H}_{20}\text{CoF}_{20}\text{N}_{12}\text{O}_3$ . The growth of single crystals for the X-ray diffraction analysis was achieved with a solvent combination of chloroform and methanol by the inter-conversion vaporization and slow diffusion of two solvents, which resulted in fine-fit single-crystallization and appropriate diffractions to refine the molecular structure (Fig. 4 and S1–S3†).

As shown in Fig. 4, each molecular origin of  $\text{A-C}_6\text{F}_5$  first formed bicyclo-pyrrolizine, whose initiation inhibited the catalytic reaction of the transition-metal ( $\text{CoCl}_2$ ) Lewis-acid, which occurred at vinylenenitrile. Consequently, the neighbouring pyrrole attached the nitrile carbon nucleophilically, resulting in the formation of a bicycle-pyrrolizine segment. Simultaneously, the conjugation pathway of  $\pi$ -electrons was reorganized for the proficient ligation of the metal-chelation. In the terminal pyrrole rings, the nucleophilic addition of water occurred, and the  $\pi$ -conjugation pathways were fulfilled by the deprotonation of hydroxyl groups, resulting in 2-pyrrolones. Herein, the cobalt-centre was associated with six coordination sites to complete the ligation, in which one terminal pyrrole out of the four still retained  $\alpha$ -free confirmation from the nucleophilic addition of water due to the restricted stabilization of the *enol*-form compared with that of the *keto*-form. It is noteworthy



to mention that two distinct terminal pyrroles took a relative orientation to each other and formed intramolecular hydrogen bonds (H-bonds), which did not contribute to metalation, but eventually stabilized the overall structure of compound **1** (dashed lines representing H-bonds in Fig. 4a).

Nevertheless, almost the same ligand was held between two ligands towards the cobalt centre; one possessing a  $\alpha$ -free pyrrole-terminal and the other that possessed all pyrrolone-terminals disordered the position of ligands in the crystalline state. The octahedral structure compromising the Co<sup>III</sup> centre holds two ligand units, associating with the planarity of two ligands, as shown in Fig. 4b and d. Mean deviations within 20 atoms involved in ligands main skeleton except for the terminal pyrrole/pyrrolone realized hydrogen-bonds and adjacent *meso*-methin carbons were calculated as 0.174 Å for (b) and 0.137 Å for (d). A ligand unit that places vertically in Fig. 4a is presented in (b) and (c), while the other unit that places horizontally in the figure is presented in (d) and (e). Furthermore, it was implied that terminal pyrroles, which were not involved in the metalation, were standing up from the mean planes of chelating parts due to the following factors. Firstly, the steric interference was induced in individual ligand units; the terminal pyrrolone that was not involved in the metal coordination was sterically hindered with the Co<sup>III</sup>-coordinating part. Secondly, hydrogen-bonding interactions between terminal pyrrolones, led to the relative orientation. As a result, dihedral angles were measured as 51.23° for the ligand unit (b) and 51.53° for the ligand unit (d) (Fig. 4c and e). Hydrogen-bond lengths for N–H...O were 2.871 and 2.909 Å.

The metal-chelating imino-/amino-Ns of fully conjugated multi-pyrrolyl ligands play as strong field ligands, in which a low spin state of Co<sup>III</sup> was compromised. The low spin, diamagnetic hexa-coordination of Co<sup>III</sup> was exhibited with <sup>1</sup>H NMR spectroscopy (Fig. 5). Two terminal pyrroles possessing H-bond interactions, where the amino-Ns of pyrroles acted as the proton donors of H-bonds, resonated two singlet-peaks of NH at a downfield shifted area, 10.48 and 10.21 ppm due to the




Fig. 5 The <sup>1</sup>H NMR spectra of complex **1** in CDCl<sub>3</sub> after (a and c) and before (b and d) the addition of D<sub>2</sub>O.




Fig. 6 The <sup>19</sup>F NMR spectrum of **1** in CDCl<sub>3</sub>.

deshielding effect of H-bond, whose peaks were assigned with the addition of D<sub>2</sub>O as disappearing peaks (spectra a and c).

Furthermore, the peaks of pyrrole-βHs were assigned to 6.00–7.81 ppm. Because of the distinct structures of the metal-chelating ligands with the Co<sup>III</sup> centre, twenty protons and twenty fluorines (four *p*-Fs and eight each for *o*- and *m*-Fs of four pentafluorophenyl groups) were assigned in the <sup>1</sup>H and <sup>19</sup>F NMR spectra, respectively (Fig. 5 and 6). A few peaks overlapped as random cases. The homonuclear HH-COSY NMR exhibits eight correlation pairs of the β-protons of pyrrole rings and one correlation peak for the  $\alpha$ - and  $\beta$ -protons of the  $\alpha$ -free terminal pyrrole with nine sets of proton–proton correlations, as shown in Fig. 7. Furthermore, <sup>19</sup>F NMR reveals individually located, twenty fluorine peaks (Fig. 6). The fluorine–fluorine correlations are demonstrated with the FF-COSY NMR analysis (Fig. S8†), whose spectrum shows fluorine sets for four *meso*-pentafluorophenyls as marked with four different symbols. The C=O stretching vibration of 2-pyrrolones of **1** expose broad bands overlapping around 1694 cm<sup>-1</sup> in CH<sub>2</sub>Cl<sub>2</sub>, which is significant for the 2-pyrrolone carbonyl group inducing a weakened energy vibration frequency as compared to common carbonyl groups (Fig. S9†).

CH<sub>2</sub>Cl<sub>2</sub> solution of **1** exhibits a vastly elongated absorption band of about 1100 nm and demonstrates strong absorbance nearly (log  $\epsilon$  = 4.01) at  $\lambda_{\max}$  = 795 nm (Fig. 8).




Fig. 7 The HH COSY NMR (F<sub>1</sub> = F<sub>2</sub> = 500 MHz) spectrum of **1** in CDCl<sub>3</sub>.






Fig. 8 The UV-vis-NIR absorption spectrum of complex **1** in  $\text{CH}_2\text{Cl}_2$ .




Fig. 9 The cyclic voltammograms of complex **1** in  $\text{CH}_2\text{Cl}_2$  ( $0.1 \text{ M } n\text{-Bu}_4\text{NPF}_6$ ). Working electrode: glassy carbon, counter electrode: Pt, reference cell:  $\text{Ag}/\text{AgCl}$ , and  $\text{Fc}/\text{Fc}^+$ : ferrocene/ferrocenium couple.

The difference in potentials for **1** between the 1st oxidation (0.722 V) and 1st reduction (0.194 V) is measured as 0.528 V with cyclic voltammetry (Fig. 9). The small HOMO–LUMO gap of 0.528 V much less than 1.0 V is in good agreement with the absorption band, largely bathochromic shifted, and tailed until 1300 nm. The first reduction process was reversible distinct from the  $\text{Co}^{\text{III}}$  octahedral complex containing macrocyclic, fewer liberty ligands that exposed irreversible waves for the first reduction process due to the increased liberty on the axial ligands of  $\text{Co}^{\text{II}}$  state *via* the transformation from  $\text{Co}^{\text{III}}$  to  $\text{Co}^{\text{II}}$ .<sup>8</sup> Furthermore, unlikely broad waves are observed between reduction potentials, which probably streamed from the liberty and propositioned to the broaden band at the absorption spectrum. The second reduction at  $-0.763 \text{ V}$  for  $\text{Co}(\text{II})/\text{Co}(\text{I})$  was reversible, and no decomposition was reflected in any window of the potential range, 1.0–1.6 V.

## Experimental

### Materials and methods

All reagents obtained from commercial suppliers were used without further purification. Products were column chromatographed on a silica gel (silica gel 300 or 400). The nuclear magnetic resonance (NMR) spectra of synthetic compounds were recorded using a Bruker Avance 300 MHz NMR

spectrometer using the  $\text{CDCl}_3$  solvent and trimethylsilane (TMS) as an internal standard. The X-ray diffraction patterns of the single crystal of  $\text{Co}^{\text{III}}$  complex **1** were collected at 93 K with a Varimax Saturn N instrument operated with the Rigaku operation software package. The molecular structure was analyzed using direct methods with the aid of SHELXS-97 software, and the refinement was accomplished using the SHELXL-97 program. All hydrogen atoms were placed in calculated positions.

### Preparation of $\text{Co}^{\text{III}}$ complex **1**

The precursor molecule **A-C6F5** was prepared following a previously method.<sup>5a</sup> A toluene solution (90 mL) of **A-C6F5** (60 mg, 86.4  $\mu\text{mol}$ ) and  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  (120 mg, 0.924 mmol) was heated to  $100 \text{ }^\circ\text{C}$  for 9 h and then cooled down. The resulting solution was uploaded on aluminium oxide in a column to separate two fractions using eluents having gradually increased polarities, from neat  $\text{CH}_2\text{Cl}_2$  to 1% MeOH in  $\text{CH}_2\text{Cl}_2$ . The first green fraction was separated to obtain the diazepino-pyrrolizine compound (last structure in Fig. 3) as a failed/retained ligand in cobalt metalation: spectral data elucidated the compound.<sup>7</sup> The second fraction with a yellowish-green colour was then collected and washed with water to remove aluminium oxide dissolved in the increased polar eluent. Using brine and  $\text{CH}_2\text{Cl}_2$ , the organic layer was washed, separated, and dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The residue was then precisely purified on a silica gel using neat  $\text{CH}_2\text{Cl}_2$ . A brownish yellow coloured fraction, fluorescent under the projection of a 365 nm UV lamp was collected to obtain compound **1** (10% yield). The  $^1\text{H}$  NMR spectrum of the compound exposes the two significant singlets (10.48 and 10.21 ppm) of amino-NHs, supporting the molecular structure that preserves pyrroles, but not the metalation. Furthermore, the growth of single crystals was achieved with a  $\text{CHCl}_3$  solution and MeOH *via* vapor diffusion.

### Spectral data of **1**‡

$\lambda_{\text{max}}$  (nm (log  $\epsilon$ ),  $\text{CH}_2\text{Cl}_2$ ): 320 (4.34), 456 (4.17), 795 (4.01);  $\delta_{\text{H}}$  (500 MHz, 298 K,  $\text{CDCl}_3$ ): 10.48 (bs, 1H, NH), 10.21 (bs, 1H, NH), 7.80 (d,  $J = 4.5 \text{ Hz}$ , 1H), 7.66 (d,  $J = 4.5 \text{ Hz}$ , 1H), 6.81 (s, 1H), 6.68 (d,  $J = 5.4 \text{ Hz}$ , 1H), 6.67–6.58 (m, 3H), 6.34 (d,  $J = 4.8 \text{ Hz}$ , 1H), 6.28 (d,  $J = 3.6 \text{ Hz}$ , 1H), 6.27 (d,  $J = 3.7 \text{ Hz}$ , 1H), 6.11 (d,  $J = 5.6 \text{ Hz}$ , 1H), 6.08 (d,  $J = 3.7 \text{ Hz}$ , 1H), 6.06 (d,  $J = 5.3 \text{ Hz}$ , 2H), 6.05 (d,  $J = 3.7 \text{ Hz}$ , 1H), 6.03 (d,  $J = 5.7 \text{ Hz}$ , 1H), 6.01 (d,  $J = 5.4 \text{ Hz}$ , 1H);  $\delta_{\text{F}}$  (470.57 MHz, 298 K, calibrated with external  $\text{CF}_3\text{COOH}$ ):  $-134.9$  (d,  $J = 23.5 \text{ Hz}$ , 1F, *o*-F),  $-135.7$  (dd,  $^{\text{d}}J = 51.8 \text{ Hz}$ ,  $^{\text{d}}J =$

‡ Crystallographic data of **1** (CCDC 1870416): ( $\text{C}_{68}\text{H}_{20}\text{CoF}_{20}\text{N}_{12}\text{O}_3$ ) ( $\text{CHCl}_3$ ), 0.38( $\text{CH}_3\text{OH}$ )  $M_r = 1623.19$ ,  $T = 93(2) \text{ K}$ , crystal size =  $0.09 \times 0.09 \times 0.01 \text{ mm}^3$ , Green prism, Mo radiation, triclinic, space group  $P1$  (#2),  $a = 11.204(5) \text{ \AA}$ ,  $b = 16.844(5) \text{ \AA}$ ,  $c = 18.858(5) \text{ \AA}$ ,  $\alpha = 63.767(5)^\circ$ ,  $\beta = 83.998(5)^\circ$ ,  $\gamma = 77.380(5)^\circ$ ,  $V = 3115.1(19) \text{ \AA}^3$ ,  $Z = 2$ ,  $P_{\text{calcd}} = 1.728 \text{ g cm}^{-3}$ ,  $R_1(F) = 0.0972$  ( $I > 2(I)$ ),  $wR_2(F^2) = 0.2944$  (all),  $\text{GoF} = 1.072$ . Disordered solvent molecules were refined in one chloroform and around 0.38 methanol molecules per one molecule of **1**. Furthermore, one of the terminal pyrrolones, positioning the closet to the terminal pyrrole of the partner ligand in the cobalt complex was also disordered: two terminal parts were disordered and placed close to each other by forming hydrogen bonds, N–H...O.





23.5 Hz, 1F, *o*-F),  $-136.2$  (dd,  ${}^{\text{dd}}J = {}^{\text{d}}J = 23.5$  Hz, 1F, *o*-F),  $-137.6$  (dd,  ${}^{\text{dd}}J = 7.1$  Hz,  ${}^{\text{d}}J = 23.5$  Hz, 1F, *o*-F),  $-139.5$  (dd,  ${}^{\text{dd}}J = 51.8$  Hz,  ${}^{\text{d}}J = 23.5$  Hz, 1F, *o*-F),  $-140.1$  (dd,  ${}^{\text{dd}}J = {}^{\text{d}}J = 23.5$  Hz, 1F, *o*-F),  $-140.7$  (d,  $J = 23.5$  Hz, 1F, *o*-F),  $-140.9$  (d,  $J = 23.5$  Hz, 1F, *o*-F),  $-149.5$  (t,  $J = 23.5$  Hz, 1F, *p*-F),  $-149.6$  (t,  $J = 23.5$  Hz, 1F, *p*-F),  $-150.3$  (t,  $J = 23.5$  Hz, 1F, *p*-F),  $-152.8$  (t,  $J = 23.5$  Hz, 1F, *p*-F),  $-159.2$  (td,  ${}^{\text{td}}J = 4.7$  Hz,  ${}^{\text{t}}J = 23.5$  Hz, 1F, *m*-F),  $-159.5$  (td,  ${}^{\text{td}}J = 4.7$  Hz,  ${}^{\text{t}}J = 23.5$  Hz, 2F, *m*-F),  $-159.9$  (td,  ${}^{\text{td}}J = 4.7$  Hz,  ${}^{\text{t}}J = 23.5$  Hz, 1F, *m*-F),  $-160.7$  (t,  $J = 23.5$  Hz, 1F, *m*-F),  $-160.9$  (t,  $J = 23.5$  Hz, 1F, *m*-F),  $-162.1$  (t,  $J = 23.5$  Hz, 1F, *m*-F),  $-163.9$  (t,  $J = 23.5$  Hz, 1F, *m*-F); MALDI TOF MS: calcd for  $[M + nH]^+$ ,  $M = C_{68}H_{19}CoF_{20}N_{12}O_3$ : 1492.087 (calcd), 1492.858 (obs.); MALDI TOF MS: calcd for  $C_{68}H_{19}CoF_{20}N_{12}O_3$  ( $[M + nH]^+$ ): 1492.087; found: 1492.858.

## Conclusions

In conclusion, the  $Co^{III}$  complex **1** was synthesized and isolated from a toluene solution of dicyanovinylene-bis(*meso*-aryl) dipyrriin *via* thermal metalation of the pyrrolo-pyrrolizine derivative. The molecular structure of **1** was elucidated by the X-ray diffraction analysis, in which two pyrrolizine-ligands chelating to the  $Co^{III}$  atom oriented in an AB-BA alignment with the octahedral  $Co^{III}$  complex. Furthermore, NMR spectroscopic and X-ray crystallographic details disclosed the complex **1** to be diamagnetic, low spin octahedral  $Co(III)$ -complex of strong-field ligands.

## Conflicts of interest

There are no conflicts of interest to declare.

## Acknowledgements

This research was supported by the JSPS KAKENHI grant number JP18K04925. J.-Y. Shin acknowledges the Department of Molecular and Macromolecular Chemistry, Graduate School of Engineering, Nagoya University for its great support for the provision of experimental and analytical instruments as well as indispensable advice.

## Notes and references

- (a) A. Kamkaew, S. H. Lim, H. B. Lee, L. V. Kiew, L. Y. Chung and K. Burgess, *Chem. Soc. Rev.*, 2013, **42**, 77; (b) C. S. Kue, S. Y. Ng, S. H. Voon, A. Kamkaew, L. Y. Chung, L. V. Kiew and H. B. Lee, *Photochem. Photobiol. Sci.*, 2018, **17**, 1691; (c) C.-L. Hou, Y. Yao, D. Wang, J. Zhang and J.-L. Zhang, *Org. Chem. Front.*, 2019, **6**, 2266; (d) J. R. Pankhurst, N. L. Bell, M. Zegke, L. N. Platts, C. A. Lamfsus, L. Maron, L. S. Natrajan, S. Sproules, P. L. Arnold and J. B. Love, *Chem. Sci.*, 2017, **8**, 106; (e) L. Lecarme, A. Kochem, L. Chiang, J. Moutet, F. Berthiol, C. Philouze, N. Leconte, T. Storr and F. Thomas, *Inorg. Chem.*, 2018, **57**, 9708; (f) J. R. Pankhurst, N. L. Bell, M. Zegke, L. N. Platts, C. A. Lamfsus, L. Maron, L. S. Natrajan, S. Sproules, P. L. Arnold and J. B. Love, *Chem. Sci.*, 2017, **8**, 108; (g) D. Perl, S. W. Bisset and S. G. Telfer, *Dalton Trans.*, 2016, **45**, 2440.
- (a) H. Fischer and H. Orth in *Die Chemie des Pyrrols*, Akademische Verlagsgesellschaft, Leipzig, 1940, vol. 2, pp.1–151; (b) H. Falk in *The Chemistry of Linear Oligopyrroles and Bile Pigments*, Springer, Vienna, 1989; (c) C. Brückner, Y. Zhang, S. J. Rettig and D. Dolphin, *Inorg. Chim. Acta*, 1997, **263**, 279; (d) R. S. Singh, R. P. Paitandi, R. K. Gupta and D. S. Pandey, *Coord. Chem. Rev.*, 2020, **414**, 213269; (e) S. A. Baudron, *Dalton Trans.*, 2020, **49**, 6161.
- (a) C. V. K. Sharma, G. A. Broker, J. G. Huddleston, J. W. Baldwin, R. M. Metzger and R. D. Rogers, *J. Am. Chem. Soc.*, 1999, **121**, 1137; (b) H. Yanagi, H. Mukai, K. Ikuta, T. Shibusani, T. Kamikado, S. Yokoyama and S. Mashiko, *Nano Lett.*, 2002, **2**, 601; (c) Q. Chen and D. Dolphin, *Can. J. Chem.*, 2002, **80**, 1668; (d) A. Thompson, S. J. Rettig and D. Dolphin, *Chem. Commun.*, 1999, 631; (e) Y. Zhang, A. Thompson, S. J. Rettig and D. Dolphin, *J. Am. Chem. Soc.*, 1998, **120**, 13537; (f) A. Thompson and D. Dolphin, *Org. Lett.*, 2000, **2**, 1315; (g) R. Taniguchi, S. Shimizu, J.-Y. Shin, H. Furuta and A. Osuka, *Tetrahedron Lett.*, 2003, **44**, 2505; (h) J.-Y. Shin, D. Dolphin and B. O. Patrick, *Cryst. Growth Des.*, 2004, **4**, 659; (i) J.-Y. Shin, B. O. Patrick and D. Dolphin, *CrystEngComm*, 2008, **10**, 960; (j) Q. Miao, J.-Y. Shin, B. O. Patrick and D. Dolphin, *Chem. Commun.*, 2009, 2541; (k) Z. Zhang and D. Dolphin, *Chem. Commun.*, 2009, 6931; (l) L. Ma, J.-Y. Shin, B. O. Patrick and D. Dolphin, *CrystEngComm*, 2008, **10**, 1539; (m) Y. Chiba, T. Nakamura, R. Matsuoka and T. Nabeshima, *Synlett*, 2020, **31**, 1663.
- (a) D. Walker and J. D. Hiebert, *Chem. Rev.*, 1967, **67**, 153; (b) J. S. Lindsey, I. C. Schreiman, H. C. Hsu, P. C. Kearney and A. M. Marguerettaz, *J. Org. Chem.*, 1987, **52**, 827; (c) L. Yu, K. Muthukumar, J. V. Sazanovich, C. Kirmaier, E. Hindin, J. R. Diers, P. D. Boyle, D. f. Bocian, D. Holten and J. S. Lindsey, *Inorg. Chem.*, 2003, **42**, 6629; (d) G. R. Geier III, J. F. B. Chick, J. B. Callinan, C. G. Reid and W. P. Auguscinski, *J. Org. Chem.*, 2004, **69**, 4159; (e) L. Capella, P. C. Montevecchi and D. Nanni, *J. Org. Chem.*, 1994, **59**, 7379; (f) C. Brückner, V. Karunaratne, S. J. Rettig and D. Dolphin, *Can. J. Chem.*, 1996, **74**, 2182; (g) J. S. Lindsey, in *The porphyrin Handbook* ed. K. M. Kadish, K. M. Smith and R. Guilard, Academic Press, San Diego, CA, 2000, vol. 1; (h) Z. Gross, N. Galili and I. Saltsman, *Angew. Chem., Int. Ed.*, 1999, **38**, 1427; (i) J.-Y. Shin, H. Furuta, K. Yoza, S. Igarashi and A. Osuka, *J. Am. Chem. Soc.*, 2001, **123**, 7190; (j) T. E. Wood and A. Thompson, *Chem. Rev.*, 2007, **107**, 1831.
- (a) J.-Y. Shin, B. O. Patrick and D. Dolphin, *Org. Biomol. Chem.*, 2009, **7**, 2032; (b) J.-Y. Shin and D. Dolphin, *New J. Chem.*, 2011, **35**, 2483.
- N. H. Faialaga, S. Ito, H. Shinokubo, Y. Kim, K. Kim and J.-Y. Shin, *Dalton Trans.*, 2017, **46**, 10802.
- J.-Y. Shin, *RSC Adv.*, 2019, **9**, 40031.
- A. Mahammed, B. Mondal, A. Rana, A. Dey and Z. Gross, *Chem. Commun.*, 2014, **50**, 2725.

