


 Cite this: *RSC Adv.*, 2021, 11, 10804

# Characterization of carotenoids and phenolics during fruit ripening of Chinese raspberry (*Rubus chingii* Hu)

 Xiaobai Li,<sup>a</sup> Jian Sun,<sup>b</sup> Zhen Chen,<sup>c</sup> Jingyong Jiang<sup>d</sup> and Aaron Jackson<sup>e</sup>

Chinese raspberry (*Rubus chingii* Hu) is a fruit valued for its health benefits, which is indigenous to China. It is a great source of antioxidants. However, the fruit phytochemicals are poorly understood. Phenolics and carotenoids attract much attention for their antioxidant capability, and they dramatically change during fruit ripening, leading to the difference in color, flavor and medicinal components. In this study, we investigated the change of carotenoids, phenolics and antioxidant activity using spectrophotometry during four different ripening phases *i.e.* mature green (MG), green yellow (GY), yellow orange (YO) and red (RE). The major components of carotenoids, anthocyanins, ellagitannins and flavonols were identified and quantified by LC-MS/MS. As a result, five carotenoids (mainly  $\beta$ -Citaurin and its esters), six anthocyanins (mainly anthocyanins covalently linked to another flavonoid unit), methyl (S)-flavogallate and rourinoside were first identified in *Rubus*. In contrast to other known raspberries, *R. chingii* had a continuous decrease in total phenolics during fruit ripening, which was due to a continuous decrease in flavonoids (including anthocyanin). Total anthocyanin and flavonoid respectively declined from 19.5 to 6.9 mg/100 g FW, and 646.2 to 128.5 mg/100 g FW during fruit maturation and coloration. Accordingly, the components of anthocyanins, ellagitannins and flavonols also declined, thus resulting in a decrease in antioxidant activity (from 41.2 to 10.1 TEAC/100 g FW in ABTS and from 35.3 to 7.7 mmol TEAC/100 g FW in FRAP). In contrast, total carotenoid increased from 184.2 to 305.4 mg/100 g FW. Accordingly, the components of carotenoids also increased, with the exception of lutein. Additionally, kaempferol and quercetin were the main flavonoid aglycones, which were linked to a variety of glycosides. These kaempferol- and quercetin-glycosides mainly accumulated in epidermal hair and placenta. Notably, carotenoids (*i.e.*  $\beta$ -citaurin esters), instead of anthocyanins, gradually accumulated during fruit ripening, imparting the reddish color to ripe fruit.

Received 9th December 2020

Accepted 12th January 2021

DOI: 10.1039/d0ra10373j

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

## 1. Introduction

*Rubus chingii* Hu with the Chinese name “Fu-Pen-Zi” is also called Chinese raspberry, which is indigenous to China. The unripe fruit is often used in Chinese medicine. It provides health promoting and protective properties against a variety of human diseases, *e.g.* improving renal function,<sup>1</sup> protecting hepatocyte function<sup>2</sup> and relieving anxiety, pain and inflammation.<sup>3</sup> The ripe fruit is a nutritional fruit, just like red or black raspberry.

The health benefits of the *Rubus* genus are believed to be mainly due to their abundance of phenolics and carotenoids. In *Rubus*, phenolics mainly consist of anthocyanins, flavonols and ellagitannins.<sup>4,5</sup> Red and black raspberry share the same profile of anthocyanins. Their anthocyanins are predominantly cyanidin glycosides (*e.g.* glucosides, sophorosides, rutinosides, sambubioside and glucosyl-rutinosides), but they only contain low to trace levels of pelargonidin glycosides.<sup>4,6–8</sup> Black raspberry has up to five-fold greater anthocyanin content than red raspberry.<sup>9,10</sup> Flavonols in red and black raspberry are mainly kaempferol/quercetin glycosides with glucosides, rutinoside and coumaroylglucoside.<sup>7,8</sup> Ellagitannins in red and black raspberry comprise dimeric HHDP (hexahydroxydiphenic) sanguin H-6 and a tetrameric HHDP lambertianin C, as well as ellagic acid.<sup>7,10,11</sup> Previous studies have focused on a few compounds in unripe fruit,<sup>1,2,12</sup> but a comprehensive analysis of phenolics throughout the whole fruit ripening process has not been done until now.

Carotenoids benefit human health for antioxidant capability of combating the “superoxide anion radical” to reduce cancer

<sup>a</sup>Zhejiang Academy of Agricultural Sciences, Hangzhou, 310021, China. E-mail: hufanfan1982815@outlook.com; lixiaobai@mail.zaas.ac.cn

<sup>b</sup>Zhejiang Research Institute of Traditional Chinese Medicine Co., Ltd., Hangzhou, 310023, China

<sup>c</sup>College of Life Sciences, Taizhou University, Taizhou, 318000, China. E-mail: chenzh@tztc.edu.cn

<sup>d</sup>Taizhou Academy of Agricultural Sciences, Linhai, 317000, China. E-mail: jiy5971@163.com

<sup>e</sup>South Oak, Stuttgart, AR 72160, USA. E-mail: sativadna@gmail.com



risk. Some components are transformed into vitamin A, which is required for healthy skin and mucus membranes, and night vision. In raspberries fruit, apocarotenoids are very abundant e.g.  $\alpha$ - and  $\beta$ -ionone, responsible for a large part of the characteristic raspberry aroma, but amounts of xanthophyll are relatively low.<sup>13</sup> However, the information on carotenoids in *R. chingii* has been very limited until now.

In *R. chingii*, the unripe and ripe fruits are used differently. The different uses are attributed to the discrepancy in phytochemicals, especially for phenolics and carotenoids. They dramatically change throughout the process of fruit ripening, which has attracted considerable research attention in other *Rubus* species. This study was undertaken to investigate the composition of phenolics and carotenoids, and their changes during fruit ripening.

## 2. Experimental

### 2.1. Plant material

*R. chingii* fruits were collected from five to six plants at different ripening phases i.e. mature green (MG), green yellow (GY), yellow orange (YO) and red (RE) during the growing season (May, 2019) at Linhai, Zhejiang, China (Fig. 1a). Ten fruits were pooled as one replicate. Three biological replicates were designed for further experiments. The whole fruit were ground with liquid nitrogen into powders, which were used for analysis of mRNA, protein and metabolites. The whole fruit tissues were “ground” in liquid nitrogen into “powder” for further analysis.

### 2.2. Total carotenoid, anthocyanin, flavonoid, and phenolic content

Fruits were ground with liquid nitrogen and into powder, of which approximately 0.3 g was mixed with 8 mL extraction solvent (ethanol : acetone = 1 : 2). The extract was sonicated and set aside in the dark for half-hour until the residues became colorless. The absorbance of the extract was measured at 440, 645 and 663 nm for carotenoid, chlorophyll a and b respectively.

$$\text{Chlorophyll a (mg g}^{-1}\text{ FW)} = 0.01 \times (12.7 \times A_{663} - 2.69 \times A_{645}) \times V/(M \times 1000);$$

$$\text{Chlorophyll b (mg g}^{-1}\text{ FW)} = 0.01 \times (22.9 \times A_{645} - 4.68 \times A_{663}) \times V/(M \times 1000);$$

$$\text{Total chlorophyll (mg g}^{-1}\text{ FW)} = (20.21 \times A_{645} + 8.02 \times A_{663}) \times V/(M \times 1000);$$

$$\text{Carotenoids (mg g}^{-1}\text{ FW)} = 4.695 A_{440} - 0.268 [\text{chlorophyll (a + b)}] \times V/(M \times 1000);$$

where  $V$  is the extract volume (mL) and  $M$  is the mass of the tissue extracted (g).

Total anthocyanin content was determined *via* spectrophotometry. Approximately, 0.3 g of ground tissue with liquid nitrogen were added to 10 mL 1% (v/v) HCl methanol and sonicated for half-hour at room temperature in the dark. After

centrifuging, supernatants were measured for absorbance at 530, 620 and 650 nm. The anthocyanin content was estimated using the following formulas:

$$\text{The anthocyanin content} = [(A_{530} - A_{620}) - 0.1 \times (A_{650} - A_{620})] \times V \times M/(\epsilon \times m).$$

where  $V$  is the extract volume (mL),  $\epsilon$  is the molar extinction coefficient of cyanidin-3-glucoside at 530 nm (29 600),  $M$  is the molecular weight of cyanidin-3-glucoside (449 g mol<sup>-1</sup>), and  $m$  is the mass of the fruit extracted. The results were expressed as cyanidin-3-glucoside equivalents (mg CG/g FW).


Total flavonoid content was quantified by a colorimetric assay method. Approximately 0.3 g of tissue powder was mixed with 10 mL ethanol for 2 h at room temperature in the dark, and centrifuged. Of supernatant, 1 mL was mixed with 2.4 mL ethanol and 0.4 mL NaNO<sub>2</sub>. After 6 min, the mixture was added to 0.4 mL 10% Al(NO<sub>3</sub>)<sub>3</sub> solution. After an additional 6 min, the mixture was added to 4 mL 4% NaOH and filled to 10 mL with 100% ethanol. After 15 min, the absorbance was determined at 510 nm and measured relative to a blank extraction solvent. Total flavonoid content was expressed as rutin equivalent (mg RE/g FW). Total phenolic content was determined using the Folin-Ciocalteu method following the procedure.<sup>14</sup> Fruit tissue was finely ground in liquid nitrogen. Of tissue powders, approximately 0.3 g was mixed with 10 mL of acidified methanol (0.1% hydrochloric acid) and sonicated in ice for half-hour in the dark and centrifuged. Two mL of supernatant was transferred to another colorimetric tube, mixed with 1 mL 0.5 N Folin-Ciocalteu's phenol reagent, and set for 5 min. The reaction was neutralized with 2 mL of 5% saturated Na<sub>2</sub>CO<sub>3</sub> and incubated for 60 min at 30 °C. The absorbance was measured at 760 nm. TPCs were expressed as gallic acid equivalent (mg GAE/g FW).

### 2.3. ABTS assay and FRAP assay

The free radical-scavenging activity was determined by ABTS radical cation decolorization method.<sup>15</sup> Approximately 0.3 g was weighed and added to H<sub>2</sub>O. ABTS radical cation (ABTS<sup>•+</sup>) was obtained by mixing 7.0 mmol L<sup>-1</sup> ABTS solution with 2.45 mmol L<sup>-1</sup> potassium persulfate at 2 : 1 (v/v) and stored in the dark for at least 16 h. The ABTS<sup>•+</sup> solution was diluted with 80% ethanol until its absorbance reached 0.700 ± 0.02 at 734 nm. The ABTS<sup>•+</sup> solution (4.0 mL, absorbance of 0.700 ± 0.02) was thoroughly mixed with 0.1 mL appropriately diluted fruit aqueous extract. The mixture was placed at room temperature for 6 min, and its absorbance was immediately measured at 734 nm. Results were expressed at Trolox equivalent (mmol TEAC/g FW).

The reducing antioxidant power of samples was determined using the FRAP method.<sup>15</sup> Fruit tissue was completely ground with liquid nitrogen, and approximately 0.3 g of tissue powder was weighed and added to H<sub>2</sub>O. The FRAP reagent (0.3 M, pH3.6 acetate buffer, 10 M TPTZ in 40 M HCl, and 20 M FeCl<sub>3</sub>, v/v/v = 10 : 1 : 1) was freshly prepared. The FRAP reagent (1.0 mL) and appropriately diluted fruit aqueous extract (0.05 mL) were mixed thoroughly. The absorbance was measured at 593 nm





**Fig. 1** Dynamic change of fruit appearance, anthocyanin, carotenoid, flavonoid and phenolic content, and antioxidant capacity during maturation in *R. chingii* Hu. (a) Fruit appearance (b) anthocyanin and carotenoid (c) phenolics and flavonoids, (d) the free radical-scavenging activity (ABTS) and reducing antioxidant power (FRAP) and (e) Pearson correlation between quality trait. Fruit maturation are MG: mature green, GY: green yellow, YO, yellow orange, RE: red. \*: significant difference when compared to MG ( $P < 0.05$ ).

after incubation at 37 °C for 30 min. Results were expressed as Trolox equivalent (mmol TEAC/g FW).

#### 2.4. LC-MS/MS analysis of carotenoids, anthocyanins and flavonoids

Carotenoids were extracted by hexane/acetone/ethanol (1 : 1 : 1) solution and sonicated in an ice bath for 30 min. The extracts was concentrated by CentriVap Refrigerated Centrifugal Concentrators at 10 °C (Labconco Models 73100 Series). The residue was dissolved with THF/acetonitrile/methanol (15 : 30 : 55) solution to a final volume of 1 mL, passed through a 0.45 μm microporous membrane filter for UPLC-MS/MS analysis. Carotenoids were separated by HPLC with YMC

Carotenoid C30 column (4.6 mm × 250 mm, YMC, Japan). The mobile phases were methanol (A) and acetonitrile/isopropanol (1 : 1) (B). The linear gradient was: 0 min, 5%B; 60 min, 95% B; 70 min, 95%B; sample injection, 5 μL; column oven temperature, 25 °C; flow rate, 1 mL min<sup>-1</sup>.

Anthocyanins were extracted with 1% (v/v) HCl methanol, concentrated by CentriVap refrigerated Centrifugal Concentrators at 8 °C (Labconco Models 73100 Series) and then re-dissolved with 1 mL 1% (v/v) HCl methanol. Flavonoids was extracted with 70% methanol for 2 h at room temperature in the dark, and refrigerated Centrifugal Concentrators at 8 °C (Labconco Models 73100 Series) and then re-dissolved it with 1 mL 70% methanol. The anthocyanin and flavonoid extracts were



passed through a 0.22  $\mu\text{m}$  microporous membrane filter for LC-MS analysis. Anthocyanin and flavonoid were separated by UPLC with an ACQUITY UPLC HSS T3 column (1.8  $\mu\text{m}$ , 2.1  $\times$  150 mm; Waters Corp.). For anthocyanin, the mobile phases were 1% formic acid-water (A) and acetonitrile (B). The linear gradient was as follows, 0/5, 25/35, 37/95 (min/B%); sample injection volume, 5  $\mu\text{L}$ ; column oven temperature, 50  $^{\circ}\text{C}$ ; flow rate, 0.4  $\text{mL min}^{-1}$ ; and the UV detector was set at 530 nm. For flavonoids and ellagitannins, the mobile phases were 0.1% formic acid-water (A) and 0.1% formic acid-acetonitrile (B). The linear gradient was as follows, 0/5, 5/10, 25/25, 37/95 (min/B%); sample injection volume, 5  $\mu\text{L}$ ; column oven temperature, 25  $^{\circ}\text{C}$ ; flow rate, 0.3  $\text{mL min}^{-1}$ ; and the UV detector was set at 280 and 360 nm.

The separated carotenoids and anthocyanin were analyzed by MS AB Triple TOF 5600<sup>plus</sup> System (AB SCIEX, Framingham, USA) in positive ion mode (source voltage was +5.5 kV, and the source temperature was 600  $^{\circ}\text{C}$ ). The separated flavonoids were analyzed in both negative ion (source voltage at -4.5 kV, and source temperature at 550  $^{\circ}\text{C}$ ) and negative ion (source voltage was +5.5 kV, and the source temperature was 600  $^{\circ}\text{C}$ ). Maximum allowed error was set to  $\pm 5$  ppm. Declustering potential (DP), 100 V; collision energy (CE), 10 V. For MS/MS acquisition mode, the parameters were almost the same except that the collision

energy (CE) was set at  $40 \pm 20$  V, ion release delay (IRD) at 67, ion release width (IRW) at 25. The IDA-based auto-MS<sup>2</sup> was performed on the 8 most intense metabolite ions in a cycle of full scan (1 s). The scan range of  $m/z$  of precursor ion and product ion were set as 100–2000 Da and 50–2000 Da. The exact mass calibration was performed automatically before each analysis employing the Automated Calibration Delivery System.

The content of anthocyanin compounds was expressed as pelargonidin 3-glucoside equivalents (mg PG/g FW). The content of flavanol, ellagitannins and hydroxybenzoic acid components was determined based on their corresponding standard subtracts except methyl (*S*)-flavogallate and casuarictin (galloyl-bis-HHDP-glucose), ellagic acid pentoside, rourinoside, kaempferol-3-*o*-rutinoside isomer, and kaempferol 3-*O*-hexoside isomer. The content of ellagitannins components was expressed as ellagic acid equivalents (mg g<sup>-1</sup> FW). The content of carotenoid compounds was estimated based on the standard curve of lutein and expressed as lutein equivalents.

## 2.5. Frozen sections for 2-aminoethyl diphenylborate (DPBA) staining *in situ*

The fresh fruits were cut into several parts, and then embedded in a dedicated embedding medium (SCEM; Section-Lab). The




Fig. 2 Analysis of the representative carotenoid and anthocyanin compounds in *R. chingii*. (a) UPLC spectra of anthocyanins listed in table, (b) HPLC spectra of carotenoids in table. The upper part: electrospray product ion mass spectra; the lower part: fragment ions mass spectra; on the right: putative molecular structure and cleavage pattern. As mass spectra showed, peak 2 and peak 4 shared another fragment (Afzelechin(4 $\alpha$ ->8) pelargonidin glucoside,  $m/z$  705) by losing one glucoside. However, peak 4 had a higher ratio in 705/543 than peak 2, which indicates that peak 4 has more chances than peak 2 to lose one glucoside, while peak 2 has more chances than peak 4 to lose two glucosides. It is highly probable that two glucosides are connected in peak 2 and separated in peak 4.



embedded tissue was immediately frozen at  $-20\text{ }^{\circ}\text{C}$ . The frozen samples in embedding medium were trimmed and then carefully sliced to produce 50–80  $\mu\text{m}$  fresh-frozen sections using a Cryostat (CM1850; Leica Microsystems) set at  $-20\text{ }^{\circ}\text{C}$ . The sections of fruit were stained in a freshly prepared aqueous solution of 0.25% (w/v) DPBA and 0.00375% (v/v) Triton X-100 for at least 30 min. A Zeiss LSM Zeiss LSM880 confocal laser scanning microscope was used to excite the roots with 30% maximum laser power at 458 nm, and the fluorescence was collected at 475–504 nm for kaempferol and 577 to 619 nm for quercetin.<sup>16</sup>

### 3. Results and discussion

#### 3.1. Composition analysis of carotenoids, anthocyanins and fruit coloration

The fruit color changed during ripening (Fig. 1a) and the content of two major pigments (anthocyanin, and carotenoids) were examined (Fig. 1b). It was surprising that carotenoids were very abundant in content while anthocyanins were relatively low. More surprisingly, total anthocyanin decreased in content while total carotenoids increased. The pattern of anthocyanin change was different from previous reports in other known berries *e.g.* red raspberry, black raspberry, blueberry and strawberry *etc.* Even in other species of *Rubus*, *i.e.* red and black raspberries, concentration of fruit anthocyanins continuously increases throughout ripening.<sup>17,18</sup> However, the pattern of carotenoid change is different among fruit crops. Tomato and loquat carotenoids progressively accumulate during fruit development and maturation.<sup>19–21</sup> Contrarily, other fruit crops, fruit carotenoids do not accumulate but gradually decrease during ripening.<sup>22</sup> The results suggest that the reddish color of mature fruit was probably associated with carotenoids rather than anthocyanins.

These anthocyanins consisted of monomeric (*i.e.* pelargonidin 3-*O*-glucoside and cyanidin-3-(6''-cinnamoyl)glucoside and polymeric anthocyanins covalently linked to another flavonoid unit (Fig. 2a). In agreement to total anthocyanin, these compounds also showed a continuous decrease in content during the fruit maturation process in *R. chingii* (Table 1). This pattern of anthocyanin change was negatively correlated with its red coloration during ripening. These polymeric anthocyanins were first identified in *R. chingii*, which have not been reported in *Rubus*. Of them, four were flavanol–anthocyanins, and pelargonidin was the main type of anthocyanin aglycones. Flavanol–anthocyanins, derived from a spontaneous condensation reaction between anthocyanins and flavanols, is usually found during storage and processing in plant-derived foods.<sup>23</sup> Also, this type of anthocyanin (purple-colored pigments) has been reported in small amounts in a few plants. For example, 5-carboxypyranocyanidin/pelargonidin glycosides have been found in red onion<sup>24</sup> and strawberries<sup>25</sup> while 5-methylpyranocyanidin/delphinidin glycosides in blackcurrant seeds.<sup>26</sup> A kind of cyanidin pigment linked to gallic acid with the C–C bonds, have been found in petals of *Rosa*.<sup>27</sup> Additionally, anthocyanin linked to (epi)catechin or (epi)afzelechin moieties have been found in strawberry,<sup>25</sup> runner beans and purple

corn.<sup>28</sup> Pelargonidin is the main type of aglycone for flavanol–anthocyanin in strawberry, while cyanidin is the main type in runner beans, and purple corn. In red raspberry, cyanidin 3-glucoside is the most prominent, followed by cyanidin 3-glucosylrutinoside and cyanidin 3-rutinoside,<sup>17</sup> and then by pelargonidin 3-(glucosyl) rutinoside, cyanidin-3-*O*-rutinoside, and cyanidin-3-glucoside.<sup>18</sup> Similarly, in black raspberry, anthocyanin increases as fruit matures,<sup>29</sup> and cyanidin-3-*O*-rutinoside is the primary anthocyanin, followed by cyanidin-3-xylosylrutinoside and cyanidin-3-sambubioside.<sup>30</sup> The results indicate that *R. chingii* has a special profile of anthocyanin, which is very different from other known species in *Rubus*.

These carotenoids primarily consisted of three apocarotenoids ( $\beta$ -citaurin,  $\beta$ -citaurin laurate, and  $\beta$ -citaurin myristate), and two xanthophylls (zeaxanthin, and lutein) (Fig. 2b).  $\beta$ -citaurin and its esters, and zeaxanthin gradually accumulated while lutein gradually decreased (Table 1). In fact, zeaxanthin was the precursor of these apocarotenoids. This is the first time these  $\beta$ -citaurin esters have been identified in *Rubus*.  $\beta$ -citaurin is a C30 apocarotenoid first discovered in Sicilian oranges,<sup>31</sup> and gives rise to the peel color of citrus fruits which can range from yellow to red.<sup>31,32</sup> However, the accumulation of  $\beta$ -citaurin is not common, and only observed in the flavedos of some varieties during fruit ripening.<sup>33</sup> Raspberry is another one of the few fruits accompanied by xanthophyll degradation but by a massive apocarotenoids production during fruit ripening.<sup>13</sup> In spite of the fact that raspberries are very rich in the apocarotenoids *i.e.*  $\alpha$ - and  $\beta$ -ionone, responsible for a large part of the characteristic raspberry aroma, they contain relatively low amounts of carotenes.<sup>13,34</sup> Raspberry carotenoids are diverse in composition. In yellow and red raspberry, ripe fruits contain considerable amounts of free lutein, esterified lutein (saturated fatty acids) and apocarotenoids ( $\alpha$ - and  $\beta$ -ionone), but a small amount of zeaxanthin, phytoene,  $\beta$ -carotene and  $\alpha$ -carotene.<sup>13,35</sup> Lutein and  $\beta$ -carotene decrease in content while phytoene, esterified lutein and apocarotenoids ( $\alpha$ - and  $\beta$ -ionone) increase in content during fruit ripening. In a wild raspberry (*Rubus palmatus*),  $\beta$ -cryptoxanthin accumulates during fruit ripening.<sup>36</sup> The results indicate that  $\beta$ -citaurin and its esters are species-specific products in *R. chingii*.

Taken together, the reddish coloration of fruit is caused by  $\beta$ -citaurin and its esters rather than any component of anthocyanins. The profile of *R. chingii* carotenoids and anthocyanins is very different from that of other known *Rubus* species.

#### 3.2. Phenolics and antioxidant capability

Total flavonoid content averaged 3.3 mg RE/100 g FW, and it decreased by 42.1%, 55.3%, and 23.3% from MG to GY, YO, and RE (Fig. 1b). Total phenolic content averaged 20.89 mg GAE/100 g FW, and dropped by 39.0%, 54.7%, and 31.6% from MG to GY, YO, and RE respectively (Fig. 1b). It was surprising to see that the total phenolics (including anthocyanins and other flavonoids) showed a continuous decrease during the fruit maturation process in *R. chingii*. The pattern of phenolic change was different from any other report in *Rubus* species including red and black raspberry. In red raspberry fruit, the total



Table 1 Dynamic change in content of carotenoid, anthocyanin ellagitannins, hydroxybenzoic derivatives, and flavonols during fruit ripening<sup>a</sup>

	Peak	Compound	MG ( $\mu\text{g g}^{-1}$ )	GY ( $\mu\text{g g}^{-1}$ )	YO ( $\mu\text{g g}^{-1}$ )	RE ( $\mu\text{g g}^{-1}$ )
Carotenoid	1	$\beta$ -Citaurin	42 $\pm$ 2	72 $\pm$ 3*	226 $\pm$ 10*	489 $\pm$ 20
	2	Lutein	1450 $\pm$ 60	1022 $\pm$ 50*	239 $\pm$ 10*	111 $\pm$ 4
	3	Zeaxanthin	375 $\pm$ 20	971 $\pm$ 50*	1136 $\pm$ 50*	1172 $\pm$ 40
	4	$\beta$ -Citaurin laurate	0 $\pm$ 0	307 $\pm$ 10*	2961 $\pm$ 100*	10 543 $\pm$ 400
	7	$\beta$ -Citaurin myristate	405 $\pm$ 20	234 $\pm$ 10*	556 $\pm$ 20*	1494 $\pm$ 50
Anthocyanin	8	Cyanidin-3-(6''-cinnamoyl)glucoside	18.8 $\pm$ 0.7	12.0 $\pm$ 0.5*	11.6 $\pm$ 0.4*	9.7 $\pm$ 0.4
	9	Afzelechin(4 $\alpha$ ->8)pelargonidin 3,5-O-diglucoside	17.6 $\pm$ 0.6	12.0 $\pm$ 0.5*	11.6 $\pm$ 0.5*	10.0 $\pm$ 0.4
	10	Catechin(4 $\alpha$ ->8)pelargonidin 3-O- $\beta$ -D-glucopyranoside	53 $\pm$ 2	44 $\pm$ 1*	31 $\pm$ 1*	16.5 $\pm$ 0.8
	11	Afzelechin(4 $\alpha$ ->8)pelargonidin 3-O-sophoroside	123 $\pm$ 4	66 $\pm$ 2*	59 $\pm$ 3*	21 $\pm$ 1
	12	Afzelechin(4 $\alpha$ ->8)pelargonidin 3-O- $\beta$ -D-glucopyranoside	73 $\pm$ 3	57 $\pm$ 2*	40 $\pm$ 2*	19.6 $\pm$ 0.9
	13	Pelargonidin 3-O-glucoside	66 $\pm$ 2	45 $\pm$ 2*	41 $\pm$ 2*	23 $\pm$ 1
Ellagitannins	14	Methyl (S)-flavogallate <sup>b</sup>	482 $\pm$ 20	279 $\pm$ 20*	125 $\pm$ 5*	95 $\pm$ 4
	15	Casuarictin (galloyl-bis-HHDP-glucose) <sup>b</sup>	803 $\pm$ 40	490 $\pm$ 30*	222 $\pm$ 10*	151 $\pm$ 6
Hydroxybenzoic derivatives, and flavonols	16	Ellagic acid pentoside <sup>b</sup>	25.4 $\pm$ 0.9	14.6 $\pm$ 0.5*	3.4 $\pm$ 0.1*	4.0 $\pm$ 0.1
	17	Rourinoside <sup>b</sup>	2.33 $\pm$ 0.08	1.34 $\pm$ 0.04*	0.49 $\pm$ 0.02*	0.32 $\pm$ 0.01
	18	Ellagic acid	50 $\pm$ 2	30 $\pm$ 1*	14.0 $\pm$ 0.8*	9.7 $\pm$ 0.5
	19	Rutin (quercetin 3-O-rutinoside)	4.7 $\pm$ 0.3	2.7 $\pm$ 0.1*	1.21 $\pm$ 0.06*	0.93 $\pm$ 0.05
	20	Isoquercitrin (quercetin 3-glucoside)	5.8 $\pm$ 0.3	3.4 $\pm$ 0.2*	1.51 $\pm$ 0.07*	1.16 $\pm$ 0.06
	21	Kaempferol-3- <i>o</i> -rutinoside isomer <sup>c</sup>	26.3 $\pm$ 0.8	15.2 $\pm$ 0.8*	6.8 $\pm$ 0.3*	3.1 $\pm$ 0.1
	22	Kaempferol 3-O-hexoside isomer <sup>c</sup>	13.3 $\pm$ 0.5	7.0 $\pm$ 0.3*	3.2 $\pm$ 0.1*	2.11 $\pm$ 0.08
	23	Nikotoflorin (kaempferol-3- <i>o</i> -rutinoside)	61 $\pm$ 3	31 $\pm$ 2*	13.5 $\pm$ 0.8*	9.7 $\pm$ 0.4
	24	Astragalin (kaempferol-3-glucoside)	31 $\pm$ 2	18 $\pm$ 1*	8.2 $\pm$ 0.4*	6.3 $\pm$ 0.3
	25	Tiliroside (kaempferol-3- <i>p</i> -coumaroylglucoside)	17 $\pm$ 1	10.4 $\pm$ 0.6*	4.7 $\pm$ 0.2*	3.2 $\pm$ 0.2

<sup>a</sup> Carotenoid was expressed as lutein equivalents. Anthocyanin was expressed as pelargonidin 3-O-glucoside equivalents. <sup>b</sup>: was expressed as Ellagic acid equivalents. <sup>c</sup>: was expressed as kaempferol-3-*o*-rutinoside equivalents. \*: *T*-test ( $P < 0.05$ ).

phenolic concentration decreases from the green to the veraison stage, and then increases until maturity.<sup>17</sup> The increasing trend in anthocyanins and "V"-type change in phenolics are ubiquitous during fruit ripening in many berries, e.g. blueberry,<sup>14</sup> cranberry,<sup>37</sup> strawberry,<sup>38</sup> and grape.<sup>39</sup> The increase in phenolics at the later stage of maturation is mainly due to substantial increases in anthocyanin after veraison.<sup>14</sup> In contrast to most known berries, *R. chingii* had the continuous decrease in anthocyanins, which contributed to the continuous decrease in flavonoids and phenolics. Antioxidant capacity of fruit was estimated by the free radical-scavenging activity (ABTS) and the reducing antioxidant power (FRAP) assay. ABTS averaged 23.3 mmol TEAC/100 g and it dropped by 33.7%, 45.8% and 32.1% from MG to GY, YO, and RE (Fig. 1d). Similarly, FRAP averaged 19.6 mmol TEAC/100 g and it dropped by 36.2%, 42.2%, and 41.0% (Fig. 1d). Anthocyanins, flavonoids, and phenolics were tightly related with antioxidant capacity (Pearson correlations ranging from 0.968 to 0.999). However, carotenoid was negatively correlated with these phenolics and antioxidant capacity (Pearson correlations ranging from -0.867 and -0.738) (Fig. 1e). The result indicates that the antioxidant capability is highly dependent on phenolic content rather than

carotenoids. It may be due to that carotenoids principally scavenge singlet molecular oxygen and peroxy radicals, and their antioxidant capability can be detected using other assays, like ORAC-L.<sup>40,41</sup>

Total content of phenolics peaked at 4026.3 (mg GAE/100 g FW) at MG (Fig. 1c). It is 10-fold higher than mature fruit in red raspberry (357.8 mg GAE/100 g FW), blackberry (850.5), strawberry (621.9), blueberry (305.4) and cherry (314.5).<sup>42,43</sup> Also, the total content of flavonoids peaked at 646.2 in MG (mg RE/100 g FW) (Fig. 1c), which was higher than that in raspberry.<sup>43</sup> ABTS peaked at 41.2 (mmol TEAC/100 g FW) or 411.8 ( $\mu\text{mol TEAC/g FW}$ ) in MG fruit and was over 20 folds higher than that in mature fruit of red raspberry (6.3  $\mu\text{mol TEAC/g FW}$ ), blackberry (13.2), strawberry (7.9), blueberry (5.9) and cherry (8.8).<sup>42</sup> The extremely high antioxidant capacity of unripe fruit may be one of the reasons for its utilization in a traditional Chinese medicine.

### 3.3. Composition analysis of flavonoids and ellagitannins

Ellagitannins and other flavonoids were identified by LC-MS/MS. Two characteristic wavelengths were used to detect the compounds based on their structural properties, i.e. 280 nm for






Fig. 3 Analysis of ellagitannins, hydroxybenzoic derivatives, and flavonols in *R. chingii*. (a) UPLC spectra of two ellagitannins, (b) UPLC spectra of ten flavonoid compounds. The upper part: electrospray ion mass spectra; the lower part: their fragment ions mass spectra; the beside part: putative molecular structure and cleavage pattern. The IDA-based auto-MS2 was used to distinguish between conjugates of quercetin and ellagic acid since they produce identical deprotonated ions ( $m/z$  301) in the MS/MS spectra.


ellagitannin, and 360 nm for hydroxybenzoic acid and flavonoids. The UPLC profiles at 280 and 360 nm are shown in Fig. 3a and 4b, respectively. The major two compounds detected at 280 nm belonged to the ellagitannins family (Fig. 3a). Ellagitannins are hydrolyzable tannins, esterified with hexahydroxydiphenic acid (HHDP) and a polyol (*i.e.* glucose). Of ellagitannins, casuarictin was predominant in content followed by methyl (*S*)-flavogallate. Both decreased in content during fruit maturation (Table 1). Of hydroxybenzoic acid and flavonoids, kaempferol-3-*o*-rutinoside was predominant in content, followed by ellagic acid and kaempferol-3-glucoside, while rourinoside was the least in content, followed by rutin and isoquercitrin. All of them also showed a trend of decrease in content.

Plants ellagic acid is present as a free compound, in glycosylated and/or acylated form, or as ellagitannin derivatives usually esterified with glucose. In raspberries, free ellagic acid constitutes only a minor part of the total ellagic acids.<sup>44</sup> In red raspberries, the most abundant ellagitannins are sanguin H-6, sanguin H-10 isomer, and lambertianin C,<sup>45,46</sup> while less abundant ellagitannins are sanguin H-2 and [galloyl-bis-HHDP-glucose]2-gallate.<sup>45</sup> Both sanguin H-2 and [galloyl-bis-HHDP-glucose]2-gallate are either present naturally or derived from degradation of lambertianin C during hot-water extraction processes.<sup>46</sup> In black raspberries, sanguin H-6 and its derivatives,

lambertianin C/D, ellagic acid and its derivatives are also found.<sup>47,48</sup> These ellagitannins can be hydrolyzed with acids or bases to release hexahydroxydiphenol units which spontaneously cyclizes into ellagic acid.<sup>48</sup> The ellagic acid released after acid hydrolysis are one of important phenolic compounds in *Rubus* fruit, accounting for approximately 80% of the total phenolics.<sup>49</sup> In *R. chingii*, the main ellagitannins (lambertianin A, sanguin H-6 and casuarictin) and ellagic acid are found in unripe fruits.<sup>50</sup> In this study, methyl (*S*)-flavogallate and casuarictin were the main components of ellagitannins, and the former was first identified in *R. chingii* fruit. Ellagitannins were much higher contents than other phenolics, which contributed to a large part of antioxidant capability. These ellagitannins all decreased as the fruit matured, which was consistent to what was observed in other *Rubus* species.<sup>17,47</sup> The high antioxidant capacity of ellagitannins are believed to have multiple health benefits, *e.g.* antiglycation activity,<sup>50</sup> lung, oesophagus function<sup>51</sup> and as a remedy for combating prostate cancer.<sup>52</sup> Additionally, the ellagitannins is always correlated with oral astringency.<sup>53</sup> The high content of ellagitannins in the unripe fruit of *R. chingii* also explains its use in traditional Chinese medicine, while the low content in ripe fruit makes it have a much less astringent taste than unripe fruit.

*R. chingii* had a varied flavonoid profile due to the occurrence of quercetin and kaempferol derivatives (Table 1; Fig. 3b). These





**Fig. 4** *In situ* flavonoid staining of fresh-fruit sections. (a) Fruit radial sections by paraffin method. (b) Fruit radial sections by frozen method. Fluorescence was collected at (c) 475–504 nm for kaempferol and (d) 577 to 619 nm for quercetin after fresh-fruit section were stained with diphenylboric acid 2-aminoethyl ester (DPBA) (e) Flavonoid localization in inflorescences combing (c) and (d), (f) original figure without fluorescence. The results showed that flavonoid mainly accumulated in epidermal hair and episperm.

components decreased during fruit ripening, which was consistent with what has been observed in red raspberry (S. Y. Wang *et al.*, 2009). Of these, nikotoflorin is predominant, followed by astragaln, ellagic acid and tiliroside, which is consistent with previous reports in *R. chingii*.<sup>54</sup> Ellagic acid and astragaln are prevalent in fruit, and were also reported in red raspberry<sup>7,17,45</sup> and blackberry<sup>47</sup> while tiliroside only exists in some varieties of Poland red raspberry<sup>8</sup> and in leaves of Bulgarian *Rubus* species.<sup>55</sup> Nikotoflorin is not found in red or black raspberry, but is found in *R. chingii* with high concentrations.<sup>56</sup> Isoquercitrin and astragaln ubiquitously exist in red and black raspberries, as well as several ellagic acid pentosides, ellagic acid acetyl pentosides, hyperoside and rutin.<sup>8,47</sup> Rourinoside was first identified in *Rubus*, which was also found in the fractionation of the antimalarial active CHCl<sub>3</sub> extract of the dried stems of *Rourea minor* (Gaertn.).<sup>57</sup> Isoquercitrin, nikotoflorin and tiliroside exhibit significant bioactivity, *e.g.* Isoquercitrin has shown bioactivity against cancer, cardiovascular disorders, diabetes and allergic reactions,<sup>58</sup> nikotoflorin protects the liver from CCl<sub>4</sub>-induced oxidative damage,<sup>59</sup> while

tiliroside possesses anti-inflammatory, antioxidant, anticarcinogenic and hepatoprotective activities.<sup>60</sup> The kaempferol- and quercetin-based flavonoids mainly accumulated in the fruit epidermal hair, and in the placentae and seed coats, but rarely in fruit pericarp (the exocarp, hypodermis and mesocarp) (Fig. 4). Thus, nikotoflorin and rourinoside are species-specific products, which could be applied to the taxonomic classification of *Rubus* species.

## 4. Conclusions

In *R. chingii*, the phenolics drastically decreased throughout fruit ripening while apocarotenoids dramatically increased, which led to discrepancy in color, flavor, and nutritional components between unripe and ripe fruits. The discrepancy determines their different uses. Unripened is extremely rich in healthy phenolic compounds, which could be useful in the development of health care products, while ripe fruit is much less astringent tasting and can be marketed as produce. Notably, *R. chingii* has very special profiles of phenolics and





carotenoids, which is totally different from other raspberries. For example, the  $\beta$ -citraurin esters rather than anthocyanin components are responsible for the fruit reddish coloration.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This research was funded by Natural Science Foundation of Zhejiang Province (LY19C160008), the Science-Technology Programs of Linqi Government, and Basic Public Welfare Projects of Zhejiang Province (2017C32082).

## References

- H. Y. Ding, *Int. J. Mol. Sci.*, 2011, **12**, 3941–3949.
- T. T. Zhang, C. L. Lu, J. G. Jiang, M. Wang, D. M. Wang and W. Zhu, *Carbohydr. Polym.*, 2015, **130**, 307–315.
- N. Sun, Y. Wang, Y. Liu, M.-L. Guo and J. Yin, *Chem. Nat. Compd.*, 2013, **49**, 49–53.
- S. P. Mazur, A. Nes, A. B. Wold, S. F. Remberg and K. Aaby, *Food Chem.*, 2014, **160**, 233–240.
- S. H. Nile and S. W. Park, *Nutrition*, 2014, **30**, 134–144.
- I. A. Ludwig, P. Mena, L. Calani, G. Borges, G. Pereira-Caro, L. Bresciani, R. D. Del, M. E. Lean and A. Crozier, *Free Radical Biol. Med.*, 2015, **89**, 758–769.
- A. V. Rao and D. M. Snyder, *J. Agric. Food Chem.*, 2010, **58**, 3871–3883.
- M. Kula, M. Majdan, D. Glód and M. Krauze-Baranowska, *J. Food Compos. Anal.*, 2016, **52**, 74–82.
- L. S. Kim, S. H. Youn and J. Y. Kim, *J. Korean Soc. Food Sci. Nutr.*, 2014, **43**, 1357–1362.
- M. Krauze-Baranowska, M. Majdan, R. Hałas, D. Glód, M. Kula, I. Fecka and A. Orzeł, *Food Funct.*, 2014, **5**, 2536–2541.
- M. Park, H. Cho, H. Jung, H. Lee and K. T. Hwang, *J. Food Biochem.*, 2014, **38**, 259–270.
- Q. L. Guo, J. Y. Gao and J. S. Yang, *Chromatographia*, 2005, **62**, 145–150.
- J. Beekwilder, I. M. van der Meer, A. Simic, J. Uitdewilligen, J. van Arkel, R. C. H. de Vos, H. Jonker, F. W. A. Verstappen, H. J. Bouwmeester, O. Sibbesen, I. Qvist, J. D. Mikkelsen and R. D. Hall, *Biofactors*, 2008, **34**, 57–66.
- X. Li, L. Jin, X. Pan, L. Yang and W. Guo, *Food Chem.*, 2019, **290**, 216–228.
- L. Jin, X.-B. Li, D.-Q. Tian, X.-P. Fang, Y.-M. Yu, H.-Q. Zhu, Y.-Y. Ge, G.-Y. Ma, W.-Y. Wang, W.-F. Xiao and M. Li, *Ind. Crops Prod.*, 2016, **87**, 198–209.
- D. R. Lewis, M. V. Ramirez, N. D. Miller, P. Vallabhaneni, W. K. Ray, R. F. Helm, B. S. J. Winkel and G. K. Muday, *Plant Physiol.*, 2011, **156**, 144–164.
- S. Y. Wang, C.-T. Chen and C. Y. Wang, *Food Chem.*, 2009, **112**, 676–684.
- J. A. Stavang, S. Freitag, A. Foito, S. Verrall, O. M. Heide, D. Stewart and A. Sønsteby, *Sci. Hortic.*, 2015, **195**, 216–225.
- P. D. Fraser, M. R. Truesdale, C. R. Bird, W. Schuch and P. M. Bramley, *Plant Physiol.*, 1994, **105**, 405–413.
- L. Zhang, Z. Zhang, T. Zheng, W. Wei, Y. Zhu, Y. Gao, X. Yang and S. Lin, *Horticultural Plant Journal*, 2016, **2**, 9–15.
- M.-J. Rodrigo, J. F. Marcos and L. Zacarías, *J. Agric. Food Chem.*, 2004, **52**, 6724–6731.
- K. Karppinen, L. Zoratti, M. Sarala, E. Carvalho, J. Hirsimäki, H. Mentula, S. Martens, H. Haggman and L. Jaakola, *BMC Plant Biol.*, 2016, **16**, 95.
- S. Remy-Tanneau, C. Le Guernevé, E. Meudec and V. Cheyner, *J. Agric. Food Chem.*, 2003, **51**, 3592–3597.
- T. Fossen and O. M. Andersen, *Phytochemistry*, 2003, **62**, 1217–1220.
- T. Fossen, S. Rayyan and Ø. M. Andersen, *Phytochemistry*, 2004, **65**, 1421–1428.
- G. J. McDougall, S. Gordon, R. Brennan and D. Stewart, *J. Agric. Food Chem.*, 2005, **53**, 7878–7885.
- Y. Fukui, T. Kusumi, K. Masuda, T. Iwashita and K. Nomoto, *Tetrahedron Lett.*, 2002, **43**, 2637–2639.
- A. M. González-Paramás, F. Lopes da Silva, P. Martín-López, G. Macz-Pop, S. González-Manzano, C. Alcalde-Eon, J. J. Pérez-Alonso, M. T. Escribano-Bailón, J. C. Rivas-Gonzalo and C. Santos-Buelga, *Food Chem.*, 2006, **94**, 428–436.
- T. K. Hyun, S. Lee, Y. Rim, R. Kumar, X. Han, S. Y. Lee, C. H. Lee and J. Y. Kim, *PLoS One*, 2014, **9**, e88292.
- M. Dossett, J. Lee and C. E. Finn, *J. Funct. Foods*, 2010, **2**, 292–297.
- A. L. Cual, *J. Food Sci.*, 1965, **30**, 13–18.
- J. N. Mi and S. Al-Babili, *Mol. Plant*, 2019, **12**, 1173–1175.
- G. Ma, L. Zhang, A. Matsuta, K. Matsutani, K. Yamawaki, M. Yahata, A. Wahyudi, R. Motohashi and M. Kato, *Plant Physiol.*, 2013, **163**, 682–695.
- D. Marinova and F. Ribarova, *J. Food Compos. Anal.*, 2007, **20**, 370–374.
- E. Carvalho, P. D. Fraser and S. Martens, *Food Chem.*, 2013, **139**, 744–752.
- K. Mizuno, T. Tokiwano and Y. Yoshizawa, *Biochem. Biophys. Res. Commun.*, 2017, **484**, 845–849.
- I. O. Vvedenskaya and N. Vorsa, *Plant Sci.*, 2004, **167**, 1043–1054.
- J. Song, L. Du, L. Li, W. Kalt, L. C. Palmer, S. Fillmore, Y. Zhang, Z. Zhang and X. Li, *J. Proteomics*, 2015, **122**, 1–10.
- M. Giribaldi, I. Perugini, F. X. Sauvage and A. Schubert, *Proteomics*, 2007, **7**, 3154–3170.
- H. Liu, J. Mao, S. Yan, Y. Yu, L. Xie, J. G. Hu, T. Li, A. M. Abbasi, X. Guo and R. H. Liu, *Int. J. Food Sci. Technol.*, 2018, **53**, 381–388.
- W. Stahl and H. Sies, *Mol. Aspects Med.*, 2003, **24**, 345–351.
- V. R. de Souza, P. A. Pereira, T. L. da Silva, L. C. de Oliveira Lima, R. Pio and F. Queiroz, *Food Chem.*, 2014, **156**, 362–368.
- L. Chen, X. Xin, H. Zhang and Q. Yuan, *J. Funct. Foods*, 2013, **5**, 508–515.
- K. R. Määttä-Riihinen, A. Kamal-Eldin and A. R. Törrönen, *J. Agric. Food Chem.*, 2004, **52**, 6178–6187.
- R. Bobinaite, P. Viskelis and P. R. Venskutonis, *Food Chem.*, 2012, **132**, 1495–1501.



- 46 M. M. Kool, D. J. Comeskey, J. M. Cooney and T. K. McGhie, *Food Chem.*, 2010, **119**, 1535–1543.
- 47 L. Kaume, L. R. Howard and L. Devareddy, *J. Agric. Food Chem.*, 2012, **60**, 5716–5727.
- 48 U. Vrhovsek, A. Palchetti, F. Reniero, C. Guillou, D. Masuero and F. Mattivi, *J. Agric. Food Chem.*, 2006, **54**, 4469–4475.
- 49 H. Sh, M. Heinonen, K. So, M. Hm and R. Trnnen, *Food Res. Int.*, 1999, **32**, 345–353.
- 50 C. Yue, X. Leilei, W. Yajie, C. Zhongqin and C. Haixia, *LWT-Food Sci. Technol.*, 2019, **121**, 108967.
- 51 L. A. Kresty, M. A. Morse, C. Morgan, P. S. Carlton, J. Lu, A. Gupta, M. Blackwood and G. D. Stoner, *Cancer Res.*, 2001, **61**, 6112–6119.
- 52 N. P. Seeram, W. J. Aronson, Y. Zhang, S. M. Henning, A. Moro, R. P. Lee, M. Sartippour, D. M. Harris, M. Rettig, M. A. Suchard, A. J. Pantuck, A. Belldegrun and D. Heber, *J. Agric. Food Chem.*, 2007, **55**, 7732–7737.
- 53 T. Hofmann, A. Glabasnia, B. Schwarz, K. N. Wisman, K. A. Gangwer and A. E. Hagerman, *J. Agric. Food Chem.*, 2006, **54**, 9503–9509.
- 54 J. M. He, N. Sun, W. D. Wu, L. J. Fan and M. L. Guo, *Zhongguo Zhongyao Zazhi*, 2013, **38**, 4351–4356.
- 55 R. Gevrenova, I. Badjakov, M. Nikolova and I. Doichinova, *Biochem. Syst. Ecol.*, 2013, **50**, 419–427.
- 56 G. Yu, Z. Luo, W. Wang, Y. Li, Y. Zhou and Y. Shi, *Front. Pharmacol.*, 2019, **10**, 1323.
- 57 Z.-D. He, C.-Y. Ma, G. T. Tan, K. Sydara, P. Tamez, B. Southavong, S. Bouamanivong, D. D. Soejarto, J. M. Pezzuto, H. H. S. Fong and H.-J. Zhang, *Phytochemistry*, 2006, **67**, 1378–1384.
- 58 K. Valentová, J. Vrba, M. Bancířová, J. Ulrichová and V. Křen, *Food Chem. Toxicol.*, 2014, **68**, 267–282.
- 59 Y. Wang, C. Tang and H. Zhang, *J. Food Drug Anal.*, 2015, **23**, 310–317.
- 60 T. Goto, A. Teraminami, J.-Y. Lee, K. Ohyama, K. Funakoshi, Y.-I. Kim, S. Hirai, T. Uemura, R. Yu, N. Takahashi and T. Kawada, *J. Nutr. Biochem.*, 2012, **23**, 768–776.

