


 Cite this: *RSC Adv.*, 2021, **11**, 30937

 Received 3rd July 2021
 Accepted 24th August 2021

DOI: 10.1039/d1ra05134b

rsc.li/rsc-advances

Vitamin B1-catalyzed aerobic oxidative esterification of aromatic aldehydes with alcohols†

 Xin-Long Luo,^{‡a} Danhua Ge,^{‡a} Zi-Lun Yu,^a Xue-Qiang Chu^{‡*a} and Pei Xu^{*b}

A straightforward aerobic oxidative esterification of aryl aldehydes with alcohols has been developed for the synthesis of substituted esters by employing vitamin B1 as a cost-effective, metal-free, and eco-friendly NHC catalyst. Air is used as a green terminal oxidant. The reaction is a useful addition to the existing NHC-catalytic oxidative esterification.

Introduction

Vitamin B1 (VB1), also known as thiamine, is one of the eight water-soluble B vitamins. In the human body, VB1 plays an essential role in the synthesis of nucleic acids (*e.g.*, DNA), the conduction of nerve impulses, and the production of energy from food.¹ The structure of VB1 contains a thiazole ring and a pyrimidine ring linked by a methylene bridge (Fig. 1). As an important subarea of biocatalysis, VB1 and its analogs have been widely used as non-toxic, low cost, and stable catalysts in diverse organic transformations² including oxidation,³ cyclization,⁴ Michael addition, C–X bond coupling,⁵ benzoin condensation,⁶ Knoevenagel condensation,⁷ *etc.*⁸ Despite this impressive progress which has been achieved, reports related to the C–O bond formation by employing this renewably biobased catalyst are still highly desirable, offering benefits from an environment point of view.

The esters have been found a myriad of applications in fields ranging from agrochemistry, pharmacology, and materials science to organic synthesis, mainly owing to their unique biological and chemical properties.⁹ The development of efficient strategies towards ester functionality has inspired chemists for more than 100 years, therefore, a variety of well-established methods are developed. Classical condensation of carboxylic acids with alcohols is carried out under strongly acidic conditions in the presence of a large excess of either substrate.¹⁰ In addition, esterification can also be achieved through the stoichiometric activation of the corresponding acids (*e.g.*, acid halides, anhydrides, or activated esters) following nucleophilic substitution with appropriate alcohols.

On the other hand, transition metal-catalyzed direct oxidative functionalization of aldehydes provides a powerful platform for ester synthesis over the past decades (Scheme 1A). Despite advances, these transformations commonly suffered from harsh reaction conditions, the use of costly metal catalysts, stoichiometric oxidants, and extra additives, which constitute drawbacks for large-scale applications and late-stage modifications.¹¹ Alternatively, N-heterocyclic carbenes (NHCs) derived from thiazolium ions have the capability to accomplish the oxidative esterification with the assistance of various oxidants, such as MnO₂,¹² nitroarene,¹³ TEMPO,¹⁴ NFSI,¹⁵ diphenone,¹⁶ azobenzene,¹⁷ phenazine,¹⁸ CCl₃CN,¹⁹ and O₂/air (Scheme 1B).²⁰ From a viewpoint of green chemistry, a metal-free NHC-catalyzed one-pot esterification of aldehydes under air has become more sought after. In this regard, most of the existing methods relied on the preformed NHC catalysts which required multi-step synthesis. Continuing with our interests in green chemistry,²¹ herein we introduce the commercially available VB1 as an efficient catalyst for oxidative esterification (Scheme 1C). As a result, a cost-effective, metal-free, eco-friendly, and catalytic route that minimizes hazardous waste is developed for the synthesis of esters by employing a variety of aryl aldehydes and alcohols as coupling partners or reaction solvents. This metal-free approach offers a valuable alternative when compared to the above-mentioned methods: (i) biocompatible VB1 is used as an NHC catalyst; (ii) transition metal and expensive oxidant are not necessary; (iii) DABCO serves as




Fig. 1 Thiamine (VB1) as a versatile catalyst in organic transformations.


^aSchool of Chemistry and Molecular Engineering, Nanjing Tech University, Nanjing 211816, China. E-mail: xueqiangchu@njtech.edu.cn

^bJiangsu Key Laboratory of New Drug Research and Clinical Pharmacy, School of Pharmacy, Xuzhou Medical University, Xuzhou 221004, China. E-mail: xupeix@xzhmu.edu.cn

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d1ra05134b

‡ X.-L. Luo and D. Ge contributed equally to this work.





Scheme 1 General routes for the oxidative esterification.

a basic promoter. In particular, aerobic oxidation using dioxygen gas as a green oxidant is attractive.

Results and discussion

We commenced the study by investigating the model reaction of 4-nitrobenzaldehyde (**1a**) with ethanol (**2a**) (Table 1). Gratifyingly, the reaction proceeded smoothly in the presence of VB1 (20 mol%) as a catalyst and triethylenediamine (DABCO, 2 equiv.) as a base at 40 °C for 6 h under air, leading to the desired coupled product **3aa** in 58% yield (entry 1). In contrast, eroded efficiency was observed when the amount of VB1 was increased from 20 mol% to 30 mol% (entry 2), or when the amount of DABCO was increased from 2 equiv. to 3 equiv. (entry 3). Based on the control experiments, the necessity of adding catalyst and base were adequately demonstrated for the present reaction (entries 4 and 5). Subsequently, variations of various reaction parameters, including base, reaction temperature, reaction time, and solvent, were also carried out. It was found that DABCO was the base of choice, as the same reactions performed with other bases (including Na₂CO₃, Et₃N, and NaOH) produced the corresponding product **3aa** in low yields (entries 6–8). Furthermore, evaluation of reaction temperature and reaction time showed that the isolated yield of product **3aa** could be increased to 65% when the reaction was performed at 80 °C for 12 h (entry 9). Interestingly, a low equivalent of ethanol (1 mmol) in CHCl₃ did not considerably reduce the yield efficiency (49% yield, entry 10). Further screening of other solvents (*e.g.*, DMSO, toluene, THF, and acetone) revealed that the reaction performance could be improved in acetone as the solvent (57% yield, entry 14). Finally, it was found that 63% yield could be obtained by carrying out the reaction at 60 °C in acetone for 12 h

Table 1 Optimization of reaction conditions^a


Entry	VB1 (x mol%)	Base (y equiv.)	Solvent	T (°C)	Yield ^b (%)
1	20	DABCO (2)	EtOH	40	58
2	30	DABCO (2)	EtOH	40	49
3	20	DABCO (3)	EtOH	40	47
4	—	DABCO (2)	EtOH	40	0
5	20	—	EtOH	40	0
6	20	Na ₂ CO ₃ (2)	EtOH	40	<10
7	20	Et ₃ N (2)	EtOH	40	10
8	20	NaOH (2)	EtOH	40	<10
9	20	DABCO (2)	EtOH	80	65 ^c
10	20	DABCO (2)	CHCl ₃	40	49 ^d
11	20	DABCO (2)	DMSO	40	22 ^d
12	20	DABCO (2)	Toluene	40	Trace ^d
13	20	DABCO (2)	THF	40	Trace ^d
14	20	DABCO (2)	Acetone	40	57 ^d
15	20	DABCO (2)	Acetone	60	63 ^d
16	20	DABCO (2)	Acetone	60	61 ^{d,e}

^a Reaction conditions: 4-nitrobenzaldehyde (**1a**, 1 mmol), VB1 (0–0.6 mmol), and base (0–3 mmol) in EtOH (**2a**, 2 mL) at 40–80 °C (oil bath) under air for 6 h. ^b Isolated yield. ^c 12 h. ^d Reaction conditions: 4-nitrobenzaldehyde (**1a**, 3 mmol), EtOH (**2a**, 1 mmol), VB1 (0.2 mmol), and DABCO (2 mmol) in solvent (2 mL) at 40–60 °C (oil bath) under air for 12 h. ^e O₂ is used as an oxidant.

(entry 15). However, further improvement of the yield of product **3aa** and diminishment of the undesired process were failed by replacing air with pure O₂ (entry 16).

With this optimized result in hand, we continued to study the substrate scope of the present esterification by employing a wide variety of structurally varied alcohols and aryl aldehydes as starting materials, and the results were summarized in Table 2. The aerobic oxidative esterification of 4-nitrobenzaldehyde (**1a**) worked equally well when the reactions were performed in different alcohols **2** as the solvents at 80 °C for 12–24 h, furnishing the corresponding esters **3aa–3aj** in 18–65% yields. Interestingly, the chain length of the alkyl alcohols only had a slight impact on the reaction outcome (**3aa–3ae**, 52–65% yields). Notably, a sterically hindered secondary alcohol could be employed as substrate with equal ease (**3ad**, 53% yield). However, 2,2,2-trifluoroethan-1-ol with a strong electron-withdrawing CF₃ group obviously reduced the reactivity presumably because of its low nucleophilicity (**3af**, 18% yield). To our delight, alcohols containing functional groups such as Cl atom (**2g**), alkene (**2h**), and alkyne (**2i**) were compatible with the mild reaction conditions, providing the anticipated products **3ag–3ai** in 44–59% yields. Moreover, aryl aldehydes **1** bearing various substituents successfully underwent oxidative coupling with ethanol. In most cases, the cross-coupled products were produced in moderate yields (**3ba–3fa**). In addition to 4-nitrobenzaldehyde, aryl aldehydes possessing F, CN, I, and Br in the




Table 2 Substrate scope of various alcohols and aryl aldehydes^a

^a Reaction conditions: aryl aldehyde **1** (1.0 mmol), VB1 (0.2 mmol), and DABCO (2.0 mmol) in alcohol **2** (2.0 mL) at 80 °C (oil bath) for 12–24 h under air; isolated yields.

benzene ring were also proven to be appropriate substrates for the reaction (**3ca–3fa**). Unfortunately, further examination revealed that 4-methoxybenzaldehyde (**1g**) and heterocyclic aldehydes **1h–1j** were reluctant to participate in the titled transformation. The failures could be attributed to the electronic effect. However, the reaction of 2-nitrobenzaldehyde (**1k**) with alcohol also failed to give the desired product, mainly owing to the negative effect of steric hindrance.


Encouraged by the above results, the generality of the substrate scope was further evaluated by applying different alcohols in acetone. As summarized in Table 3, benzyl alcohols **2** possessing either electron-donating group (e.g., Me, Me) or electron-withdrawing group (e.g., F, Cl, Br, and I) smoothly underwent the present organic transformations to give the desired esters **3ak–3ar** in moderate to good yields. Notably, halogen functionality was satisfactorily compatible with the mild reaction conditions, which could be allowed for further

Table 3 Substrate scope of various alcohols 2^a

^a Reaction conditions: 4-nitrobenzaldehyde (**1a**, 3.0 mmol), alcohol **2** (1.0 mmol), VB1 (0.2 mmol), and DABCO (2.0 mmol) in acetone (2.0 mL) at 60 °C (oil bath) for 24 h under air; isolated yields.

divergent late-stage modifications (**3ao–3ar**). Similarly, thiophen-2-ylmethanol (**2s**), furan-2-ylmethanol (**2t**), and pyridin-2-ylmethanol (**2v**) were proven to be good candidates, affording the anticipated products **3as**, **3at**, and **3av** in 63–69% yields. However, other heterocyclic substrate such as (1*H*-indol-3-yl)methanol (**2u**) could not participate into the reactions with 4-nitrobenzaldehyde (**1a**). In addition, we were able to accomplish the conversion of alkyl alcohols such as 2-(4-methoxyphenyl)ethan-1-ol (**3aw**), 2-(naphthalen-1-yl)ethan-1-ol (**3ax**), and cyclohexylmethanol (**3ay**) to the coupled products **3aw–3ay** in 46–66% yields. Remarkably, (*E*)-3-phenylprop-2-en-






Scheme 2 Scale-up synthesis of product 3aa and control experiment.

1-ol can be easily converted into 3az in 62% yield. During the course of our studies, other types of nucleophiles including phenol 2a' and amine 2b' failed to react under the optimized reaction conditions.

Subsequently, a scale-up reaction was also applied to the model reaction (Scheme 2A). It was found that 10 mmol scale reaction worked with equally high efficiency under the well-established reaction conditions to produce the desired product 3aa in 61% yield. Additionally, when the reaction of 4-nitrobenzaldehyde and ethanol was performed under N₂ atmosphere, the formation of product 3aa was almost completely inhibited (Scheme 2B). This result suggests that the presence of O₂ is important for the intermolecular oxidative coupling reaction.²² Furthermore, a radical trapping experiment by using 2,2,6,6-tetramethylpiperidinoxy (TEMPO, 1.0 equiv.) as a radical inhibitor provided clear evidence that the reaction might not proceed in a radical manner (Scheme 2C).

Based on the control experiment and a literature survey,^{12–20} a proposed mechanism is depicted in Scheme 3. Initially, deprotonation of VB1 under the basic condition gives rise to carbene catalyst A. Subsequently, the resulting species reacts with aryl aldehyde to produce the Breslow intermediate B,



Scheme 3 Proposed mechanism.

which in the presence of air can be further oxidized to acyl azolium intermediate C. Next, an intermolecular nucleophilic attack from the alcoholic OH occurred to give the desired product and regenerate the NHC catalyst.

Conclusions

In summary, we have developed a vitamin B1-catalyzed aerobic oxidative esterification of aryl aldehydes with alcohols, providing easy access to synthetically important ester derivatives in moderate yields. The reaction proceeds in the air without an external oxidant. Moreover, commercially available VB1 was used as a cost-effective, metal-free, eco-friendly, and green catalyst, complementing the previously reported NHC-catalytic methods for the esterification.

Experimental

General procedure of vitamin B1-catalyzed aerobic oxidative esterification for the synthesis of products 3

Method 1. A solution of aldehyde (1 mmol), vitamin B1 (60.2 mg, 0.2 mmol), and triethylenediamine (DABCO, 224.3 mg, 2 mmol) in alcohol (2 mL) was stirred at 80 °C under air for 12–24 h. The reaction was then quenched by saturated NH₄Cl solution (20 mL) and extracted with EtOAc (20 mL × 3). The organic layer was washed with saturated brine twice, dried over MgSO₄, filtered, and concentrated under reduced pressure. The crude product was purified by flash column chromatography (300–400 mesh) using petroleum ether/ethyl acetate (100/1–2/1) as eluent to afford the pure products 3.

Method 2. A solution of 4-nitrobenzaldehyde (453.4 mg, 3 mmol), alcohol (1 mmol), vitamin B1 (60.2 mg, 0.2 mmol), and triethylenediamine (DABCO, 224.3 mg, 2 mmol) in acetone (2 mL) was stirred at 60 °C under air for 24 h. The reaction was then quenched by saturated NH₄Cl solution (20 mL) and extracted with EtOAc (20 mL × 3). The organic layer was washed with saturated brine twice, dried over MgSO₄, filtered, and concentrated under reduced pressure. The crude product was purified by flash column chromatography (300–400 mesh) using petroleum ether/ethyl acetate (100/1–2/1) as eluent to afford the pure products 3.

General procedure for the scale-up synthesis of pyridine 3aa

A solution of 4-nitrobenzaldehyde (1.51 g, 10 mmol), vitamin B1 (601.6 mg, 2 mmol), and triethylenediamine (DABCO, 2.24 g, 20 mmol) in EtOH (20 mL) was stirred at 80 °C under air for 24 h. The reaction solvent was removed under reduced pressure, then the reaction mixture was quenched by saturated NH₄Cl solution (20 mL) and extracted with EtOAc (20 mL × 3). The organic layer was washed with saturated brine twice, dried over MgSO₄, filtered, and concentrated under reduced pressure. The crude product was purified by flash column chromatography (300–400 mesh) using petroleum ether/ethyl acetate (100/1–2/1) as eluent to afford the pure products 3aa (1.19 g, 61% yield).



Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We gratefully acknowledge the financial support from Xuzhou Medical University (Start-up Grant no. RC20552038), Nanjing Tech University (start-up grant no. 39837146), National Students' platform for innovation and entrepreneurship training program (202110291019Z), Natural Science Foundation of Jiangsu Province (BK20180690), and National Natural Science Foundation of China (22001121).

Notes and references

- (a) C. J. Abularrage, A. N. Sidawy, P. W. White, G. Aidinian, K. J. DeZee, J. M. Weiswasser and S. Arora, Effect of Folic Acid and Vitamins B₆ and B₁₂ on Microcirculatory Vasoactivity in Patients with Hyperhomocysteinemia, *Vascular and Endovascular Surgery*, 2007, **41**, 339–345; (b) W. G. Christen, R. J. Glynn, E. Y. Chew, C. M. Albert and J. E. Manson, Folic Acid, Pyridoxine, and Cyanocobalamin Combination Treatment and Age-Related Macular Degeneration in Women The Women's Antioxidant and Folic Acid Cardiovascular Study, *Arch. Intern. Med.*, 2009, **169**, 335–341.
- R. Singh, S. A. Ganaie and A. Singh, Vitamin B1: A versatile organocatalyst for organic synthesis, *Curr. Organocatal.*, 2017, **4**, 84–103.
- Y. Liu, C. Wang, D. Xue, M. Xiao, J. Liu, C. Li and J. Xiao, Reactions Catalysed by a Binuclear Copper Complex: Relay Aerobic Oxidation of *N*-Aryl Tetrahydroisoquinolines to Dihydroisoquinolones with a Vitamin B1 Analogue, *Chem.–Eur. J.*, 2017, **23**, 3062–3066.
- M. Lei, L. Ma and L. Hu, One-Pot Synthesis of 1*H*-Benzimidazole Derivatives Using Thiamine Hydrochloride as a Reusable Organocatalyst, *Synth. Commun.*, 2012, **42**, 2981–2993.
- (a) V. B. Purohit, S. C. Karad, K. H. Patel and D. K. Raval, Palladium *N*-heterocyclic carbene catalyzed regioselective thiolation of 1-aryl-3-methyl-1*H*-pyrazol-5(4*H*)-ones using aryl thiols, *Tetrahedron*, 2016, **72**, 1114–1119; (b) V. B. Purohit, S. C. Karad, K. H. Patel and D. K. Raval, Palladium *N*-heterocyclic carbene catalyzed expected and unexpected C–C and C–N functionalization reactions of 1-aryl-3-methyl-1*H* pyrazol-5(4*H*)-ones, *RSC Adv.*, 2016, **6**, 111139–111143.
- (a) T. Lu, X. Li, L. Gu and Y. Zhang, Vitamin B1-Catalyzed Acetoin Formation from Acetaldehyde: A Key Step for Upgrading Bioethanol to Bulk C₄ Chemicals, *ChemSusChem*, 2014, **7**, 2423–2426; (b) Y. Chen, W. Shan, M. Lei and L. Hu, Thiamine hydrochloride (VB1) as an efficient promoter for the one-pot synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones, *Tetrahedron Lett.*, 2012, **53**, 5923–5925.
- (a) I. R. Siddiqui, P. Rai, Rahila, A. Srivastava, A. Srivastava and A. Srivastava, Synthesis of fused pyridines in the presence of thiamine hydrochloride as an efficient and reusable catalyst in aqueous conditions, *New J. Chem.*, 2013, **37**, 3798–3804; (b) K. Pradhan, P. Bhattacharyya, S. Paul and A. R. Das, Synthesis of 3,4-dihydropyridin-2-one derivatives in convergent mode applying bio catalyst vitamin B1 and polymer supported catalyst PEG–SO₃H from two different sets of building blocks, *Tetrahedron Lett.*, 2012, **53**, 5840–5844; (c) M. Lei, L. Ma and L. Hu, Highly Chemoselective Condensation of β-Naphthol, Aldehyde, and Urea Catalyzed by Thiamine Hydrochloride, *Synth. Commun.*, 2011, **41**, 3424–3432; (d) M. Lei, L. Ma and L. Hu, VB1–Al₂O₃-catalyzed one-pot condensation of aromatic ketone, aromatic aldehyde, and amide, *Tetrahedron Lett.*, 2010, **51**, 4746–4749; (e) J. Liu, M. Lei and L. Hu, Thiamine hydrochloride (VB1): an efficient promoter for the one-pot synthesis of benzo[4,5]imidazo[1,2-*a*]pyrimidine and [1,2,4]triazolo[1,5-*a*] pyrimidine derivatives in water medium, *Green Chem.*, 2012, **14**, 840–846.
- (a) M. Lei, L. Ma and L. Hu, A convenient one-pot synthesis of formamide derivatives using thiamine hydrochloride as a novel catalyst, *Tetrahedron Lett.*, 2010, **51**, 4186–4188; (b) K. Azizi and A. Heydari, Vitamin B1 supported on silica-encapsulated γ-Fe₂O₃ nanoparticles: design, characterization and application as a greener biocatalyst for highly efficient acylation, *RSC Adv.*, 2014, **4**, 8812–8816.
- J. Otera and J. Nishikido, *Esterification: Methods, Reactions, and Applications*, Wiley-VCH, Weinheim, 2nd edn, 2010.
- For selected examples, see: (a) K. Ishihara, S. Ohara and H. Yamamoto, Direct Condensation of Carboxylic Acids with Alcohols Catalyzed by Hafnium(IV) Salts, *Science*, 2000, **290**, 1140–1142; (b) K. Manabe, X.-M. Sun and S. Kobatashi, Dehydration Reactions in Water. Surfactant-Type Brønsted Acid-Catalyzed Direct Esterification of Carboxylic Acids with Alcohols in an Emulsion System, *J. Am. Chem. Soc.*, 2001, **123**, 10101–10102.
- (a) X.-F. Wu and C. Darcel, Iron-Catalyzed One-Pot Oxidative Esterification of Aldehydes, *Eur. J. Org. Chem.*, 2009, **2009**, 1144–1147; (b) C. Liu, J. Wang, L.-K. Meng, Y. Deng, Y. Li and A.-W. Lei, Palladium-Catalyzed Aerobic Oxidative Direct Esterification of Alcohols, *Angew. Chem., Int. Ed.*, 2011, **50**, 5144–5148; (c) M.-L. Zhang, S.-H. Zhang, G.-Y. Zhang, F. Chen and J. Cheng, Palladium/NHC-catalyzed oxidative esterification of aldehydes with phenols, *Tetrahedron Lett.*, 2011, **52**, 2480–2483; (d) L. Fu, C.-J. Yao, N.-J. Chang, J.-R. Chen, L.-Q. Lu and W.-J. Xiao, Oxidative cross-esterification of dithiolanes with alcohols through a cross-dehydrogenative coupling (CDC)/deprotection sequence, *Org. Biomol. Chem.*, 2012, **10**, 506–508; (e) W.-J. Yoo and C.-J. Li, Copper-catalyzed oxidative esterification of alcohols with aldehydes activated by Lewis acids, *Tetrahedron Lett.*, 2007, **48**, 1033–1035.
- B. Maji, S. Vedachalan, X. Ge, S.-T. Cai and X.-W. Liu, *N*-Heterocyclic Carbene-Mediated Oxidative Esterification of



- Aldehydes: Ester Formation and Mechanistic Studies, *J. Org. Chem.*, 2011, **76**, 3016–3023.
- 13 J. Castells, F. Pujol, H. Llitjós and M. Moreno-Mañas, Oxidative benzoin reactions, *Tetrahedron*, 1982, **38**, 337–346.
- 14 S. D. Sarkar, S. Grimme and A. Studer, NHC Catalyzed Oxidations of Aldehydes to Esters: Chemoselective Acylation of Alcohols in Presence of Amines, *J. Am. Chem. Soc.*, 2010, **132**, 1190–1191.
- 15 X. Dong, W. Yang, W. Hu and J. Sun, N-Heterocyclic Carbene Catalyzed Enantioselective α -Fluorination of Aliphatic Aldehydes and α -Chloro Aldehydes: Synthesis of α -Fluoro Esters, Amides, and Thioesters, *Angew. Chem., Int. Ed.*, 2015, **54**, 660–663.
- 16 S. De Sarkar, S. Grimme and A. Studer, NHC Catalyzed Oxidations of Aldehydes to Esters: Chemoselective Acylation of Alcohols in Presence of Amines, *J. Am. Chem. Soc.*, 2010, **132**, 1190–1191.
- 17 C. Noonan, L. Baragwanath and S. J. Connon, Nucleophilic carbene-catalysed oxidative esterification reaction, *Tetrahedron Lett.*, 2008, **49**, 4003–4006.
- 18 X. Zhao, K. E. Ruhl and T. Rovis, N-Heterocyclic-Carbene-Catalyzed Asymmetric Oxidative Hetero-Diels–Alder Reactions with Simple Aliphatic Aldehydes, *Angew. Chem., Int. Ed.*, 2012, **51**, 12330–12333.
- 19 Z. Wu, D. Jiang and J. Wang, Carbene-catalyzed oxidative acylation promoted by an unprecedented oxidant CCl_3CN , *Org. Chem. Front.*, 2019, **6**, 688–693.
- 20 W. Harnying, P. Sudkaow, A. Biswas and A. Berkessel, N-Heterocyclic Carbene/Carboxylic Acid Co-Catalysis Enables Oxidative Esterification of Demanding Aldehydes/Enals, at Low Catalyst Loading, *Angew. Chem., Int. Ed.*, 2021, **60**, 19631–19636.
- 21 (a) T. Xie, G.-Q. Wang, Y.-W. Wang, W. Rao, H. Xu, S. Li, Z.-L. Shen and X.-Q. Chu, Selective Quadruple $\text{C}(\text{sp}^3)\text{-F}$ Functionalization of Polyfluoroalkyl Ketones, *iScience*, 2020, **23**, 101259; (b) Q.-D. Wang, Y.-W. Wang, T. Xie, Y.-Y. Cui, M. Ma, Z.-L. Shen and X.-Q. Chu, Three-Component Bisannulation for the Synthesis of Trifluoromethylated Tetracyclic Aza-Aromatics through Six $\text{C}(\text{sp}^3)\text{-F}$ Bond Cleavage and Four C-N Bond Formation, *J. Org. Chem.*, 2021, **86**, 8236–8247; (c) D. Ge, X.-L. Luo, X. Tang, C.-B. Pang, X. Wang and X.-Q. Chu, Metal-Free [3 + 2 + 1] Annulation of Allylic Alcohols, Ketones, and Ammonium Acetate: Radical-Involving Synthesis of 2,3-Diarylpyridine Derivatives, *Org. Biomol. Chem.*, 2021, **19**, 2277–2283; (d) D. Ge, X. Wang and X.-Q. Chu, SOMophilic Alkynylation Using Acetylenic Sulfones as Functional Reagents, *Org. Chem. Front.*, 2021, **8**, 5145–5164; (e) S.-Z. Cai, D. Ge, L.-W. Sun, W. Rao, X. Wang, Z.-L. Shen and X.-Q. Chu, Three-Component Heteroannulation for Tetrasubstituted Furan Construction Enabled by Successive Defluorination and Dual Sulfonylation Relay, *Green Chem.*, 2021, **23**, 935–941; (f) X.-Q. Chu, D. Ge, Y.-Y. Cui, Z.-L. Shen and C.-J. Li, Desulfonylation *via* Radical Process: Recent Developments in Synthetic Applications, *Chem. Rev.*, 2021, DOI: 10.1021/acs.chemrev.1c00084.
- 22 For selected examples of oxidation reactions with air as the oxidant, see: (a) K.-J. Liu, Z. Wang, L.-H. Lu, J.-Y. Chen, F. Zeng, Y.-W. Lin, Z. Cao, X. Yu and W.-M. He, Synergistic cooperative effect of $\text{CF}_3\text{SO}_2\text{Na}$ and bis(2-butoxyethyl)ether towards selective oxygenation of sulfides with molecular oxygen under visible-light irradiation, *Green Chem.*, 2021, **23**, 496–500; (b) K.-J. Liu, J.-H. Deng, T.-Y. Zeng, X.-J. Chen, Y. Huang, Z. Cao, Y.-W. Lin and W.-M. He, 1,2-Diethoxyethane catalyzed oxidative cleavage of *gem*-disubstituted aromatic alkenes to ketones under minimal solvent conditions, *Chin. Chem. Lett.*, 2020, **31**, 1868–1872.

