


 Cite this: *RSC Adv.*, 2021, **11**, 28934

 Received 15th July 2021  
 Accepted 13th August 2021

DOI: 10.1039/d1ra05443k

[rsc.li/rsc-advances](https://rsc.li/rsc-advances)

## Synthesis of xanthohumol and xanthohumol-d<sub>3</sub> from naringenin†

 Joanna Andrusiak,<sup>ab</sup> Kinga Mylkie,<sup>cd</sup> Małgorzata Wysocka,<sup>b</sup> Jacek Ścianowski,<sup>id a</sup> Andrzej Wolan<sup>ab</sup> and Marcin Budny<sup>id \*b</sup>

A six-step synthesis of xanthohumol (**1a**) and its d<sub>3</sub>-derivative (**1b**) from easily accessible naringenin is reported. The prenyl side chain was introduced by Mitsunobu reaction followed by the europium-catalyzed Claisen rearrangement and base-mediated opening of chromanone gave access to an α,β-conjugated ketone system. Compound **1b** was used as an internal standard in stable isotope dilution assays of **1a** in two Polish beers.

Xanthohumol (**1a**, Scheme 1A) is a naturally occurring prenylated chalcone produced by lupulin glands in female inflorescences of hop plants.<sup>1</sup> In recent years, **1a** has attracted significant attention due to its vast range of biological activities including cancer-preventive, antioxidative, anti-inflammatory, and antiviral.<sup>2–8</sup> Such properties combined with low toxicity to the human body make **1a** a prospective therapeutic agent, diet supplement, or ingredient of cosmetics.<sup>9</sup> Although **1a** was isolated from natural sources in 1913,<sup>10</sup> the first synthesis of this compound was reported as late as in 2007.<sup>11</sup> Several other syntheses have been reported since then, but only minor improvements have been achieved.<sup>12–14</sup>

Bioactive compounds labeled with stable isotopes (deuterium, carbon-13) are widely applied in metabolomic studies for tracking metabolic pathways and as internal standards in stable isotope dilution assays.<sup>15,16</sup> Deuterated compounds are also considered as attractive drug candidates due to the influence of the kinetic isotope effect on pharmacokinetics.<sup>17–19</sup> Although approaches to <sup>13</sup>C-enriched xanthohumol<sup>20,21</sup> and hydrogen/deuterium exchange in **1a**<sup>22</sup> were reported, no scalable and cost-effective synthesis of the deuterium-labeled derivative of **1a** (*i.e.* **1b**) has been disclosed to date.

Two main challenges have to be faced in the synthesis of **1a**: (i) construction of a pentasubstituted aromatic ring containing a prenyl side chain and (ii) selection of suitable protecting groups for phenols. In the case of (i), phloroglucinol is used as a precursor and an acyl-substituent is introduced by Friedel–

Crafts acylation with a subsequent Claisen–Schmidt condensation. The prenyl side-chain is introduced by Mitsunobu alkylation, followed by Claisen rearrangement. In the case of (ii), acid-sensitive alkoxymethyl protecting groups, removable under conditions in which **1a** does not cyclize to isoxanthohumol (**2a**), are used most often (Scheme 1A).

In this study, we have developed a synthetic approach for the formation of **1a** and its deuterated analog **1b**. We envisioned that both **1a** and **1b** can be directly obtained by the base-promoted chromanone ring-opening of **2a** or **2b**, which in turn can be obtained from easily accessible naringenin (**3**) (*ca.* 1 \$/1 g) *via* two-step prenylation and Williamson etherification of the phenolic OH (Scheme 1B). The use of **3** as the starting material is beneficial as only one prenyl substituent has to be introduced.

The synthetic route leading to **1a** and **1b** is depicted in Scheme 2. Our synthesis commenced from naringenin (**3**), which was selectively converted to diester **4** (Ac<sub>2</sub>O, pyridine). *O*-Alkylation of **4** under Mitsunobu conditions (3-methyl-2-butene-1-ol, Ph<sub>3</sub>P, DIAD), followed by the catalytic Claisen rearrangement of **5** (Eu(fod)<sub>3</sub>, 1,2-dichloroethane, 80 °C) afforded prenyl-derivative **6**. Notably, performing the latter reaction in 1,2-dichloroethane above its boiling point was superior in comparison to earlier reports.<sup>23–26</sup> Alkylation of **6** (CH<sub>3</sub>I, Ag<sub>2</sub>O or CD<sub>3</sub>I, Ag<sub>2</sub>O) afforded **7a/7b** in good yields. An alternative approach to **7b** involving the alkylation of phenolic OH under Mitsunobu conditions (CD<sub>3</sub>OD, Ph<sub>3</sub>P, DIAD) required a large excess of reagents and afforded the product in moderate yield. Basic hydrolysis (KOH, MeOH) of esters afforded isoxanthohumols **2a/2b**. Although the chromane ring was stable during the hydrolysis, it could be opened under more harsh conditions (DBU, DMF, 70 °C),<sup>27,28</sup> leading to **1a/1b** in good yields after a mild acidic workup.

With **1a** and **1b** in hand, we investigated their MS-fragmentation patterns in electrospray ionization in positive and negative ion modes. The MRM transitions were found by an automatic procedure and they are listed in Table 1 (see ESI† for

<sup>a</sup>Department of Organic Chemistry, Faculty of Chemistry, Nicolaus Copernicus University, Gagarina 7, 87-100 Toruń, Poland

<sup>b</sup>Synthex Technologies Sp. z o.o., Gagarina 7/134B, 87-100 Toruń, Poland. E-mail: budny@synthex.com.pl


<sup>c</sup>Department of Biomedical and Polymer Chemistry, Faculty of Chemistry, Nicolaus Copernicus University, Gagarina 7, 87-100 Toruń, Poland

<sup>d</sup>Noctiluca S.A., Gagarina 7/41B, 87-100 Toruń, Poland

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d1ra05443k




## A) First synthetic approach to xanthohumol (ref. 11):



## Reagents and conditions:

i. Lewis acid, Ac<sub>2</sub>O or AcCl; ii. MOMCl, DIPEA, DCM, rt, 60%; iii. 3-methyl-2-butene-1-ol, DEAD, (Ph)<sub>3</sub>P, toluene/THF, rt, 80%; iv. *N,N*-dimethylaniline, 200 °C, 64%; v. dimethylsulfate, K<sub>2</sub>CO<sub>3</sub>, acetone, reflux, 82%; vi. 4-MOMO-C<sub>6</sub>H<sub>4</sub>-CHO, NaOH, MeOH reflux, 60%; vii. HCl (pH = 1), MeOH/H<sub>2</sub>O, rt, 72%


## B) This work:



Scheme 1 The background of the study.

details). In positive ion mode, the most intensive product ions in fragmentation of **1a** were ions with *m/z* values of 178.9, 299.0, 113.0, and 150.9. Corresponding ions with *m/z* +3 values can be found in the fragmentation of **1b**. On the other hand, in the negative ion mode, the same product ions are observed both in the case of **1a** and **1b**, indicating that the CH<sub>3</sub>/CD<sub>3</sub> groups were lost during fragmentations.

One of the criteria for an effective internal standard is the coelution of the labeled and non-labeled compounds during the HPLC analysis. This is particularly important in case of deuterium-labeled compounds as with the increase in the number of deuterons in the molecule, retention times may be extended. The retention times of **1a** and **1b** under different HPLC conditions are listed in Table 2. Notably, coelution of **1a**



## Reagents and conditions:

i. Ac<sub>2</sub>O (2 equiv.), pyridine, rt, 4 h, 80%; ii. 3-methyl-2-buten-1-ol (1.3 equiv.), Ph<sub>3</sub>P (1.2 equiv.), DIAD (1.2 equiv.), THF, 0 °C → rt, 12 h, 78%; iii. Eu(fod)<sub>3</sub> (10 mol%), (CH<sub>2</sub>Cl)<sub>2</sub>, 80 °C, 12 h, 61%; iv. CH<sub>3</sub>I (6.0 equiv.), Ag<sub>2</sub>O (4.0 equiv.), Et<sub>2</sub>O, reflux, 3 h, 62% for **7a** or CD<sub>3</sub>I (6.0 equiv.), Ag<sub>2</sub>O (4.0 equiv.), Et<sub>2</sub>O, reflux, 3 h, 86% for **7b**; v. KOH (2.0 equiv.), MeOH, 1 h, 99% for **2a**, 99% for **2b**; vi. DBU (2.0 equiv.), DMF, 70 °C, 12 h, 85% for **1a**, 72% for **1b**.

Scheme 2 The synthetic route to **1a** and **1b**.

Table 1 The MRM transitions of **1a** and **1b**

Ionization mode	<b>1a</b>		<b>1b</b>	
	Precursor ion	Product ion	Precursor ion	Product ion
ESI(+)	355.0	178.9	358.0	182.0
		299.0		302.0
		113.0		115.9
		150.9		107.9
		93.0		154.0
ESI(-)	353.0	119.1	356.0	119.1
		233.0		236.0
		295.1		295.2
		218.2		175.0
		175.0		218.1
		189.2		168.2

and **1b** is observed in case of the XB-C18 stationary phase (entry 1). Minor differences were observed when separation was attempted on C18-PFP (entry 2) and polar-C18 (entry 3) stationary phases.

As an example of applications, compound **1b** was used as an internal standard in a stable isotope dilution assay of xanthohumol (**1a**) in two Polish beers (determined concentrations: 0.4069 mg L<sup>-1</sup> and 0.5488 mg L<sup>-1</sup>, respectively). The developed MRM method allowed for the direct analysis of **1a** and any preconcentration of the analyte was not needed.

In conclusion, we have developed a six-step synthesis of xanthohumol (**1a**) and its deuterated analog **1b** from naringenin (**3**) in total 19.8% yield for **1a** and 23.3% for **1b**. In a key step, isoxanthohumols **2a/2b** were converted to the target compounds under basic conditions. The overall synthetic route was scalable and was used in the synthesis of **1a** on a 5 g scale. The MRM transitions of **1b** and its coelution with **1a** makes **1b** a suitable internal standard for the stable isotope dilution assay.

Table 2 Comparison of the retention times of **1a** and **1b** under different HPLC conditions

Entry	Conditions	Retention time [min]	
		<b>1a</b>	<b>1b</b>
1	Column: XB-C18, 100 × 3.0 mm, 2.6 μm, 100 Å; flow: 0.55 mL min <sup>-1</sup> ; oven: 35 °C; gradient MeOH/0.1% HCO <sub>2</sub> H(aq): from 5% MeOH to 95% MeOH	20.030	20.034
2	Column: Ace 5 C18-PFP, 250 × 4.6 mm; flow: 1.0 mL min <sup>-1</sup> , oven: 35 °C; isocrat. MeOH/0.1% HCO <sub>2</sub> H(aq): 80 : 20	19.615	19.700
3	Column: polar-C18, 100 × 3.0 mm, 2.6 μm, 100 Å; flow: 0.55 mL min <sup>-1</sup> ; oven: 35 °C; isocrat.: MeOH/0.1% HCO <sub>2</sub> H(aq): 65 : 35	6.590	6.520

## Experimental section

Tetrahydrofuran (THF) and *N,N*-dimethylformamide (DMF) were dried over activated molecular sieves 4 Å. Other solvents and reagents were of analytical grade and were used as received without further purification. NMR spectra were recorded on Bruker Avance III 300 MHz, 400 MHz, and 700 MHz spectrometers. Chemical shifts are reported as δ values in parts per million relative to the residual solvent signal (CDCl<sub>3</sub>: δ = 7.24 ppm for <sup>1</sup>H and 77.23 ppm for <sup>13</sup>C; CD<sub>3</sub>OD: δ = 3.31 ppm for <sup>1</sup>H and 49.15 ppm for <sup>13</sup>C). Coupling constants are in hertz (Hz). The following abbreviations are used for spin multiplicity: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, and br = broad. HPLC-ESI MS analyses were performed on a triple quadrupole Shimadzu LCMS 8030. Infrared spectra were recorded on a PerkinElmer UATR two instrument and are reported in cm<sup>-1</sup>. Melting points were determined in open glass capillaries and are uncorrected. Silica Gel 60, Merck 230–400, was used for preparative column chromatography. Sigma-Aldrich TLC plates (silica gel on Al foil with fluorescent indicator 254 nm) were used for analytical TLC. UV lamp (λ = 254 nm) and solution of phosphomolybdic acid in ethanol were used for the visualization of TLC plates.

### 4-(7-Acetoxy-5-hydroxy-4-oxochroman-2-yl)phenyl acetate (**4**)

To a suspension of naringenin (**3**) (1.0 equiv., 73.5 mmol, 20.0 g) in pyridine (60 mL), acetic anhydride (2.0 equiv., 147.0 mmol, 15.0 g) was added portionwise. The resulting solution was stirred until the complete consumption of the starting material. The reaction was quenched with conc. HCl (60 mL) and water (200 mL), extracted with EtOAc (3 × 100 mL), and washed with water (3 × 100 mL) and brine (100 mL). The extract was dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated under reduced pressure. The residue was recrystallized from MeOH to afford **4** (20.94 g, 80%) as a white solid.

Mp. 143–144 °C; IR (neat, cm<sup>-1</sup>): 2964, 1742, 1650, 1370, 1276, 1210, 1076, 1014, 840, 769; <sup>1</sup>H NMR (700 MHz, CDCl<sub>3</sub>), δ (ppm): 11.81 (s, 1H), 7.46–7.43 (m, 2H), 7.16–7.13 (m, 2H), 6.30 (d, *J* = 2.1 Hz, 1H), 6.29 (d, *J* = 2.1 Hz, 1H), 5.44 (dd, *J*<sub>1</sub> = 13.4 Hz, *J*<sub>2</sub> = 2.9 Hz, 1H), 3.08 (dd, *J*<sub>1</sub> = 17.2 Hz, *J*<sub>2</sub> = 13.4 Hz, 1H), 2.85 (dd, *J*<sub>1</sub> = 17.2 Hz, *J*<sub>2</sub> = 2.9 Hz, 1H), 2.30 (s, 3H), 2.27 (s, 3H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>), δ (ppm): 196.9, 169.4, 168.4, 163.6, 162.4, 158.7, 151.3, 135.8, 127.5, 122.3, 106.4, 103.6, 101.9, 79.0, 43.8, 21.4, 21.3; anal. calcd for C<sub>19</sub>H<sub>16</sub>O<sub>7</sub>: C, 64.04; H, 4.53. Found: C, 64.17; H, 4.60.

### 4-(7-Acetoxy-5-((3-methylbut-2-en-1-yl)oxy)-4-oxochroman-2-yl)phenyl acetate (**5**)

To a solution of **4** (1 equiv., 30.3 mmol, 10.792 g) in anhydrous THF (270 mL), 3-methylbut-2-en-1-ol (1.3 equiv., 39.4 mmol, 3.39 g) and triphenylphosphine (1.2 equiv., 36.4 mmol, 9.55 g) were added. The resulting solution was cooled to 0 °C, and then DIAD (1.2 equiv., 36.4 mmol, 7.36 g) was added portionwise. The resulting mixture was stirred at 0 °C for 30 min and then at room temperature (rt) overnight. The reaction mixture was concentrated under reduced pressure. The residue was



dissolved in MTBE (250 mL), and the resulting solution was stirred at 0 °C for 30 min. The precipitated triphenylphosphine oxide was filtered off and the solution was concentrated under a reduced pressure. The residue was purified by flash chromatography (petroleum ether-ethyl acetate = 70 : 30 → 60 : 40) affording **5** (10.03 g, 78%) as a pale yellow solid.

Mp. 113–116 °C; IR (neat, cm<sup>-1</sup>): 2979, 1764, 1680, 1596, 1373, 1197, 1108, 1076, 1031, 904, 843; <sup>1</sup>H NMR (700 MHz, CDCl<sub>3</sub>), δ (ppm): 7.46–7.43 (m, 2H), 7.13–7.11 (m, 2H), 6.40 (d, *J* = 2.1 Hz, 1H), 6.29 (d, *J* = 2.1 Hz, 1H), 5.53–5.49 (m, 1H), 5.41 (dd, *J*<sub>1</sub> = 13.2 Hz, *J*<sub>2</sub> = 2.7 Hz, 1H), 4.62–4.56 (m, 2H), 2.99 (dd, *J*<sub>1</sub> = 16.4 Hz, *J*<sub>2</sub> = 13.2 Hz, 1H), 2.80 (dd, *J*<sub>1</sub> = 16.4 Hz, *J*<sub>2</sub> = 2.7 Hz, 1H), 2.29 (s, 3H), 2.28 (s, 3H), 1.77 (br s, 3H), 1.72 (br s, 3H). <sup>13</sup>C NMR (175 MHz, CDCl<sub>3</sub>), δ (ppm): 189.2, 169.6, 168.6, 163.9, 161.4, 156.6, 151.0, 138.5, 136.3, 127.5, 122.2, 119.1, 109.8, 103.4, 100.1, 78.9, 66.3, 46.0, 26.0, 21.4, 21.3, 18.6; anal. calcd for C<sub>24</sub>H<sub>24</sub>O<sub>7</sub>: C, 67.91; H, 5.70. Found: C, 68.01; H, 5.68.

#### 4-(7-Acetoxy-5-hydroxy-8-(3-methylbut-2-en-1-yl)-4-oxochroman-2-yl)phenyl acetate (**6**)

In a 50 mL pressure vial, **5** (1 equiv., 6.96 mmol, 2.95 g), Eu(fod)<sub>3</sub> (0.1 equiv., 0.696 mmol, 722 mg) and 1,2-dichloroethane (6 mL) were placed. The resulting mixture was stirred at 80 °C (temp. of the oil bath) overnight. Then, the mixture was cooled to rt and directly purified by flash chromatography (petroleum ether-ethyl acetate = 90 : 10 → 70 : 30) to afford **6** (1.80 g, 61%) as a white solid.

Mp. 143–144 °C; IR (neat, cm<sup>-1</sup>): 2975, 1763, 1637, 1428, 1372, 1189, 1066, 1012, 896; <sup>1</sup>H NMR (700 MHz, CDCl<sub>3</sub>), δ (ppm): 11.70 (s, 1H), 7.46–7.43 (m, 2H), 7.16–7.13 (m, 2H), 6.29 (s, 1H), 5.43 (dd, *J*<sub>1</sub> = 13.3 Hz, *J*<sub>2</sub> = 2.9 Hz, 1H), 5.06–5.01 (m, 1H), 3.18–3.10 (m, 2H), 3.06 (dd, *J*<sub>1</sub> = 17.1 Hz, *J*<sub>2</sub> = 13.3 Hz, 1H), 2.87 (dd, *J*<sub>1</sub> = 17.1 Hz, *J*<sub>2</sub> = 2.9 Hz, 1H), 2.30 (s, 3H), 2.28 (s, 3H), 1.64–1.62 (m, 3H), 1.58–1.56 (m, 3H). <sup>13</sup>C NMR (175 MHz, CDCl<sub>3</sub>), δ (ppm): 197.3, 169.4, 168.4, 161.1, 160.0, 157.0, 151.2, 136.1, 132.1, 127.4, 122.2, 121.8, 113.8, 106.8, 104.3, 78.9, 43.8, 25.8, 22.9, 21.3, 21.1, 18.0; anal. calcd for C<sub>24</sub>H<sub>24</sub>O<sub>7</sub>: C, 67.91; H, 5.70; N, 14.42. Found: C, 67.94; H, 5.82.

#### 4-(7-Acetoxy-5-methoxy-8-(3-methylbut-2-en-1-yl)-4-oxochroman-2-yl)phenyl acetate (**7a**)

To a suspension of **6** (1 equiv., 9.46 mmol, 4.012 g) and freshly prepared Ag<sub>2</sub>O (4 equiv., 37.85 mmol, 8.770 g) in Et<sub>2</sub>O (95 mL), MeI (6 equiv., 56.772 mmol, 8.058 g, 3.66 mL) was added. The resulting mixture was refluxed for 3 h. Then, the reaction mixture was cooled to rt, silver salts were filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography (petroleum ether-ethyl acetate = 60 : 40) affording **7a** (2.59 g, 62%) as a white solid.

Mp. 140–141 °C; IR (neat, cm<sup>-1</sup>): 2962, 1753, 1649, 1592, 1369, 1203, 1098, 1055, 902, 834; <sup>1</sup>H NMR (700 MHz, CDCl<sub>3</sub>), δ (ppm): 7.46–7.42 (m, 2H), 7.14–7.10 (m, 2H), 6.28 (s, 1H), 5.41 (dd, *J*<sub>1</sub> = 13.3 Hz, *J*<sub>2</sub> = 2.8 Hz, 1H), 5.07–5.02 (m, 1H), 3.86 (s, 3H), 3.21–3.15 (m, 2H), 2.98 (dd, *J*<sub>1</sub> = 16.4 Hz, *J*<sub>2</sub> = 13.3 Hz, 1H), 2.83 (dd, *J*<sub>1</sub> = 16.4 Hz, *J*<sub>2</sub> = 2.8 Hz, 1H), 2.31 (s, 3H), 2.30 (s, 3H), 1.64–1.63 (m, 3H), 1.58–1.56 (m, 3H). <sup>13</sup>C NMR (175 MHz, CDCl<sub>3</sub>), δ (ppm): 190.1, 169.6, 168.9, 161.9, 159.6, 154.8, 150.9,

136.5, 132.2, 127.4, 122.1, 121.7, 115.4, 109.9, 99.6, 78.8, 56.5, 45.8, 25.9, 23.2, 21.4, 21.1, 18.00; anal. calcd for C<sub>25</sub>H<sub>26</sub>O<sub>7</sub>: C, 68.48; H, 5.98. Found: C, 68.21; H, 6.11.

#### 4-(7-Acetoxy-5-(methoxy-d<sub>3</sub>)-8-(3-methylbut-2-en-1-yl)-4-oxochroman-2-yl)phenyl acetate (**7b**)

The same procedure as for compound **7a** was used. Starting from **6** (2.36 mmol, 1.0 g) **7b** (0.89 g, 86%) as a pale yellow solid was obtained.

Mp. 138–139 °C; IR (neat, cm<sup>-1</sup>): 1754, 1685, 1593, 1362, 1204, 1091, 902. <sup>1</sup>H NMR (700 MHz, CDCl<sub>3</sub>), δ (ppm): 7.46–7.42 (m, 2H), 7.14–7.10 (m, 2H), 6.27 (s, 1H), 5.41 (dd, *J*<sub>1</sub> = 13.1 Hz, *J*<sub>2</sub> = 2.8 Hz, 1H), 5.07–5.03 (m, 1H), 3.21–3.14 (m, 2H), 2.98 (dd, *J*<sub>1</sub> = 16.5 Hz, *J*<sub>2</sub> = 13.2 Hz, 1H), 2.83 (dd, *J*<sub>1</sub> = 16.5 Hz, *J*<sub>2</sub> = 2.8 Hz, 1H), 2.30 (s, 3H), 2.29 (s, 3H), 1.64–1.62 (m, 3H), 1.58–1.56 (m, 3H). <sup>13</sup>C NMR (175 MHz, CDCl<sub>3</sub>), δ (ppm): 189.9, 169.4, 168.8, 161.8, 159.7, 154.9, 151.0, 136.6, 132.2, 127.3, 122.1, 121.8, 115.4, 110.0, 99.6, 78.8, 78.8, 45.8, 25.8, 23.2, 21.3, 21.1, 18.0; anal. calcd for C<sub>25</sub>H<sub>23</sub>D<sub>3</sub>O<sub>7</sub>: C, 68.01; H, 6.62. Found: C, 68.07; H, 6.60.

#### 7-Hydroxy-2-(4-hydroxyphenyl)-5-methoxy-8-(3-methylbut-2-en-1-yl)chroman-4-one (**2a**)

To a solution of **7a** (1 equiv., 4.349 mmol, 1.905 g) in MeOH (30 mL), KOH (2 equiv., 8.699 mmol, 488 mg) was added. The reaction mixture was stirred until the complete consumption of the starting material. The reaction mixture was neutralized with 2 M HCl, extracted with EtOAc (3 × 60 mL), washed with water (2 × 30 mL) and brine (30 mL), dried over anhydrous MgSO<sub>4</sub>, filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (petroleum ether-ethyl acetate = 50 : 50) to afford **2a** (1.524 g, 99%) as a pale yellow solid.

Mp. 157–161 °C; IR (neat, cm<sup>-1</sup>): 3150, 1646, 1590, 1452, 1410, 1349, 1270, 1088, 827; <sup>1</sup>H NMR (700 MHz, CD<sub>3</sub>OD), δ (ppm): 7.33–7.28 (m, 2H), 6.83–6.79 (m, 2H), 6.11 (s, 1H), 5.28 (dd, *J*<sub>1</sub> = 12.8 Hz, *J*<sub>2</sub> = 3.0 Hz, 1H), 5.15–5.12 (m, 1H), 3.79 (s, 3H), 3.24–3.17 (m, 2H), 2.97 (dd, *J*<sub>1</sub> = 16.7 Hz, *J*<sub>2</sub> = 12.8 Hz, 1H), 2.66 (dd, *J*<sub>1</sub> = 16.7 Hz, *J*<sub>2</sub> = 3.0 Hz, 1H), 1.62 (s, 3H), 1.56 (s, 3H). <sup>13</sup>C NMR (700 MHz, CD<sub>3</sub>OD), δ (ppm): 193.0, 164.4, 164.0, 159.0, 131.8, 131.7, 129.0, 124.1, 116.4, 110.2, 106.1, 93.8, 80.2, 56.1, 46.4, 26.0, 22.9, 18.0; anal. calcd for C<sub>21</sub>H<sub>22</sub>O<sub>5</sub>: C, 71.17; H, 6.26. Found: C, 71.38; H, 6.42.

#### 7-Hydroxy-2-(4-hydroxyphenyl)-5-(methoxy-d<sub>3</sub>)-8-(3-methylbut-2-en-1-yl)chroman-4-one (**2b**)

The same procedure as for compound **2a** was used. Starting from **7b** (4.33 mmol, 1.799 g) **2b** (1.530 g, 99%) as a pale yellow solid was obtained.

Mp. 150–151 °C; IR (neat, cm<sup>-1</sup>): 3164, 1591, 1509, 1422, 1359, 1259, 1095, 823; <sup>1</sup>H NMR (700 MHz, CD<sub>3</sub>OD), δ (ppm): 7.33–7.28 (m, 2H), 6.83–6.78 (m, 2H), 6.10 (s, 1H), 5.27 (dd, *J*<sub>1</sub> = 12.8 Hz, *J*<sub>2</sub> = 2.9 Hz, 1H), 5.18–5.11 (m, 1H), 3.24–3.17 (m, 2H), 2.97 (dd, *J*<sub>1</sub> = 16.5 Hz, *J*<sub>2</sub> = 12.8 Hz, 1H), 2.65 (dd, *J*<sub>1</sub> = 16.5 Hz, *J*<sub>2</sub> = 2.9 Hz, 1H), 1.61 (s, 3H), 1.56 (s, 3H). <sup>13</sup>C NMR (100 MHz, CD<sub>3</sub>OD), δ (ppm): 193.0, 164.3, 164.0, 162.0, 158.9, 131.8, 131.7, 129.0, 124.0, 116.4, 110.1, 106.1, 93.6, 80.1, 46.4, 26.1, 22.9, 18.0; anal. calcd for C<sub>21</sub>H<sub>19</sub>D<sub>3</sub>O<sub>5</sub>: C, 70.57; H, 7.05. Found: C, 70.46; H, 6.96.



**(E)-1-(2,4-Dihydroxy-6-methoxy-3-(3-methylbut-2-en-1-yl)phenyl)-3-(4-hydroxyphenyl)prop-2-en-1-one (1a)**

To a solution of **2a** (1 equiv., 4.349 mmol, 1.544 g) in DMF (20 mL), DBU (2 equiv., 8.699 mmol, 1.324 g) was added dropwise and the reaction mixture was stirred at 70 °C for 12 h. Then, the reaction mixture was neutralized with 2 M HCl, extracted with EtOAc (4 × 60 mL), washed with water (3 × 30 mL) and brine (30 mL). The combined extracts were dried over anhydrous MgSO<sub>4</sub>, filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (petroleum ether-ethyl acetate = 60 : 40) affording **1a** (1.31 g, 85%) as orange solid.

Mp. 151–152 °C; IR (neat, cm<sup>-1</sup>): 3184, 2916, 1596, 1511, 1437, 1340, 1100, 824, 804; <sup>1</sup>H NMR (700 MHz, CD<sub>3</sub>OD), δ (ppm): 7.80 (d, *J* = 15.6 Hz, 1H), 7.67 (d, *J* = 15.6 Hz, 1H), 7.51–7.49 (m, 2H), 6.85–6.81 (m, 2H), 6.02 (s, 1H), 5.22–5.18 (m, 1H), 3.09 (s, 3H), 3.23 (d, *J* = 7.1 Hz, 2H), 1.76 (s, 3H), 1.65 (s, 3H). <sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD), δ (ppm): 194.2, 166.3, 163.6, 161.2, 143.4, 131.5, 131.4, 128.6, 126.0, 124.4, 117.0, 109.6, 106.7, 91.8, 56.3, 26.1, 22.4, 18.0; anal. calcd for C<sub>21</sub>H<sub>22</sub>O<sub>5</sub>: C, 71.17; H, 6.26. Found: C, 71.03; H, 6.20.

**(E)-1-(2,4-Dihydroxy-6-(methoxy-d<sub>3</sub>)-3-(3-methylbut-2-en-1-yl)phenyl)-3-(4-hydroxyphenyl)prop-2-en-1-one (1b)**

The same procedure as for compound **1b** was used.

Starting from **2b** (4.43 mmol, 1.581 g) **1b** (1.141 g, 72%) as a pale yellow solid was obtained.

Mp. 149–151 °C, IR (neat, cm<sup>-1</sup>): 3302, 2915, 1521, 1439, 1337, 1213, 1157, 1098, 830; <sup>1</sup>H NMR (700 MHz, CD<sub>3</sub>OD), δ (ppm): 7.79 (d, *J* = 15.5 Hz, 1H), 7.66 (d, *J* = 15.5 Hz, 1H), 7.51–7.46 (m, 2H), 6.85–6.79 (m, 2H), 6.01 (s, 1H), 5.22–5.18 (m, 1H), 3.23 (d, *J* = 7.0 Hz, 2H), 1.76 (s, 3H), 1.65 (s, 3H). <sup>13</sup>C NMR (100 MHz, CD<sub>3</sub>OD), δ (ppm): 194.3, 166.3, 163.8, 162.6, 161.1, 143.4, 131.5, 131.4, 128.7, 126.1, 124.4, 117.0, 109.6, 106.7, 91.9, 26.1, 22.4, 18.0; anal. calcd for C<sub>21</sub>H<sub>19</sub>D<sub>3</sub>O<sub>5</sub>: C, 70.57; H, 7.05. Found: C, 70.55; H, 6.97.

## Conflicts of interest

The authors declare no conflict of interest.

## Acknowledgements

Financial support from Synthex Technologies Sp. z o.o. and Noctiluca S.A. is kindly acknowledged. Financial support from the Polish Ministry of Science and Higher Education (“Industrial Doctorates 1<sup>st</sup> Edition”) for Joanna Andrusiak is kindly acknowledged.

## References

- J. F. Stevens and J. E. Page, Xanthohumol and Related Prenylflavonoids from Hops and Beer: To Your Good Health!, *Phytochemistry*, 2004, **65**(10), 1317–1330, DOI: 10.1016/j.phytochem.2004.04.025.
- C. Gerhäuser, A. Alt, E. Heiss, A. Gamal-Eldeen, K. Klimo, J. Knauff, I. Neumann, H.-R. Scherf, N. Frank, H. Bartsch

and H. Becker, Cancer Chemopreventive Activity of Xanthohumol, a Natural Product Derived from Hop 1 Support for This Work Has Been Provided by Verein Zur Förderung Der Krebsforschung in Deutschland e.V. and by Wissenschaftsförderung Der Deutschen Brauwirtschaft e.V., *Mol. Cancer Ther.*, 2002, **1**(11), 959–969.

- M. Liu, P. E. Hansen, G. Wang, L. Qiu, J. Dong, H. Yin, Z. Qian, M. Yang and J. Miao, Pharmacological Profile of Xanthohumol, a Prenylated Flavonoid from Hops (*Humulus Lupulus*), *Molecules*, 2015, **20**(1), 754–779, DOI: 10.3390/molecules20010754.
- M. Stompor, K. Danciewicz, B. Gabryś and M. Anioł, Insect Antifeedant Potential of Xanthohumol, Isoxanthohumol, and Their Derivatives, *J. Agric. Food Chem.*, 2015, **63**(30), 6749–6756, DOI: 10.1021/acs.jafc.5b02025.
- Z. A. Nowakowska, Review of Anti-Infective and Anti-Inflammatory Chalcones, *Eur. J. Med. Chem.*, 2007, **42**(2), 125–137, DOI: 10.1016/j.ejmech.2006.09.019.
- C. Gerhäuser, Beer Constituents as Potential Cancer Chemopreventive Agents, *Eur. J. Cancer*, 2005, **41**(13), 1941–1954, DOI: 10.1016/j.ejca.2005.04.012.
- B. Orlikova, D. Tasdemir, F. Golais, M. Dicato and M. Diederich, Dietary Chalcones with Chemopreventive and Chemotherapeutic Potential, *Genes Nutr.*, 2011, **6**(2), 125–147, DOI: 10.1007/s12263-011-0210-5.
- Y. Zhou, J. Zheng, Y. Li, D.-P. Xu, S. Li, Y.-M. Chen and H.-B. Li, Natural Polyphenols for Prevention and Treatment of Cancer, *Nutrients*, 2016, **8**(8), 515, DOI: 10.3390/nu8080515.
- D. I. Batovska and I. T. Todorova, Trends in Utilization of the Pharmacological Potential of Chalcones, *Curr. Clin. Pharmacol.*, 2010, 1–29, DOI: 10.2174/157488410790410579.
- F. B. Power, F. Tutin and H. Rogerson, CXXXV.—The Constituents of Hops, *J. Chem. Soc., Trans.*, 1913, **103**, 1267–1292, DOI: 10.1039/CT9130301267.
- R. S. Khupse and P. W. Erhardt, Total Synthesis of Xanthohumol, *J. Nat. Prod.*, 2007, **70**(9), 1507–1509, DOI: 10.1021/np070158y.
- S. Vogel, S. Ohmayer, G. Brunner and J. Heilmann, Natural and Non-Natural Prenylated Chalcones: Synthesis, Cytotoxicity and Anti-Oxidative Activity, *Bioorg. Med. Chem.*, 2008, **16**(8), 4286–4293, DOI: 10.1016/j.bmc.2008.02.079.
- J. Yao, B. Zhang, C. Ge, S. Peng and J. Fang, Xanthohumol, a Polyphenol Chalcone Present in Hops, Activating Nrf2 Enzymes To Confer Protection against Oxidative Damage in PC<sub>12</sub> Cells, *J. Agric. Food Chem.*, 2015, **63**(5), 1521–1531, DOI: 10.1021/jf505075n.
- B. Zhang, D. Duan, C. Ge, J. Yao, Y. Liu, X. Li and J. Fang, Synthesis of Xanthohumol Analogues and Discovery of Potent Thioredoxin Reductase Inhibitor as Potential Anticancer Agent, *J. Med. Chem.*, 2015, **58**(4), 1795–1805, DOI: 10.1021/jm5016507.
- E. Ciccimaro and I. A. Blair, Stable-Isotope Dilution LC–MS for Quantitative Biomarker Analysis, *Bioanalysis*, 2010, **2**(2), 311–341, DOI: 10.4155/bio.09.185.
- A. Chokkathukalam, D.-H. Kim, M. P. Barrett, R. Breitling and D. J. Creek, Stable Isotope-Labeling Studies in



- Metabolomics: New Insights into Structure and Dynamics of Metabolic Networks, *Bioanalysis*, 2014, **6**(4), 511–524, DOI: 10.4155/bio.13.348.
- 17 C. Schmidt, First Deuterated Drug Approved, *Nat. Biotechnol.*, 2017, **35**(6), 493–494, DOI: 10.1038/nbt0617-493.
- 18 J. F. Liu, S. L. Harbeson, C. L. Brummel, R. Tung, R. Silverman and D. Doller, Chapter Fourteen – A Decade of Deuteration in Medicinal Chemistry, in *Platform Technologies in Drug Discovery and Validation*, ed. R. A. Goodnow, Academic Press, 2017, vol. 50, pp. 519–542, DOI: 10.1016/bs.armc.2017.08.010.
- 19 S. Cargnin, M. Serafini and T. Pirali, A Primer of Deuterium in Drug Design, *Future Med. Chem.*, 2019, **11**(16), 2039–2042, DOI: 10.4155/fmc-2019-0183.
- 20 D. C. Ellinwood, M. F. El-Mansy, L. S. Plagmann, J. F. Stevens, C. S. Maier, A. F. Gombart and P. R. Blakemore, Total Synthesis of [<sup>13</sup>C]<sub>2</sub>, [<sup>13</sup>C]<sub>3</sub>, and [<sup>13</sup>C]<sub>5</sub>-Isotopomers of Xanthohumol, the Principal Prenylflavonoid from Hops, *J. Labelled Compd. Radiopharm.*, 2017, **60**(14), 639–648, DOI: 10.1002/jlcr.3571.
- 21 S. Berwanger, J. Zapp and H. Becker, <sup>13</sup>C-Labeling of Xanthohumol in Hop Cones (*Humulus Lupulus*), *Planta Med.*, 2005, **71**(6), 530–534.
- 22 L. Buckett, S. Schinko, C. Urmann, H. Riepl and M. Rychlik, Stable Isotope Dilution Analysis of the Major Prenylated Flavonoids Found in Beer, Hop Tea, and Hops, *Front. Nutr.*, 2020, 319.
- 23 S. Gester, P. Metz, O. Zierau and G. Vollmer, An Efficient Synthesis of the Potent Phytoestrogens 8-Prenylnaringenin and 6-(1,1-Dimethylallyl)Naringenin by Europium(III)-Catalyzed Claisen Rearrangement, *Tetrahedron*, 2001, **57**(6), 1015–1018, DOI: 10.1016/S0040-4020(00)01078-4.
- 24 S. Tischer and P. Metz, Selective C-6 Prenylation of Flavonoids via Europium(III)-Catalyzed Claisen Rearrangement and Cross-Metathesis, *Adv. Synth. Catal.*, 2007, **349**(1–2), 147–151, DOI: 10.1002/adsc.200600454.
- 25 A. Keßberg and P. Metz, Utilizing an *O*-Quinone Methide in Asymmetric Transfer Hydrogenation: Enantioselective Synthesis of Brosimine A, Brosimine B, and Brosimacutin L, *Angew. Chem., Int. Ed.*, 2016, **55**(3), 1160–1163, DOI: 10.1002/anie.201507269.
- 26 O. Schaefer, R. Bohlmann, W.-D. Schleuning, K. Schulze-Forster and M. Hümpel, Development of a Radioimmunoassay for the Quantitative Determination of 8-Prenylnaringenin in Biological Matrices, *J. Agric. Food Chem.*, 2005, **53**(8), 2881–2889, DOI: 10.1021/jf047897u.
- 27 H. Zhang, Z. Wang, I. Ghiviriga, G. G. Pillai, F. Jabeen, J. A. Arami, W. Zhou, P. J. Steel, C. Dennis Hall and A. R. Katritzky, Synthesis, Characterization and Energetic Properties of Novel 1-Methyl-1,2,4-Triazolium *N*-Aryl/*N*-Pyridinyl Ylids, *Tetrahedron Lett.*, 2017, **58**(12), 1079–1085, DOI: 10.1016/j.tetlet.2016.12.016.
- 28 N. Jun, G. Hong and K. Jun, Synthesis and Evaluation of 2',4',6'-Trihydroxychalcones as a New Class of Tyrosinase Inhibitors, *Bioorg. Med. Chem.*, 2007, **15**(6), 2396–2402, DOI: 10.1016/j.bmc.2007.01.017.

