


 Cite this: *RSC Adv.*, 2021, **11**, 39058

Nanomaterial catalyzed green synthesis of tetrazoles and its derivatives: a review on recent advancements

 Suman Swami, ^a Satya Narayan Sahu ^b and Rahul Shrivastava ^{*a}

Tetrazoles are indispensable nitrogen containing heterocyclic scaffolds that offer a broad spectrum of applications in various domains such as medicinal chemistry, high energy material science, biochemistry, pharmacology etc. Owing to their useful applications, a wide range of catalysts have been explored for green synthesis of tetrazole derivatives. In recent times, nanomaterials have emerged as extremely efficient catalysts for different organic transformations because of their high surface area-to-volume ratio, easy surface modification, simple fabrications, easy recovery and reusability. In this article, we have presented an overview of utilization of various nano-catalysts, nanocomposites and other solid-supported nanomaterials as an efficient environmental benign catalytic system for green synthesis of tetrazoles and derivatives. This review will provide an exclusive emphasis on boehmite, magnetic, copper, carbon, MCM-41, and composite based nanomaterials that have been developed since the year 2010 for the synthesis of tetrazole derivatives. In addition, we have briefly discussed the fabrication, functionalization and characterization of some novel nanomaterials and their advantages in the synthesis of tetrazole and its derivatives along with the reaction mechanism that involves synthesis of tetrazole derivatives via nanomaterials catalysed reactions.

 Received 6th August 2021
 Accepted 11th November 2021

DOI: 10.1039/d1ra05955f

rsc.li/rsc-advances

^aDepartment of Chemistry, Manipal University Jaipur, VPO-Dehmi-Kalan, Off Jaipur-Ajmer Express Way, Jaipur, Rajasthan, India-303007. E-mail: chem.rahul@gmail.com

^bSchool of Chemistry, Sambalpur University, Jyoti Vihar, Burla, Sambalpur, Orissa, India-768019

1. Introduction

Chemicals are necessary for production of various essential products, pharmaceutically relevant moieties, food additives, pesticides, fertilizers and many more. The production of these necessary products involves well established chemical routes



Dr Suman Swami is currently a CSIR research associate at the Department of Chemistry Manipal University Jaipur, India. Dr Swami did her master's in Chemistry from St. Wilfred's College (University of Rajasthan) in 2011. She carried out her doctoral studies in organic chemistry at the Manipal University Jaipur, India, and was awarded a PhD in 2018. Subsequently, she moved for

a professional course in sanitation to IHE Delft, The Netherlands. Her research areas include nano-catalyst, heterocyclic synthesis, multicomponent reaction, chemosensors, and nanosensors development for hazardous metal ions and anion sensing.



Dr Satya Narayan Sahu earned his MSc degree in Chemistry from Utkal University, Odisha, in 2001 and received his MTech degree in Modern Methods of Chemical Analysis in 2003 and PhD degree in Organic Chemistry in 2009 from Indian Institute of Technology, Delhi. After completion of his PhD he joined Sambalpur University, Odisha, as an Assistant Professor in Chemistry, and then moved to

Bielefeld University, Germany for post-doctoral work after receiving the BOYSCAST fellowship in 2010 from the Department of Science and Technology, Government of India. His current research interest focuses on the synthesis of chromofluorescent molecular sensors for ionic and molecular recognitions.






Fig. 1 Tetrazole derivatives possessing devised important pharmacological properties.

that have not been upgraded or modified over the years by chemical industries. Many of these chemical processes accomplished with high temperature, pressure, toxic solvent, catalyst, chemicals and generate enormous amount of toxic chemical waste which impose hazardous impacts on environment and health.^{1,2} In recent years, environmental policies have become more stringent because of increased awareness about harmful impacts of generated by-products and reagents on environment

and living species. Further, owing to global health crisis generated by industrial toxic chemical wastes, various regulatory bodies demand to adopt eco-compatible greener synthetic processes against the conventional synthetic procedures in chemical industries. Among the various aspects for advancement in chemical process, use of green and efficient catalyst played a key role with aim of reducing temperature, pressure, reagent-based waste with minimum unwanted toxic by-products. In this context, designing and development of sustainable environmental benign catalysts are one of the urgently indispensable challenge facing by chemists. The idea of green chemistry has become an integral part of sustainability which makes catalysis science even more innovative. From the viewpoint of green chemistry, sustainable catalyst must possess a series of distinct advantages like high activity, selectivity and efficiency, reasonable recovery, high stability and excellent recyclability.

Catalysts can be generally divided into two types based on reaction phase: (i) homogeneous (ii) heterogeneous catalyst. Homogenous catalysts have several advantages such as high selectivity, better yield, and high turnover numbers due to its solubility in reaction medium which makes catalytic sites of homogeneous catalyst more accessible. Despite its significant advantages, separation of homogenous catalyst from reaction medium is time-consuming and laborious task and needs



Dr Rahul Shrivastava is Associate professor of organic chemistry at the Manipal University Jaipur, Jaipur, India. Dr Shrivastava received his Master's degree from Indian Institute of Technology, Delhi in 2002. He obtained his PhD degree from Indian Institute of Technology Delhi, India in 2009. His research interest includes molecular recognition, calixarene and cyclo-triveratrylene based receptor molecules, nano sensors, chemosensor, nano-catalysts and multi-component reactions.



specific techniques. Also, homogenous catalyst derived chemical process are not compatible with green chemistry principles in modern-day catalysis science, especially from environmental and economic standpoint. Therefore, in recent years, chemists have focused their effort to transform homogenous catalytic process by heterogeneous catalysts so that they can avoid indorsed limitation. During the last two decades, chemists all over the world are fascinated toward nanomaterials that can be used as an excellent alternative of conventional catalytic system thereby can reduce gap between homogenous and heterogeneous catalysis. The nanomaterials could provide incredible solid support for immobilization of homogeneous catalysts to produce combine benefits of homogeneous and heterogeneous catalysis. Nano catalysts or nanomaterials support can assist chemists in development of sustainable catalyst with outstanding activity, selectivity and stability by fine-tuning their morphologies, shapes and sizes. These nanomaterials based catalytic systems are useful for numerous reasons such as:

- The interaction between nano catalyst and reactant are significantly higher due to the large surface area of nano catalyst which facilitates excellent rate of reaction in comparison to its homogeneous counterpart.
- Due to nano meter size, their surface area-to-volume ratio are very high.
- The sizes, shapes, and morphology of nano catalysts can be easily tuned as per the necessity of catalytic applications which would generally be tricky in their homogeneous counterparts.
- The recovery of magnetic nano catalysts from reaction mixtures with help of an external magnet is easy and convenient.
- Nanomaterials based catalysts are recyclable and can be reused many times which could bring them for more industrial applications, especially from economic standpoint.

To develop innovative pharmaceutical entities, number of novel heterocyclic scaffolds have been explored based on their intrinsic toxicity towards numerous pathogens. Amongst the various nitrogen-containing heterocycles, tetrazole based molecules have attracted considerable attention due to their potential biological applications towards antifungal,³⁻⁷ anti-HIV,⁸ antitubercular,⁹ antibacterial,¹⁰ antiulcer,^{11,12} anti-allergic,¹³ antihypertensive,¹⁴ antileishmanial,¹⁵ anticancer,¹⁶⁻¹⁹ antibacterial and antiproliferative²⁰ activities (Fig. 1). Further, tetrazole derivatives are leading bio-isosteres of carboxylic acids/cis-amide which play important roles in enhancement of lipophilicity, bioavailability, pharmacokinetic properties and reduce side effects of the drug molecules.¹⁶ Besides medicinal utility, tetrazoles are also widely employed in various fields of research such as high energy material science, photography, bioimaging agents, explosives and organometallic chemistry *etc.*^{21,22}

Consequently, owing to the potent applications of tetrazole scaffolds in diverse areas, different methodology for synthesis of tetrazole entities have been attempted. In general, tetrazoles are directly prepared either by: (a) [3+2] cycloaddition reaction between isocyanides and hydrazoic acid/trimethyl azide²³⁻²⁵ or (b) one-pot multicomponent reaction (MCR) between amine/organic nitrile, triethyl orthoformate/aldehyde and sodium azide²¹ under influence of an appropriate catalyst and solvent conditions as shown in Fig. 2.

It was observed that despite several advantages of conventional synthetic route [3+2] cycloaddition reaction of isocyanides and trimethyl azide/hydrazoic acids, this methodology also suffers some limitations such as uses of hazardous acids, toxic solvents, high temperatures and low to moderate yields and longer reaction times. Therefore, various research group offered multicomponent reaction (MCR) approaches with




Fig. 2 Synthetic routes for tetrazole derivatives; (a) conventional [3+2] cycloaddition reaction of isocyanides and trimethyl azide/hydrazoic acid and (b) one-pot multicomponent reaction (MCR) of amine/organic nitrile, triethyl orthoformate/aldehyde and sodium azide.






Fig. 3 Progress in research papers: synthesis of tetrazoles and its derivatives by using various nanomaterials. Data collected from various search engines (Google Scholar, Science Direct, Research Gate, and PubMed Central etc.) using key words: tetrazole, nanomaterials catalyzed synthesis of tetrazole.

suitable catalysts, solvents, and reaction conditions to overcome limitations associated with conventional methodology. In this context, number of acidic homogenous and heterogenous catalysts were investigated for synthesis of tetrazole derivatives *via* multicomponent reaction. It was found that despite high selectivity and activity of homogenous catalyst, it suffers from

few drawbacks such as tedious workup and purification procedure, difficulties in separation of catalysts and high temperature whereas using traditional heterogeneous catalyst, low yield and long reaction time are main disadvantages for synthesis of tetrazoles derivatives. Therefore, considering limitations associated with both homogenous and heterogeneous catalyst systems, researcher devoted massive efforts to achieve green synthetic methodology for synthesis of tetrazole *via* introducing nanostructured catalyst systems.

As a result, several nano-catalysts or nanomaterials supported catalysts have been explored to avoid drawbacks associated with the conventional catalytic system for synthesis of tetrazole derivatives. It is observed that interest in synthesis of tetrazole derivatives *via* environmental benign process by using various nanomaterials have continuously risen as depicted by Fig. 3. As shown in figure, ample magnetic and non-magnetic nanomaterials-based research articles are published since 2010 for tetrazoles synthesis comprising the green chemical processes.

The detail literature survey reflects that no systematic and comprehensive overview approaching nanomaterials catalyzed tetrazoles synthesis have been reported so far. The published reviews on nanoparticles catalyzed heterocyclic compounds are majorly discussed on synthesis of numerous diverse heterocyclic ring system rather than emphasis only on tetrazole and




Fig. 4 Class of nanostructures green catalyst engaged in the tetrazole synthesis.



derivatives. Thus, we believe that an insightful overview of the established nano-catalysts employed for synthesis of tetrazole derivatives, would be a driving step to the scientific community working in this domain. In the present review, we have emphasized mainly on exploring applications of different nanoparticle-based catalysts used in tetrazole synthesis through multicomponent reaction so far. Examples described in the present review are grouped under different sections (Fig. 4) based on boehmite, magnetic, non-magnetic, copper, ZnO, MCM-41 and carbon nanoparticles catalyzed synthesis of tetrazole derivatives with comprehensive analysis on synthesis and advantages thereof.

2. Boehmite based nano-catalyzed synthesis of 5-substitute tetrazoles

Boehmite is an aluminum oxide hydroxide (γ -AlOOH) having large number of hydroxyl groups on its surface (Fig. 5).^{26–28} These terminal hydroxyl groups provide high reactivity and hydrophilic environment to boehmite which enhances catalytic activities. In view of high reactivity and hydrophilic nature of boehmite, it was envisaged that transforming boehmite from molecular scale level to nanoscale level makes them even more attractive and efficient catalyst for various organic transformations. It was found that at nanoscale, boehmite devises several appealing features such as high thermal, mechanical, and non-toxicity, high dispersion of the active phases, chemical stability, high specific surface area, readily and easily available and favorable biocompatibility.^{26,29–35}




Fig. 5 Framework of boehmite (aluminum oxide hydroxide).


The high specific surface area (more than $120 \text{ m}^2 \text{ g}^{-1}$), hydroxyl groups on surface, and ease of surface modification of boehmite^{29,36} make them suitable green nano-catalyst for several organic transformation. The overwhelming performance of boehmite@nanostructure as catalyst in tetrazole synthesis motivates researchers to explore it at large extends. In this connection, a boehmite supported Pd-SMTU@boehmite nano-catalyst was prepared and examined their catalytic utility in 5-substituted-1*H*-tetrazole synthesis (Table 1, entry 1).³⁷ Optimization of reaction conditions involved a reaction between nitrile (1 mmol) and sodium azide (1.4 mmol) in the presence of Pd-SMTU@boehmite at 120°C in poly(ethylene glycol)-400 as green solvent. The optimized condition reveals that 0.3 mol% catalyst loading (Pd-SMTU@boehmite) are sufficient to achieve maximum yield. Similarly, boehmite@SiO₂@Tris-Cu(I) nano-catalyst was prepared for synthesis of 5-substituted-1*H*-tetrazole derivatives (Table 1, entry 2).³⁸ The boehmite@SiO₂@Tris-Cu(I) synthesis comprises immobilization of a Cu(I) complex on modified-surface of boehmite nanoparticles and characterized through FT-IR, XRD, EDS, ICP, X-ray mapping, TGA and SEM techniques. Further, nickel-anchored curcumin-functionalized boehmite nanoparticle (BNPs@Cur-Ni) was prepared *via* the sequential process as depicted in the Scheme 1 (Table 1, entry 3). The synthesized nano-catalyst was successfully employed in 5-substituted-1*H*-tetrazole synthesis.³⁹ The BNPs@Cur-Ni NPs exhibited excellent catalytic activity for 5-substituted-1*H*-tetrazole synthesis with magnificent turnover number (TON) and turnover frequency (TOF) results.



In another attempt, boehmite based nano catalyst was synthesized through three-step process as shown in Scheme 2. The synthesized Ni-SMTU@boehmite was effectively used for synthesis of tetrazole derivatives with good yield (Table 1, entry 4).³² However organic moiety functionalized nanostructure always demanding in organic transformation, thus a Schiff base functionalized boehmite nanostructure was synthesized and utilized for the same (Table 1, entry 5).³⁶

A palladium-arginine complex immobilized on boehmite nanoparticles (Pd-Arg@boehmite) was synthesized to serve as a green catalyst in tetrazole synthesis (Table 1, entry 6).⁴⁰ The

Table 1 Boehmite based nano-catalyzed for synthesis of 5-substituted-1*H*-tetrazole containing scaffolds *via* [3+2] cycloaddition reaction

Entry	Boehmite@nano-structures	Catalyst loading (mg or mol%)	Temperature ($^\circ \text{C}$)	Solvent	Time (h)	Yield (%)	Ref.
1	Pd-SMTU@boehmite	0.30 mol%	120	PEG-400	2.5	95	37
2	Boehmite@SiO ₂ @Tris-Cu(I)	0.20 mol%	120	PEG-400	2	97	38
3	Boehmite Nps@Cur-Ni	40 mg	120	PEG-400	1.25	97	39
4	Ni-SMTU@boehmite	25 mg	120	PEG-400	1.33	94	32
5	Pd-isatin@boehmite	35 mg	120	PEG-400	7	96	36
6	Pd-Arg@boehmite	25 mg	120	PEG-400	7	97	40



Scheme 1 Synthesis of BNPs@Cur-Ni nano-catalyst.³⁹Scheme 2 Synthesis of Ni-SMTU@boehmite nanoparticles.³²

reported boehmite supported catalytic systems offers several advantages like high substrate competency, use of environmental-friendly and commercially accessible starting materials, operational simplicity, shorter reaction time, ease of catalyst separation and reusability over several times without leaching of the braced metal ions (Cu, Pd, Ni) and high yield.

3. Magnetic nano-catalyzed synthesis of tetrazoles

Magnetic nano particles have received tremendous attention in the last few decades as an excellent catalyst or supporting materials for the organic transformation. Magnetic nano-materials have number of remarkable features like easy preparation, reusability, high catalytic efficiency, and handy separation from reaction mixtures through an external magnet which makes them ideal and efficient nano-catalyst.^{37–39} For example, Zarghania and Akhlaghinia have explored magnetite-chitin (Fe_3O_4 @chitin) as an eco-friendly magnetic nano-catalyst for the synthesis of 5-substituted-1*H*-tetrazoles under

solvent-free conditions (Table 2, entry 1).⁴¹ The magnetite-chitin was prepared *via* a hydrothermal synthesis and characterized with XRD, TEM, SEM, FT-IR, TGA and VSM analytical techniques. It was observed that Fe_3O_4 @chitin catalyst deals with the synthesis of a different tetrazoles from reaction of 1-butyl-3-methylimidazolium azide ([bmim][N₃]) with nitriles under solvent-free condition. The excellent yield, shorter reaction times (15–120 minutes), easy preparation of the catalyst, abating chemical waste and good reusability without any significant loss of catalytic efficiency are some of the advantages of the present protocol.

In another interesting report, magnetic Fe_3O_4 @tryptophan-La catalyst was prepared *via* anchoring lanthanum on the surface of Fe_3O_4 nanoparticles and characterized through FT-IR, SEM, TGA, XRD, EDX, and ICP techniques. The catalytic behavior of Fe_3O_4 @tryptophan-La was investigated as recyclable catalytic system for synthesis of 5-substituted tetrazoles using various substituted aromatic nitrile and NaN_3 in neat reaction condition (Table 2, entry 2). Along with substrate generality, La leaching was studied by La loading amount before



and after recycling of catalyst by ICP-OEIS technique which shows a very negligible leaching of La metal from catalyst surface. This catalyst offers excellent product yield, high reusability cycles, negligible leaching of La and easy separation of catalyst *via* an external magnetic field.⁴² A similar catalyst ($\text{Fe}_3\text{O}_4@\text{Nd}$) was synthesized by T. Tamoradi *et al.* for purpose of synthesis of 5-substituted-1*H*-tetrazoles (Table 2, entry 3). The catalyst was synthesized through surface functionalization of Fe_3O_4 nanoparticles with tryptophan ligand followed by Nd immobilization. The prepared $\text{Fe}_3\text{O}_4@\text{Nd}$ nano-catalyst exhibits remarkable catalytic activity toward the synthesis of aromatic tetrazoles with excellent yield.⁴³

Further, in another report, Fe_3O_4 functionalized with zinc(II)-adenine complex was successfully synthesized and employed for synthesis of 5-substituted tetrazoles (Table 2, entry 4). The catalytic activity of Fe_3O_4 -adenine-Zn was investigated with model reactants 4-chlorobenzonitrile and sodium azide in PEG at 120 °C. It was observed that Fe_3O_4 -adenine-Zn nano-catalyst offers 96% yield within 80 minutes. Additionally, nano-catalyst Fe_3O_4 -adenine-Zn gave excellent product yield with variety of aromatic and aliphatic nitriles.⁴⁴


A magnetic nano-catalyst, $\text{Fe}_3\text{O}_4@\text{MCM-41}@\text{Cu-P2C}$ was developed by Nikoorazm and Erfani and examined its catalyst efficiency for synthesis of 5-substituted-1*H*-tetrazoles derivatives using wide variety of nitriles (Table 2, entry 5). The $\text{Fe}_3\text{O}_4@\text{MCM-41}@\text{Cu-P2C}$ nano-catalyst offers many advantages like simple handling, excellent yield, easy recovery by an external magnet and reusability up to five times without any significant loss of its catalytic activity.⁴⁵ Similarly, $\text{Fe}_3\text{O}_4@L$ -aspartic-Gd nano-catalyst was prepared and characterized. The $\text{Fe}_3\text{O}_4@L$ -aspartic-Gd nano-catalyst was utilized for synthesis of

5-substituted-1*H*-tetrazoles (Table 2, entry 6). The reaction conditions were optimized with model reaction of sodium azide (1.2 mmol) and 4-chlorobenzonitrile (1 mmol). The solvent and temperature influence studies revealed that 100 °C with PEG solvent provided maximum yield of product. Further, Gd leaching studies was also examined by ICP-OEIS technique which indicated that traces of Gd was leached from the initial load to the next run. The beneficial features associated with the protocol were easy separation of catalyst, high catalytic activity, chemical and physical stability, easy workups, short reaction-time and good yields.⁴⁶

An efficient $\text{Fe}_3\text{O}_4@$ tryptophan@Ni (Table 2, entry 7) nano-catalyst was prepared by immobilization of Ni metal on tryptophan coated Fe_3O_4 nanoparticles as depicted in Scheme 3. The prepared $\text{Fe}_3\text{O}_4@$ tryptophan@Ni was characterized using various spectroscopic techniques. The synthesized $\text{Fe}_3\text{O}_4@$ tryptophan@Ni nano-catalyst was successfully furnished desired tetrazole derivatives in excellent yield and shorter reaction time (20 minutes). The reusability and metal leaching studies revealed that catalyst can be used up to seven cycles without significant leaching of Ni metal. The easy recovery, reusability and thermal stability are features advantages of $\text{Fe}_3\text{O}_4@$ tryptophan@Ni nano-catalyzed synthesis of tetrazole derivatives.⁴⁷


In another attempt, $\text{Fe}_3\text{O}_4@$ SiO₂-APTES-TFA nano-catalyst was developed by Fatahi *et al.* for synthesis of 5-substituted-1*H*-tetrazoles from various nitriles and sodium azide in EtOH at 80 °C (Table 2, entry 8). The synthesized nano-catalyst produced target tetrazoles in excellent yields in short reaction time of 4 h. Along with this, nano catalyst was easily recovered from external magnet and reused several times without significant loss of

Table 2 Magnetic nano-catalyst catalyzed protocol for the synthesis of 5-substituted tetrazole containing scaffolds *via* [3+2] cycloaddition reaction



Entry	Magnetic@nanocatalyst	Solvent	Catalyst loading mg or mol%	Temperature (°C)	Time (h)	Yield (%)	Ref.
1	$\text{Fe}_3\text{O}_4@$ chitin	—	0.03 g	110	0.33	95	41
2	$\text{Fe}_3\text{O}_4@$ tryptophan-La	H_2O	8 mg	80	0.1	96	42
3	$\text{Fe}_3\text{O}_4@$ Nd	H_2O	4 mg	80	0.16	95	43
4	Fe_3O_4 -adenine-Zn	PEG-400	50 mg	120	1.33	96	44
5	$\text{Fe}_3\text{O}_4@\text{MCM-41}@\text{Cu-P2C}$	PEG	20 mg	120	3	96	45
6	$\text{Fe}_3\text{O}_4@L$ -aspartic-Gd	PEG	70 mg	100	2.08	93	46
7	$\text{Fe}_3\text{O}_4@$ tryptophan@Ni	PEG-400	0.08 g	120	0.33	97	47
8	$\text{Fe}_3\text{O}_4@$ SiO ₂ -APTES-TFA	Ethanol	0.1 g	80	4	97	48
9	$\text{Fe}_3\text{O}_4@L$ -lysine-Pd(0)	H_2O	0.30%	100	1	99	49
10	$\text{Fe}_3\text{O}_4/\text{HT-GLYMO-TA(IV)}$	H_2O	0.005 g	70	0.33	95	50
11	Methionine@ Fe_3O_4	DMSO	0.05 g	120	0.166	100	51
12	Co $\text{Fe}_2\text{O}_4@$ glycine-Yb	PEG	0.07 g	120	2.33	93	52
13	Chitosan supported magnetic ionic liquid nanoparticles (CSMIL)	—	2.5 mol%	70	6	92	54



Scheme 3 Preparation of Fe₃O₄@tryptophan@Ni nanoparticles.⁴⁷

catalytic activity.⁴⁸ Similarly, structurally diverse wide library of 5-substituted-1*H*-tetrazoles drug-like molecules was synthesized by Ashraf *et al.* by introducing new magnetic nano-catalyst Fe₃O₄@L-lysine-Pd(0) catalyzed protocol (Table 2, entry 9). The Fe₃O₄@L-lysine-Pd(0) catalyzed green sustainable protocol involved [3+2] cycloaddition reactions of various aryl nitriles (1.0 mmol) possessing either electron-withdrawing or electron-donating substituents with sodium azide (1.3 mmol) in water. It was found from optimization study that 0.30 mol% amount of catalyst is sufficient to yield maximum yield of desired tetrazole derivatives. Furthermore, it was noticed that nano catalyst could be reused up to eight cycles without losing appreciable activity and leaching of Pd in solution.⁴⁹ In another study, Ghasemzadeh and Akhlaghinia was prepared Fe₃O₄/HT-GLYMO-TA(IV) as magnetic nano-catalyst for synthesis of 1- and 5-substituted tetrazole synthesis (Tables 2 and 3, entries 10 & 8). The catalytic activity of Fe₃O₄/HT-GLYMO-TA(IV) was studied for both 1-substituted-tetrazole and 5-substituted-tetrazoles synthesis under similar reaction condition. It was observed that 1-substituted-1*H*-tetrazoles were synthesized using triethyl orthoformate, amines, and [bmim]N₃ whereas 5-substituted-1*H*-tetrazoles, using various nitriles and [bmim][N₃]. Both electron-donating and electron-withdrawing groups containing aromatic nitriles were successfully gave target product in high to excellent yields within 20 minutes. However, main advantages reported for nano catalyst are high yield, easy work-up procedure and recyclable up to eight cycle without losing significant activity.⁵⁰

Methionine-coated Fe₃O₄ nanoparticles was successfully employed in 5-substituted-1*H*-tetrazoles synthesis by Karimian *et al.* (Table 2, entry 11). Moreover, they found that synthesized catalyst was easily separate from reaction mixture and re-claimed in many cycles without loss of activity. This nano

catalyst also offers several other benefits such as cost-effective, quick reaction, low loading of catalyst, clean process, high yields, and simple operation.⁵¹ Similarly, CoFe₂O₄@glycine-M (Pr, Tb and Yb) (Table 2, entry 12) have been prepared and utilized in the 5-substituted tetrazole synthesis.⁵²

It has been found that synthesis of various 5-substituted tetrazole derivatives have been widely documents, their 1-substituted analogs also have been subject of investigation in view their useful medical and biological applications. Most reported synthetic protocols for 1-substituted analogs involve cyclization among triethyl orthoformate, primary amines and sodium azide by using suitable catalyst. In this context and to follow advancement in catalyst, Naeimi and Mohamadabadi carried out functionalization at Fe₃O₄ as core structure to form sulfonic acid-functionalized silica-coated magnetic nanoparticles. The catalyst was characterized by FTIR, SEM, TGA, XRD, and VSM. The catalytic activity of Fe₃O₄@silica sulfonic acid was investigated in model reaction of triethyl orthoformate, sodium azide and 4-chloroaniline at 100 °C (Table 3, entry 1). The Fe₃O₄@silica sulfonic acid offers excellent yield in a shorter time (50 minutes) and efficiently apply for a range of substrates. It was observed that anilines containing electron-withdrawing and electron-donating groups gave complete reaction in lesser reaction times in comparison to unsubstituted amines. The *para*-substituted anilines produce better results compared to *ortho*-substituted aniline due to steric hindrance at *ortho* position.⁵³ In an alternative method, chitosan biopolymer supported magnetic ionic liquid nanoparticles (CSMIL) were developed for both 1- and 5-substituted tetrazole synthesis as novel heterogeneous catalyst (Tables 2 and 3, entries 13 & 2). Authors explored this new environmental-friendly nano-catalyst in tetrazole synthesis under solvent-free condition at 70 °C. Furthermore, they compared catalytic




Table 3 Magnetic nano-catalyst catalyzed synthesis of 1-substituted tetrazole containing scaffolds *via* one pot MCR between amine with triethyl orthoformate and azides

Entry	Catalyst	Solvent	Catalyst loading	Temperature (°C)	Time (h)	Yield (%)	Ref.
1	Fe ₃ O ₄ @silica sulfonic acid	—	0.02 g	100	50	97	53
2	Chitosan supported magnetic ionic liquid nanoparticles (CSMIL)	—	2.5 mol%	70	1	92	54
3	Fe ₃ O ₄ @Quinol@Cu	—	0.01 g	100	40	92	55
4	Fe ₃ O ₄ @Phen@Mn	—	50 mg	100	40	92	56
5	Fe ₃ O ₄ @WO ₃ -EAE-SO ₃ H(III)	H ₂ O	0.01 g	60	30	95	57
6	Fe ₃ O ₄ /HT-NH ₂ -Cu ^{II}	H ₂ O	0.005 g	90	1	94	58
7	Fe ₃ O ₄ @HT@AEPH ₂ -Co ^{II}	H ₂ O	0.005 g	90	60	95	59
8	Fe ₃ O ₄ /HT-GLYMO-TA	H ₂ O	0.02 g	90	45	95	50

activity of this catalyst with other reported catalyst like In(OTf)₃, SSA, [HBIm]BF₄, and natrolite zeolite. In the comparison studies, they found that 2.5 mol% of this nano-catalyst is sufficient to give maximum yield in shorter time compared to other reported catalyst. Moreover, the reported synthetic protocol offered other several advantages such as operation


easiness, high competence, low cost, easily recyclable and competency for large-scale synthesis.⁵⁴

To carry out nano-catalyst development for 1-substituted-tetrazole synthesis, Habibi *et al.* have prepared Fe₃O₄@5,10-dihydropyrido[2,3-*b*]quinoxaline-7,8-diol copper complex as a heterogeneous magnetic nano-catalyst (Table 3, entry 3). The



Scheme 4 Synthesis of Fe₃O₄@WO₃-EAE-SO₃H(III) NPs.⁵⁷



Scheme 5 Synthesis of the catalyst $\text{Fe}_3\text{O}_4/\text{HT-NH}_2\text{-Cu}^{\text{II}}$ NPs.⁵⁸

catalyst successfully furnished tetrazole formation (84–92%) in one-pot reaction from various substituted anilines with triethyl orthoformate and electron-donating or electron-withdrawing groups with sodium azide. Further, ICP analysis revealed that amount of Cu leaching was about 0.25 after first run and 2.5% after five consecutive runs while no leaching was observed for iron from nanocatalyst.⁵⁵ Similarly, Habibi *et al.* also developed $\text{Fe}_3\text{O}_4@1,10\text{-phenanthroline-5,6\text{-diol}@Mn}$ nano-catalyst for 1-substituted-tetrazole synthesis (Table 3, entry 4).⁵⁶

A functionalized magnetic $\text{Fe}_3\text{O}_4@WO_3\text{-EAE-SO}_3\text{H}$ nano-catalyst was prepared by Ghasemzadeh and Akhlaghinia as depicted in the Scheme 4. This nano-catalyst serves as a significant replacement of Brønsted acids and used in rapid preparation of 1-substituted-1*H*-tetrazoles *via* cyclization reaction between 1-butyl-3-methylimidazolium azide ($[\text{bmim}][\text{N}_3]$), triethyl orthoformate and primary amines (Table 3, entry 5). This protocol offered characteristics advantages including operational simplicity, excellent yield, high purity, gracefully separation of the catalyst *via* an external magnet and its potential reusability.⁵⁷


A bifunctional magnetite nano-catalyst ($\text{Fe}_3\text{O}_4/\text{HT-NH}_2\text{-Cu}^{\text{II}}$) was prepared and characterized by Salimi and Zamanpour research group and examined its catalytic efficiency for synthesis of 1-substituted-1*H*-tetrazoles derivatives (Table 3, entry 6). The catalyst was prepared *via* five consecutive steps as depicted in the Scheme 5. The catalytic activity of $\text{Fe}_3\text{O}_4/\text{HT-NH}_2\text{-Cu}^{\text{II}}$ was monitored through reaction of different aromatic amine, (1 mmol) triethyl orthoformate, (1.4 mmol) and sodium azide (1.1 mmol) 90 °C in H_2O solvent. Authors also extend

range of diverse substrates like arylamines with electron-releasing or electron-withdrawing group that gracefully succeeded 1-substituted-1*H*-tetrazoles.⁵⁸ Similarly, Salimi's research group was prepared $\text{Fe}_3\text{O}_4@HT@AEP\text{H}_2\text{-Co}^{\text{II}}$ nano-catalyst for 1-substituted-tetrazole synthesis (Table 3, entry 7). This catalyst also exhibits catalytic competence as previous one in preparation of 1-substituted-1*H*-tetrazoles from triethyl orthoformate (1.2 mmol), sodium azide, (1 mmol) and amine (1.0 mmol).⁵⁹

To utilize non-toxic and cost-effective starting reactants for synthesis of 5-substituted tetrazoles, one-pot multicomponent reaction approach using easily available, lower toxic and diverse aldehydes with active methylene compound and azides were explored in presence of suitable catalyst systems. For example, fibroin based magnetic nano-catalyst ($\text{Fe}_3\text{O}_4@$ fibroin) was reported by Parouch's research group for synthesis of tetrazole derivatives (Table 4, entry 1). The $\text{Fe}_3\text{O}_4@$ fibroin- SO_3H was synthesized by coating of Fe_3O_4 NPs with fibroin followed by sulfonation. The catalytic activity of $\text{Fe}_3\text{O}_4@$ fibroin- SO_3H was investigated in a reaction between malononitrile (1.00 mmol) sodium azide (2.00 mmol) and aryl aldehyde (1.00 mmol) to furnish tetrazoles derivatives. It was observed that 10 mol% amount of catalyst is sufficient to yield 86% of tetrazole product.⁶⁰ Further, Khodamorady *et al.* have prepared magnetic boehmite based nano-catalyst immobilized with palladium chitosan complex ($\text{Fe}_3\text{O}_4@$ BNPs-CPTMS-chitosan-Pd(0)) (Table 4, entry 2) for development of green synthetic protocol for synthesis of 5-substituted tetrazole derivatives by reaction between from aldehyde, malononitrile and sodium azide.



Table 4 Magnetic nano-catalyst catalyzed protocol for the synthesis of 5-substituted tetrazole containing scaffolds *via* one-pot MCR of aldehydes with malononitrile and azides



Entry	Catalyst	Solvent	Catalyst loading	Temperature (°C)	Time (h)	Yield (%)	Ref.
1	Fe ₃ O ₄ @fibroin-SO ₃ H	—	10 mol%	100	2	86	60
2	Fe ₃ O ₄ @BNPs-CPTMS-chitosan Pd(0)	—	2.8 mol%	70	50	96	65
3	Fe ₃ O ₄ -CNT-SO ₃ H	—	0.02 g	80	2.5	90	66
4	Nano-Fe ₃ O ₄	—	20 mol%	80	4.5	90	67




Fig. 6 (a) FESEM of the Fe₃O₄@BNPs and (b) FESEM of the Fe₃O₄@BNPs-CPTMS-chitosan-Pd(0). Adapted with consent from ref. 65. Copyright John Wiley and Sons.

However, literature reports stated that crystallinity of Fe₃O₄ NPs was altered during functionalization with various coating/immobilization.^{61–64} In the current instance, crystallinity of Fe₃O₄@BNPs was not altered with binding of CPTMS linker and chitosan–palladium complex on the surface of catalyst which was confirmed by SEM and XRD results as shown in Fig. 6 and 7 respectively. The Fe₃O₄@BNPs-CPTMS-chitosan-Pd(0) nano-catalyst was initially catalyzed for chemo-selective oxidation of alcohol in presence of H₂O₂ in ethanol to furnish corresponding aldehydes. Then resulting aldehydes were utilized in the homo-selective synthesis of tetrazoles. The reaction condition was optimized from mixture of malononitrile (1 mmol), benzaldehyde (1 mmol), sodium azide (1.5 mmol) using different amounts of nano-catalyst in solvent-free conditions at 70 °C which showed the maximum yield (96%) of tetrazole derivatives at 2.8 mol% of catalyst. The Fe₃O₄@BNPs-CPTMS-chitosan-Pd(0) offered numerous advantages such as green ligand (Chitosan), thermal and mechanical stability, high-grade magnetic strength, high surface area, reusability (5 cycles), an easy and fast separation.⁶⁵

Akbarzadeh *et al.* embedded chlorosulfonic acid functionalized carbon nanotubes on Fe₃O₄ nanoparticles to produce

Fe₃O₄-CNT-SO₃H nano catalyst as depicted in Scheme 6. The Fe₃O₄-CNT-SO₃H nano catalyst was employed for 2-(1-*H*-tetrazole-5-yl) acrylonitrile synthesis *via* multicomponent domino Knoevenagel condensation/1,3-dipolar cycloaddition





Fig. 7 (a) XRD of Fe₃O₄@BNPs and (b) XRD of Fe₃O₄@BNPs-CPTMS-chitosan-Pd(0). Adapted with consent from ref. 65. Copyright John Wiley and Sons.





Scheme 6 Preparation of the $\text{Fe}_3\text{O}_4\text{-CNT-SO}_3\text{H}$ nano-catalyst.⁶⁶

reaction among aromatic aldehydes, (1.0 mmol) malononitrile, (1.0 mmol) and sodium azide (1.5 mmol) under solvent-free conditions (Table 4, entry 3). Further authors compared catalytic performance of $\text{Fe}_3\text{O}_4\text{-CNT-SO}_3\text{H}$ with previously reported heterogeneous catalysts including, nano ZrP_2O_7 , OPNSA, Nano NiO, Fe_3O_4 . They revealed that this catalytic system produced high product yield within a shorter reaction time and eliminates toxic and volatile solvents. Further tempting to green catalyst, recyclability of $\text{Fe}_3\text{O}_4\text{-CNT-SO}_3\text{H}$ is much easier compared to other stated.⁶⁶


Similarly, it was found that, unmodified nano Fe_3O_4 exhibited remarkable catalytic activity for 5-substituted tetrazole synthesis from malononitrile (1 mmol), aromatic aldehyde (1 mmol), sodium azide (1 mmol) as tabulated in Table 4, entry 4. They observed that nano Fe_3O_4 gave best result at 20 mol%

amount. Prominent features of the presented protocol comprise of reactions under microwave condition, short reaction time, exclusion of toxic solvents, reasonable and reclaimable catalyst, mild reaction condition and easy workup.⁶⁷

4. Copper-based nanomaterial catalyzed synthesis of tetrazoles

Development of copper-based nanomaterials received huge attention in recent years specially in domain of nano-catalysis organic transformations due to its extensive range of oxidation states which can enables reactivity through both one- and two-electron pathways.⁶⁶ Also, copper-based nano-materials found applications in various other areas like nanotechnology, electrocatalysis, and photocatalysis.⁶⁸⁻⁷⁰ In this context,

Table 5 Copper-based nano-catalyst catalyzed protocol for the synthesis of 5-substituted tetrazole containing scaffolds via [3+2] cycloaddition reaction



Entry	Catalyst	Solvent	Catalyst loading	Temperature (°C)	Time (h)	Yield (%)	Ref.
1	$\text{Cu(II)/Fe}_3\text{O}_4\text{@APTMS-DFXn}$	DMSO	2.5 mol%	120	1	98	71
2	$\text{Cu-MOFs-1 (Cu-TMU-17-NH}_2\text{)}$	PEG	0.05 g	110	5	94	72
3	$\text{Cu-MOFs-2 (Cu-TMU-17-NH}_2\text{)}$	PEG	0.05 g	110	5	95	72
4	$\text{CuO/aluminosilicate}$	DMF	35 mg	110	2–13	80–93	73
5	Cu-TBA@biochar	PEG	0.78 mol%	130	7	98	74
6	SBA-15@glycine-Ni	PEG	0.04 g	100	110	98	78
7	SBA-15@glycine-Cu	PEG	0.04 g	100	110	98	78
8	$\text{SBA-15/thioamide-Cu(I)}$	H_2O	1 mol%	80	240	97	79
9	Cu(II)-DCC-CMK-3	PEG	20 mg	120	90	97	80
10	Cu(II) immobilized on $\text{Fe}_3\text{O}_4\text{@SiO}_2\text{@L-arginine}$	PEG	1.0 mol%	120	3	95	81
11	$\text{Fe}_3\text{O}_4\text{@SiO}_2\text{-DAQ-Cu(II)}$	DMF	0.9 mol%	110	2.5	95	82
12	$\text{Fe}_3\text{O}_4\text{@SiO}_2\text{/ligand/Cu(II)}$	DMF	0.4 mol%	110	4	92	83
13	$\text{Fe}_3\text{O}_4\text{@SiO}_2\text{/salen Cu(II)}$	DMF	0.4 mol%	120	6	92	84
14	$\text{Fe}_3\text{O}_4\text{-AMPD-Cu}$						85
15	Cu(II) immobilized on aminated epichlorohydrin activated silica (CAES)	DMSO	1.0 mol%	130	1	95	86
16	Cu(II) immobilized on $\text{Fe}_3\text{O}_4\text{@SiO}_2\text{@L-histidine}$	PEG	0.05 g	120	60	95	87







Fig. 8 SEM image of Cu-TBA@biochar nanoparticles. Adapted with consent from ref. 74. Copyright Elsevier.

Cu-immobilized heterogenous nano-catalysts have been extensively utilized in synthesis of tetrazoles derivatives (Tables 5–8). For example, Cu(II) immobilized magnetic Cu(II)/Fe₃O₄@-APTMS-DFX nano-catalyst was developed and examined for [3+2] cycloaddition reactions of sodium azide and various organic nitriles to provided biologically active 5-substituted-1*H*-tetrazoles (Table 5, entry 1). It was found that 2.5 mol% of nano-catalyst produced a series of 5-substituted-1*H*-tetrazoles in DMSO at 120 °C. This nano-catalyzed protocol offered

advantage of recyclability up to five runs with insignificant leaching of copper from catalyst.⁷¹


Two analogues of Cu-TMU-17-NH₂'s framework entitled Cu-MOF-1 and Cu-MOF-2 were prepared by Salahshournia *et al.* as efficient and recoverable heterogeneous nano-catalyst for the 5-substituted-1*H*-tetrazoles synthesis (Table 5, entries 2–3). The Cu-MOF-1 and Cu-MOF-2 were obtained *via* post-synthetic metalation of a Zn-MOF (TMU-17-NH₂) with Cu(OAc)₂ and CuCl₂·2H₂O respectively as copper precursors. The synthesized catalysts offer excellent yield of 5-substituted-1*H*-tetrazoles (more than 94%) in PEG as a green solvent probably due to the porous nature of nano Cu-MOFs. Further, it also provide several advantages such as feasible synthesis, smooth recovery, and reusability (five cycles) without dropping its catalytic activity.⁷² In this line, CuO/aluminosilicate, as an effective heterogeneous nano-catalyst was developed by Movaheditabar *et al.* for preparation of 5-substituted-1*H*-tetrazoles (Table 5, entry 4). The nano-catalyst affords good to excellent yield of tetrazoles from a series of aliphatic and aromatic nitriles.⁷³

In the progress for sustainable nano-catalyst development, Moradi *et al.* have explored the use of biochar as a solid support material for preparation of functionalized biochar-based nano-catalyst (Table 5, entry 5). Biochar is a carbon-rich solid having carbonyl, carboxylic acid, and hydroxyl functional groups on its surfaces. In the process of preparation of nano catalyst, initially biochar nanoparticles were synthesized *via* pyrolysis of chicken manure and then functionalized with 3-chloropropyltrimethoxy silane to obtain CPTMS@biochar. Subsequently, 2-(thiophene-2-yl)-1*H*-benzo[*d*]imidazole was embedded on CPTMS@biochar to form TBA@biochar for Cu immobilization. Finally, copper



Scheme 7 Preparation of the SBA-15/thioamide-Cu(I) NPs.⁷⁹



Scheme 8 Preparation of Cu(II)-DCC-CMK-3 nano-catalyst.⁸⁰

was immobilized on TBA@biochar. The prepared Cu-TBA@biochar characterized by TGA, XRD, SEM, EDS, N₂ adsorption-desorption isotherms, AAS and FT-IR techniques which indicated that particle formed in less than 100 nm range (Fig. 8). It was observed that 0.78 mol% of Cu-TBA@biochar was sufficient to give excellent yield (98%) of tetrazoles derivatives with reusability of several runs without loss of copper leaching and catalytic activity.⁷⁴

In view of stringent demand of green nano-catalyst for heterocyclic transformation, different solid supports like zeolites, silica, carbon nanotubes, organic polymers, alumina and other metal oxides have been used for development of nano-catalyst.⁷⁵⁻⁷⁷ Among different solid supports, mesoporous silica nanoparticles especially SBA-15 have extensively used in different catalytic systems due to their hexagonally packed arrays and hierarchical structure.²³ Tamoradi *et al.* utilized SBA-15 as catalyst support for preparation of two heterogeneous catalysts SBA-15@glycine-Ni and SBA-15@glycine-Cu (Table 5, entries 6-7) by immobilization of Ni and Cu metal on ordered SBA-15 coated with glycine. The structure of SBA-15@glycine-Ni and SBA-15@glycine-Cu was analyzed by FT-IR, TGA, BET, ICP-OES, XRD and EDX analysis and successfully employed for synthesis of 5-substituted tetrazoles derivatives. The high yield (83-98%) of desired tetrazole derivatives revealed high catalytic activity of prepared catalysts. It was observed that improved Lewis's acid characteristic of SBA-15@glycine-Cu is mainly responsible for its high catalytic performance than SBA-15@glycine-Ni. Further, the present catalysts offers some exclusive advantages such as high catalytic nature, use of a green solvent (ethanol and PEG) and metals (Ni & Cu) and ligand (glycine).⁷⁸


Similarly, an SBA-15 modified thioamide-Cu(I) (Table 5, entry 8) catalyst was developed by Pourhassan and Eshghi as illustrated in Scheme 7. In synthesis of SBA-15/thioamide-Cu(I), SBA-15 channels was modified with tris(2-aminoethyl)amine (TAEA) groups and then reacted with S₈ and phenylacetylene to form thioamide groups. This synthesized porous material was found

to be an active host for immobilization of economical Cu(II) ions. The 1.0 mol% amount of catalyst gave 95% yield of tetrazoles derivative at room temperature. Further, practical application of catalyst was examined by recovery and reusability of catalyst, and it was observed that catalyst can be reused consecutively up to eleven runs in a model reaction without losing its catalytic efficiency. The copper ion leaching was monitored by ICP-OES analysis. It was reported that after eleven successful runs of tetrazole synthesis only 0.6 wt% of Cu species leached in solution.⁷⁹

Another nano-catalyst Cu(II)-DCC-CMK-3 was synthesized using SBA-15 as a solid template. The synthesis of Cu(II)-DCC-CMK-3 was achieved *via* multi-step process as depicted in Scheme 8. In the first step mesoporous carbon CMK-3 was synthesized from SBA-15 silica which was used for preparation of Cu(II)-DCC-CMK-3 catalyst. The catalyst was analyzed by FT-IR, TGA, BET, XRD, SEM, ICP-OES and EDX techniques. The SEM image of the nano-catalyst indicated that CMK-3 carbon particles maintain rod-like structures, whereas, TEM image of Cu(II)-DCC-CMK-3 reveals well-ordered mesostructured with hexagonal symmetry. The synthesized catalyst was successfully employed for synthesis of tetrazole derivatives (Table 5, entry 9).⁸⁰

A non-corrosive Cu(II) immobilized Fe₃O₄@SiO₂@L-arginine, nano catalyst was developed for synthesis of 5-substituted-1*H*-tetrazoles *via* cycloaddition reaction of various organic nitriles with sodium azide (Table 5, entry 10). The optimization results with aryl nitrile (1 mmol), NaN₃ (1 mmol) in the presence of Cu(II) immobilized Fe₃O₄@SiO₂@L-arginine catalyst revealed that 1.020 mol% of catalyst gave 95% yield at 120 °C in PEG solvent. Furthermore, high competence, ease of operation, easy recovery, low cost and reusability are some significant beneficial features of this catalytic system.⁸¹ In an another report, 1,4-dihydroxyanthraquinone-copper(II) was immobilized on superparamagnetic Fe₃O₄@SiO₂ NPs to developed Fe₃O₄@SiO₂-DAQ-Cu(II) nano-catalyst for synthesis of 1- and 5-substituted-1*H*-tetrazoles as illustrated in Scheme 9 (Tables 5 and 6, entry 11 & 4). It was reported that 0.9 mol% of this catalyst was sufficient





Scheme 9 Preparation of the $\text{Fe}_3\text{O}_4@ \text{SiO}_2\text{-DAQ-Cu(II)}$ nano-catalyst.⁸²

for 5-substituted-1-*H*-tetrazole and 0.8 mol% for 1-substituted-1-*H*-tetrazoles synthesis. It was observed that different aromatic nitriles containing electron-donating and electron-withdrawing groups were successfully employed for tetrazole synthesis with excellent product yields and the substituents on benzonitriles play significant impact on the reaction time. For example, model reaction with electron-poor aromatic and heteroaromatic nitriles completed within a few hours whereas nitriles having electron-donating substituents took longer reaction time. The $\text{Fe}_3\text{O}_4@ \text{SiO}_2\text{-DAQ-Cu(II)}$ catalyzed synthetic protocol serves many advantages like operational simplicity, reusability up to six runs and negligible copper leaching and excellent product yield.⁸²

Similarly, Esmaeilpour *et al.* prepared $\text{Fe}_3\text{O}_4@ \text{SiO}_2/\text{ligand}/\text{Cu(II)}$ magnetic nano-catalyst for 1- and 5-substituted-1-*H*-tetrazoles synthesis such as illustrated in Scheme 10 (Tables 5 and

6, entry 12 & 3). It was observed that 0.4 mol% of the nano catalyst are sufficient to yield more than 90% of the targeted product. Furthermore, super magnetic nature of the catalyst eliminates catalyst filtration after completion of reaction which represents its main advantage in the expression of energy-consuming and environmental-friendly.⁸³

In another method from Dehghani *et al.*, 1- and 5-substituted-1-*H*-tetrazoles were synthesized from nitriles and amines using, $\text{Fe}_3\text{O}_4@ \text{SiO}_2/\text{salen}$ of Cu(II) based magnetite nanoparticles was developed by (Tables 5 and 6, entry 13 & 2). The structural and magnetic properties of $\text{Fe}_3\text{O}_4@ \text{SiO}_2/\text{Salen}$ of Cu(II) was elucidated by TEM, SEM, FTIR, and VSM analysis. The reported catalyst successfully furnished synthesis of various 1- and 5-substituted tetrazole derivative with wide substrate scope in good to excellent. Furthermore, it was observed that 0.4 mol% of catalyst is best for both 1- and 5-substituted



Table 6 Copper-based nano-catalyst catalyzed protocol for the synthesis of 1-substituted tetrazole containing scaffolds *via* one-pot MCR of triethyl orthoformate, amines and azide


$$\text{R-NH}_2 + \text{EtO}-\underset{\text{OEt}}{\overset{\text{H}}{\text{C}}}-\text{OEt} + \text{NaN}_3 \xrightarrow[\text{Solvent, temperature}]{\text{Nanocatalyst}} \text{R-N} \begin{array}{c} \diagup \text{N} \\ \diagdown \text{N} \\ \text{N} \end{array}$$

Entry	Catalyst	Solvent	Catalyst loading	Temperature (°C)	Time (h)	Yield (%)	Ref.
1	Fe ₃ O ₄ /SiO ₂ /CPTMS/MT/Cu	—	50 mg	40	2	94	88
2	Fe ₃ O ₄ @SiO ₂ /salen Cu(II)	—	0.4 mol%	100	75	92	84
3	Fe ₃ O ₄ @SiO ₂ /ligand/Cu(II)	—	0.4 mol%	100	0.75	93	83
4	Fe ₃ O ₄ @SiO ₂ -DAQ-Cu(II)	—	0.8 mol%	100	1	96	82
5	Fe ₃ O ₄ /SiO ₂ /CPTMS/MT/Cu	H ₂ O	20 mg	40	2	94	89
6	Fe ₃ O ₄ /SiO ₂ /CPTMS/AT/Cu	Ethanol	20 mg	100	1	97	90

tetrazole derivatives. Authors also compared catalytic competence of reported catalyst with other previously reported catalysts such as nano natrolite zeolite, ZnHAP, ZnO/Co₃O₄ and Zn/Al-HT *etc.* which showed that Fe₃O₄@SiO₂/Salen of Cu(II) gave better product yield in shorter reaction time compare to other ones.⁸⁴


Another copper immobilized Fe₃O₄-AMPD-Cu catalyst was prepared and employed for 5-substituted tetrazole synthesis (Table 5, entry 14). The nano-catalyst offers several advantages

including environmental-benign condition, inexpensive, chemically inert reagents, shorter reaction time (28 minutes), easy separation *via* an external magnet, and reusability.⁸⁵ Analogously, Cu(II) immobilized on aminated epichlorohydrin activated silica (CAES) as efficient heterogeneous nano-catalyst was synthesized *via* five-step synthetic method for synthesis of 5-substituted-1H-tetrazoles as depicted in Scheme 11 (Table 5, entry 15). The synthesized CAES catalyst in 0.1 mol% gave excellent yield of tetrazoles *via* [3+2] cycloaddition reaction of



Scheme 10 Preparation of the Fe₃O₄@SiO₂/ligand/Cu(II) nano-catalyst.⁸⁵





Scheme 11 Preparation of Cu(II) immobilized on aminated epichlorohydrin activated silica (CAES)⁸⁶

sodium azide and nitriles with wide substrate scope.⁸⁶ Further, Azadi *et al.* reported synthesis of copper(II) immobilization on $\text{Fe}_3\text{O}_4@\text{SiO}_2$ -L-histidine as a reusable nano-catalyst for the synthesis of 5-substituted tetrazoles (entry 16, Table 5). This synthetic protocol provide excellent yield in a shorter time with efficient reusability.⁸⁷

A Cu(II) immobilized $\text{Fe}_3\text{O}_4/\text{SiO}_2/\text{CPTMS}/\text{MT}/\text{Cu}$ magnetic catalyst was prepared and characterized by distinctive methods includes TEM, SEM, XRD, ICP, EDX, VSM and TG-DTA analysis *etc.* The reported catalyst successfully utilized for synthesis of tetrazole derivatives by reaction of aromatic amines with sodium azide in presence of triethyl orthoformate in a solvent-free condition at 100 °C (Table 6, entry 1). The catalyst exhibits excellent competence in the expression of 94% yield within 2 h⁸⁸

In some report's amino acids was utilized for synthesis of tetrazole derivatives *via* one-pot MCR pathway using triethyl orthoformate, and azide. For example, Cu(II) immobilized nano catalyst ($\text{Fe}_3\text{O}_4/\text{SiO}_2/\text{CPTMS}/\text{MT}/\text{Cu}$) was prepared *via* five-step synthetic protocol. The prepared magnetic nano-catalyst was effectively utilized for synthesis of tetrazoles using several amino acids in aqueous medium at 40 °C (entry 5, Table 6).⁸⁹ Similarly, in another alternative method, $\text{Fe}_3\text{O}_4/\text{SiO}_2/\text{CPTMS}/\text{AT}/\text{Cu}$ nano catalyst was developed for tetrazole synthesis from triethyl orthoformate, sodium azide and amino acids in ethanol at reflux condition (Table 6, entry 6).⁹⁰

A copper-based magnetic heterogeneous catalyst was prepared *via* functionalization surface of nano $\text{Fe}_3\text{O}_4@\text{SiO}_2$ with polyvinyl alcohol and the resultant $\text{Fe}_3\text{O}_4@\text{SiO}_2$ -PVA was then immobilized with copper(II) complex. The nano-catalyst $\text{Fe}_3\text{O}_4@\text{SiO}_2$ -PVA/Cu(II) was well-characterized and employed for green synthesis of 5-substituted-1*H*-tetrazoles using hydroxylamine hydrochloride (1.5 mmol), aldehydes (1.0 mmol) and sodium azide (1.5 mmol) in aqueous medium (Table 7, entry 1). It was observed that 0.5 mol% of $\text{Fe}_3\text{O}_4@\text{SiO}_2$ -PVA/Cu(II) nano-catalyst was found to be enough for the maximum yield. Further, easy handling, easy separation of catalyst, reusability (seven successive runs), and yielding high to excellent yields of the anticipated products were few beneficial features of present protocol.⁹¹

5. Zinc oxide derived nanomaterials catalyzed synthesis of tetrazoles


Some inorganic metal oxide nanoparticles are well known to display both Lewis acid–base and redox properties on their surfaces. In this context, zinc oxide nanoparticles exhibits Lewis acidic sites at their surfaces and outstanding physical and chemical properties like wide bandgaps energy,^{92,93} large surface area, high pores volume, reusability and environmental sustainability⁹⁴ which was widely employed as heterogeneous

Table 7 Copper-based nano-catalyst catalyzed protocol for the synthesis of 5-substituted tetrazole containing scaffolds *via* one-pot MCR of aldehydes with hydroxylamine and azide

Entry	Catalyst	Solvent	Catalyst loading	Temperature (°C)	Time (h)	Yield (%)	Ref.
1	$\text{Fe}_3\text{O}_4@\text{SiO}_2$ -PVA/Cu(II)	H_2O	0.5 mol%	100	4	92	91




Table 8 ZnO derived nano-catalyst catalyzed for synthesis of 5-substituted tetrazole containing scaffolds via [2+3] cycloaddition



Entry	Catalyst	Solvent	Catalyst loading	Temperature (°C)	Time (h)	Yield (%)	Ref.
1	ZnO	DMF	0.1 g	120–130	14	69–82	100
2	ZnO/Co ₃ O ₄	DMF	50 mg	120	12	90	104

Table 9 ZnO derived nano-catalyst catalyzed synthesis of 1-substituted tetrazole containing scaffolds via one-pot MCR of amine, triethyl orthoformate and azide



Entry	Catalyst	Solvent	Catalyst loading	Temperature (°C)	Time (h)	Yield (%)	Ref.
1	ZnO	CH ₃ CN/H ₂ O	1 wt%	200 W	1	90	106


acid catalyst. These interesting physical properties of ZnO nanoparticles prompts to be deployed in various fields such as photodegradation of microorganisms, electrical engineering, catalysis, optical and optoelectronic devices and solar cells.^{95–98} Further, ZnO nanoparticles offers excellent catalytic activity for acid catalyzed organic transformations.⁹⁹ A nanocrystalline ZnO as a heterogeneous green catalyst was employed for preparation of 5-substituted-1*H*-tetrazoles through [3+2] cycloaddition of sodium azide and nitriles (Table 8, entry 1). Nanocrystalline ZnO effectively yielded 5-substituted-1*H*-tetrazoles from various nitriles at 120–130 °C with product yields of 69–82%. Furthermore, catalyst was graciously recovered and reused in the consecutive run.¹⁰⁰ It was established in literature that role of zinc oxide in tetrazole synthesis is to activate the –CN triple bond of nitriles.^{101–103} The acidity of zinc oxide (ZnO) could be enhanced by mixing it with other metal oxides like TiO₂, CeO₂ and Co₃O₄. A mixed metal oxide (ZnO/Co₃O₄) was employed as a pioneering catalyst for 5-substituted-1*H*-tetrazoles (Table 8, entry 2). The mixed nano metal oxide exhibits excellent catalytic activity compared to solitary ZnO or Co₃O₄. This protocol offers significant advantages comprising easy applicability, catalyst recyclability, simple work-up, and high yield.¹⁰⁴ Further, in another report, silver doped ZnO nanorods catalyzed photo-triggered synthesis of 1,5-disubstituted tetrazoles was reported. This synthetic approach involved one-pot MCR of aldehyde, chromenocyl bromide, sodium azide in presence of triethylamine using ZnO nanorods and Ag-doped ZnO nanocomposites (NCs) as photocatalysts. Optimized reaction conditions revealed that at room temperature in dark, 10 mol%

silver doped ZnO nanorods gave 97% yield of 1,5-disubstituted tetrazole derivatives.¹⁰⁵ Similarly, in another attempt, green protocol was developed for tetrazole synthesis using ZnO nanoparticles as nano catalyst under ultrasound irradiation through reaction of triethyl orthoformate, sodium azide, ethyl esters and a series of α -amino acid (Table 9, entry 1). It was observed that under irradiation at 200 W, reaction time shortened to 30 min with excellent yield (88–96%). The major advantages of this protocol are use of mild reaction conditions that eliminated formation of HN₃, no side products formation, high yields, and easy work-up procedure. The recyclability and reusability of nano-catalysts were adequate which showed cost proficiency and green aspect of the methodology.¹⁰⁶

6. Carbon-based nanomaterials catalyzed synthesis of tetrazoles


Carbon nanostructures like fullerene, graphene, graphene oxide, carbon nanotubes, carbon nanowires, carbon nanoparticles (CNPs) and nano-diamonds widely used by researchers in form of compatible catalyst support for various organic transformations due to their inevitable exclusive physical and chemical properties like high thermal, mechanical and chemical stability, large surface area, inertness, low cost, easy functionalization and tunable carbon framework.^{107,108} In line of this, a multi-walled carbon nanotube supported AMWCNTs-O-Cu(II)-PhTPY nano-catalyst was reported for facile preparation of 1-substituted and 5-substituted-1*H*-tetrazole derivatives. The AMWCNTs-O-Cu(II)-PhTPY preparation involved



Scheme 12 Schematic synthesis route for GO/Fe₃O₄@PAA-Cu-complex preparation.¹¹⁰

immobilization of copper(II) complex of 4'-phenyl-2,2':6',2''-terpyridine on activated multi-walled carbon nanotubes. The optimization of the 5-substituted-1*H*-tetrazole synthesis was carried out by taking model reactants such as nitrile, azide, ammonium acetate and DMF as solvent. Optimization studies revealed that 4.0 mol% of catalyst loading is sufficient to achieve maximum product yield at 70 °C s. The AMWCNTs-O-Cu(II)-PhTPY nano-catalyst also examined for wide substrates scope with hetero-aryl nitriles substituted with the electron-withdrawing as well as electron donating groups. It was noticed that electron-deficient nitriles furnished the target

products in shorter reaction time than the electron-rich counterparts with comparatively higher yields. The lesser reaction time, easy recovery of catalyst, wide substrate scope, catalyst reusability up to five cycles without showing copper leaching are some advantages of the reported protocol.¹⁰⁴ Similarly, a carbon nanotube supported heterogeneous Fe₃O₄-CNT-TEA-Cu(II) nano-catalyst was prepared and characterized (FTIR, XRD, FESM, TGA, and VSM) for the 5-substituted-1*H*-tetrazoles synthesis. The Fe₃O₄-CNT-TEA-Cu(II) nano-catalyst was prepared by growing metallic copper(II) nanoparticles on the surface of magnetic carbon nanotube and supported with

Fig. 9 Cu/AC/r-GO nanohybrid.¹¹¹


triethanolamine (TEA) that acts as a non-toxic ligand to capture the copper nanoparticles. The prepared $\text{Fe}_3\text{O}_4\text{-CNT-TEA-Cu(II)}$ successfully employed in synthesis of 5-substituted-1*H*-tetrazoles through one-pot MCR of aldehyde, hydroxylamine, and sodium azide. The authors revealed that protocol furnish product in high yield with low catalyst loading without leaching copper ions in the solution. Moreover, it was found that catalyst performed the recyclability up to eight cycles with same extent without losing noticeable catalytic activity. The reusability, separation with a magnet, substrate scope, high yield and lesser reaction time are some of aids with this protocol.¹⁰⁹

Among different carbon nanostructures, graphene oxide has potentials to behave as excellent catalyst support for fabrication of an ideal nano-catalyst for organic transformations because of their distinctive properties such as high thermal and chemical stability, nano-size dimension with high surface area, sheet structure, and possibility of functionalization with different organic entities *etc.* In this context, a graphene oxide (GO) supported $\text{GO/Fe}_3\text{O}_4\text{@PAA-Cu}$ -complex, nano-catalyst was prepared for 1- and 5-substituted tetrazoles synthesis by one-pot multicomponent reaction. The $\text{GO/Fe}_3\text{O}_4\text{@PAA-Cu}$ -complex nanostructure synthesis comprises multi-step protocol as depicted in Scheme 12. Graphene oxide synthesis commenced through modified Hummer's method and $\text{GO-Fe}_3\text{O}_4$ hybrid by chemical co-precipitation. The final step involved immobilization of the poly(α -amino acid)- Cu(II) complex on $\text{GO-Fe}_3\text{O}_4$ hybrid. The prepared $\text{GO/Fe}_3\text{O}_4\text{@PAA-Cu}$ -complex were characterized through XRD, EDS, FESEM, TEM, TGA, VSM and DLS analysis and utilized in tetrazole synthesis. The present method offered diversity in reaction, recyclability of catalyst over six successive turns with insignificant metal leaching.¹¹⁰

In another report, GO supported Cu/AC/r-GO nano-hybrid nano-catalyst (Fig. 9) was prepared for 5-substituted-1*H*-tetrazoles synthesis bearing bioactive *N*-heterocyclic cores. Different types of nitriles with diverse *N*-heterocyclic cores reacted with sodium azide in water/*i*-PrOH (50 : 50, v/v) system under catalytic effect of Cu/AC/r-GO nano-hybrid at reflux condition gave corresponding 5-substituted-1*H*-tetrazoles. The nano-hybrid exhibited chemical and thermal stability, inexpensive, and environmentally benign nano-catalyst that significantly endorsed cycloaddition of the azide and nitrile. Further, the advantage of recyclability and reused for many consecutive runs without significant loss of its activity makes this system more effective.¹¹¹


7. Composite nanomaterial catalyzed synthesis of tetrazoles

Composites are superintended or naturally transpiring solid materials in which two or more different constituent materials are combined to produce modified materials with superior physical and chemical properties.¹¹² In recent years, to overcome limitations associated with various engineering materials, nanocomposites has emerged as beneficial alternatives. Therefore, numerous nanocomposites were explored in different organic transformation as heterogeneous catalyst. For instance, a biosynthesized Pd/MnO_2 nanocomposite catalyst explored for the 5-aryl-1*H*-tetrazoles synthesis from various aryl halides. It was observed that this strategy suit to a range of aryl halides having electron withdrawing groups on benzene and heterocyclic ring. Further authors also clarified that MnO_2 play important role in stabilizing Pd NPs, by preventing agglomeration of



Scheme 13 Preparation of the $\text{Fe}_3\text{O}_4\text{@MCM-41-SB-Cu}$ nano-catalyst.¹¹⁴





Scheme 14 Synthesis of 5-substituted-1*H*-tetrazoles catalyzed by Fe₃O₄@MCM-41-SB-Cu nanocomposite.¹¹⁴

Pd nanoparticles and leaching during reactions. The biocatalyst (Pd/MnO₂) offers several advantages like high yields, easygoing preparation, use of K₄[Fe(CN)₆] as a non-toxic cyanide source, simple work-up procedure, escaping formation of destructive hydrazoic acid, and accessibility of MnO₂ as support.¹¹³

A magnetic Fe₃O₄@MCM-41-SB-Cu nanocomposite as a catalyst was prepared for 5-substituted-1*H*-tetrazoles synthesis. The Fe₃O₄@MCM-41-SB-Cu nanocomposite synthesis involved the several steps shown in Scheme 13. In the first step magnetic mesoporous silica nanocomposite was prepared and functionalized with *N*-(2-aminoethyl)-3-aminopropyltrimethoxysilane. Thereafter Schiff base grafted NPs were prepared by condensation reaction between 5,5'-methylene bis (salicylaldehyde) and *N*-(2-aminoethyl)-3-aminopropyltrimethoxysilane which was then reacted with benzhydrazide followed by (CH₃COOH)₂ · H₂O to give Fe₃O₄@MCM-41-SB-Cu. The new organic-inorganic hybrid nanocomposite characterized by various spectroscopic analytical techniques. The catalytic performance of Fe₃O₄@MCM-41-SB-Cu was assessed in one-pot MCR of using aldehydes, hydroxylamine hydrochloride and sodium azide for the synthesis of 5-substituted-1*H*-tetrazoles (Scheme 14.) The Fe₃O₄@MCM-41-SB-Cu nano catalyst offers easy separation due to magnetic nature and reused several times without dropping catalytic stability and activity.¹¹⁴ Similarly, a magnetic nanocomposite (Fe₃O₄@SiO₂/aza-crown ether-Cu(II)) was prepared for the synthesis of 1,2,3-triazoles, 1-substituted-1*H*-tetrazoles and 5-substituted-1*H*-tetrazoles synthesis. The fabrication of the nanocomposite catalyst could be achieved *via* covalent grafting of an aza-crown ether Cu(II) complex on a silica-coated iron oxide support. The resulted nanocomposite characterized by various techniques like TEM, SEM, XRD, TGA, FT-IR and ICP. The nanocomposite catalyzed the 1,2,3-triazoles, 5-phenyl-1*H*-tetrazoles and 1-phenyl-1*H*-tetrazoles synthesis with several advantages including thermal stability, easygoing preparation, convenient separation, short reaction time, reusability (five runs) and admirable yields.¹¹⁵

Further, Cu NPs@Fe₃O₄-chitosan nano-catalyst was prepared and characterized for different tetrazoles synthesis using sodium azide and various secondary or tertiary cyanamides. The Cu NPs@Fe₃O₄-chitosan nano-catalyst preparation involved immobilization of copper nanoparticles on the magnetic chitosan using *Euphorbia falcata* leaf extract as reducing and stabilizing agent such as depicted in the Scheme 15. The tetrazole formation was optimized using 2,4-dimethylphenylcyanamide and NaN₃ as model reactants. It was observed that Cu NPs@Fe₃O₄-chitosan display high catalytic activity for

reaction of various *N*-arylcyanamides possessing diverse functionalities with NaN₃ in aqueous medium with high yields. Further, numerous advantages also recognized with this protocol such as easy work-up procedure, minimum use of toxic hydrazoic acid, use of green solvent (H₂O), catalyst stability, reusability, and high yields.¹¹⁶

A biosynthesized Ag/sodium borosilicate nanocomposite was explored by Nasrollahzadeh *et al.* for 1-substituted-1*H*-tetrazoles synthesis. The Ag/sodium borosilicate nanocomposite (ASBN) catalyst was prepared using *Aleurites moluccana* leaf extract as stabilizing and reducing agent. The formation of the catalyst was characterized *via* various spectroscopical analysis such as FESEM, TEM, EDS, FT-IR, XRD. This synthetic protocol offered high yields, easy work-up, short reaction time, green synthesis without using harmful reducing agents. Further, reported catalyst was recycled and reused multiple times without any significant loss of activity.¹¹⁷

Another nanocomposite RuO₂/MMT was synthesized and employed in one-pot three-component (3-CR) synthesis for tetrazole using different sodium azide, amine and triethyl orthoformate under solvent-free condition. The RuO₂/MMT catalyzed synthesis of tetrazole derivatives offered excellent yield (84–97%), high reusability (five cycles), short reaction time and simple work up procedure. The enhanced catalytic activity of reported bifunctional nanocomposite is attributed by the uniform dispersion of RuO₂ NPs on surface of montmorillonite (MMT). In this protocol RuO₂ sites are accountable for the coordination of isocyanide intermediate while strong acidic character of MMT induces condensation and cyclization steps in a synergic mode.¹¹⁸

8. MCM-41 based nanomaterials catalyzed synthesis of tetrazoles

MCM-41 offered as auspicious catalyst support in liquid phase reactions due to its well-defined mesoporous structure with a high specific surface area (>1000 m² g⁻¹) and thermal stability.¹¹⁹ MCM-41 was first time reported in 1992 by scientists of Mobil Company during their project for finding new highly porous materials. Since then, mesoporous MCM-41 fascinated cumulative research attention due to their chemical versatility, large specific surface area with huge silanol groups which can be functionalized with other suitable functional groups. The 1.3 ml g⁻¹ pore volume of MCM-41 allows anchoring of diverse molecules such as metal complexes and organic ligands into highly ordered hexagonal channels.^{96,120,121} In recent years MCM-41





Scheme 15 Preparation of the Cu NPs@Fe₃O₄-chitosan nano-catalyst.¹¹⁶

Table 10 MCM-41@nano-catalyst catalyzed protocol for the synthesis of 5-substituted tetrazole containing scaffolds via [2+3] cycloaddition




Entry	MCM-41@nanostructure	Catalyst loading (mg or mol%)	Temperature (°C)	Solvent	Time (h)	Yield (%)	Ref.
1	SO ₃ H@MCM-41	50 mg	80	DMF	1.6	90	122
2	Cu-cytosine@MCM-41	3.1%	120	PEG-400	0.5	96	123
3	Ni-cytosine@MCM-41	0.1%	120	PEG-400	1	92	123
4	L-Cysteine@MCM-41	2.9%	100	PEG-400	2.5	97	124
5	Pd-SBT@MCM-41	35 mg	120	PEG-400	6.75	97	125

widely utilized as catalyst support in organic transformation like SO₃H functionalized MCM-41 nonordered heterogenous catalyst was prepared for 1- and 5-substituted-1*H*-tetrazole synthesis (Table 10, entry 1).¹²²

Cu and Ni immobilized on cytosine@MCM-41 nano catalysts (Cu-cytosine@MCM-41 & Ni-cytosine@MCM-41) were synthesized and monitored for 5-substituted-1*H*-tetrazoles synthesis (Table 10, entries 2–3) as depicted in the Scheme 16. The

synthesized organic–inorganic hybrid nano-catalysts characterized by TGA, BET, XRD, TEM, EDS, SEM, WDX, AAS and FT-IR techniques. The SEM images of synthesized nano catalyst exhibited that nano catalysts are nearly spherical in shape whereas TEM image exemplified ordered hexagonal mesoporous structure less than 5 nm and uniform quasi-spherical morphology (Fig. 10 and 11). The reaction condition was optimized by varying catalyst loading, solvents, sodium azide

Scheme 16 Synthesis of the Ni-cytosine@MCM-41 and Cu-cytosine@MCM-41 nano-catalyst.¹²³




Fig. 10 SEM images of the (a) Cu-cytosine@MCM-41 and (b) Ni-cytosine@MCM-41 (Adapted with consent from ref. 123. Copyright John Wiley and Sons).

concentration and temperature in the [3+2] cycloaddition of benzonitrile and sodium azide. It was observed that optimum results were obtained with 3.1 mol% Cu-cytosine@MCM-41 and 0.1 mol% of Ni-cytosine@MCM-41 catalyst loading in PEG-400 at 120 °C using 1.3 mmol of sodium azide. Further, to establish the role of Cu-cytosine@MCM-41 and Ni-cytosine@MCM-41 in the tetrazoles synthesis, authors also demonstrated comparative catalytic study of nickel nitrate ($\text{Ni}(\text{NO}_3)_2$), copper nitrate ($\text{Cu}(\text{NO}_3)_2$), MCM-41, and cytosine as a catalyst. It was noticed that compared to Cu-cytosine@MCM-41 and Ni-cytosine@MCM-41, model reaction was not completed in presence of either $\text{Ni}(\text{NO}_3)_2$, $\text{Cu}(\text{NO}_3)_2$, MCM-41, cytosine even

when reaction time was prolonged. However, reported organic-inorganic hybrid catalyst system offers several beneficial features such as high catalytic activity due to porous large surface area, reusability, low metal leaching in the reaction solution and high yield.¹²³

In another research, L-cysteine-Pd@MCM-41 organic-inorganic hybrid nano-catalyst was synthesized by the functionalization of L-cysteine on mesoporous MCM-41 channel followed by addition of palladium particles on surface of L-cysteine@MCM-41. The L-cysteine-Pd@MCM-41 organic-inorganic hybrid nano-catalyst was used for synthesis of 5-substituted-1*H*-tetrazoles (Table 10, entry 4). The catalyst was characterized by various spectroscopic techniques such as FTIR, XRD, TGA, EDS, WDX, SEM ICP, and N_2 adsorption-desorption measurement isotherms. The collected data suggest that palladium complex immobilized onto MCM-41 pores. Additionally, protocol offered several advantages such as simple methodology, easy separation of catalyst, shorter reaction times, high product yields and catalyst retrieval up to six times without losing significant activity and selectivity.¹²⁴

Similarly, Pd-SBT@MCM-41 nano-catalyst was prepared and examined as catalyst for 5-substituted-1*H*-tetrazoles synthesis (Table 10, entry 5). The Pd-SBT@MCM-41 nano-catalyst was prepared *via* sequential process depicted in the Scheme 17. The characterization of reported nanomaterial was done through XRD, SEM, TGA, ICP-OES and N_2 absorption-desorption isotherms techniques. The SEM analysis revealed that Pd-SBTU@MCM-41 are spherical with a mean diameter of 80 ± 20 nm. Further, TEM image exhibited that catalyst had well-ordered hexagonal type of array of regular hexagonal arrangement before and after coordination of palladium. The TEM image also reveals most of Pd particles supported on MCM-41 with an average diameter of about 4 ± 1 nm. The synthesized Pd-SBT@MCM-41 was successfully employed in 5-substituted-1*H*-tetrazole synthesis *via* reaction of various nitriles with sodium azide. This protocol offered several advantages like high





Fig. 11 TEM images of Cu-cytosine@MCM-41. Adapted with consent from ref. 123. Copyright John Wiley and Sons.




Scheme 17 Preparation of the Pd-SMTU@MCM-41 nano-catalyst.¹²⁵

stability, easy handling, simple application procedure and reusability of the catalyst.¹²⁵

9. Miscellaneous nano-catalyst for the synthesis of tetrazole

Besides these broad sorts of nanostructured catalyst some miscellaneous nano-catalysts were also explored by various research groups. Such as Movaheditabar and Javaherian explored nano-silica melamine trisulfonic acid as heterogeneous catalyst for synthesis of 5-substituted-1*H*-tetrazoles. The

reaction was optimized *via* model reaction of benzonitrile with sodium azide. It was observed that nitrile activity is critical toward azide ion in [3+2] cycloaddition reactions. Thus, in shorter time better yield was obtain when aliphatic nitriles were reacted with sodium azide contrary to aromatic nitriles. The lower activity of aromatic nitriles may be due to their significant resonance between the aromatic ring and cyano group, which decreases its electrophilicity. Additionally, authors reported that if electron-donating and withdrawing substituents were located at *para*- or *meta*-positions in aromatic compounds, no major difference reaction times or in product yields were

Scheme 18 Preparation of $\text{Fe}_3\text{O}_4@SiO_2@CPTMS@AMTDA@Co$ nano-catalyst.¹²⁹

observed. It has also observed that synthesized catalyst efficiently recovered after reaction completion and could be applicable at the industrial scale.¹²⁶ Similarly, a nickel zirconium phosphate (NiZrP) based heterogeneous catalyst was prepared and characterized for synthesis of 5-substituted-1*H*-tetrazoles *via* the [3+2] cycloaddition reaction of sodium azide with nitriles. They found that 5 mol% of nickel zirconium phosphate is optimum to achieve maximum yield of the tetrazoles. Additionally, short reaction time with excellent product formation are attractive advantages of this method.¹²⁷ The β -Ni(OH)₂ nanoparticles was reported as a green heterogeneous catalyst for synthesis of versatile substituted tetrazoles. Authors found that 4.32 mol% of β -Ni(OH)₂ catalyst loading in water is sufficient for tetrazoles synthesis *via* simple, one-pot reaction. Owing to nanocrystalline nature, good thermal stability, large surface area and small particle size of β -Ni(OH)₂ offer excellent yield (98%) of the 5-substituted-1*H*-tetrazoles. Another heterogeneous nano catalyst TiCl₄·SiO₂ was developed by Zamani *et al.* for 5-substituted-1*H*-tetrazoles synthesis. They noticed that 0.1 mol% catalyst loading is sufficient for the maximum yield of the product.¹²⁸ In this line, magnetic Fe₃O₄@SiO₂@CPTMS@AMTDA@Co heterogeneous catalyst was developed *via* sequential process for synthesis of diverse tetrazoles as depicted in Scheme 18. The Fe₃O₄@SiO₂@CPTMS@AMTDA@Co activity optimized in a model reaction of aniline with sodium azide and triethyl orthoformate (TEOF) to form 1-phenyl-1*H*-tetrazole. Moreover, the reusability of catalyst also performed up to five runs without losing noticeable catalytic activity.¹²⁹

10. Generalized reaction mechanism of the nano-catalyzed tetrazole synthesis

In this section, general mechanism involves in synthesis of tetrazoles in both [3+2] cycloaddition reaction and one pot

multicomponent reaction are described in detail using suitable reactants.

11. Mechanism for tetrazoles synthesis *via* [3+2] cycloaddition

The [3+2] cycloaddition nano-catalyzed tetrazole synthesis was initiated with interaction of nitrile group with nano catalyst and forms an intermediate which accelerates [3+2] cycloaddition step. Certainly, nano-catalyst activates nitrile groups *via* coordination to nitrogen and/or triple bond which enhances electrophilic character of cyanide group (intermediate **I**). Thereafter, sodium azide reacts with this complex and produces second intermediate (**II**). Finally, 1,3-*H*-shift produce tetrazole product with acidic work up (Fig. 12).

12. One-pot multicomponent reaction of amine/amino acids, triethyl orthoformate and sodium azide

One pot multicomponent reaction of amine/amino acids with triethyl orthoformate and azide is another alternative route for synthesis of tetrazole derivatives. The reaction mechanism of nanomaterials catalyzed synthesis of tetrazole derivatives *via* multicomponent reaction approach is well described in literature in which nanoparticle attack to oxygen atom of ethoxy group of triethyl orthoformate thereby facilitate nucleophilic attack of amino group on intermediate **A**. Successive removal of two molecule of ethanol followed by nucleophilic attack of azide to give intermediate **B** which than cyclize to produce final tetrazole (Fig. 13).

Similarly, the reaction mechanism of nanomaterial catalyzed multicomponent reaction between aldehyde, malononitrile and azide is depicted in Fig. 14. Initially, carbonyl oxygen atom of




Fig. 12 Reaction mechanism for tetrazole *via* [3+2] cycloaddition.






Fig. 13 Reaction mechanism for tetrazole via one-pot MCR of amine/amino acids, triethyl orthoformate and sodium azide.




Fig. 14 Reaction mechanism for tetrazole via aldehyde, active methylene compound (malononitrile) and sodium azide.

aldehyde was activated by nanoparticle to produce intermediate A. The activated carbonyl compound (A) reacts with malononitrile to form arylidene malononitrile (B) via Knoevenagel condensation. The nitrogen atom of arylidene malononitrile (B) activated by nanoparticles forms complex (C) which subsequently activates towards the attack of the azide ion. The [3+2] cycloaddition reaction between arylidene malononitrile and azide ion gave intermediate (D). Finally, addition of acid produces the target tetrazole.

13. Conclusions

Tetrazoles and its derivatives are vital heterocyclic scaffolds having wide applications in various fields such as biochemistry, medicinal chemistry, high energy material science and many more. The synthesis of tetrazole and its derivatives using conventional homogeneous and heterogeneous catalytic systems

are associated with several constraints such as long reaction time, moderate yield, tedious workup, and purification *etc.* Therefore, the prime focus of chemists working in this domain is to develop efficient, environmental, benign, and catalytic systems for the synthesis of tetrazoles and its derivatives with the aim of high yield, minimum by-products, low temperature conditions, catalyst reusability, easy workup procedure, and economic and eco-friendly methodology. In this context, different nanomaterial-derived catalytic systems have undoubtedly proven as an efficient and eco-friendly catalyst for tetrazole synthesis. This review article demonstrated the synthesis, characterization, and benefits of various classes of nanomaterial-derived catalysts for example boehmite, magnetic and nonmagnetic nanoparticles, nanomaterials supported copper, zinc, carbon-derived nanomaterials, MCM-41, and composite nanostructures under different reaction conditions in tetrazole synthesis. Along with



this, we have also focused on the fabrication, functionalization, and characterization of some novel nanomaterials. We hope that this review would enrich the process chemists to design and develop eco-friendly and novel synthetic methodology for tetrazole and their derivatives considering environmental and chemical industrial apprehensions and area of nanomaterials derived synthesis of tetrazole will continue to flourish.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

S.S. and R.S. also would like to acknowledge Manipal University Jaipur for the laboratory facility.

References

- 1 A. Corma and H. Garcia, *Adv. Synth. Catal.*, 2006, **348**, 1391–1412.
- 2 M. B. Gawande, Y. Monga, R. Zboril and R. Sharma, *Coord. Chem. Rev.*, 2015, **288**, 118–143.
- 3 R. S. Upadhyaya, S. Jain, N. Sinha, N. Kishore, R. Chandra and S. K. Arora, *Eur. J. Med. Chem.*, 2004, **39**, 579–592.
- 4 E. Łukowska-Chojnacka, J. Mierzejewska, M. Milner-Krawczyk, M. Bondaryk and M. Staniszevska, *Bioorg. Med. Chem.*, 2016, **24**, 6058–6065.
- 5 M. Asif, *Pharm. Methods*, 2014, **5**, 39–46.
- 6 A. Qian, Y. Zheng, R. Wang, J. Wei, Y. Cui, X. Cao and Y. Yang, *Bioorg. Med. Chem. Lett.*, 2018, **28**, 344–350.
- 7 S. Vembu, P. Parasuraman and M. Gopalakrishnan, *Pharma Chem.*, 2014, **6**, 35–44.
- 8 P. Zhan, Z. Li, X. Liu and E. De Clercq, *Mini-Rev. Med. Chem.*, 2009, **9**, 1014–1023.
- 9 K. Chauhan, M. Sharma, P. Trivedi, V. Chaturvedi and P. M. Chauhan, *Bioorg. Med. Chem. Lett.*, 2014, **24**, 4166–4170.
- 10 R. M. DeMarinis, J. R. Hoover, G. L. Dunn, P. Actor, J. V. Uri and J. A. Weisbach, *J. Antibiot.*, 1975, **28**, 463–470.
- 11 M. Uchida, M. Komatsu, S. Morita, T. Kanbe and K. Nakagawa, *Chem. Pharm. Bull.*, 1989, **37**, 322–326.
- 12 M. Uchida, M. Komatsu, S. Morita, T. Kanbe, K. Yamasaki and K. Nakagawa, *Chem. Pharm. Bull.*, 1989, **37**, 958–961.
- 13 R. E. Ford, P. Knowles, E. Lunt, S. M. Marshall, A. J. Penrose, C. A. Ramsden, A. J. Summers, J. L. Walker and D. E. Wright, *J. Med. Chem.*, 1986, **29**, 538–549.
- 14 S. Hayao, H. J. Havera, W. G. Strycker, T. Leipzig and R. Rodriguez, *J. Med. Chem.*, 1967, **10**, 400–402.
- 15 P. Purohit, A. K. Pandey, D. Singh, P. S. Chouhan, K. Ramalingam, M. Shukla, N. Goyal, J. Lal and P. M. Chauhan, *MedChemComm*, 2017, **8**, 1824–1834.
- 16 C. G. Neochoritis, T. Zhao and A. Dömling, *Chem. Rev.*, 2019, **119**, 1970–2042.
- 17 F. Gao, J. Xiao and G. Huang, *Eur. J. Med. Chem.*, 2019, **184**, 111744.
- 18 V. Bhaskar and P. Mohite, *J. Optoelectron. Biomed. Mater.*, 2010, **2**, 249–259.
- 19 L. Myznikov, A. Hrabalek and G. Koldobskii, *Chem. Heterocycl. Compd.*, 2007, **43**, 1–9.
- 20 K. Dileep, S. Polepalli, N. Jain, S. K. Buddana, R. Prakasham and M. Murty, *Mol. Diversity*, 2018, **22**, 83–93.
- 21 A. K. Gupta, C. H. Song and C. H. Oh, *Tetrahedron Lett.*, 2004, **45**, 4113–4116.
- 22 D. R. Patil, Y. B. Wagh, P. G. Ingole, K. Singh and D. S. Dalal, *New J. Chem.*, 2013, **37**, 3261–3266.
- 23 J. Dhainaut, J.-P. Dacquin, A. F. Lee and K. Wilson, *Green Chem.*, 2010, **12**, 296–303.
- 24 F. G. Fallon and R. M. Herbst, *J. Org. Chem.*, 1957, **22**, 933–936.
- 25 T. Jin, S. Kamijo and Y. Yamamoto, *Tetrahedron Lett.*, 2004, **45**, 9435–9437.
- 26 K. Bahrami, M. M. Khodaei and M. Roostaei, *New J. Chem.*, 2014, **38**, 5515–5520.
- 27 E. Carbonell, E. Delgado-Pinar, J. Pitarch-Jarque, J. Alarcón and E. García-España, *J. Phys. Chem. C*, 2013, **117**, 14325–14331.
- 28 A. Keivanloo, M. Bakherad, E. Imanifar and M. Mirzaee, *Appl. Catal., A*, 2013, **467**, 291–300.
- 29 M. Mirzaee, B. Bahramian and A. Amoli, *Appl. Organomet. Chem.*, 2015, **29**, 593–600.
- 30 Z. Wu, Y. Mao, M. Song, X. Yin and M. Zhang, *Catal. Commun.*, 2013, **32**, 52–57.
- 31 S.-M. Kim, Y.-J. Lee, K.-W. Jun, J.-Y. Park and H. Potdar, *Mater. Chem. Phys.*, 2007, **104**, 56–61.
- 32 A. Ghorbani-Choghamarani, P. Moradi and B. Tahmasbi, *RSC Adv.*, 2016, **6**, 56638–56646.
- 33 A. Ghorbani-Choghamarani, P. Moradi and B. Tahmasbi, *RSC Adv.*, 2016, **6**, 56458–56466.
- 34 V. Vatanpour, S. S. Madaeni, L. Rajabi, S. Zinadini and A. A. Derakhshan, *J. Membr. Sci.*, 2012, **401**, 132–143.
- 35 B. Tahmasbi, A. Ghorbani-Choghamarani and P. Moradi, *New J. Chem.*, 2020, **44**, 3717–3727.
- 36 A. Jabbari, B. Tahmasbi, M. Nikoorazm and A. Ghorbani-Choghamarani, *Appl. Organomet. Chem.*, 2018, **32**, e4295.
- 37 P. Moradi and A. Ghorbani-Choghamarani, *Appl. Organomet. Chem.*, 2017, **31**, e3602.
- 38 A. Ghorbani-Choghamarani, H. Aghavandi and M. Mohammadi, *Appl. Organomet. Chem.*, 2020, **34**, e5804.
- 39 M. A. Jani and K. Bahrami, *Appl. Organomet. Chem.*, 2020, **34**, e6014.
- 40 B. Tahmasbi and A. Ghorbani-Choghamarani, *Appl. Organomet. Chem.*, 2017, **31**, e3644.
- 41 M. Zarghani and B. Akhlaghinia, *RSC Adv.*, 2016, **6**, 31850–31860.
- 42 T. Tamoradi, A. Irandoust and M. Ghadermazi, *J. Iran. Chem. Soc.*, 2019, **16**, 1723–1733.
- 43 T. Tamoradi, S. Taherabadi and M. Ghadermazi, *Polyhedron*, 2019, **171**, 305–311.
- 44 T. Tamoradi, A. Ghorbani-Choghamarani and M. Ghadermazi, *New J. Chem.*, 2017, **41**, 11714–11721.



- 45 M. Nikoorazm and Z. Erfani, *Chem. Phys. Lett.*, 2019, **737**, 136784.
- 46 M. Nemati, T. Tamoradi and H. Veisi, *Polyhedron*, 2019, **167**, 75–84.
- 47 N. Moeini, T. Tamoradi, M. Ghadermazi and A. Ghorbani-Choghamarani, *Appl. Organomet. Chem.*, 2018, **32**, e4445.
- 48 H. Fatahi, M. Jafarzadeh and Z. Pourmanouchehri, *J. Heterocycl. Chem.*, 2019, **56**, 2090–2098.
- 49 M. A. Ashraf, Z. Liu, C. Li and D. Zhang, *Appl. Organomet. Chem.*, 2020, e6133.
- 50 M. S. Ghasemzadeh and B. Akhlaghinia, *ChemistrySelect*, 2020, **5**, 6440–6452.
- 51 A. Karimian, M. Namvar-Mhaboub and R. Abbasi, *Russ. J. Org. Chem.*, 2020, **56**, 1646–1653.
- 52 T. Tamoradi, A. Ghorbani-Choghamarani and M. Ghadermazi, *Solid State Sci.*, 2019, **88**, 81–94.
- 53 H. Naeimi and S. Mohamadabadi, *Dalton Trans.*, 2014, **43**, 12967–12973.
- 54 A. Khalafi-Nezhad and S. Mohammadi, *RSC Adv.*, 2013, **3**, 4362–4371.
- 55 D. Habibi, N. Pakravan, A. Arabi and Z. Kaboudvand, *Appl. Organomet. Chem.*, 2018, **32**, e3988.
- 56 D. Habibi, S. Heydari, A. Gil, M. Afsharfarnia, A. Faraji, R. Karamian and M. Asadbegy, *Appl. Organomet. Chem.*, 2018, **32**, e4005.
- 57 M. S. Ghasemzadeh and B. Akhlaghinia, *Bull. Chem. Soc. Jpn.*, 2017, **90**, 1119–1128.
- 58 M. Salimi and A. Zamanpour, *Appl. Organomet. Chem.*, 2020, **34**, e5682.
- 59 M. Salimi, F. Esmaeli-nasrabadi and R. Sandaroods, *Inorg. Chem. Commun.*, 2020, **122**, 108287.
- 60 A. N. Parouch, N. Koukabi and E. Abdous, *Res. Chem. Intermed.*, 2020, **46**, 3295–3310.
- 61 R. Gao, Z. Yang, L. Zheng, L. Gu, L. Liu, Y. Lee, Z. Hu and X. Liu, *ACS Catal.*, 2018, **8**, 1955–1963.
- 62 N. Zhou, Q. An, Z. Xiao, S. Zhai and Z. Shi, *ACS Sustainable Chem. Eng.*, 2017, **5**, 5394–5407.
- 63 W. Xie, Z. Guo, F. Gao, Q. Gao, D. Wang, B.-s. Liaw, Q. Cai, X. Sun, X. Wang and L. Zhao, *Theranostics*, 2018, **8**, 3284.
- 64 X. Sun, M. Sui, G. Cui, L. Li, X. Li, X. Lv, F. Wu and G. Gu, *RSC Adv.*, 2018, **8**, 17489–17496.
- 65 M. Khodamorady and K. Bahrami, *ChemistrySelect*, 2019, **4**, 8183–8194.
- 66 P. Akbarzadeh, N. Koukabi and M. M. Hosseini, *J. Heterocycl. Chem.*, 2020, **57**, 2455–2465.
- 67 P. Akbarzadeh, N. Koukabi and E. Kolvari, *Res. Chem. Intermed.*, 2019, **45**, 1009–1024.
- 68 B. C. Ranu, R. Dey, T. Chatterjee and S. Ahammed, *ChemSusChem*, 2012, **5**, 22–44.
- 69 S. E. Allen, R. R. Walvoord, R. Padilla-Salinas and M. C. Kozlowski, *Chem. Rev.*, 2013, **113**, 6234–6458.
- 70 K. Pan, H. Ming, H. Yu, Y. Liu, Z. Kang, H. Zhang and S. T. Lee, *Cryst. Res. Technol.*, 2011, **46**, 1167–1174.
- 71 F. Taghavi, M. Gholizadeh, A. S. Saljooghi and M. Ramezani, *MedChemComm*, 2017, **8**, 1953–1964.
- 72 B. Salahshournia, H. Hamadi and V. Nobakht, *Appl. Organomet. Chem.*, 2018, **32**, e4416.
- 73 P. Movaheditabar, M. Javaherian and V. Nobakht, *React. Kinet., Mech. Catal.*, 2017, **122**, 217–228.
- 74 P. Moradi, M. Hajjami and B. Tahmasbi, *Polyhedron*, 2020, **175**, 114169.
- 75 J. Y. Kim, K. Park, S. Y. Bae, G. C. Kim, S. Lee and H. C. Choi, *J. Mater. Chem.*, 2011, **21**, 5999–6005.
- 76 T. Tamoradi, M. Ghadermazi and A. Ghorbani-Choghamarani, *New J. Chem.*, 2018, **42**, 5479–5488.
- 77 L. Wei, S. Yan, H. Wang and H. Yang, *NPG Asia Mater.*, 2018, **10**, 899–911.
- 78 T. Tamoradi, A. Ghorbani-Choghamarani, M. Ghadermazi and H. Veisi, *Solid State Sci.*, 2019, **91**, 96–107.
- 79 F. Pourhassan and H. Eshghi, *Catal. Lett.*, 2020, **150**, 1287–1300.
- 80 R. Ghafouri-Nejad and M. Hajjami, *React. Kinet., Mech. Catal.*, 2020, **129**, 371–389.
- 81 A. Ghorbani-Choghamarani, L. Shiri and G. Azadi, *RSC Adv.*, 2016, **6**, 32653–32660.
- 82 M. Esmailpour, J. Javidi and S. Zahmatkesh, *Appl. Organomet. Chem.*, 2016, **30**, 897–904.
- 83 M. Esmailpour, J. Javidi, F. N. Dodeji and M. M. Abarghoui, *J. Mol. Catal. A: Chem.*, 2014, **393**, 18–29.
- 84 F. Dehghani, A. R. Sardarian and M. Esmailpour, *J. Organomet. Chem.*, 2013, **743**, 87–96.
- 85 M. Darabi, T. Tamoradi, M. Ghadermazi and A. Ghorbani-Choghamarani, *Transition Met. Chem.*, 2017, **42**, 703–710.
- 86 N. Razavi and B. Akhlaghinia, *RSC Adv.*, 2015, **5**, 12372–12381.
- 87 G. Azadi, A. Ghorbani-Choghamarani and L. Shiri, *Transition Met. Chem.*, 2017, **42**, 131–136.
- 88 V. Khorramabadi, D. Habibi and S. Heydari, *Green Chem. Lett. Rev.*, 2020, **13**, 50–59.
- 89 V. Khorramabadi, D. Habibi and S. Heydari, *Org. Prep. Proced. Int.*, 2020, **52**, 139–146.
- 90 M. Ariannezhad, D. Habibi and S. Heydari, *Polyhedron*, 2019, **160**, 170–179.
- 91 A. R. Sardarian, H. Eslahi and M. Esmailpour, *ChemistrySelect*, 2018, **3**, 1499–1511.
- 92 T. Sun, J. Qiu and C. Liang, *J. Phys. Chem. C*, 2008, **112**, 715–721.
- 93 E. S. Jang, J. H. Won, S. J. Hwang and J. H. Choy, *Adv. Mater.*, 2006, **18**, 3309–3312.
- 94 Y. A. Attia, *Mater. Express*, 2016, **6**, 211–219.
- 95 T.-J. Hsueh, C.-L. Hsu, S.-J. Chang, P.-W. Guo, J.-H. Hsieh and I.-C. Chen, *Scr. Mater.*, 2007, **57**, 53–56.
- 96 C. Huo, J. Ouyang and H. Yang, *Sci. Rep.*, 2014, **4**, 1–9.
- 97 R. S. Mane, W. J. Lee, H. M. Pathan and S.-H. Han, *J. Phys. Chem. B*, 2005, **109**, 24254–24259.
- 98 O. Seven, B. Dindar, S. Aydemir, D. Metin, M. Ozinel and S. Icli, *J. Photochem. Photobiol., A*, 2004, **165**, 103–107.
- 99 P. Chhattise, S. Saleh, V. Pandit, S. Arbuji and V. Chabukswar, *Mater. Adv.*, 2020, **1**, 2339–2345.
- 100 M. Lakshmi Kantam, K. S. Kumar and C. Sridhar, *Adv. Synth. Catal.*, 2005, **347**, 1212–1214.



- 101 F. Himo, Z. P. Demko and L. Noodleman, *J. Org. Chem.*, 2003, **68**, 9076–9080.
- 102 F. Himo, Z. P. Demko, L. Noodleman and K. B. Sharpless, *J. Am. Chem. Soc.*, 2002, **124**, 12210–12216.
- 103 F. Himo, Z. P. Demko, L. Noodleman and K. B. Sharpless, *J. Am. Chem. Soc.*, 2003, **125**, 9983–9987.
- 104 S. M. Agawane and J. M. Nagarkar, *Catal. Sci. Technol.*, 2012, **2**, 1324–1327.
- 105 Y. A. Attia, Y. M. Mohamed, M. M. Awad and S. Alexeree, *J. Organomet. Chem.*, 2020, **919**, 121320.
- 106 Y. M. A. Mohamed and Y. A. Attia, *Appl. Organomet. Chem.*, 2020, **34**, e5758.
- 107 C. Testa, A. Zammataro, A. Pappalardo and G. T. Sfrazzetto, *RSC Adv.*, 2019, **9**, 27659–27664.
- 108 A. V. Vakhrushev, V. I. Kodolov, A. Haghi and S. C. Ameta, *Carbon Nanotubes and Nanoparticles: Current and Potential Applications*, CRC Press, 2019.
- 109 P. Akbarzadeh, N. Koukabi and E. Kolvari, *Mol. Diversity*, 2019, 1–15.
- 110 M. Kazemnejadi, B. Mahmoudi, Z. Sharafi, M. A. Nasser, A. Allahresani and M. Esmaeilpour, *Appl. Organomet. Chem.*, 2020, **34**, e5273.
- 111 M. N. Soltani Rad, S. Behrouz, V. Sadeghi Dehchenari and S. J. Hoseini, *J. Heterocycl. Chem.*, 2017, **54**, 355–365.
- 112 M. Sen, in *Nanotechnology and the Environment*, IntechOpen, 2020.
- 113 M. Nasrollahzadeh, F. Ghorbannezhad and S. M. Sajadi, *Appl. Organomet. Chem.*, 2019, **33**, e4698.
- 114 A. Ahmadi, T. Sedaghat, H. Motamedi and R. Azadi, *Appl. Organomet. Chem.*, 2020, **34**, e5572.
- 115 F. Rezaei, M. A. Amrollahi and R. Khalifeh, *Inorg. Chim. Acta*, 2019, **489**, 8–18.
- 116 N. Motahharifar, M. Nasrollahzadeh, A. Taheri-Kafrani, R. S. Varma and M. Shokouhimehr, *Carbohydr. Polym.*, 2020, **232**, 115819.
- 117 M. Nasrollahzadeh, M. Sajjadi, M. R. Tahsili, M. Shokouhimehr and R. S. Varma, *ACS Omega*, 2019, **4**, 8985–9000.
- 118 H. R. Pawar and R. C. Chikate, *J. Mol. Struct.*, 2021, **1225**, 128985.
- 119 J. Choi, D. Kim, S. Chang and W. Ahn, *Appl. Catal., A*, 2003, **254**, 225–237.
- 120 A. S. Diez, M. Alvarez and M. A. Volpe, *J. Braz. Chem. Soc.*, 2015, **26**, 1542–1550.
- 121 S. Alahmad, *Orient. J. Chem.*, 2012, **28**, 1.
- 122 F. Matloubi Moghaddam, M. Eslami and N. Ghadirian, *Sci. Iran.*, 2019, **26**, 1463–1473.
- 123 M. Nikoorazm, B. Tahmasbi, S. Gholami and P. Moradi, *Appl. Organomet. Chem.*, 2020, **34**, e5919.
- 124 M. Nikoorazm, P. Moradi and N. Noori, *J. Porous Mater.*, 2020, **27**, 1159–1169.
- 125 M. Nikoorazm, A. Ghorbani-Choghamarani, M. Ghobadi and S. Massahi, *Appl. Organomet. Chem.*, 2017, **31**, e3848.
- 126 P. Movaheditabar and M. Javaherian, *J. Org. Chem. Res.*, 2019, **5**, 174–189.
- 127 F. Abrishami, M. Ebrahimikia and F. Rafiee, *Iran. J. Catal.*, 2016, **6**, 245–251.
- 128 M. Halder, M. M. Islam, P. Singh, A. Singha Roy, S. M. Islam and K. Sen, *ACS Omega*, 2018, **3**, 8169–8180.
- 129 F. Sarrafioun, S. Jamehbozorgi and M. Ramezani, *Russ. J. Org. Chem.*, 2019, **55**, 1777–1784.

