


 Cite this: *RSC Adv.*, 2021, 11, 14260

## Correction: Metal-free oxidative coupling of arylmethyamines with indoles: a simple, environmentally benign approach for the synthesis of 3,3'-bis(indolyl)methanes

 Vikas D. Kadu,<sup>\*a</sup> Sankala Naga Chandrudu,<sup>b</sup> Mahesh G. Hublikar,<sup>a</sup> Dattatraya G. Raut<sup>a</sup> and Raghunath B. Bhosale<sup>a</sup>

DOI: 10.1039/d1ra90099d

[rsc.li/rsc-advances](https://rsc.li/rsc-advances)

 Correction for 'Metal-free oxidative coupling of arylmethyamines with indoles: a simple, environmentally benign approach for the synthesis of 3,3'-bis(indolyl)methanes' by Vikas D. Kadu *et al.*, *RSC Adv.*, 2020, 10, 23254–23262, DOI: 10.1039/D0RA03221B.


The authors regret several errors throughout this *RSC Advances* manuscript. These include an incorrect description of ref. 23 (ref. 1) which starts at line 11 of paragraph 3 in the Introduction and should read as: "Previously, the Gopalaiah research group reported the synthesis of 3,3'-benzylidenebis(1*H*-indole), using an iron(II) triflate catalyst, from benzylamine with indoles.<sup>23</sup>" as opposed to "Previously, the Sankala research group reported the synthesis of 3,3'-benzylidenebis(1*H*-indole), using an iron(II) triflate catalyst, from benzylamine with indoles.<sup>23</sup>"

The authors regret an error in Scheme 2 where 4-methylbenzylamine was incorrectly labelled as compound **1a** and should have been labelled as compound **1b**. The correct version of Scheme 2 is shown below.

The authors regret the inclusion of the following sentence starting at line 3 of paragraph 1 in the Experimental section: "Iron salts were purchased from Sigma-Aldrich and were used as received." This sentence should be removed as it does not relate to the experimental procedures in this *RSC Advances* article.

The authors also regret that the wrong temperature was stated in line 4 of paragraph 2 in the Experimental section, it should read as "The round bottom flask was equipped with an O<sub>2</sub> balloon, and the reaction mixture was stirred at 120 °C until the complete consumption of indole **2** occurred, as monitored *via* TLC."

The authors regret that 4-methyl-*N*-(4-methylbenzylidene)benzylamine was incorrectly described in lines 10 and 11 of paragraph 3 in the Experimental section. 4-Methylbenzylamine was also incorrectly labelled as compound **1a** in line 2 and the wrong temperature was stated in line 5. Paragraph 3 in the Experimental section should therefore read as: "4-Methylbenzylamine (**1b**)



Scheme 2 Control experiments.

<sup>a</sup>School of Chemical Sciences, Punyashlok Ahilyadevi Holkar Solapur University, Solapur-413255, Maharashtra, India. E-mail: vikaskadu1@gmail.com

<sup>b</sup>Department of Chemistry, College of Engineering, Rayalseema University, Kurnool-518002, Andhra Pradesh, India


(350 mg, 2.89 mmol), AcOH (10 mol%), and dry chlorobenzene (2 mL) were added to a 25 mL round bottom flask. The round bottom flask was equipped with an O<sub>2</sub> balloon, and the reaction mixture was stirred at 120 °C for 3.4 h. The reaction mixture was cooled to room temperature and adsorbed on basic alumina. It was purified *via* column chromatography over basic alumina using a hexane/ethyl acetate (9 : 1) mixture as the eluent to afford 4-methyl-*N*-(4-methylbenzylidene)benzylamine (**4**) (232 mg, 72% yield) as a pale yellow semisolid.”

The authors have also updated some <sup>1</sup>H and <sup>13</sup>C NMR spectra in the ESI to differentiate them from the NMR spectra in their previous *Synthesis* article.<sup>1</sup> The updated spectra in the revised ESI are the <sup>1</sup>H and <sup>13</sup>C NMR spectra for compounds **3ba**, **3da**, **3ea**, **3fa**, **3ga**, **3ha**, **3ia**, **3ja**, **3na**, **3oa**, **3pa**, **3qa**, **3ra** and **3sa**. The <sup>1</sup>H NMR and <sup>13</sup>C NMR signal values given in the Experimental section in the main article have been corrected to reflect the updated NMR spectra. An expert viewed the corrected data and was satisfied that the conclusions are not affected. The corrected NMR data are given below:

### Synthesis of 3,3'-(4-methylbenzylidene)bis(1*H*-indole) (**3ba**)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.88 (br s, 2H, NH), 7.40 (d, *J* = 8.0 Hz, 2H), 7.36–7.34 (m, 2H), 7.24 (d, *J* = 8.4 Hz, 2H), 7.20–7.16 (m, 2H), 7.09 (d, *J* = 8.0 Hz, 2H), 7.04–7.00 (m, 2H), 6.64 (d, *J* = 2.4 Hz, 2H), 5.86 (s, 1H), 2.33 (s, 3H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 140.9, 136.7, 135.5, 128.9, 128.6, 127.1, 123.6, 121.9, 119.9, 119.8, 119.2, 111.0, 39.7, 21.1 ppm.

### Synthesis of 3,3'-(4-hydroxybenzylidene)bis(1*H*-indole) (**3da**)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.68 (br s, 2H, NH), 8.17 (br s, 1H, OH), 7.32–7.24 (m, 4H), 7.10–7.04 (m, 4H), 6.90–6.87 (m, 2H), 6.70–6.67 (m, 2H), 6.59 (d, *J* = 1.2 Hz, 2H), 5.72 (s, 1H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 154.1, 136.7, 135.2, 129.5, 127.0, 123.7, 121.3, 119.8, 119.7, 118.6, 115.1, 111.1, 39.3 ppm.

### Synthesis of 3,3'-(4-*tert*-butylbenzylidene)bis(1*H*-indole) (**3ea**)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.04 (br s, 2H, NH), 7.41 (d, *J* = 8.0 Hz, 2H), 7.35 (d, *J* = 8.0 Hz, 2H), 7.27–7.26 (m, 4H), 7.18–7.13 (m, 2H), 7.01–6.97 (m, 2H), 6.90 (d, *J* = 2.0 Hz, 2H) 5.86 (s, 1H), 1.29 (s, 9H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 148.7, 140.8, 136.6, 128.2, 127.1, 125.0, 123.5, 121.7, 119.9, 119.8, 119.0, 111.0, 39.6, 34.4, 31.4 ppm.

### Synthesis of 3,3'-(4-fluorobenzylidene)bis(1*H*-indole) (**3fa**)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.94 (br s, 2H, NH), 7.38–7.28 (m, 6H), 7.19 (t, *J* = 7.6 Hz, 2H), 7.04–6.95 (m, 4H), 6.47 (d, *J* = 1.6 Hz, 2H), 5.88 (s, 1H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 139.7, 136.7, 130.1, 130.0, 126.9, 123.5, 122.0, 119.9, 119.6, 119.3, 115.1, 114.9, 111.1, 39.4 ppm.

### Synthesis of 3,3'-((4-chlorophenyl)methylene)bis(1*H*-indole) (**3ga**)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.95 (br s, 2H, NH), 7.38–7.23 (m, 8H, Ar-H), 7.20–7.16 (m, 2H, Ar-H), 7.04–6.99 (m, 2H, Ar-H), 6.66 (t, *J* = 1.2 Hz, 2H, Ar-H), 5.87 (s, 1H, Ar-CH) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 142.5, 136.7, 131.8, 130.1, 128.4, 126.9, 123.6, 122.1, 119.8, 119.3, 119.2, 111.1, 39.6 ppm.

### Synthesis of 4-(di(1*H*-indol-3-yl)methyl)benzylidene (**3ha**)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.86 (br s, 2H, NH), 7.26–7.11 (m, 7H, Ar-H), 6.98–6.81 (m, 5H, Ar-H), 6.71 (t, *J* = 7.6 Hz, 2H, Ar-H), 6.09 (s, 1H, CH) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 144.2, 135.1, 132.4, 129.3, 128.7, 128.2, 126.6, 120.2, 118.8, 118.6, 111.1, 110.2, 38.3 ppm.

### Synthesis of 3,3'-(2-methoxybenzylidene)bis(1*H*-indole) (**3ia**)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.79 (br s, 2H, NH), 7.43 (d, *J* = 7.6 Hz, 2H), 7.31 (d, *J* = 8.0 Hz, 2H), 7.21–7.14 (m, 4H), 7.03–6.83 (m, 4H), 6.60 (s, 2H), 6.37 (s, 1H), 3.82 (s, 3H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 156.9, 136.7, 132.3, 129.7, 127.2, 127.1, 123.7, 123.5, 122.3, 121.7, 120.4, 120.0, 119.8, 119.6, 119.0, 110.9, 110.6, 55.7, 32.0 ppm.

### Synthesis of 3,3'-(2-chlorobenzylidene)bis(1*H*-indole) (**3ja**)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.90 (br s, 2H), 7.44–7.35 (m, 5H), 7.24–7.01 (m, 7H), 6.63 (dd, *J* = 2.4 & 0.8 Hz, 2H), 6.35 (s, 1H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 141.3, 136.7, 133.9, 130.3, 129.4, 127.5, 127.0, 126.6, 123.7, 122.0, 119.8, 119.3, 118.3, 111.0, 36.6 ppm.



## Synthesis of 3,3'-[3,5-bis(trifluoromethyl)benzylidene]bis(1*H*-indole) (3na)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.98 (br s, 2H, NH), 7.83 (s, 2H), 7.78 (s, 1H), 7.40–7.35 (m, 4H), 7.23 (t, *J* = 7.6 Hz, 2H), 7.06 (t, *J* = 7.2 Hz, 2H), 6.62 (d, *J* = 2.4 Hz, 2H), 6.04 (s, 1H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 146.7, 136.7, 131.6, 131.2, 128.8, 126.5, 124.8, 123.7, 122.4, 122.1, 120.5, 119.7, 119.4, 117.9, 111.3, 40.0 ppm.

## Synthesis of 3,3'-(1-naphthylmethylene)bis(1*H*-indole) (3oa)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.85 (br s, 2H), 8.11 (d, *J* = 8.4 Hz, 1H), 7.80 (d, *J* = 8.0 Hz, 1H), 7.65 (d, *J* = 7.2 Hz, 1H), 7.38–7.19 (m, 8H), 7.09–7.05 (m, 2H), 6.91–6.87 (m, 2H), 6.58 (s, 1H), 5.51 (s, 2H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 136.8, 128.5, 126.7, 126.1, 125.6, 125.4, 125.1, 124.5, 121.4, 119.5, 118.7, 111.2, 35.7 ppm.

## Synthesis of 3,3'-(4-pyridylmethylene)bis(1*H*-indole) (3pa)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.45 (d, *J* = 6.0 Hz, 2H), 8.00 (br s, 2H, NH), 7.40 (d, *J* = 6.0 Hz, 2H), 7.31 (d, *J* = 8.4 Hz, 2H), 7.05 (t, *J* = 7.2 Hz, 2H), 6.93–6.90 (m, 5H), 6.74 (t, *J* = 7.2 Hz, 2H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 174.1, 148.9, 136.7, 127.1, 124.5, 121.7, 121.6, 119.2, 111.0, 36.9 ppm.

## Synthesis of 3,3'-(pyridin-3-ylmethylene)bis(1*H*-indole) (3qa)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.67 (d, 1H, *J* = 2.4 Hz, ArH), 8.48 (dd, 1H, *J* = 4.8 & 1.6 Hz, ArH), 8.26 (brs, 2H, NH), 7.62–7.59 (m, 1H, ArH), 7.38–7.34 (m, 4H, ArH), 7.21–7.16 (m, 3H, ArH), 7.04–7.00 (m, 2H, ArH), 6.62 (d, 2H, *J* = 2.0 Hz, ArH), 5.92 (s, 1H, Ar-CH) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 150.3, 147.5, 139.5, 136.7, 136.1, 126.7, 123.7, 123.3, 122.2, 119.7, 119.4, 118.5, 111.2, 60.4 ppm.

## Synthesis of 3,3'-(2-thienylmethylene)bis(1*H*-indole) (3ra)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.92 (br s, 2H, NH), 7.48 (d, *J* = 8.0 Hz, 2H), 7.37 (d, *J* = 8.4 Hz, 2H), 7.21–7.16 (m, 3H), 7.06–7.02 (m, 2H), 6.94–6.91 (m, 2H), 6.84 (dd, *J* = 2.4 & 0.8 Hz, 2H), 6.17 (s, 1H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 148.6, 136.6, 126.7, 126.4, 125.1, 123.6, 123.1, 122.0, 119.8, 119.7, 119.6, 119.4, 111.1, 35.3 ppm.

## Synthesis of 3,3'-benzylidenebis(5-bromo-1*H*-indole) (3sa)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.23 (br s, 2H, NH), 7.38 (d, *J* = 1.6 Hz, 2H), 7.25–7.13 (m, 9H), 6.59 (d, *J* = 2.0 Hz, 2H), 5.68 (s, 1H) ppm; <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 143.5, 135.5, 128.6, 128.5, 128.3, 126.3, 125.2, 124.3, 121.9, 118.4, 112.9, 112.0, 39.8 ppm.

The authors also regret errors and the misspelling of names in ref. 1, 2, 3, 10 and 12 in the original article which are given correctly as ref. 1–5 below, respectively.

## References

- 1 K. Gopalaiah, S. Naga Chandrudu and A. Devi, *Synthesis*, 2015, **47**, 1766–1774.
- 2 R. B. Tjalkens and S. Safe, *US Pat.*, 8580843, 2013.
- 3 A. Ramirez and S. Garcia-Rubio, *Curr. Med. Chem.*, 2003, **10**, 1891–1915.
- 4 M. A. Zeligs, *J. Med. Food.*, 1998, **1**, 67–85.
- 5 C. H. Thomas, *US Pat.*, 4500689, 1985.

