

Cite this: *Chem. Sci.*, 2021, 12, 6726

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 10th March 2021  
Accepted 10th April 2021

DOI: 10.1039/d1sc01410b

rsc.li/chemical-science

## Light-driven release of cucurbit[8]uril from a bivalent cage†

Pim J. de Vink, Tim van der Hek and Luc Brunsveld\*

Temporal control over supramolecular systems has great potential for the modulation of binding and assembly events, such as providing orthogonal control over protein activity. Especially light controlled triggering provides unique entries for supramolecular systems to interface in a controlled manner with enzymes. Here we report on the light-induced release of cucurbit[8]uril (CB[8]) from a bivalent cage molecule and its subsequent activation of a proteolytic enzyme, caspase-9, that itself is unresponsive to light. Central to the design is the bivalent binding of the cage with high affinity to CB[8], 100-fold stronger than the UV-inactivated products. The affinity switching occurs in the (sub-)micromolar concentration regime, matching the concentration characteristics required for dimerizing and activating caspase-9 by CB[8]. The light-responsive caged CB[8] concept presented offers a novel platform for tuning and application of switchable cucurbiturils and beyond.

## Introduction

Supramolecular systems have shown great potential for the modulation of biomolecular assemblies, providing orthogonal and additive control to natural recognition and switching events.<sup>1–4</sup> Several supramolecular host molecules, such as molecular tweezers, calixarenes, cyclodextrins, and cucurbit[*n*]urils, are able to selectively recognize specific peptide and protein elements through, for example, combinations of side chain residues.<sup>5–10</sup> In particular cucurbit[*n*]urils have shown to be successful in protein binding, enabling for example protein activity modulation, because of their relatively strong affinity to biomolecular epitopes in water.<sup>11–14</sup> Cucurbit[8]uril (CB[8]), specifically, is able to incorporate two N-terminal phenylalanine-glycine-glycine (FGG) motifs simultaneously in its hydrophobic cavity and therewith acts as a supramolecular glue<sup>5</sup> allowing to control the dimerization and assembly of biomolecules such as caspases<sup>15</sup> and larger complexes.<sup>16</sup>

Light controlled triggering systems offer the possibility to perturb and modulate supramolecular over biological systems with spatio or temporal control.<sup>17,18</sup> Beautiful examples of supramolecular light stimulus-responsive systems include: light induced pH jumps,<sup>19</sup> photo redox switches<sup>20</sup> and photo-caged guests to release drug molecules from CB[8].<sup>21</sup> Of particular note are diarylethenes-based switches featuring low to high nanomolar affinity switches between their two states,<sup>22</sup> which

has been used for example for induction of light-mediated self-sorting.<sup>23</sup> Notwithstanding, there is a need for novel switchable CB[8]-based systems for controlling biomolecular assemblies, including those with affinities matching the concentrations and assembly characteristics of the biomolecules involved. Here we




Fig. 1 General concept and design elements for light-triggered supramolecular activation of enzymes. CB[8] is temporarily caged by bivalent FGG-peptide **1** that cleaves upon UV-light radiation, which substantially reduces the affinity of the individual cage elements for CB[8]. This liberates the CB[8] to bind and activate two monomers of caspase-9. Also shown is control cage **2** that cannot be light-inactivated.

Laboratory of Chemical Biology, Department of Biomedical Engineering, Institute for Complex Molecular Systems, Eindhoven University of Technology, P. O. Box 513, 5600 MB Eindhoven, The Netherlands. E-mail: l.brunsveld@tue.nl

† Electronic supplementary information (ESI) available: Figures and tables, experimental details of peptide synthesis, protein expression, NMR, fluorometric assays, ITC. See DOI: 10.1039/d1sc01410b



report on a supramolecular caged CB[8] capable of undergoing light-triggered release using a photo-cleavable bivalent FGG-cage (Fig. 1). We illustrate its application for subsequent activation in downstream processes, such as fluorescent dye aggregation assembly and control over enzyme activity. For the light triggered CB[8] to work in unison with downstream protein effectors, the affinities were balanced such that the cage completely blocks the CB[8] prior to the stimulus, yet releases the host molecule after the stimulus for efficient binding to the proteins. Central to this design is the usage of a bivalent FGG-cage with a high affinity for CB[8] through an enhanced effective molarity<sup>24</sup> and the usage of FGG-tagged enzymes, whose affinity for dimerization is enhanced upon two-fold binding to the CB[8] platform.<sup>25</sup> In our design, two FGG-epitopes are bridged by an amino acid linker including two photo-cleavable amino-3-(2-nitrophenyl)propanoic acid (Anp) residues (1).<sup>26,27</sup> Upon irradiation with 365 nm light, the amino acid backbone cleaves, resulting in two monovalent FGG-peptide fragments that have significant lower overall affinity, releasing the CB[8] to interact with other guests such as the FGG-tagged caspase-9.

## Results and discussion

The peptide cages were synthesized using Fmoc-based solid-phase peptide synthesis. The two Anp residues are positioned within the backbone of the supramolecular cage, to ensure a high photo-cleavage yield. The second FGG-tag was introduced from a branching lysine side chain, temporarily protected *via* a 4-methoxytrityl group. After assembly of the complete backbone, this protecting group was selectively removed, followed by the subsequent installation of the two FGG-tags. The two N-terminal phenylalanines are 11 amino acids apart, giving it sufficient flexibility to encompass the CB[8] and bind the host in a bivalent manner without excessive strain on the peptide backbone (ESI Fig. S1†). As a control, also a non-cleavable variant (2) was made that has  $\beta$ -alanines instead of the Anp-residues.

The photo-cleavage of FGG-cage 1 was investigated using LC-MS, UV-Vis and <sup>1</sup>H-NMR. The peptide cage was mixed with an internal standard as reference for LC-MS quantification and samples were analyzed at different time points of exposure to irradiation with UV light (365 nm) (Fig. 2A). Within 5 minutes of UV light exposure most of cage 1 was cleaved, with a half-life of around 40 seconds. The control  $\beta$ -alanine-cage 2 remained intact, also after prolonged exposure. Gratifyingly, the presence of CB[8] had no significant influence on the cleavage rate.

The photo-cleavage of 1 was also followed by UV-Vis spectra at various time points during irradiation (Fig. 2B). Upon radiation, the characteristic peak at 260 nm of the *ortho*-nitro group rapidly disappears, while two new peaks at 230 and 300 nm appeared, indicating the successful photo-conversion. The presence of CB[8] had no significant influence on the optical properties of 1 (ESI Fig. S4†). <sup>1</sup>H-NMR spectra of 1 before and after UV radiation (Fig. 2C and ESI Fig. S5†) revealed that after 10 minutes of irradiation the signals characteristic of the Anp groups had all disappeared.




Fig. 2 Cage-cleavage by UV-light. (A) Cleavage followed over time by LC-MS of 1 and 2 (control) with an internal standard as reference. The cleavage half-life ( $\tau_{1/2}$ )  $\sim$  40 s. The presence of CB[8] has no significant influence on the cleavage rate. (B) UV-Vis absorption spectra of 1 upon UV-radiation at different time points. Inset: absorbance at 230 nm over time. (C) <sup>1</sup>H-NMR spectra of peptide cage 1 (500  $\mu$ M) before (blue) and after (red) UV-radiation in D<sub>2</sub>O. Spectrum between 5 and 3.5 ppm is omitted for clarity (see ESI Fig. S3† for full spectra).

To verify complexation of the bivalent cage 1 with CB[8], <sup>1</sup>H-NMR complex formation studies were performed. The aromatic signals belonging to the (N-terminal) phenylalanines, shifted up-field upon addition of CB[8] from 7.2 to 6.4 ppm, indicative of the de-shielding inside the CB[8] cavity for FGG-peptides (ESI Fig. S6†).<sup>5</sup> Binding studies were performed to explore whether photo-cleavage of cage 1 would result in differential binding properties between both chemical states. Initial experiments were performed using a fluorometric assay *via* the displacement of acridine orange (AO) from the cavity of CB[8]. The AO dye is



auto-quenched inside the cavity of CB[8], and becomes fluorescent again when another guest displaces it from the cavity (Fig. 3A).<sup>28</sup> The FGG-cages **1** and **2** were titrated to a solution of pre-complexed CB[8] and AO. In previous studies,<sup>28</sup> a relative high concentration of the FGG-mono peptide was required to

displace AO from the CB[8] cavity ( $IC_{50}$  62  $\mu$ M). In comparison, the bivalent FGG-cages **1** and **2** had a much stronger affinity and displayed AO with  $IC_{50}$ 's of 400 and 500 nM respectively. Upon UV irradiation of the titration plate, the affinity of the light-sensitive **1** weakened towards the value of the monovalent




Fig. 3 (A) Schematic representation of fluorometric assay. (B) Titration of bivalent cage **1** into a solution of acridine orange (1  $\mu$ M) and titration with bivalent control **2** in HBS (10 mM HEPES, pH 7.4, 150 mM NaCl) at various time points after UV-radiation, performed in triplicate; error bars represent the standard deviation. (C) Apparent  $K_D$  determined by isothermal titration calorimetry before and after photo cleavage of cage **1** and control **2**; error bars represent the confidence interval of the fit. (D) ITC trace of **1** before and after UV-radiation; (see ESI Table S1† for an overview of the isothermal titration calorimetry characteristics).



peptide (Fig. 3B on the left, blue to red). In contrast, the  $IC_{50}$  of non-cleavable **2** was not affected (Fig. 3B on the right).

Isothermal titration calorimetry (ITC) studies were performed to provide insights into the thermodynamic parameters of the supramolecular caging and light induced uncaging (Fig. 3C, D; ESI Table S1; Fig. S7 and S8†). Titration of FGG-cage **1** to CB[8] resulted in a dissociation constant ( $K_D$ ) of 22 nM. Irradiation of **1** resulted in a 100-fold weakening of the apparent  $K_D$  to 2.4  $\mu$ M. Interestingly, the  $\Delta H$  of the binding interactions before and after UV-light induced cleavage remained similar and the lowering of the  $\Delta G$  results from the change of the  $\Delta S$ . These results strongly support a binding mode that is molecularly very similar in both states (same molecular interactions), but that has changed from bivalent to monovalent.<sup>29</sup> In line with previous results, the affinity of the non-cleavable cage **2** was not affected significantly upon UV irradiation.

With the light-controllable CB[8] in hand, we investigated its utility as downstream control element on enzyme activity. Protein dimerization is a ubiquitous mechanism to regulate protein activity in a broad range of biological processes including apoptosis. Molecular control over these processes is critical to elucidate and perturb the molecular mechanisms of the proteins involved.<sup>30</sup> Caspase-9 (casp-9) is a critical cysteine protease and initiator of the apoptosis pathway and is responsible for cleaving proteins at specific aspartate residues. In the cell, caspase-9 exists primarily in its inactive monomeric form,<sup>31</sup> becoming activated only upon induced dimerization by regulatory factors as part of controlled cell death.<sup>32</sup> For example extensive UV-damage to the DNA<sup>33</sup> leads to mitochondrial release of cytochrome-C which triggers the assembly of the apoptosome that acts a scaffolding platform that recruits multiple copies of caspase-9, initiating apoptosis.<sup>34</sup> This scaffolding can be mimicked by bringing two caspase-9 proteins in close proximity by virtue of dual binding of CB[8] to N-terminal FGG-tags on the caspase-9.<sup>15</sup> The intrinsic affinity between two caspase monomers ( $K_D \sim 100 \mu$ M)<sup>35</sup> causes the caspases to act as a supramolecular bidentate.<sup>28</sup> As a consequence, the caspases primarily form active dimers, even when CB[8] is in 10-fold excess over the caspase monomers. We employed a proteolytic activity assay for caspase-9 using the synthetic substrate Ac-LEHD-AFC (where AFC is 7-amino-4-(trifluoromethyl) coumarin), that becomes fluorescent after cleavage by caspase-9 (Fig. 4A). Kinetic traces were recorded by following the fluorescence over time. The optimal concentration of CB[8] for caspase-9 activation, independent of the bivalent cage, was determined to be 10  $\mu$ M.

We screened for the optimal concentration of cage **1** that had the highest on/off ratio before and after UV cleavage (ESI Fig. S10†). We then investigated the effect of photo-cleavage of cage **1** in comparison with caspase with and without CB[8] as maximum and minimum effect (Fig. 4B). Without UV radiation the caspase-9 activity in the presence of CB[8] and cage **1** is similar to the characteristic caspase-9 background activity, around 1  $\mu$ M  $\text{min}^{-1}$  (Fig. 4C). After irradiation the activity increases dramatically to an initial slope of 11  $\mu$ M  $\text{min}^{-1}$  (the rate decreases over time due to depletion of the AFC substrate).




Fig. 4 (A) Light-triggered dimerization of caspase-9 monomers by CB[8] induces caspase-9 activity, resulting in the conversion of Ac-LEHD-AFC into a fluorescent product. (B) Time-activity traces of increasing molarity corresponding to cleavage of synthetic substrate Ac-LEHD-AFC (200  $\mu$ M) by FGG-caspase-9 (1  $\mu$ M) in 100  $\mu$ L activity assay buffer (20 mM  $\text{Na}_2\text{HPO}_4$ , 150 mM NaCl, 1 mM EDTA, 2 mM TCEP, pH 7.0, RT). 10  $\mu$ M **1** or **2** and 10  $\mu$ M CB[8] was used; the 'minimum' contains only casp-9 the 'maximum' only casp-9 and CB[8]. Measurements were performed in triplicate. (C) Initial rate ( $v_0$ ) of substrate cleavage in  $\mu\text{M min}^{-1}$ .

The UV-light activated caspase-9 reaches a comparable level activity as the CB[8]-only positive control (13  $\mu\text{M min}^{-1}$ ). As expected, reference inhibitor **2** effectively blocked caspase-9 activity, also after UV-light treatment. Even though the caspase-9 monomers feature a similar FGG-tag as the monovalent fragments of the inhibitor, the intrinsic interaction between two caspase monomers results in self-sorting and hetero complexation in the CB[8] only starts to play a more dominant role when concentrations exceed the effective molarity.<sup>36</sup>



## Conclusions

In conclusion, a versatile photo-triggered release system for CB [8] is presented, as well as its application for subsequent downstream modulation of fluorescent guest binding and binding and activation of caspase-9 enzymes. The UV-sensitive cage exhibits a 100-fold higher affinity in its intact bivalent, inhibitory mode as compared to its cleaved state. The UV-induced switching occurs in the submicromolar regime, highly favourable for supramolecular chemical biology applications, with proteins and alike that typically act in a similar concentration regime.<sup>37</sup> Light-activatable supramolecular systems are in high demand, especially those that work at low concentrations and those that can interface with biomolecules.<sup>38</sup> The caged CB[8] concept presented here offers many entries for tuning and application to the specific biomolecular system at hand, for example by varying the concentration of cage and CB[8] or by tuning of the affinities by utilizing stronger or weaker binding FGG analogues.<sup>39</sup> Similarly, the conceptual approach of a cleavable bivalent inhibitor, as presented here, can be extended to other types of triggers, which are currently being explored.

## Author contributions

P. J. d. V. and T. v. d. H. performed experiments, P. J. d. V. and L. B. conceptualized the studies, all authors contributed to writing the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This research was funded by the Netherlands Organization for Scientific Research (NWO) through Gravity program 024.001.035 and VICI grant 016.150.366. We thank Bas Rosier for the discussions about caspases.

## Notes and references

- 1 L. Zhang, Y. Wu and L. Brunsveld, *Angew. Chem., Int. Ed.*, 2007, **46**, 1798–1802.
- 2 S. Sakamoto and K. Kudo, *J. Am. Chem. Soc.*, 2008, **130**, 9574–9582.
- 3 D. Bier, R. Rose, K. Bravo-Rodriguez, M. Bartel, J. M. Ramirez-Anguita, S. Dutt, C. Wilch, F.-G. Klärner, E. Sanchez-Garcia, T. Schrader and C. Ottmann, *Nat. Chem.*, 2013, **5**, 234–239.
- 4 S. C. Zimmerman, *Beilstein J. Org. Chem.*, 2016, **12**, 125–138.
- 5 L. M. Heitmann, A. B. Taylor, P. J. Hart and A. R. Urbach, *J. Am. Chem. Soc.*, 2006, **128**, 12574–12581.
- 6 H. D. Nguyen, D. T. Dang, J. L. J. van Dongen and L. Brunsveld, *Angew. Chem., Int. Ed.*, 2010, **49**, 895–898.
- 7 D. A. Uhlenheuer, K. Petkau and L. Brunsveld, *Chem. Soc. Rev.*, 2010, **39**, 2817–2826.
- 8 R. E. McGovern, H. Fernandes, A. R. Khan, N. P. Power and P. B. Crowley, *Nat. Chem.*, 2012, **4**, 527–533.
- 9 S. Sonzini, A. Marcozzi, R. J. Gubeli, C. F. van der Walle, P. Ravn, A. Herrmann and O. A. Scherman, *Angew. Chem., Int. Ed.*, 2016, **55**, 14000–14004.
- 10 S. H. Hewitt and A. J. Wilson, *Chem. Commun.*, 2016, **52**, 9745–9756.
- 11 J. M. Chinai, A. B. Taylor, L. M. Ryno, N. D. Hargreaves, C. A. Morris, P. J. Hart and A. R. Urbach, *J. Am. Chem. Soc.*, 2011, **133**, 8810–8813.
- 12 K. L. Kim, G. Sung, J. Sim, J. Murray, M. Li, A. Lee, A. Shrinidhi, K. M. Park and K. Kim, *Nat. Commun.*, 2018, **9**, 1712–1721.
- 13 F. Guagnini, P. M. Antonik, M. L. Rennie, P. O'Byrne, A. R. Khan, R. Pinalli, E. Dalcanale and P. B. Crowley, *Angew. Chem., Int. Ed.*, 2018, **57**, 7126–7130.
- 14 L. A. Logsdon, C. L. Schardon, V. Ramalingam, S. K. Kwee and A. R. Urbach, *J. Am. Chem. Soc.*, 2011, **133**, 17087–17092.
- 15 D. T. Dang, H. D. Nguyen, M. Merckx and L. Brunsveld, *Angew. Chem., Int. Ed.*, 2013, **52**, 2915–2919.
- 16 C. Si, J. Li, Q. Luo, C. Hou, T. Pan, H. Li and J. Liu, *Chem. Commun.*, 2016, **52**, 2924–2927.
- 17 A. Gautier, C. Gauron, M. Volovitch, D. Bensimon, L. Jullien and S. Vríz, *Nat. Chem. Biol.*, 2014, **10**, 533–541.
- 18 M. J. Hansen, W. A. Velema, M. M. Lerch, W. Szymanski and B. L. Feringa, *Chem. Soc. Rev.*, 2015, **44**, 3358–3377.
- 19 C. Parente Carvalho, V. D. Uzunova, J. P. D. Silva, W. M. Nau and U. Pischel, *Chem. Commun.*, 2011, **47**, 8793–8795.
- 20 F. Tian, D. Jiao, F. Biedermann and O. A. Scherman, *Nat. Commun.*, 2012, **3**, 1207.
- 21 M. A. Romero, N. Basilio, A. J. Moro, M. Domingues, J. A. González-Delgado, J. F. Arteaga and U. Pischel, *Chem.–Eur. J.*, 2017, **23**, 13105–13111.
- 22 P. Ferreira, B. Ventura, A. Barbieri, J. P. Da Silva, C. A. T. Laia, A. J. Parola and N. Basilio, *Chem.–Eur. J.*, 2019, **25**, 3477–3482.
- 23 P. Remón, D. González, S. Li, N. Basilio, J. Andréasson and U. Pischel, *Chem. Commun.*, 2019, **55**, 4335–4338.
- 24 M. Ramaekers, S. P. W. Wijnands, J. L. J. van Dongen, L. Brunsveld and P. Y. W. Dankers, *Chem. Commun.*, 2015, **51**, 3147–3150.
- 25 R. P. G. Bosmans, J. M. Briels, L.-G. Milroy, T. F. A. de Greef, M. Merckx and L. Brunsveld, *Angew. Chem., Int. Ed.*, 2016, **55**, 8899–8903.
- 26 E. Profft and F.-J. Becker, *J. Prakt. Chem.*, 1965, **30**, 18–38.
- 27 B. B. Brown, D. S. Wagner and H. M. Geysen, *Mol. Diversity*, 1995, **1**, 4–12.
- 28 P. J. de Vink, J. M. Briels, T. Schrader, L.-G. Milroy, L. Brunsveld and C. Ottmann, *Angew. Chem., Int. Ed.*, 2017, **56**, 8998–9002.
- 29 G. Wu, M. Olesińska, Y. Wu, D. Matak-Vinkovic and O. A. Scherman, *J. Am. Chem. Soc.*, 2017, **139**, 3202–3208.
- 30 J. D. Klemm, S. L. Schreiber and G. R. Crabtree, *Annu. Rev. Immunol.*, 1998, **16**, 569–592.
- 31 S. J. Riedl and G. S. Salvesen, *Nat. Rev. Mol. Cell Biol.*, 2007, **8**, 405–413.



- 32 B. J. H. M. Rosier, A. J. Markvoort, B. Gumí Audenis, J. A. L. Roodhuizen, A. den Hamer, L. Brunsveld and T. F. A. de Greef, *Nat. Catal.*, 2020, **3**, 295–306.
- 33 L. A. Sitailo, S. S. Tibudan and M. F. Denning, *J. Biol. Chem.*, 2002, **277**, 19346–19352.
- 34 P. Li, L. Zhou, T. Zhao, X. Liu, P. Zhang, Y. Liu, X. Zheng and Q. Li, *Oncotarget*, 2017, **8**, 23996–24008.
- 35 M. Rénatus, H. R. Stennicke, F. L. Scott, R. C. Liddington and G. S. Salvesen, *Proc. Natl. Acad. Sci. U. S. A.*, 2001, **98**, 14250–14255.
- 36 S. Di Stefano and G. Ercolani, in *Advances in Physical Organic Chemistry*, ed. I. H. Williams and N. H. Williams, Academic Press, 2016, vol. 50, pp. 1–76.
- 37 L.-G. Milroy, T. N. Grossmann, S. Hennig, L. Brunsveld and C. Ottmann, *Chem. Rev.*, 2014, **114**, 4695–4748.
- 38 O. Dagliyan and K. M. Hahn, *Curr. Opin. Struct. Biol.*, 2019, **57**, 17–22.
- 39 A. R. Urbach and V. Ramalingam, *Isr. J. Chem.*, 2011, **51**, 664–678.

