


 Cite this: *RSC Adv.*, 2022, 12, 25587

 Received 16th July 2022
Accepted 1st September 2022

DOI: 10.1039/d2ra04387d

rsc.li/rsc-advances

Aromatic fluorine atom-induced highly amine-sensitive trimethine cyanine dye showing colorimetric and ratiometric fluorescence change†

 Ryunosuke Kani,^a Yasuhiro Kubota,^a Toshiyasu Inuzuka^b
and Kazumasa Funabiki^{*,a}

Herein, introducing multiple fluorine atoms into aromatic rings of trimethine cyanine dyes is proposed as a powerful method for dramatically increasing sensitivity to amines. The highly sensitive ratiometric fluorescence properties previously available only by intramolecular addition can be exploited in reactions with intermolecular amines or other nucleophiles.

Introduction

Ratiometric fluorescence, a method wherein the ratio of fluorescence intensities at two or more wavelengths are used, has attracted much attention in analytical sensing and optical imaging because it can reduce the effects of many factors such as dye concentration, surrounding pH, polarity, and changes in environmental temperature compared to when single fluorescence intensity is measured.^{1–4}

Amines could cause serious damage in public health and the environment owing to their toxic nature, so, simple, fast, low-cost, and accurate amine monitoring systems are becoming highly urgent.⁵ The detection of amines using ratiometric fluorescence through utilized intramolecular charge transfer (ICT),^{6–17} twisted ICT,¹⁸ aggregation-induced emission,^{19–23} and excited state intramolecular proton transfer²⁴ has been reported.

Polymethine cyanine dyes have several features such as good fluorescence quantum yield and the ability to easily prepare the fluorescence wavelength by changing the methine chain length.^{25–27} However, they lack reactivity with amines, and the reported examples are limited to reactions with intramolecular amines.^{28–34} Herein, we report that the prepared ring-perfluorinated trimethine cyanine dye **2a** (Fig. 1) has a significantly higher response to *n*-hexylamine than the non-fluorinated dye **2b**. The dye **2a** exhibited a dual change in the solution and fluorescence color at widely shifted wavelengths, visible to the naked eye.

Furthermore, dye **2a** adsorbed on the filter paper could be repeatedly used because the orange emission was instantly converted into blue by the amine vapor and returned to its original color after drying rapidly in the air. These results are the first example of the highly sensitive ratiometric fluorescence properties of polymethine cyanine dyes with intermolecular amines and advance the development of rapid and reversible vapochromic response materials to amines with two changes in papers and emission color.

Results and discussion

The responsivity of the ring-perfluorinated trimethine cyanine dyes **2a** and **2b** to *n*-hexylamine in tetrahydrofuran (THF) stored overnight in the dark at 25 °C were investigated; the results are summarized in Fig. 2 and Table 1. *n*-Hexylamine was used as a representative example of a primary aliphatic amine because it is not easily volatilized. As the amount of *n*-hexylamine gradually increased, the orange fluorescence at 596 nm from dye **2a** decreased, and the blue emission at 404 nm increased at an excitation of 257 nm. This phenomenon led to the disappearance of the solution color and a shift in the fluorescence color from orange through pink to blue, according to the CIE color system. Thus, dye **2a** was highly sensitive, completing the reaction with only 2 equiv. of the amine. For dye **2b**, the red fluorescence at 612 nm diminished, and the blue fluorescence




Fig. 1 Structure of trimethine cyanine dyes **2a** and **2b**.

^aDepartment of Chemistry and Biomolecular Science, Gifu University, 1-1 Yanagido, Gifu 501-1193, Japan

^bDivision of Instrumental Analysis, Life Science Research Centre, Gifu University, 1-1 Yanagido, Gifu 501-1193, Japan

† Electronic supplementary information (ESI) available. See <https://doi.org/10.1039/d2ra04387d>






Fig. 2 (a) Ultraviolet-visible (UV-vis), (b) fluorescence spectra, and (c) CIE color systems of dye **2a** (5×10^{-6} M) in THF stored overnight in the dark at 25 °C after adding *n*-hexylamine. (d) UV-vis, (e) fluorescence spectra, and (f) CIE color systems of dye **2b** under similar conditions. The excitation wavelengths were determined by the excitation spectra (Fig. S2†) and were as follows: 257 nm (dye **2a**), 250 nm (dye **2b**), and 365 nm (CIE color systems). Inset photos were obtained under daylight and UV light ($\lambda = 365$ nm), and the order of amine concentration is left to right.

at 380 nm increased significantly at an excitation of 250 nm, although 6400 equiv. of the amine was required to complete the reaction with the non-fluorinated dye **2b**. These results suggest that replacing all the hydrogen atoms with fluorine on the aromatic rings dramatically improved the sensitivity of trime-thine cyanine dye to the amine.

The fluorescence quantum yields for dyes **2a** and **2b** were 14 and 10%, respectively. When the dyes reacted with the amine, the maximum excitation wavelength overlapped with the absorption band of THF; therefore, excluding this band, the dyes were excited at 290 nm, with fluorescence quantum yields of 4 and 24%, respectively. Furthermore, a calibration curve for the fluorescence intensity ratio of dye **2a** ($I_{404 \text{ nm}}/I_{596 \text{ nm}}$) versus the concentration of *n*-hexylamine demonstrated good linearity ($R^2 = 0.988$), while the estimated detection limit ($3\sigma/\text{slope}$) was 17 nM, as shown in Fig. 3. The obtained value shows sensitivity as high as that of an activated ester and aldehydes.^{8,10,11}

Next, absorbance changes in various solvents were followed to examine the sensitivity of dye **2a** to *n*-hexylamine (Fig. S1†). The results revealed that this reaction was primarily promoted by THF, followed by other solvents in the order dimethyl sulfoxide (DMSO), methyl alcohol (MeOH), acetone, chloroform, and dichloromethane (DCM). The large difference in the reactivity was related to the Gutmann donor number (DN) and acceptor number (AN) of solvent molecules, which are the empirical measures of the nucleophilic and electrophilic properties of the solvents.^{35,36}

Solvents with high DN can stabilize cations,³⁷ while an opposite phenomenon is observed for AN. Therefore, the reaction of the dye with the amine is more likely to proceed in THF and DMSO, which have larger DN and/or smaller AN than acetone, MeOH, chloroform, and DCM, because the cationic dye is most involved in this system. The DN and AN of these solvents are listed in Table S1.†

Table 1 Optical properties of dyes **2a** and **2b** in THF with *n*-hexylamine

Dye ^a	<i>n</i> -Hexylamine (equiv.)	λ_{max} (nm) (ϵ (M ⁻¹ cm ⁻¹))	λ_{ex} (nm)	F_{max} (nm) ^b	$\phi_{\text{f}}^{\text{c}}$	τ_{s} (ns) ^e
2a	0	579 (120 000)	580	596	0.14	0.7
	0.5	579 (91 000)	—	—	—	—
	1	579 (30 000)	—	—	—	—
	2	579 (200)	257	404	0.04 ^d	6.1
2b	0	591 (110 000)	591	612	0.10	0.5
	6400	254	250	380	0.24	12.5

^a Measured dye **2** (5×10^{-6} M) in THF stored overnight in the dark at 25 °C after adding *n*-hexylamine. ^b Measured using excitation wavelength (λ_{ex}) at low sensitivity. ^c Measured using an integrating sphere method excited at λ_{ex} . ^d Measured by excitation at 290 nm. ^e Measured using a single-photon-counting method.






Fig. 3 Calibration curve for the fluorescence intensity ratio ($I_{404 \text{ nm}}/I_{596 \text{ nm}}$) of dye **2a** vs. the concentration of *n*-hexylamine ($0-4 \times 10^{-6}$ M).

Other analytes, such as not only other amines (aniline, triethylamine, and diethylamine) but also anions (perchlorate, acetate, bromide, chloride, and fluoride) in THF were also examined. The results are summarized in Fig. 4. The reactions of dye **2a** with 2 equiv. of amines such as triethylamine and diethylamine occurred smoothly, and the color of the fluorescent changed from orange to blue, as in the case of *n*-hexylamine, as illustrated in Fig. 4(c). Although, accurate measurement for aniline was not possible due to its fluorescence, the dye-derived maximum absorption was slightly reduced.

Using 2 equiv. of a couple of anions, such as acetate and fluoride,³⁸ also resulted in observation to change the fluorescent color from orange to blue, as in the case of *n*-hexylamine, as

illustrated in Fig. 4(c). The detection of other anionic species, such as bromide and chloride, demonstrated only moderate changes in the fluorescent color and reacted slightly with perchlorate.

To observe the difference in the detection of these neutral and anionic nucleophiles, UV-vis and fluorescence spectra of dye **2a** were measured in detail in THF with different equiv. of not only diethylamine and triethylamine but also acetate and fluoride anions, were carried out. The results are summarized in Fig. 5. It was found that the highly nucleophilic anion, such as acetate and fluoride are more reactive than *n*-hexylamine. Detection of diethylamine appears to be similar to that of *n*-hexylamine, but somewhat lower for the tertiary triethylamine.

Importantly, the ring-perfluorinated trimethine cyanine dye **2a** has a significantly higher response to very small amounts of neutral amines than the non-fluorinated dye **2b**, and underwent a dual change in the solution and fluorescent. The reason can seem to be that the introduction of fluorine atoms into the aromatic ring significantly lowers the LUMO of trimethine cyanine dye **2a** and promotes the addition reaction of even neutral amines, as shown in Fig. 6.³⁹

For the reaction mechanism, the results of nuclear magnetic resonance (NMR) and high-resolution mass spectrometry illustrated in Fig. 7, S3, and S4† suggest the 1,2-addition of *n*-hexylamine to the iminium carbon of dye **2a**. In particular, ¹H NMR confirmed that the doublet signal split in two and the triplet signal changed to a double doublet, along with an up-field shift in all signals regarding the methine chain protons. Furthermore, the amine-added dyes **2a** and **2b** were characterized by MS analysis (Fig. S4†), where peaks at m/z 774.2719 (calculated = 774.2718, $[C_{39}H_{36}F_{12}N_3]^+$) and m/z 558.3849




Fig. 4 (a) UV-vis, (b) fluorescence spectra, and (c) the photos of dye **2a** (5×10^{-6} M) in THF stored overnight in the dark at 25°C after adding various analytes (2 equiv.). The excitation wavelength was 257 nm. (d) Absorbance at 579 nm and (e) fluorescence intensity ratio ($I_{404 \text{ nm}}/I_{596 \text{ nm}}$) were calculated from the UV-vis and fluorescence spectra, respectively. Analytes are as follows: (1) **2a** only, (2) ClO_4^- , (3) AcO^- , (4) Br^- , (5) Cl^- , (6) F^- , (7) PhNH_2 , (8) NEt_3 , (9) NHEt_2 , and (10) *n*-hexylamine. Photos were captured under daylight and UV light ($\lambda = 365 \text{ nm}$) with various analytes. The counter cations of these anionic species are all Bu_4N^+ .




Fig. 5 UV-vis and fluorescence spectra of dye **2a** (5×10^{-6} M) in THF stored overnight in the dark at 25 °C after adding analytes which are as follows: (a) AcO^- , (b) F^- , (c) NEt_3 , and (d) NHEt_2 . The counter cations of these anionic species are all Bu_4N^+ .

(calculated = 558.3848, $[\text{C}_{39}\text{H}_{48}\text{N}_3]^+$) were detected, respectively. Although it was difficult to measure the non-fluorinated dye **2b** by NMR owing to its poor reactivity with amines, it is expected to react with amines in the manner similar to that as the fluorinated dye **2a** because of its very similar structure and the MS result.

In order to assess the versatility of application, the repeatability of dye **2a** was investigated using filter paper, as shown in Fig. 8.

The filter paper for dye adsorption was fabricated by soaking a white filter paper overnight in the DCM solution of dye **2a** (5×10^{-4} M) and then drying in the dark for 1 h at 25 °C. Exposure to *n*-hexylamine vapor resulted in rapid disappearance of the pink of the filter paper and change in the fluorescence from orange to blue. Subsequently, drying in the air caused the pink of the filter paper to return to pink and emit orange fluorescence. This reversible change in fluorescent color was particularly

pronounced, with the blue fluorescence reverting to orange in only 5 s when kept in the air after exposure to the amine (see Video†).

Conclusions

We reported that the ring-perfluorinated trimethine cyanine dye **2a** was highly sensitive to *n*-hexylamine and underwent a dual change in the solution and fluorescent color with a large shift. The methodology of introducing many fluorine atoms proved valuable for efficiently enhancing the sensitivity of the dye to amines. In addition, the fluorinated dye **2a** adsorbed on the filter paper exhibited reversible ratiometric fluorescence properties, such that the orange emission was instantly changed to blue by the amine vapor and returned to its original color when dried in the air for 5 s.






Fig. 6 Energy diagrams and molecular orbitals of the cationic parts of dyes obtained using DFT calculations at the B3LYP/6-31+G(d,p) level.

Experimental

Materials

Acetone, methanol, and ethyl acetate were purchased from Kanto Chemical Industry Co., Inc. Tetrahydrofuran (THF; stabilizers not included), dimethyl sulfoxide, chloroform, triethylamine, aniline, and tetrabutylammonium chloride were purchased from FUJIFILM Wako Pure Chemical Co. *n*-Hexylamine, diethylamine, tetrabutylammonium fluoride, and tetrabutylammonium acetate were purchased from TCI Co., Ltd. Tetrabutylammonium bromide was purchased from Sigma-Aldrich. Tetrabutylammonium perchlorate and dichloromethane were purchased from Nacalai Tesque Inc. Dyes **2a** and **2b** were synthesized by our method.⁴⁰

Measurement

¹H nuclear magnetic resonance (NMR) spectra were measured at 392 or 400 MHz in hexadeuterodimethyl sulfoxide ((CD₃)₂SO) solutions with residual solvents as internal standards using




Fig. 7 ¹H NMR analysis for the reaction of dye **2a** (1×10^{-5} mol) with *n*-hexylamine (20 equiv.) in DMSO-*d*₆.




Fig. 8 Repeatability of fluorescence using filter paper adsorbed with dye **2a** upon alternative exposure to *n*-hexylamine vapor and drying in the air. The inset photos were captured under UV light ($\lambda = 365$ nm).

a JEOL ECS-400 or ECX-400P FT-NMR spectrometer. ¹³C NMR spectra were obtained at 99 or 101 MHz in (CD₃)₂SO solution with residual solvents as internal standards using a JEOL ECS-400 or ECX-400P FT-NMR spectrometer. ¹⁹F NMR spectra were recorded at 369 or 376 MHz in (CD₃)₂SO solutions using a JEOL ECS-400 or ECX-400P FT-NMR spectrometer. The data were reported as (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br s = broad singlet, coupling constant(s), integration). Electrospray ionization mass spectroscopy (ESI-MS) analysis using MeOH was performed with a JEOL JMS-T100LP (AccuTOF LC-plus). Ultraviolet-visible absorption spectra were obtained using a Hitachi U-4100. The fluorescence spectra were acquired using a Jasco FP-8600 spectrofluorometer. The absolute fluorescence quantum yields were obtained on Hamamatsu Quantaurus-QY C11347-01. The fluorescence lifetimes were measured using Hamamatsu Quantaurus-Tau compact fluorescence lifetime spectrometer C11367-01.

Determination of limit of detection (LOD)

The fluorescence intensity ratio at 404 nm and 596 nm of the ring-perfluorinated dye **2a** in THF was collected 10 times to obtain the background noise σ . The fluorescence intensity ratios were measured overnight after adding different concentrations of *n*-hexylamine. LOD ($3\sigma/\text{slope}$) was calculated as three times the background noise σ divided by the slope of the fluorescence intensity ratio fitted to a calibration curve as a function of the amine concentration.

Author contributions

KF and RK conceived and designed the project. RK performed all experiments, analysed the data, and wrote the manuscript. YK and TI contributed to useful discussions. KF reviewed and modified the manuscript.

Conflicts of interest

There are no conflicts to declare.



Acknowledgements

This work is supported by the “Interdisciplinary Frontier Next-Generation Researcher Program of the Tokai Higher Education and Research System” to RK from JST SPRING, grant number JPMJSP2125. This work was partially supported by JSPS KAKENHI grant numbers JP20K05647 to KF. We would like to thank Editage (www.editage.com) for English language editing.

Notes and references

- 1 R. Gui, H. Jin, X. Bu, Y. Fu, Z. Wang and Q. Liu, *Coord. Chem. Rev.*, 2019, **383**, 82–103.
- 2 M. H. Lee, J. S. Kim and J. L. Sessler, *Chem. Soc. Rev.*, 2015, **44**, 4185–4191.
- 3 J. Fan, M. Hu, P. Zhan and X. Peng, *Chem. Soc. Rev.*, 2013, **42**, 29–43.
- 4 T. Doussineau, A. Schulz, A. Lapresta-Fernandez, A. Moro, S. Korsten, S. Trupp and G. J. Mohr, *Chem.–Eur. J.*, 2010, **16**, 10290–10299.
- 5 L. Wang, X. Ran, H. Tang and D. Cao, *Dyes Pigm.*, 2021, **194**, 109634.
- 6 S. Zhang, Z. Li, B. Zhang, F. Dong, B. Han, J. Lv, Y. Sun, H. Lu, Y. Yang and H. Ma, *New J. Chem.*, 2021, **45**, 20441–20446.
- 7 Z. Quan, H. He, H. Zhou, Y. Liang, L. Wang, S. Tian, H. Zhu and S. Wang, *Sens. Actuators, B*, 2021, **333**, 129535.
- 8 S. Sawminathan, S. Munusamy, D. Jothi and S. K. Iyer, *ChemistrySelect*, 2021, **6**, 858–864.
- 9 H. Nawaz, J. Zhang, W. Tian, K. Jin, R. Jia, T. Yang and J. Zhang, *J. Hazard. Mater.*, 2020, **387**, 121719.
- 10 M. Saravanakumar, B. Umamahesh, R. Selvakumar, J. Dhanapal, S. K. A. Kumar and K. I. Sathiyarayanan, *Dyes Pigm.*, 2020, **178**, 108347.
- 11 S. Jeon, T.-I. Kim, H. Jin, U. Lee, J. Bae, J. Bouffard and Y. Kim, *J. Am. Chem. Soc.*, 2020, **142**, 9231–9239.
- 12 R. Jia, W. Tian, H. Bai, J. Zhang, S. Wang and J. Zhang, *Nat. Commun.*, 2019, **10**, 795.
- 13 D. Kim, S. W. Cho, Y. W. Jun and K. H. Ahn, *Bull. Korean Chem. Soc.*, 2017, **38**, 995–996.
- 14 S. Mallick, F. Chandra and A. L. Koner, *Analyst*, 2016, **141**, 827–831.
- 15 D. Meng, S. Yang, D. Sun, Y. Zeng, J. Sun, Y. Li, S. Yan, Y. Huang, C. W. Bielawski and J. Geng, *Chem. Sci.*, 2014, **5**, 3130–3134.
- 16 J.-T. Hou, K. Li, B.-Y. Liu, Y.-X. Liao and X.-Q. Yu, *Tetrahedron*, 2013, **69**, 2118–2123.
- 17 S.-i. Sasaki, Y. Kotegawa and H. Tamiaki, *Tetrahedron Lett.*, 2006, **47**, 4849–4852.
- 18 T. Han, W. Wei, J. Yuan, Y. Duan, Y. Li, L. Hu and Y. Dong, *Talanta*, 2016, **150**, 104–112.
- 19 M. Kataria, J. Hwang, H. D. Chau, N. Y. Kwon, D. W. Lee, T. Kim, K. Lee, Y. Kim, S. Park and D. H. Choi, *Sens. Actuators, B*, 2021, **343**, 130110.
- 20 R. Roy, N. R. Sajeev, V. Sharma and A. L. Koner, *ACS Appl. Mater. Interfaces*, 2019, **11**, 47207–47217.
- 21 S. Pramanik, H. Deol, V. Bhalla and M. Kumar, *ACS Appl. Mater. Interfaces*, 2018, **10**, 12112–12123.
- 22 S. Lohar, D. A. Safin, A. Sengupta, A. Chattopadhyay, J. S. Matalobos, M. G. Babashkina, K. Robeyns, M. P. Mitoraj, P. Kubisiak, Y. Garcia and D. Das, *Chem. Commun.*, 2015, **51**, 8536–8539.
- 23 A. Ghosh, A. Sengupta, A. Chattopadhyay and D. Das, *Chem. Commun.*, 2015, **51**, 11455–11458.
- 24 S. Zheng, Y. Fang, Y. Chen, Q. Kong, F. Wang and X. Chen, *Spectrochim. Acta, Part A*, 2022, **267**, 120616.
- 25 K. Ilina and M. Henary, *Chem. - Eur. J.*, 2021, **27**, 4230–4248.
- 26 W. Sun, S. Guo, C. Hu, J. Fan and X. Peng, *Chem. Rev.*, 2016, **116**, 7768–7817.
- 27 A. Mishra, R. K. Behera, P. K. Behera, B. K. Mishra and G. B. Behera, *Chem. Rev.*, 2000, **100**, 1973–2012.
- 28 A. Mukherjee, P. C. Saha, R. S. Das, T. Bera and S. Guha, *ACS Sens.*, 2021, **6**, 2141–2146.
- 29 H. Mu, K. Miki, H. Harada, K. Tanaka, K. Nogita and K. Ohe, *ACS Sens.*, 2021, **6**, 123–129.
- 30 M. Oe, K. Miki and K. Ohe, *Org. Biomol. Chem.*, 2020, **18**, 8620–8624.
- 31 M. Oe, K. Miki, H. Mu, H. Harada, A. Morinibu and K. Ohe, *Tetrahedron Lett.*, 2018, **59**, 3317–3321.
- 32 H. Mu, K. Miki, Y. Takahashi, N. Teshima, M. Oe, K. Kojima and K. Ohe, *Chem. Lett.*, 2018, **47**, 1147–1150.
- 33 K. Miki, K. Kojima, K. Oride, H. Harada, A. Morinibu and K. Ohe, *Chem. Commun.*, 2017, **53**, 7792–7795.
- 34 L. He, W. Lin, Q. Xu, M. Ren, H. Wei and J.-Y. Wang, *Chem. Sci.*, 2015, **6**, 4530–4536.
- 35 E. V. Vetrova, N. I. Makarova, N. I. Omelichkin, L. G. Minyaeva, V. V. Chernyavina, R. N. Borisenko and A. V. Metelitsa, *J. Photochem. Photobiol., A*, 2019, **370**, 127–134.
- 36 V. Gutmann, *The Donor-Acceptor Approach to Molecular Interactions*, Springer, New York, 1978.
- 37 E. A. Ribeiro, T. Sidooski, L. G. Nandi and V. G. Machado, *Spectrochim. Acta, Part A*, 2011, **81**, 745–753.
- 38 There is a report where an anionic fluoride ion can be detected *via* addition to non-fluorinated cyanine dye, see Y. Wang, S.-Y. Gwon, Y.-A. Son and S.-H. Kim, *Mol. Cryst. Liq. Cryst.*, 2012, **566**, 61–62.
- 39 Introduction CN groups also effective for detection of amines *via* addition to the fluorescent dyes, see Z. Li, R. Toivola, J. Yang, P.-N. Lai, T. Howie, S.-H. Jang, B. D. Flinn and A. K.-Y. Jen, *Adv. Mater.*, 2016, **28**, 6592–6597.
- 40 K. Funabiki, Y. Saito, T. Kikuchi, K. Yagi, Y. Kubota, T. Inuzuka, Y. Miwa, M. Yoshida, O. Sakurada and S. Kutsumizu, *J. Org. Chem.*, 2019, **84**, 4372–4380.

