



Cite this: *RSC Adv.*, 2022, 12, 33584

Received 12th September 2022

Accepted 9th November 2022

DOI: 10.1039/d2ra05739e

rsc.li/rsc-advances

B(C₆F₅)₃-catalyzed oxidation of α -diazoesters using DMF and molecular oxygen as oxygen sources†

Bei Wang,^{†abc} Guo-Min Zhang,^{†abc} Hua Zhang^{abc} and Ji-Yu Wang^{ID*ab}


A metal-free catalytic oxidation of α -diazoesters via a green environmental-friendly route was developed. The α -diazoesters were converted to α -ketoesters using DMF and molecular oxygen as oxygen sources and B(C₆F₅)₃ as the catalyst, without any additives. This protocol has a broad adaptability of substrates and good compatibility with a range of functional groups, and it offers new insight into reactions catalyzed by B(C₆F₅)₃.

Introduction


α -Diazoesters are an important class of organic synthesis synthon.¹ Diazo compounds can form metal carbenes by transition metal catalysis, and the reactions of insertion or cyclopropanation based on metal carbenes have been developed in the past few decades.² The oxidation reaction of α -diazoesters can generate α -ketoesters, which usually have biological activity, and can also transform into a variety of functional groups.³ Various methods for this oxidation have been reported, although many require the use of harsh oxidants like dimethyldioxirane (DMDO) or *t*-BuOCl,⁴ or expensive transition metal catalysts like Rh.⁵ In 2016, Stoltz's group found that dimethyl sulfoxide (DMSO) could also serve as an oxidant to achieve oxidation of aryl α -diazoesters (Fig. 1a).⁶ But the method applied only to the electron-rich diazo compounds. According to green and sustainable chemistry principles, molecular oxygen is considered to be an ideal oxidant due to its natural, inexpensive, and environmentally friendly nature, and therefore it has attractive academic and industrial prospects.⁷ The reactions of preformed stable metal carbene compounds with oxygen are known.⁸ In 2021, Xu *et al.* reported a highly efficient and catalytic procedure for the aerobic oxidation of α -diazoesters to α -ketoesters via a copper carbene intermediate (Fig. 1a).⁹ In addition, Liu *et al.* used cheap, readily available Eosin Y as a photocatalyst and O₂ (air) as a green oxidant, and achieved the aerobic oxidation of α -diazoesters under visible light in air at room temperature (Fig. 1a).¹⁰ Research on the aerobic oxidation of α -diazoesters especially by non-metal catalysts still has space for exploration.

In recent years, tris(pentafluorophenyl)borane has been widely used in the reduction or addition reactions of unsaturated compounds, and in other reactions.¹¹ However, few efforts have been devoted to B(C₆F₅)₃-catalyzed oxidation reactions. For instance, the borane-mediated hydride abstraction of amines results in the generation of reactive iminium hydridoborate salts that participate in a variety of stoichiometric and catalytic processes.¹² In 2021, Basak *et al.* provided a detailed overview of the borane-mediated dehydrogenation functionalization of amine compounds.¹³ Moreover, in 2017, Babu's group reported an efficient one-pot oxidative esterification and amidation of aldehydes using B(C₆F₅)₃ as the catalyst and TBHP as the oxidant.¹⁴ In 2018, Ling *et al.* reported a B(C₆F₅)₃-catalyzed

a. Synthesis of α -ketoesters via the oxidation of α -diazoesters



b. Boron-Catalyzed Oxidation of α -Diazoesters



c. This work




Fig. 1 The oxidation reactions of α -diazoesters.

^aDepartment of Chemistry, Asymmetric Synthesis and Chiral Technology Key Laboratory of Sichuan Province, Xihua University, Chengdu 610041, P. R. China

^bChengdu Institute of Organic Chemistry, Chinese Academy of Sciences, Chengdu 610041, P. R. China

^cUniversity of Chinese Academy of Sciences, Beijing 100049, P. R. China

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2ra05739e>

‡ B. Wang and G.-M. Zhang contributed equally to this work.




oxidative deamination/cyclization cascade reaction of benzyl amines and ketones for the synthesis of 2,4,6-triarylpyridines in an oxygen atmosphere.¹⁵ In addition, several reports have been published on the $B(C_6F_5)_3$ -catalyzed activation of diazo compounds;¹⁶ for example, Tang's group reported a boron-catalyzed O–H bond insertion of α -aryl α -diazoesters in water.¹⁷ In previous works, our group has also conducted research on $B(C_6F_5)_3$ -catalyzed reactions and achieved some results.¹⁸ During our research, we found that $B(C_6F_5)_3$ could catalyze the oxidation of α -diazoesters under an O_2 atmosphere. Subsequently, Zhou's group described a $B(C_6F_5)_3$ -catalyzed oxidation reaction of α -diazoesters with DMSO as an oxygen source (Fig. 1b).¹⁹ But, different to this work, we report a $B(C_6F_5)_3$ -catalyzed oxidation of α -diazoesters to obtain α -ketoesters by activating O_2 directly (Fig. 1c). And this protocol features a wide substrate scope including aromatic heterocycle α -diazoesters and good functional group tolerance.

Results and discussion


The ethyl phenyldiazoacetate **1a** was selected as the model substrate to investigate the oxidation reaction under various reaction conditions (Table 1). Initially, the reaction was carried out using different catalysts at 100 °C for 2 hours in DMF in an oxygen atmosphere. First, only a small amount of oxidation product **2-1a** was obtained without any catalyst (entry 1). To our delight, $B(C_6F_5)_3$ catalyzed the oxidation reaction and the yield of **2-1a** increased to 57% (entry 2). 2,4,6- BAr^F , 3,4,5- BAr^F , $Cu(OAc)_2$ and $CoCl_2$ also had certain catalytic effects on the

reaction, but the results were not as good as those with $B(C_6F_5)_3$ (entries 3–6). Furthermore, the catalytic effects of metal Lewis acid catalysts with weaker acidity such as $La(OTf)_3$ or $Sc(OTf)_3$ were worse (entries 7 and 8). The byproduct ethyl mandelate was reduced when using dry DMF as solvent (entry 9). Next, we tried to increase the amount of catalyst. When the amount of catalyst was increased to 20 mol%, the yield of **2-1a** was 86% (entries 10–12). Then, the temperature was adjusted to 80 °C, and the yield of the expected product decreased significantly (entry 13). Further screening of solvents showed that the type of solvent had effects on the reaction. When the reaction was performed in CH_3CN , THF or 1, 4-dioxane, the yields of **2-1a** were very low (entries 14–16). Finally, the best conditions for the reaction were determined: the reaction was run with **1a** (0.3 mmol) and

Table 1 Optimization of reaction conditions^{a,b}

					
Entry	Catalyst	Solvent	Catalyst (x mol%)	2-1a (%)	2-2a (%)
1	—	DMF	5	5	Trace
2	$B(C_6F_5)_3$	DMF	5	57	20
3	2,4,6- BAr^F	DMF	5	50	18
4	3,4,5- BAr^F	DMF	5	47	22
5	$Cu(OAc)_2$	DMF	5	42	18
6	$CoCl_2$	DMF	5	39	20
7	$La(OTf)_3$	DMF	5	20	25
8	$Sc(OTf)_3$	DMF	5	24	23
9 ^c	$B(C_6F_5)_3$	DMF	5	68	7
10 ^c	$B(C_6F_5)_3$	DMF	10	63	4
11 ^c	$B(C_6F_5)_3$	DMF	15	70	6
12 ^c	$B(C_6F_5)_3$	DMF	20	86	1
13 ^{c,d}	$B(C_6F_5)_3$	DMF	20	52	7
14 ^c	$B(C_6F_5)_3$	CH_3CN	20	15	12
15 ^c	$B(C_6F_5)_3$	THF	20	19	13
16 ^c	$B(C_6F_5)_3$	1,4-Dioxane	20	17	16

^a Reaction conditions: **1a** (54.0 mg, 0.3 mmol) and catalyst (x mol%) in solvent (1 mL) for 2 h at T °C. ^b Yield of GC. ^c The dry solvent. ^d The reaction temperature was 80 °C.



Scheme 1 Substrate scope for α -diazoesters. Reaction conditions: **1** (0.3 mmol, 1 equiv.), $B(C_6F_5)_3$ (20 mol%), and DMF (1 mL) at 100 °C for 2 h. Yield of isolated product. Scale-up reaction conditions: **1** (10 mmol, 1.9 g, 1 equiv.), $B(C_6F_5)_3$ (20 mol%), and DMF (30 mL) at 100 °C for 2 h. Yield of isolated product.

$\text{B}(\text{C}_6\text{F}_5)_3$ (20 mol%) under an oxygen atmosphere (balloon) in 1 mL of dry DMF at 100 °C for 2 hours.

Next, using the optimized conditions, the applicability of the protocol for a broad range of substrates was explored, as shown in Scheme 1. Generally, phenyl α -diazooesters with alkyl, benzyl, allyl, or propargyl substituents at R_2 all afforded moderate to high yields of the desired α -ketoesters (**2-1a–2-1h**). The position and electrical properties of the substituents on the benzene ring have a great influence on the reaction. With substituents at the *ortho* position, the yield of products substantially declined, and the decrease is more obvious when the methyl group was substituted (**2-1i–2-1j**). This is probably due to steric hindrance and the donating property of the group, because the carbene intermediate that formed during the reaction was unstable. The yield of the products was moderate when a *meta* substituent was located on the benzene ring (**2-1k–2-1l**). The yield of the products was lower when the benzene ring had an electron donating substituent at the *para* position (**2-1m–2-1n**), and the yield of the products was moderate to good with halogen substituents (**2-1o–2-1q**). Multisubstituted aryl α -diazooesters such as ethyl 2-diazo-2-(2,3-dichlorophenyl)acetate could be converted to the corresponding oxidation product **2-1r** with a yield of 48%. Unsaturated substituents, such as alkyne groups, also tolerated the reaction conditions (**2-1s**). In the case of polycyclic substrates, such as naphthalene and 2-anthraquinone, the desired products **2-1t** and **2-1u** could also be formed in good yields; the oxidation product of ethyl-2-(benzo[d][1,3]dioxol-5-yl)-2-diazoacetate could be obtained with a yield of 40% (**2-1v**). In addition, the substrate bearing a quinoline moiety also gave the corresponding product **2-1w**, albeit in a slightly diminished yield. Aryl α -diazooesters containing tocopherol moieties could also smoothly undergo the reaction to give the corresponding product with a yield of 43% (**2-1x**). Finally, diphenyl-diazomethane was converted to benzophenone in a moderate yield (**2-1y**) and an oxindole scaffold was compatible with the reaction, allowing for the formation of isatin in 48% yield (**2-1z**).

Subsequently, we increased the amount of ethyl phenyl-diazoacetate **1a** to 10 mmol and the yield of the product **2-1a** did not decrease remarkably, demonstrating that the reaction has potential for industrial scale-up (Scheme 1).

In order to explore the mechanism, we conducted some control experiments. First, we carried out ^{19}F NMR titration experiments using $\text{B}(\text{C}_6\text{F}_5)_3$ and DMF; when DMF was added to the CDCl_3 solution of $\text{B}(\text{C}_6\text{F}_5)_3$, the ^{19}F NMR spectra of $\text{B}(\text{C}_6\text{F}_5)_3$ showed two new peaks at chemical shifts of -158 and -165 ppm (Fig. 2), which were caused by $\text{B}(\text{C}_6\text{F}_5)_3$ coordinating with DMF. Then, the oxidation reaction was carried out in a pure argon atmosphere, and the yield of **2-1a** reduced remarkably, while the results of the control experiment with $^{18}\text{O}_2$ confirmed that the oxygen source of the product was mainly DMF, and a small portion of it came from $^{18}\text{O}_2$ (Scheme 2).

According to the related literature reports^{16b,20–22} and the results of control experiments, we can shed light on the reaction mechanism (Scheme 3). On the one hand, the diazoester compound might be activated by $\text{B}(\text{C}_6\text{F}_5)_3$ and then release N_2 to generate carbene intermediate **B**. Then, the carbene intermediate may directly react with DMF or molecular oxygen to form






Fig. 2 ^{19}F NMR titration experiments.



Scheme 2 Control experiments.



Scheme 3 Proposed reaction mechanism.

the corresponding oxidation product **2-1a** (path A and path C). On the other hand, $\text{B}(\text{C}_6\text{F}_5)_3$ might coordinate with DMF, and its coordination intermediate **G** will then react with the resonance structure **F** of **1a** to form intermediate **H**. Next, intermediate **H** will further react to generate the product **2-1a** (path B).

Conclusions

In summary, a metal-free $\text{B}(\text{C}_6\text{F}_5)_3$ -catalyzed oxidative reaction of α -diazooesters in dry DMF under an oxygen atmosphere was



reported. This reaction features a broad substrate scope, good compatibility of functional groups and a green environment-friendly nature. Importantly, the control experiments confirmed that the oxygen sources of the product **2-1a** were DMF and molecular oxygen. The protocol offers new insight on reactions catalyzed by $B(C_6F_5)_3$.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We are grateful for the financial support from CAS "Light of West China" Program and the Applied Basic Research Programs of Sichuan province, China (No. 20YYJC0492) and Technological Innovation Program of Chengdu, Sichuan province, China (No. 2021-YF05-00776-SN).

Notes and references

- (a) H. M. L. Davies and R. E. J. Beckwith, *Chem. Rev.*, 2003, **103**, 2861–2904; (b) H. M. L. Davies and D. Beckwith, *Chem. Soc. Rev.*, 2011, **40**, 1857–1869.
- (a) Q. Xiao, Y. Zhang and J.-B. Wang, *Acc. Chem. Res.*, 2012, **46**, 236–247; (b) W.-H. Hu, X.-F. Xu, J. Zhou, W.-J. Liu, X.-X. Huang, J. Hu, L.-P. Yang and L.-Z. Gong, *J. Am. Chem. Soc.*, 2008, **130**, 7782–7783; (c) M. P. Doyle, R. Duffy, M. Ratnikov and L. Zhou, *Chem. Rev.*, 2009, **110**, 704–724; (d) T. Hashimoto and K. Maruoka, *Org. Biomol. Chem.*, 2008, **6**, 829–835; (e) D. C. Cruz, F. Yuste, M. R. Martín, A. Tito and J. L. G. Ruano, *J. Org. Chem.*, 2009, **74**, 3820–3826; (f) A. V. Dubrovskiy, N. A. Markina and R. C. Larock, *Org. Biomol. Chem.*, 2013, **11**, 191–218; (g) Y. Zhang and J.-B. Wang, *Chem. Commun.*, 2009, **36**, 5350–5361; (h) T. Hashimoto, H. Kimura, N. Nakatsu and K. Maruoka, *J. Org. Chem.*, 2011, **76**, 6030–6037.
- For select reviews, see: (a) C. Zhang, P. Feng and N. Jiao, *J. Am. Chem. Soc.*, 2013, **135**, 15257–15262; (b) Y. Kumar, Y. Jaiswal and A. Kumar, *J. Org. Chem.*, 2016, **81**, 12247–12257; (c) P.-J. Zhou, C.-K. Li, S.-F. Zhou, A. Shoberu and J.-P. Zou, *Org. Biomol. Chem.*, 2017, **15**, 2629–2637; (d) P. Saha, S. Kumar Ray and V. K. Singh, *Tetrahedron Lett.*, 2017, **58**, 1765–1769; (e) P. Gu, X.-P. Wu, Y. Su, X.-Q. Li, P. Xue and R. Li, *Synlett*, 2014, **25**, 535–538.
- (a) M. Ma, C. Li, L. Peng, F. Xie, X. Zhang and J. Wang, *Tetrahedron Lett.*, 2005, **46**, 3927–3929; (b) P. Truong, C. S. Shanahan and M. P. Doyle, *Org. Lett.*, 2012, **14**, 3608–3611; (c) M. B. Rubin and R. Gleiter, *Chem. Rev.*, 2000, **100**, 1121–1164.
- (a) Z. Guo, H. Huang, Q. Fu and W. Hu, *Synlett*, 2006, **15**, 2486–2488; (b) Y. Yu, Q. Sha, H. Cui, K. S. Chandler and M. P. Doyle, *Org. Lett.*, 2018, **20**, 776–779.
- N. R. O'Connor, P. Bolgar and B. M. Stoltz, *Tetrahedron Lett.*, 2016, **57**, 849–851.
- (a) S. E. Allen, R. R. Walvoord, R. Padilla-Salinas and M. C. Kozlowski, *Chem. Rev.*, 2013, **113**, 6234–6458; (b) Z. Shi, C. Zhang, C. Tang and N. Jiao, *Chem. Soc. Rev.*, 2012, **41**, 3381–3430.
- (a) K. P. Kornecki, J. F. Briones, V. Boyarskikh, F. Fullilove, J. Autschbach, K. E. Schrote, K. M. Lancaster, H. M. L. Davies and J. F. Berry, *Science*, 2013, **342**, 351–354; (b) K. H. Dötz, *Angew. Chem., Int. Ed.*, 1984, **23**, 587–608; (c) J. Barluenga, F. Andina, M. A. Fernandez-Rodríguez, P. García-García, I. Merino and E. Aguilar, *J. Org. Chem.*, 2004, **69**, 7352–7354.
- C.-M. Xu, Y.-C. Wang and L. Bai, *J. Org. Chem.*, 2020, **85**, 12579–12584.
- R.-S. Liu, Q.-S. Liu, H.-R. Meng, H.-Y. Ding, J.-D. Hao, Z.-Y. Ji, H.-L. Yue and W. Wei, *Org. Chem. Front.*, 2021, **8**, 1970–1975.
- For select reviews, see: (a) K. Nagy, H. Mehdi, I. Pápai, P. Nagy, P. Király, G. Tárkányi, E. Gábor and S. Tibor, *Chem.-Eur. J.*, 2012, **18**, 574–585; (b) P. Oña-Burgos, A. Kubas, F. C. Falk, F. Breher, K. Fink, G. Lutz and J. Paradies, *Dalton Trans.*, 2012, **41**, 9056–9060; (c) D. J. Scott, M. J. Fuchter and A. E. Ashley, *J. Am. Chem. Soc.*, 2014, **136**, 15813–15816; (d) I. Blanca, D. Palomas, S. Holle, S. Steinberg, J. A. Nicasio and M. Alcarazo, *Angew. Chem., Int. Ed.*, 2012, **51**, 12367–12369; (e) Y. Hoshimoto, T. Kinoshita, S. Hazra, M. Ohashi and S. Ogoshi, *J. Am. Chem. Soc.*, 2018, **140**, 7292–7300; (f) F. G. M. Alexander, T. Sebastian, T. Schneider, U. Flörke, Z.-W. Qu, S. Grimme and J. Paradies, *Angew. Chem., Int. Ed.*, 2016, **55**, 12219–12223; (g) D. G. Sravanthi, R. C. Alpina and B. N. Babu, *New J. Chem.*, 2017, **41**, 2328–2332; (h) Y.-X. Han, S.-T. Zhang, J.-H. He and Y.-T. Zhang, *J. Am. Chem. Soc.*, 2017, **139**, 7399–7407.
- For select reviews, see: (a) M. Cao, A. Yesilcimen and M. Wasa, *J. Am. Chem. Soc.*, 2019, **141**, 4199–4203; (b) M. Hellal, F. C. Falk, E. Wolf, M. Dryzhakov and J. Moran, *Org. Biomol. Chem.*, 2014, **12**, 5990–5994; (c) M. Dryzhakov, M. Hellal, E. Wolf, F. C. Falk and J. Moran, *J. Am. Chem. Soc.*, 2015, **137**, 9555–9558; (d) G. Ku-maraswamy and M. Gangadhar, *ChemistrySelect*, 2019, **4**, 8973–8977; (e) M. Hatano, K. Hayashi, T. Sakamoto, Y. Makino and K. Ishihara, *Synlett*, 2016, **27**, 1061–1067; (f) M. Hatano, T. Sakamoto, T. Mizuno, Y. Goto and K. Ishihara, *J. Am. Chem. Soc.*, 2018, **140**, 16253–16263; (g) M. Hellal, C. F. Florian, E. Wolf, M. Dryzhakov and J. Moran, *Org. Biomol. Chem.*, 2014, **12**, 5990–5994; (h) D. Marian and J. Moran, *ACS Catal.*, 2016, **6**, 3670–3673.
- S. Basak, L. Winfrey, B. A. Kustiana, R. L. Melen, L. C. Morrill and A. P. Pulis, *Chem. Soc. Rev.*, 2021, **50**, 3720–3737.
- D. G. Sravanthi, A. R. Chari, A. Nagarsenkar, D. K. Sigalapallia and B. N. Babu, *New J. Chem.*, 2017, **41**, 2328–2332.
- F. Ling, L.-X. Shen and W.-H. Zhong, *Tetrahedron Lett.*, 2018, **59**, 3678–3682.
- (a) G.-Y. Xu, S.-B. Tang, B. Shao and J.-T. Sun, *Chem. Commun.*, 2019, **55**, 9096–9099; (b) A. Dasgupta, K. Stefkova, R. Babaahmadi, L. Gierlichs, A. Ariafard and R. L. Melen, *Angew. Chem.*, 2020, **59**, 15492–15496; (c) A. Dasgupta, R. Babaahmadi, B. Slater, B. F. Yates, A. Ariafard and R. L. Melen, *Chem*, 2020, **6**, 2364–2381; (d)



- J. P. Mancinelli and S. M. Wilkerson-Hill, *ACS Catal.*, 2020, **10**, 11171–11178; (e) J. Fang and M. Brewer, *Org. Lett.*, 2018, **20**, 7384–7387; (f) M. J. Hensinger, N. J. Dodge and M. Brewer, *Org. Lett.*, 2020, **22**, 497–500; (g) H.-H. San, C.-Y. Wang, H.-P. Zeng, S.-T. Fu, M. Jiang and X.-Y. Tang, *J. Org. Chem.*, 2019, **84**, 4478–4485.
- 17 H.-H. San, S.-J. Wang, M. Jiang and X.-Y. Tang, *Org. Lett.*, 2018, **20**, 4672–4676.
- 18 (a) Y. Dong, H. Zhang, J. Yang, S. He, Z.-C. Shi, X.-M. Zhang and J.-Y. Wang, *ACS Omega*, 2019, **4**, 21567–21577; (b) X.-Y. Zhan, H. Zhang, Y. Dong, J. Yang, S. He, Z.-C. Shi, L. Tang and J.-Y. Wang, *J. Org. Chem.*, 2020, **85**, 6578–6592; (c) H. Zhang, X.-Y. Zhan, Y. Dong, J. Yang, S. He, Z.-C. Shi, X.-M. Zhang and J.-Y. Wang, *RSC Adv.*, 2020, **10**, 16942–16948; (d) G.-M. Zhang, H. Zhang, B. Wang and J.-Y. Wang, *RSC Adv.*, 2021, **11**, 17025–17031; (e) B. Wang, H. Xu, H. Zhang, G.-M. Zhang, F.-Y. Li, S. He, Z.-C. Shi and J.-Y. Wang, *Org. Chem. Front.*, 2021, **8**, 6670–6677.
- 19 X.-Y. Wu, W.-X. Gao, Y.-B. Zhou, M.-C. Liu and H.-Y. Wu, *Adv. Synth. Catal.*, 2022, **364**, 750–754.
- 20 (a) W.-B. Liu, C. Chen and P. N. Zhou, *J. Org. Chem.*, 2017, **82**, 2219–2222; (b) J.-J. Da, H.-Y. Ji Jeon, H.-J. Kim, Y. Kim and S. Lee, *Org. Lett.*, 2014, **16**, 2208–2211; (c) W. Ai, Y. Liu, Q. Wang, Z. Lu and Q. Liu, *Org. Lett.*, 2018, **20**, 409–412.
- 21 D. P. Higley and R. W. Murray, *J. Am. Chem. Soc.*, 1974, **96**, 3330–3332.
- 22 (a) M. Santi, D. M. C. Ould, J. Wenz, Y. Soltani, R. L. Melen and T. Wirth, *Angew. Chem., Int. Ed.*, 2019, **58**, 7861–7865; (b) S. Rao, R. Kapanaiah and K. R. Prabhu, *Adv. Synth. Catal.*, 2019, **361**, 1301–1306.

