RSC Advances



View Article Online

View Journal | View Issue

PAPER



Cite this: RSC Adv., 2022, 12, 35215

DTBP-mediated cross-dehydrogenative coupling of 3-aryl benzofuran-2(3*H*)-ones with toluenes/ phenols for all-carbon quaternary centers†

Zhou Tong, Xinju Peng, Zhi Tang, D Weijun Yang, * Wei Deng,* Shuang-Feng Yin, Nobuaki Kambe and Renhua Qiu *

Received 4th October 2022 Accepted 2nd December 2022 ben DOI: 10.1039/d2ra06231c

rsc.li/rsc-advances

We have developed a transition-metal free protocol for efficient cross-dehydrogenative coupling of 3-aryl benzofuran-2(3*H*)-ones and toluenes/phenols using DTBP as an oxidant. A diverse range of 3-aryl benzofuran-2(3*H*)-ones, toluenes, and phenols undergo C–H bond cleavage to generate all-carbon quaternary centers in good yields, making this protocol useful for the synthesis of complex molecules. A gram scale experiment was performed in good yield.

Introduction

Cross-dehydrogenative coupling (CDC) is a straightforward strategy for carbon–carbon bond formation.¹ Many pioneering efforts have been made to develop CDC methodologies to access useful motifs by using a variety of transition metal catalysts.² As for the transition-metal free methods for CDC reactions, there are only a few pioneering works that have been developed,³ including oxidative promotion, photocatalysis and electrochemical catalysis. Although a transition-metal free CDC reaction has been developed, its application in all-carbon quaternary center construction has never been disclosed.⁴

All-carbon quaternary centers are broadly present in various bioactive natural products, artificially synthesized pharmaceutical and medicinal molecules,5 such as hopeahainol A,5b oxyphenisatin, TOP216 analogue,^{5c} yuccaol,^{5d} estrogen receptor modulator,^{5e} cyclophilin inhibitor,^{5f} hamigeran B (Fig. 1).^{5g} These molecules show unique bioactivities due to the rigid three-dimensional structure. Therefore, the efficient and selective construction of all-carbon quaternary stereocenters has received intensive attention in recent decades.6 However, the construction of quaternary carbon centers via CDC reaction is rare.7 Previously, we have developed a Cu-catalyzed CDC reaction of 3-aryl benzofuranones with heteroatom arenes, which furnishes highly functionalized 3,3-diaryl benzofuranones bearing a three aryls quaternary carbon center at C3 position in vields.71 These good complex compounds can be

decarbonylation to be triaryl methane easily.⁸ However, this protocol can be only applied to the coupling between $C(sp^3)$ -H bond and $C(sp^2)$ -H bond.

Herein, we disclosed the first example of the construction of all-carbon quaternary centers *via* a transition-metal free CDC reaction of 3-aryl benzofuran-2(3H)-ones with toluenes/phenols by using DTBP (di-*t*-butyl peroxide) as an oxidant, which the two different $C(sp^3)$ -H bonds can be cross-coupling site-selectively in good yields. This protocol shows a broad substrate scope and gram-scalable ability. While toluenes were changed into phenols with methyl group in this protocol, the *ortho*- $C(sp^2)$ -H bond of phenol was activated site selectively.

Results and discussion

Initially, we explored the reactions using 4-methylbenzofuranone (1a) and toluene (2a) as substrates under different reaction conditions (Table 1). CuI, NiCl₂, and CoCl₂ were then



Fig. 1 Bioactive all-carbon quaternary centers.

State Key Laboratory of Chemo/Biosensing and Chemometrics, Advanced Catalytic Engineering Research Center of the Ministry of Education, College of Chemistry and Chemical Engineering, Hunan University, Changsha, 410082, P. R. China. E-mail: renhuaqiu1@hnu.edu.cn

[†] Electronic supplementary information (ESI) available. CCDC 1949691. For ESI and crystallographic data in CIF or other electronic format see DOI: https://doi.org/10.1039/d2ra06231c

Table 1 Survey on condition for 2a formation^a



^a Reaction conditions: 1a (0.2 mmol), cat. (mol%), oxidant (0.1 mmol) and **2a** (1.0 mL), 12 h, under N₂, sealed tube, isolated yield. ^{*b*} Under air. ^{*c*} DTBP 0.2 equiv. ^{*d*} DTBP 1.0 equiv. TBHP = *t*-butylhydroperoxide, CHP = cumyl hydroperoxide, mCPBA = meta-cholorperoxybenzoicacid, $DMF = N_N$ -dimethyl-formicaci, DMSO = dimethyl sulfoxide.

examined in the presence of DTBP as an oxidant under N₂ atmosphere, the yields of 3a were 83%, 88%, 88%, respectively (entries 1-3) and changed the amount of catalyst resulting in a decrease of yield (entries 4-5). However, without a metal catalyst, the yield of 3a was also up to 88% (entry 6), indicating only DTBP is enough for this protocol. While the reaction atmosphere was changed into the air, the yield of 3a was decreased to 77% (entry 7). When DTBP was replaced by THBP, CHP and mCPBA resulted in the decrease of the yield of 3a to 10%, 27% and no product (entry 8-10). When mixed solvent was added, the target product could not be obtained (entry 11-12). Both increasing and decreasing the amount of oxidant would have a negative impact on the reaction (entry 13-14). The adoption of other temperatures also resulted in lower yields (entries 15-16). The molecular structure of 3a were being isolated as racemic mixtures and has been unambiguously confirmed by single crystal X-ray diffraction study (figure in Scheme 1, CCDC 1949691[†]).

With the optimized reaction conditions in hand, we next investigated the substrate scope (Scheme 1). A variety of substituted benzofuranones (1) could proceed well in this CDC reaction to deliver corresponding products in moderate to good yields, and valuable groups such as 4-H (3b, 91%), 3-tert-butyl (3c, 96%), 2,4-di-tert-butyl (3d, 88%), 2-tert-butyl (3e, 94%), 4ethyl (3f, 97%), 3,5-dimethyl (3g, 72%), 2,4-dimethyl (3h, 85%), 4-isopropyl (3i, 95%), 4-phenyl (3j, 82%), 4-fluorine (3k, 73%), 4chloro (3l, 82%. 3m, 53%), and 4-bromo (3n, 76%) were tolerated. Overall, the adaptability of the reaction system to



N₂, sealed tube, isolated yield. ^b 2k 0.4mmol.

Scheme i Substrates scope of benzoraranone.	Scheme 1	Substrates	scope	of be	enzofuranon	ie. ^a
---	----------	------------	-------	-------	-------------	------------------

functional groups is good, only when electron-withdrawing groups are attached to the benzofuranones, a slight inhibitory effect on the reaction was observed. And strong electron withdrawing group would completely inhibit the reaction (30). Furthermore, compared with the reactivity of HP-136 (3d), three benzofuranones (3c, 3e, 3f) have higher carbon radical trapping ability, which may have some potential to be used as a benzyl radical anti-oxidant.

On the other hand, the scope of toluene analogues was also tested (Scheme 2). A series of aryl-substituted toluenes were subjected to this CDC reaction to react with benzofuranone 1a, and various products were obtained in moderate to good yields. The substituted groups with electron-donating and electronwithdrawing groups at the para or ortho position of the aryl ring could proceed smoothly to afford the desired products (4a-4i). In addition, the radical trapping reagent 2, 6-di-tert-butyl-4methylphenol (BHT) was also applicable in this CDC reaction to furnish **4h** in 61% yield, showing that this reaction may involve a carbon radical center. These results showed that the method is suitable for multi-functionalized steric substrates. Nevertheless, for the large steric biphenyl methane, it could also get the target product 4j in 35% yield. The reaction system could also be applied to heterocyclic systems. 6-Methylquinoline (2k) could also undergo CDC cross-coupling to get the product (4k). But 2methyindole (21) could only get the product substituted at the 3position as a triaryl methane molecule (41). Other indoles also gave the same 3-substituted indole products (4m, 3n) in 76% and 55%, respectively. And the furan has two products (40, 40')which have been isolated respectively.8 It should be noted that, for the synthesis of **4p** and **4q** in Pd or Cu systems,^{7c,1} there are

1



 a **1a** (0.2 mmol), oxidant (0.1 mmol), **2** (1.0 mL), 140 °C, 12 h under N₂, sealed tube, isolated yield. b **2k** 0.4 mmol.

Scheme 2 Substrates scope of toluene.^a

no products obtained. While this protocol can afford **4p** and **4q** in 34 and 55% yield, respectively (Scheme 2b). When we chose *n*-propyl benzene as substrate which bearing benzylic C–H bond, methylene C–H bond and methyl C–H, we can get the benzylic product **4r** in 75% yield, while only trace (<1%) amount of side-products **4r**' and **4r**'' were detected (Scheme 2c). This result shows a good site-selectivity of our method.

When the steric *o*-*t*Bu substituted groups were removed from BHT, the *ortho*-C–H functionalization of phenols could engage in this process to deliver the triaryl all-carbon quaternary centers (Scheme 3). A series of phenols, such as electrondonating groups and halogen groups, were tolerated in this CDC reaction to react with benzofuranone **1a** to give **5a**–**5g** in good yields. Side-products of C–O construction were detected



^o**1** (0.2 mmol), **4** (0.4 mmol), oxidant (0.1 mmol), 140°C, 12 h, under air, sealed tube, isolated yield of **5**. ^bDetected by GC. ND: No detected.

Scheme 3 Substrates scope of phenols.^a

especially in the target product (5') with low yield (the ratio of 5: 5' is shown in the parenthesis). When there were electron donating groups in the para position of phenols, it was beneficial to the generation of radicals, so the yield would be better. When there were other groups in the ortho position of phenols, radicals were not easy to transfer, so the yield of 5 would be reduced. Halogenated phenol, especially iodophenol, were more active, and would get side reactions, leading to low yield. However, when **1a** reacted with the thiophenol, only the C–S cross coupling product was obtained (**6a**, 30%). This result was consistent with the literature,⁹ which the ortho position of phenol is indeed easy to activate as C–C bond formation rather than C–O bond formation, while that of thiophenol is more difficult to be activated, and the direct construction of C–S bond was observed.

Gram scale experiments were performed for the synthesis of **3a** and **5a**, and the yields are up to 65% and 71%, respectively (Scheme 4a). Further amplify the amount of reaction may have the risk of explosion, and a special reactor would be needed. Compound **5a** contains lactone and phenol groups, making its transformation to other useful motifs possible. For example, it gave the target decarbonyl product 7 with two phenols in the yield of 85%. And it could be trifluoroacetylated by trifluoroacetic anhydride to **8** in 60% yield. Also compound **8** could undergo dehydroxylation to form product **9** with 83% yield (Scheme 4b).⁸

To probe the possible reaction mechanism, some control experiments were conducted. An intermolecular competition experiment between toluene and the deuterated toluene was carried out to investigate the kinetic isotope effect (KIE), which was found to be significant ($k_{\rm H}/k_{\rm D} = 5.6$) (Scheme 5a). In the hydrocarbon activation reaction, when $k_{\rm H}/k_{\rm D} > 2$, the cleavage of the C–H bond may be considered to be the rate-determining step of the reaction.¹⁰ The cleavage of the C–H bond may be the rate-determining step in this system. It was observed that the reaction was suppressed by the addition of a radical scavenger 2,2,6,6-tetramethyl-1-piperidinyloxy (Tempo) and 1,4-

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence

(a) Gram-scale synthesis of 3a and 5a Tol. 25 ml DTBP 0.5 eg 500 uL Detected by GC-MS 10.0 mmol 1.0 mL 1a 5.0 mmol 1.12 g Ó٢ DTBP 0.5 eg 500 uL 5a, 1.17 g, 71% (b) Transformation of 5a нсоон Pd(OAc)₂ (0.1 eq.) dppf (0.2 eq.) Tf₂O Et₃N, DMSO, 60 °C, 12 I Et₂N, 80 °C, 12 h Vield 83% Yield, 60% CH₃ONa DME 160 °C 12 h

Scheme 4 Gram-scale synthesis and transformation.

DTBP 0.5 eq

140 °C

DTBP 0.5 eq

140 °C

10

DTBP 0.5 eq

DTBP 0.5 eq

140 °C

DTBP 0.5 eq

140 °C

Yield, 85%

a) KIE Experiment



2a

DTBP 0.5 eq

140

coupling Experi

d) 1,4-Naphthoguinone Experiments



benzoquinone (BQ), implying the involvement of a radical species (Scheme 5b). And butylated hydroxytoluene (BHT) has trapped the radical of **1a** instead of toluene radicals. In addition, the toluene homo-coupling product was detected by GC-MS. Based on the observation, the reaction appears to go through a plausible radicals pathway. When **1a** was subjected to the reaction system without toluene, only self-coupling product **10** was obtained under the action of DTBP. And **3a** could be obtained in 66% yield from **10** after reacting with toluene (Scheme 5c). At the same time, the reaction of **1a** with naphthoquinone (**11**) did not produce naphthol, which may indicate that the reaction with phenol does not involve intermediates of quinone (Scheme 5d).

Based on the above results and those reported in the literature,^{9,11} a possible pathway for the cross-dehydrogenative coupling reaction was proposed (Scheme 6). 3-Aryl benzofuran-2(3*H*)-ones and toluene/phenols respectively generated free carbon radicals with DTBP, and then two radicals were cross-coupling to obtain **3a** and **4a**, respectively. Probably, the 3aryl benzofuran-2(3*H*)-ones could also generate relative radical by reacting with benzyl radical species, which can be easily trapped by another benzyl radical to target products.

Conclusions

3a-d₇, 15%

viold

0

ND

radical scavenger

(2 eq)

none

TEMPO BQ

4h. 57%

NiCl₂ 0.1 eq K₂S₂O₈ 2.0 eq

CH_CN, 80 °C

Ref. 7I

3a, 669

12, 63%

In summary, we have disclosed the first transition-metal free synthesis of all-carbon quaternary centers *via* a robust, efficient, and practical cross-dehydrogenative coupling of benzofuranones and toluenes/phenols. A diverse array of functional groups are tolerated in this transformation. A reaction mechanism with radical participation is proposed. This simple method would be useful for the synthesis of complex molecules.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors thank the Natural Science Foundation of China (21676076, 21878081, 21971060 and 22278120), Hu-Xiang High Talent in Hunan Province (2018RS3042) and Recruitment Program for Foreign Experts of China (WQ20164300353 and G2022160001) for financial support.

8

Notes and references

1 Selected reviews on CDC reaction: (a) M. K. Lakshman and P. K. Vuram, Chem. Sci., 2017, 8, 5845; (b) L. Lv and Z. Li, Top. Curr. Chem., 2016, 374, 38; (c) S. A. Girard, T. Knauber and C.-J. Li, Angew. Chem., Int. Ed., 2014, 53, 74; (d) C. Zhang, C. Tang and N. Jiao, Chem. Soc. Rev., 2012, 41, 3464; (e) C. Liu, H. Zhang, W. Shi and A. Lei, Chem. Rev., 2011, 111, 1780; (f) C. S. Yeung and V. M. Dong, Chem. Rev., 2011, 111, 1215; (g) C.-L. Sun, B.-J. Li and Z.-J. Shi, Chem. Rev., 2011, 111, 1293; (h) S. H. Cho, J. Y. Kim, J. Kwak and S. Chang, Chem. Soc. Rev., 2011, 40, 5068; (i) C.-J. Li, Acc. Chem. Res., 2009, 42, 335; (j) C.-J. Li and Z. Li, Pure Appl. Chem., 2006, 78, 935; (k) X.-X. Guo, D.-W. Gu, Z. Wu and W. Zhang, Chem. Rev., 2015, 115, 1622; (1) L. C. M. Castro and N. Chatani, Chem. Lett., 2015, 44, 410; (m) X. Cen, K. M. Engle, D.-H. Wang and J.-Q. Yu, Angew. Chem., Int. Ed., 2009, 48, 5094.

- 2 Selected examples of transition metal catalyzed CDC reaction, palladium catalyzed: (a) J. Zhang, Y. Zhuang, Y. Ma, X. Yang and M. Szostak, Adv. Synth. Catal., 2019, 361, 5709; (b) C. Dai, Z.-B. Huang, L. Liu, Y. Han, D.-Q. Shi and Y. Zhao, Eur. J. Org. Chem., 2020, 2020, 826; (c) E. Kianmehr, M. Torabi, M. Rezazadeh Khalkhali, N. Faghih and K. M. Khan, Eur. J. Org. Chem., 2015, 2015, 2796; (d) Ruthenium catalyzed: S. Dana, D. Chowdhury, A. Mandal, F. A. S. Chipem and M. Baidya, ACS Catal., 2018, 8, 10173; (e) C. Heitz, A. W. Jones, B. S. Oezkaya, C. L. Bub, M.-L. LouillatHabermeyer, V. Wagner and F. W. Patureau, Chem.-Eur. J., 2016, 22, 17980; (f) Q. Huang, S. Fu, S. Ke, H. Xiao, X. Zhang and S. Lin, Eur. J. Org. Chem., 2015, 2015, 6602; (g) Cobalt catalyzed: B. Lin, S. Shi, R. Lin, Y. Cui, M. Fang, G. Tang and Y. Zhao, J. Org. Chem., 2018, 83, 6754; (h) M. Khrizanforov, S. Strekalova, V. Khrizanforova, A. Dobrynin, K. Kholin, T. Gryaznova, V. Grinenko, A. Gubaidullin, M. K. Kadirov and Y. Budnikova, Top. Catal., 2018, 61, 1949; (i) T. Li, Y. Yang, B. Li and P. Yang, Chem. Commun., 2019, 55, 353; (j) Copper catalyzed: G. S. Grandhi, S. Dana, A. Mandal and M. Baidya, Org. Lett., 2020, 22, 2606; (k) R. M. Gorman, T. E. Hurst, W. F. Petersen and R. J. K. Taylor, Tetrahedron, 2019, 75, 130711; (l) M. Kumar, Raziullah, A. A. Khan, A. Ahmad, H. S. Dutta, R. Kant and D. Koley, J. Org. Chem., 2019, 84, 13624; (m) Other transition catalyzed: L. K. Jin, L. Wan, J. Feng and C. Cai, Org. Lett., 2015, 17, 4726; (n) K. Yang and Q. Song, Org. Lett., 2015, 17, 548; (o) S. S. Kotha, S. Chandrasekar, S. Sahu and G. Sekar, Eur. J. Org. Chem., 2014, 2014, 7451.
- 3 Selected examples of metal-free CDC reaction, oxidative promoted: (a) H. Jiang, X. Tang, S. Liu, L. Wang, H. Shen, J. Yang, H. Wang and Q.-W. Gui, Org. Biomol. Chem., 2019, 17, 10223; (b) Q. Chen, G. Yu, X. Wang, Y. Ou and Y. Huo, Green Chem., 2019, 21, 798; (c) F. Jafarpour and M. Darvishmolla, Org. Biomol. Chem., 2018, 16, 3396; (d) Q. Yang, X. Han, J. Zhao, H.-Y. Zhang and Y. Zhang, J. Org. Chem., 2019, 84, 11417; (e) Photocatalysis: G. Kibriya,

A. K. Bagdi and A. Hajra, J. Org. Chem., 2018, 83, 10619; (f)
W. Wei, L. Wang, H. Yue, P. Bao, W. Liu, C. Hu, D. Yang and H. Wang, ACS Sustainable Chem. Eng., 2018, 6, 17252; (g) Electrochemical catalysis: M.-J. Luo, Y. Li, X.-H. Ouyang, J.-H. Li and D.-L. He, Chem. Commun., 2020, 56, 2707; (h)
Y.-Z. Yang, R.-J. Song and J.-H. Li, Org. Lett., 2019, 21, 3228; (i) Z. J. Wu and H.-C. Xu, Angew. Chem., Int. Ed., 2017, 56, 4734.

- 4 D. J. C. Constable, P. J. D. Lorenz, J. D. Hayler,
 G. R. Humphrey, J. L. Leazer Jr, R. J. Linderman,
 K. J. Manley, B. A. Pearlman, A. Wells, A. Zaks and
 T. Y. Zhang, *Green Chem.*, 2007, 9, 411.
- 5 Selected all-carbon quaternary centers as bioactive compounds: (a) T. Ling and F. Rivas, Tetrahedron, 2016, 72, 6729; (b) D. J. Newman and G. M. Cragg, J. Nat. Prod., 2016, 79, 629; (c) K. C. Nicolaou, S. A. Snyder, X.-H. Huang, K. B. Simonsen, A. E. Koumbis and A. Bigot, J. Am. Chem. Soc., 2004, 126, 10162; (d) M. K. Christensen, K. D. Erichsen, C. Trojel-Hansen, J. Tjørnelund, S. J. Nielsen, K. Frydenvang, T. N. Johansen, B. Nielsen, M. Sehested, P. B. Jensen, M. Ikaunieks, A. Zaichenko, E. Loza, I. Kalvinsh and F. Björkling, J. Med. Chem., 2010, 53, 7140; (e) S. Piacente, P. Montoro, W. Oleszek and C. Pizza, J. Nat. Prod., 2004, 67, 882; (f) X. Peng, Z. Hu, J. Zhang, W. Ning, S. Zhang, C. Dong, X. Shi and H.-B. Zhou, Chem. Commun., 2019, 55, 14570; (g) S. Daum, M. Schumann, S. Mathea, T. Aumuller, M. A. Balsley, S. L. Constant, B. F. de Lacroix, F. Kruska, M. Braun and C. Schiene-Fischer, Biochemistry, 2009, 48, 6268; (h) K. D. Wellington, R. C. Cambie, P. S. Rutledge and P. R. Bergquist, J. Nat. Prod., 2000, 63, 79.
- 6 Selected examples of the all-carbon quaternary centers formation: (a) R. Long, J. Huang, J. Gong and Z. Yang, Nat. Prod. Rep., 2015, 32, 1584; (b) I. Marek, Y. Minko, M. Pasco, T. Mejuch, N. Gilboa, H. Chechik and P. J. Das, J. Am. Chem. Soc., 2014, 136, 2682; (c) B. M. Trost and C. Jiang, Synthesis, 2006, 369; (d) K. W. Quasdorf and L. E. Overman, Nature, 2014, 516, 181; (e) C. G. Watson, A. Balanta, T. G. Elford, S. Essafi, J. N. Harvey and V. K. Aggarwal, J. Am. Chem. Soc., 2014, 136, 17370; (f) R. Alam, T. Vollgraff, L. Eriksson and K. J. Szabj, J. Am. Chem. Soc., 2015, 137, 11262.
- 7 Selected examples of constructing all-carbon quaternary centers by CDC reactions: (a) G. Zhang, Y. Zhang and R. Wang, Angew. Chem., Int. Ed., 2011, 50, 10429; (b) Tanaka, K. Hashiguchi, T. Tanaka, R. Yazaki and т. Ohshima, ACS Catal., 2018, 8, 8430; (c) G. Hong, T. D. Nahide, U. K. Neelam, P. Amadeo, A. Vijeta, Ρ. J. M. Curto, C. E. Hendrick, K. F. VanGelder and M. C. Kozlowski, ACS Catal., 2019, 9, 3716; (d) J. E. M. N. Klein, A. Perry, D. S. Pugh and R. J. K. Taylor, Org. Lett., 2010, 12, 3446; (e) C. L. Moody, V. Franckevičius, P. Drouhin, J. E. M. N. Klein and R. J. K. Taylor, Tetrahedron Lett., 2012, 53, 1897; (f) T. E. Hurst, R. M. Gorman, P. Drouhin, A. Perry and R. J. K. Taylor, Chem.-Eur. J., 2014, 20, 14063; (g) T. E. Hurst and R. J. K. Taylor, Eur. J. Org. Chem., 2017, 2017, 203; (h)

Y.-X. Jia and E. P. Kündig, Angew. Chem., Int. Ed., 2009, 48, 1636; (i) C. Dey, E. Larionov and E. P. Kündig, Org. Biomol. Chem., 2013, 11, 6734; (j) J. R. Donald, R. J. K. Taylor and W. F. Petersen, J. Org. Chem., 2017, 82, 11288; (k) Y. Liu, J. Li, X. Ye, X. Zhao and Z. Jiang, Chem. Commun., 2016, 52, 13955; (l) Z. Tang, Z. Liu, Z. Tong, Z. Xu, C. T. Au, R. Qiu and N. Kambe, Org. Lett., 2019, 21, 5152.

- 8 Z. Tang, L. Peng, Y. Yuan, T. Li, R. Qiu and N. Kambe, *J. Org. Chem.*, 2020, **85**, 5300.
- 9 2-Position of phenol: (a) Z. Liu, Y. Jiang, C. Liu, L. Zhang, J. Wang, T. Li, H. Zhang, M. H. Li and X. Yang, J. Org. Chem., 2020, 85, 7386; (b) D. Glavač, N. Topolovčan and

M. Gredičak, *J. Org. Chem.*, 2020, **85**, 14253; (c) C–S construction: W. Jiang, J. Zhuge, J. Li, G. Histand and D. Lin, *J. Org. Chem.*, 2020, **85**, 2415; (d) A. J. Basson and M. G. McLaughlin, *J. Org. Chem.*, 2020, **85**, 5615.

- 10 (a) E. M. Simmons and J. F. Hartwig, Angew. Chem., Int. Ed., 2012, 51, 3066; (b) P. K. Pramanick, Z. Zhou, Z.-L. Hou and B. Yao, J. Org. Chem., 2019, 84, 5684.
- 11 (a) S. Jin, B. Xie, S. Lin, C. Min, R. Deng and Z. Yan, Org. Lett.,
 2019, 21, 3436; (b) H. Peng, J.-T. Yu, Y. Jiang, H. Yang and
 J. Cheng, J. Org. Chem., 2014, 79, 9847; (c) C. Pan, Q. Ni,
 Y. Fu and J.-T. Yu, J. Org. Chem., 2017, 82, 7683.

35220 | RSC Adv., 2022, 12, 35215-35220