




 Cite this: *RSC Adv.*, 2022, 12, 34463

The role of polyplexes in developing a green sustainable approach in agriculture

 Pratyush K. Das,  Gyanendra Panda, Kananbala Patra, Nivedita Jena and Mamoni Dash *

Rise in global population has increased the food demands and thus the competition among farmers to produce more and more. In the race to obtain higher productivity, farmers have resorted to injudicious farming practices that include the reckless use of nitrogenous fertilizers and intensive cropping on farmlands. Such practices have paved the path for large scale infestations of crops and plants by pests thus affecting the plant productivity and crop vigour. There are several traditional techniques to control pest infestations in plants such as the use of chemical or bio-pesticides, and integrated pest management practices which face several drawbacks. Delivery of gene/nucleic acid in plants through genetic engineering approaches is a more sustainable and effective method of protection against pests. The technology of RNA interference (RNAi) provides a sustainable solution to counter pest control problems faced by other traditional techniques. The RNAi technique involves delivery of dsDNA/dsRNA or other forms of nucleic acids into target organisms thereby bringing about gene silencing. However, RNAi is also limited to its use because of their susceptibility to degradation wherein the use of cationic polymers can provide a tangible solution. Cationic polymers form stable complexes with the nucleic acids known as "polyplexes", which may be attributed to their high positive charge densities thus protecting the exogenous nucleic acids from extracellular degradation. The current paper focuses on the utility of nucleic acids as a sustainable tool for pest control in crops and the use of cationic polymers for the efficient delivery of nucleic acids in pests thus protecting the plant from infestations.

 Received 17th October 2022
 Accepted 15th November 2022

DOI: 10.1039/d2ra06541j

rsc.li/rsc-advances

1. Introduction

Sustainability in agriculture is brought about by effective practices and technologies that consider minimizing the environmental footprint.¹ The use of nucleic acids as a sustainable means for control of pests in crops and plants has started to gain foothold. Nucleic acids which include either DNA or RNA can act as an active ingredient in pest control formulations. Upon exposure of an organism to these nucleic acids, can either act to cause mutagenesis or can cause gene silencing thereby altering the expression in a particular gene or a set of genes.² RNA is the most common active ingredient used in topical pesticidal applications. Targeting specific gene/mRNA sequences in pathogens and pests of different plants inducing RNA silencing *via* specifically designed nucleic acids seems to be quite promising. Special interests of researchers have been drawn towards the use of dsRNAs, and siRNAs to induce RNAi mediated gene silencing.³ Exogenously supplied dsRNAs could induce gene silencing in an organism by means of injecting, soaking, or feeding and can cause heritable effects in certain cases.⁴ An RNAi study in *C. elegans* was a first of its kind

affecting gene silencing in the organism *via* exposure to environmental RNAi. The resultant gene silencing as a result was evident from the systemic spread of the silencing signals to all the cells in the organism. Environmental RNAi offers to be a cost-effective and simple method to deliver dsRNA into organisms. This particular method has led to a large scale RNAi screening studies thus paving way for several innovative pest control approaches in agriculture.⁵⁻⁸

Chemical pesticides are mostly used to control pest infestations in farmlands owing to their low cost-nature but have several negative implications on the crop⁹ as well as its environment.¹⁰ Moreover, aggressive uses of such chemicals have resulted in the occurrence of resistance amongst pests towards the particular pesticide. There arises an immediate need to look for alternative options that not only will help effectively control the pests but also simultaneously act as a cheap, biodegradable, and non-toxic product. Conventional gene transfer methods employing genetic engineering approaches have proved to be successful but face several drawbacks including public acceptance issues.¹¹ As such nucleic acid based environmental RNAi approaches could be instrumental in controlling the pests of plants thus substantially improving the productivity as well as preventing large scale crop loss.

Institute of Life Sciences, DBT-ILS, Bhubaneswar, Odisha, India. E-mail: mamoni.dash@ils.res.in



Development of nucleic acid based topical RNAi strategies for protection against pests pose several bottle necks. A major issue is the lack of amplification of silencing signal mostly due to the degradation of the active ingredient (dsRNA) post ingestion.¹² The availability of RNAi effectors and their silencing inducing ability has been found to decrease with an increase in the distance from the site of its exogenous application.¹³ Therefore, the current research activities in relation to the development of RNAi based topical pest control products are mostly focussed on two aspects. The first is to identify specific target genes in a pest to induce high mortality rates and the other is to ensure the stability of the topically applied dsRNAs.¹ Protection of the dsRNA from environmental degradation like change in pH within the pest gut is a major aspect for feeding assays. Nanotechnology offers a wide range of applications in different fields of research which includes pharmaceuticals, cosmetics, electronics, food products, sensor-based devices, and environment.^{14,15} Similar to the drug delivery capacity, nanosystems also act as an important gene delivery agent which is now mostly being explored in the field of agriculture for disease resistance and pest control. Nanoparticles have proven to play an important role in delivery of genes or dsRNA into pests thus conferring gene silencing with high specificity and efficiency.¹⁶ Cationic polymer based nanoparticles offer an effective option for safe delivery of these dsRNAs into the pests as non-viral delivery agent. They can be easily produced, stored, and literally pose to be non-pathogenic which is an advantage of such polymers over other traditional carriers like viruses.¹⁷ In one of the approaches, nucleic acids are loaded into the nanoparticle either upon encapsulation into the matrix or chemically conjugated by appropriate surface modification. In the other approach, strong positive charges of cationic polymers efficiently bind to and condense the nucleic acids to form a polyplex. This condensation renders the polyplex to become a nanosystem, thus facilitating

the entry through the cellular membranes. The polyplexes are formed upon spontaneous electrostatic condensation between the nucleic acid and the cationic polymer. These polymers are also known to enhance the endosomal escape ability. Several mechanisms of their interaction with the cell membranes have already been documented in other reviews and is beyond the scope of this review.¹⁸

The current review emphasizes on the problem of pest infestations in agricultural crops and its impact on the economy. It highlights the role of nucleic acids like dsDNA, dsRNA as an active ingredient in pest control applications over traditional pest control systems. The authors also categorically focus on the use of cationic polymers over other agents for safe and efficient delivery of nucleic acids in pests *via* inducing gene silencing through RNAi approach.

2. Nucleic acid delivery in plants

Genetic engineering in plants is an attractive and efficient way to improve crops, biosynthesize different plant products and bring about sustainability in agriculture. Several biological, chemical, and physical approaches have been successfully used in the context. The delivery of exogenous nucleic acids/genes in plants can be either through vector-mediated gene transfer (indirect method) and vector-less gene transfer (direct method) (Fig. 1).

Vector-mediated gene transfer is carried out either by *Agrobacterium*-mediated transformation or by the use of plant viruses as vectors. This approach involves the pairing of the transgene with a vector (Ti-plasmid) that delivers it to the target cells for integration. *Agrobacterium tumefaciens* is a Gram-negative soil bacterium that infects dicotyledonous plants and induces crown gall tumors at affected sites. This exceptional ability of *Agrobacterium tumefaciens* aids in the advancement of




Fig. 1 Conventional methods of gene transfer in plants.



plant transformation techniques.¹⁹ According to the researchers, the segregation of GFP signals revealed that a variety of wild *Oryza* species can be genetically transformed by utilising modified immature embryo technique thereby confirming the transmission of T-DNAs to the following generation.²⁰ The bacterium because of its capability to transfer DNA into other organisms is a potential vector to produce transgenic plants that can confer resistance to some pests.

Non-*Agrobacterium* mediated gene transfer involves recruiting improved non-*Agrobacterium* strains such as *Rhizobium*, *Ensifer*, *Ochrobactrum haywardense* equipped with subtle mechanisms to deliver the gene into plant cells. The Transbacter technology can hasten the search for non-*Agrobacterium* species capable of transforming plants.²¹ Research findings demonstrated the propensity of trans-kingdom gene transfer with associated rhizobia; when equipped with a disarmed pEHA105 Ti plasmid and a binary plasmid (pCAMBIA1105.1R) that can be used to transfer T-DNA to a variety of plant species including *Arabidopsis thaliana* (model plant), *Nicotiana tabacum* (non-food crop), and *Oryza sativa* (food crop).²²

Vector-less gene transfer is carried out by physical (electroporation, biolistic, microinjection, silicon carbide fibre-mediated) and chemical (polyethylene glycol-induced and DEAE-dextran mediated) gene transfer methods.

In essence, electroporation entails the employment of electrical impulses with high field strength to reversibly permeate cell membranes for DNA absorption. DNA can be delivered into intact plant cells and protoplasts using this method. The oil palm cell of the *Elaeis guineensis* species was successfully transformed by the electroporation technique, which led to the formation of explants with increased growth rates.²³

Particle bombardment (or biolistic) commonly known as microprojectile or gene gun is a direct gene delivery technique that uses high-velocity micro-projectiles to deliver foreign DNA into plants. DNA-coated gold or tungsten microcarriers or microprojectiles are accelerated toward the target plant in order to pierce the cell wall. The transferred DNA separates from the microprojectiles after entering the cells, where it can be partially expressed or may be permanently incorporated into the host genome.²⁴

Microinjection follows a mechanical approach to transfer the desired DNA into the target plant cells. It is employed for chromosomal modification and gene transfer. In this method, the gene is introduced into a protoplast's cytoplasm using a glass micropipette and a transgenic plant is generated by culturing the modified cell.²⁵ In fact, this method has led to the development of transgenic tobacco and *Brassica napus*. *Agrobacterium tumefaciens* strain EHA105 containing the uidA gene was microinjected into shoot apex explants of cotton (*Gossypium hirsutum* L. 'KC3') and cultivated. In this experiment, the bacterial cell suspension was carefully microinjected 1–5 times into the pre-cultured apical meristem areas of shoot apices. Microinjections of an *Agrobacterium tumefaciens* cell density up to three times produced better results, but more than three injections caused severe meristematic damage and decreased the explants' survival rates.²⁶

Sonoporation is a technique for delivering genes to targeted cells by employing ultrasound that creates small pores in the plasma membrane thereby transferring the gene of interest into the cell. In this approach, the gene–microbubble combination improves the transfer efficiency while micro bubbles lower the threshold for cavity formation.²⁷ The expression of the hCTLA4Ig gene was suppressed using siRNA in transgenic cell cultures. The chemically synthesized siRNA duplex was coupled with polyethyleneimine and the cells were exposed to sonoporation at 40 kHz and 419 W for 90 s to enhance the delivery process. The sonoporation-delivered siRNA complexes down-regulated the synthesis of hCTLA4Ig by 73%. Therefore, it can be inferred that sonoporation may improve the delivery of siRNA complexes into plant cells.²⁸

Silicon carbide fibre-mediated transfer is a technique identical to microinjection in which the DNA is transported into the cell by using silicon carbide fibres. The silicon carbide fibres with DNA coating are vortexed with plant sample (suspension culture, calluses).²⁹ DNA attached to the fibres penetrates the cells during mixing and is successfully integrated with the host genome.³⁰ Silicon carbide whiskers with callus, plasmid harboring chitinase, and hygromycin genes were vortexed to deliver genes in peanut (*Arachis hypogaea*). In order to transform 2 g of 20 day old callus with the maximum transformation efficiency (6.88%), 200 mg of whiskers and 5 g of plasmid were employed. Hygromycin-resistant calli were grown into complete plants that produced seeds and had a far higher level of resistance to the leaf spot disease than control plants.³¹

In polyethylene glycol-mediated gene transfer the plasma membrane of protoplasts is destabilized by the Ca²⁺ ions and becomes permeable to DNA. Hence the naked DNA enters the nucleus and gets incorporated into the genome. The technique involves the protoplast isolation and suspension, addition of plasmid DNA, followed by gradual addition of 40% PEG-4000 (w/v) dissolved in calcium nitrate and mannitol solution. Protoplasts get transformed during incubation. Polyethylene glycol-mediated protoplast transfection was carried out with ribonucleoproteins comprising LbCas12a and a single guide RNA. Analysis of T1 offspring confirmed that DNA-free edits resulted at 40% frequency and the modifications are heritable.³²

Delivery of exogenous biomolecules into plants is quite a difficult process due to the barrier posed by the rigid plant cell wall. The conventional nucleic acid delivery methods for crop improvement and protection pose several drawbacks which include high cost of upstream production, difficulty in plant regeneration, and propagation of elite varieties.³³ Transgenic crops have issues related to non-acceptance among the public due to concerns regarding safety of human health, animals, and the environment.³⁴

Protection of crop from infestations by pests is an uphill task and needs utmost priority. Among all the crops, the ones belonging to the *Poaceae* family are considered economically most important which may be attributed to the large scale dependence of the human population to meet the food demands.



3. Susceptibility of plants to pests and conventional pest control measures

Nitrogen is very much essential for crops and helps in increasing the yield.³⁵ The high leachability of several forms of nitrogen in the soil makes them unavailable for absorption by crops and thus the need for exogenous application of the element through fertilizers is a common practice.^{36,37} Availability of nitrogen to plants makes them more succulent and thereby more prone to be fed upon by the pests.³⁸ Excessive use of fertilizers is also responsible for creating disturbances in the crop canopy as well as the balance between the plant and animal communities.³⁹ Increase in crop biomass also leads to increase in the crop density thereby providing suitable breeding habitat for the pests. Pest infestations are a major concern for cultivation of crops and there has been a surge in the use of chemical pesticides in recent times which may be attributed to the large scale cultivation of crops. Increased and reckless use of these pesticides has resulted in a significant loss in the biodiversity along with pollution of the water resources.¹⁰ Pesticides affect crops and plants to a great extent.⁹ Pest control forms an integral part of agricultural practices and is much important as far as crop productivity is concerned.^{40,41} Rising populations, increased demand for food, aided with the onset of green revolution in the early 1970s led to an increase in pest infestations.⁴² Practices of monoculture farming along with the use of chemical fertilizers, and pesticides became more frequent. Fertilizers and pesticides tend to lose their efficiency in the environment due to alterations in their chemical composition under high temperatures, and wash off due to heavy rainfall.⁴³ To counter the reduction in the effectiveness

the farmers tend to bring about an increase in the dosage and frequency of application of the agrochemicals, thus leading to development of resistance in pests.⁴⁴

There are several types of pesticides used in the agricultural practices which differ from each other based upon their chemical and physical nature. Drum (1980)⁴⁵ proposed the categorization of pesticides in three different ways mainly based on the origin, the target pest, and the pesticidal function as well. Natural or organic pesticides include plant phytochemicals, essential oils, and plant extracts that have been proven to be effective against several pests.⁴⁶ These compounds pose negligible toxicity towards mammalian cells, short environmental persistence, and prevent the development of pest resistance due to their chemical complexity.⁴⁷ Organic pesticides are although environment friendly but due to low persistence, can have varying effectiveness on targeted pests. High cost of organic pesticides is also a major issue that hinders its use on a field scale. Inorganic pesticides are generally composed of simple inorganic salts that have higher solubility in water as compared to the organic ones. Examples include sulphur, sulphates of metals like copper and iron, and lime.⁴⁸ The major issue with use of inorganic pesticides is their long environmental persistence and chances of being carried further in the food chain. These pesticides being rich in metal salts tend to be more toxic to the living biota especially the soil microbiota that play a major role in crop health and vigor (Fig. 2).

Man-made or synthetic pesticides are the most commonly used pest control agents in the field of agriculture. These include the organochlorines, organophosphates, carbamates, and pyrethroids. The mode of action of these pesticides on crop pests includes alteration of nervous functions, disruption of




Fig. 2 Impact of inorganic pesticides on soil microbes. (A) Inorganic pesticides are known to affect soil microbial population to a great extent. These pesticides have been found to negatively impact metabolic activities of the microbes thereby leading to death. Pesticides are known to sometimes promote the growth of a particular microbial species and retard others. This creates a disturbance in the diversity of microbial populations in the soil. Retardation of microbial growth due to application of inorganic pesticides also leads to reduced microbial biomass. The toxic nature of the pesticides not only hampers general soil microbes but also certain rhizospheric microbes colonizing the roots of the crops. The pesticidal toxicity reduces the nitrogen mineralization ability in the microbes thereby rendering the soil low on nutrient. (B) Holmsgaard *et al.* (2017)⁴⁹ studied the responses of a bacterial community to pesticides used over an agricultural season (March to September) in a farm. The microbial diversity of the soil was significantly reduced between the month of March to July as quite evident from the reduction in the operational taxonomic units (OTUs) and the Shannon diversity index thus signifying the negative impact of the pesticides on the soil microbial community of the agricultural farm. (B has been reproduced from Holmsgaard *et al.* (2017)⁴⁹ with permission from Elsevier, copyright 2017).






Fig. 3 Deleterious effect of synthetic pesticides on the biotic and abiotic components of the environment.

sodium channels, paralysis, and death. These synthetic pesticides, although have a pronouncing effect on a broad range of pests, but bear a long lasting deleterious effect on the environment as well as its components (Fig. 3).

4. Greener possibilities of pest control

Since their introduction in the late 1940s, synthetic pesticides have been widely used which may be attributed to their high efficacy, ease of application, and cost friendly nature. However, the large scale uses of pesticides have led to several deleterious effects on the environment. Some of the major effect includes reduction in soil quality, contamination of ground water, accumulation of toxic chemicals in the food chain, health disorders in humans, and destruction of the biodiversity.⁵⁰ The synthetic pesticides being non-specific in nature also tend to harm beneficial soil microbes, and other organisms along with the development of resistance among the pests.⁵¹ The increased resistance paves way for large scale destruction of crops thereby greatly affecting the production. Pests of the Lepidoptera family are one of the most harmful, accounting for approximately 10 million mega grams of loss in crop yield.⁵⁰

Green pest control technologies offer an alternative to the conventional chemical pesticides thereby preventing any sort of damage to the environment and its components. This is where the need to employ integrated pest management (IPM) comes into the limelight. IPM includes several strategies to control pests in agricultural fields. This may include either a single or combination of techniques involving genetic, mechanical, cultural, biological, and chemical tools.⁵²

4.1. Biological based techniques

Biological based IPM strategies are most feasible and environmental friendly way to control pests of crops. The bio-based techniques are mostly dependent on the environmental conditions and utilize a broad range of bio-agents like bacteria, viruses, fungi, and other predators.⁵⁰ Natural predators like spiders play a major role in keeping the pest population under control.⁵³ Natural chemicals obtained from living organisms like plants and microbes can also be instrumental in controlling the growth and development of the pests.⁵⁴ These chemicals are used as a major ingredient in the formulation of biopesticides and hold an extremely high value in a sustainable agricultural system. Plant extracts of *Azadirachta indica* has been successfully used as a bio-control agent for the control of brown plant hopper, thus leading to 82% of mortality.⁵⁵ Bt agent is a popular bio-insecticide mostly recommended and used in China to control the outbreaks of stem borer and leaf folders in rice plants.

4.2. Development of resistant varieties

Improving the resistance of crop varieties by conventional breeding approaches is also a very useful method of pest control in crops and is generally termed as the host-plant resistance mechanism. Researchers have identified 29 genes in rice plants resistance to the brown plant hopper.⁵⁶ The resistance of the conventionally breed varieties however weaken after certain generations leading to further development of resistance among the pests.⁵⁷

Genetically modified plant varieties have been found to overcome the disadvantages posed by the conventional breed varieties. Introduction of several genes into the plants have provided breakthrough results in pest control and management.



4.3. Non-biological techniques

Traps designed to kill pests utilizing specific frequencies of light have also been used to trap pests like adult stem borers and plant hoppers.⁵⁸ The major problem with these pest traps is that these lack specificity and will cause high mortality among other beneficial insects.

Post-harvest conditions are a major factor involved in pest outbreaks. Rice stubbles form the main breeding grounds for pests like stem borers and their population in the subsequent season is mostly dependent on the existing stubble environment. Mechanical harvesting of the stubble helps in reducing the pest populations. Reduction in the stubble height has been found to bring about a 70–90% reduction in the surviving pests.⁵⁸

Ecological engineering methods like growing nectar rich flowering plants can harbour several natural enemies of the pests.⁵⁹ These enemies will help control the pest population and their outbreaks thus preventing damage or yield loss.

5. RNA interference (RNAi) as a sustainable technique for pest control

Insect pests are a major threat to plants that directly increase the pressure on global food supply which already remains

affected due to the rise in population and other environmental problems. Chemical pesticides are although effective but have been found to cause collateral damage to the environment and its biotic counterparts. Moreover, they tend to kill other beneficial and non-target pests. Development of transgenic plants can counter these limitations however can lead to emergence of resistance among the pests which is also another major concern.⁶⁰ In such a scenario, pest control technologies based on RNA interference (RNAi) seems more promising due to their target specific nature.⁶¹ Three different types of RNAi pathways have been identified in insects which includes the siRNA pathway (involves dsRNA/siRNA), the miRNA pathway, and the piRNA pathway.^{62–64} The different pathways play different roles in the insects. The siRNA pathway protects the insect from viruses and transposons^{65,66} while the miRNA pathway plays a major role in the regulation of genes,⁶⁷ and the piRNA pathway suppresses the expression of germ line transposons.⁶⁸ Insect pests have been found to take dsRNA more rather than the siRNA through the process of clathrin mediated endocytosis.^{69,70}

The process of RNAi is comprised of two important steps. The first step involves the uptake of dsRNA by the cells of the pest followed by the second step which involves the processing of the same by the central RNAi machinery of the cell (Fig. 4). Hence, the cellular uptake is a major factor to be considered in RNAi based pest control strategies. dsRNA synthesized chemically can be applied directly to the leaf of a targeted plant in the




Fig. 4 Uptake of dsRNA and the RNAi machinery in pests. This figure has been reproduced from Liu *et al.* (2020)⁷² under a creative commons license (CC BY 4.0).



form of a foliar spray. Pests ingesting upon the leaf of the plant simultaneously intake the dsRNA which is thus directed into the lumen of the pests gut. Inside the gut, uptake of the dsRNA into the cells mostly occurs *via* the mechanism of clathrin dependent receptor mediated endocytosis. Besides, SID1 like (SIL) proteins, extracellular vesicles (EV), and RNA binding proteins (RBPs) that are secreted post the fusion of multi-vesicular bodies with the plasma membrane of the insect cells may also possibly help in the uptake process. Moreover, these molecules are also thought of as a probable mechanism for facilitating the movement of the dsRNA/silencing signal from one cell to the others thereby causing systemic RNAi based silencing in certain pests. However, the actual role of these molecules in the uptake of dsRNA is yet to be clearly elucidated.⁷¹ The cells lining the gut of the insect uptake the dsRNA

which is further cleaved by DICER enzyme into sRNA and then loaded into some specific members of the AGO protein family. This leads to formation of a RNA induced silencing complex (RISC) following which the guide strand of the sRNA promotes the binding of the RISC complex to the complementary target RNA. Recognition of the target results in post transcriptional gene silencing in the cytoplasm of the insect cell either by degradation of the target mRNA or by inhibiting its translation process. Sometimes transcriptional gene silencing in the nucleus of the pest cells may occur by modifications of chromatin.⁷²

Besides foliar application, RNAi can also be induced directly in the host plant (host induced gene silencing) or through viruses (viral induced gene silencing).⁷³ Host induced gene silencing involves the development of transgenic crops that

Table 1 Some targeted genes in RNAi based gene silencing in pests

Plant/crop	Pest	Target gene	RNAi type	References
Transgenic corn crop	<i>Diabrotica virgifera virgifera</i>	snf7	Host induced	61
Transgenic rice	<i>Lepidopteran</i> sp.	Cry (<i>Bacillus thuringiensis</i>)	Host induced	75
Transgenic rice	Brown plant hopper	Bph38(t), Bph37, Bph36, Bph34	Host induced	76
Maize	<i>Diabrotica virgifera virgifera</i>	Troponin I	Host induced	77
Corn	<i>O. furnacalis</i>	CHT10	Spray induced	78
Rice	<i>S. exigua</i>	Chitin synthase B	Spray induced	79
<i>Nicotiana attenuata</i>	<i>Manduca sexta</i>	MsCYPs	Virus induced	80
—	<i>Drosophila melanogaster</i>	Vha26, RPS13, and alpha COP	Virus induced	81




Fig. 5 Different types of RNAi based upon their silencing effects. In the cell autonomous RNAi, the dsRNA of a gene is generally applied to or expressed in a particular cell thus limiting the silencing effect to the cell. The non-cell autonomous RNAi includes systemic and environmental RNAi. Environmental RNAi involves uptake of the dsRNA from the environment and the effect can be observed in all the cells that are able to uptake the dsRNA. This type of exposure occurs by either soaking or feeding the targeted pest. In case of systemic RNAi the silencing signal is transported from the cell where the dsRNA is applied or expressed to other different cells or tissues of the organism thus spreading the silencing effect (this figure has been reproduced from Huvenne and Smaghe (2010)⁸² with permission from Elsevier, copyright 2010).



have the ability to express dsRNA against a specific pest. RNAi induced gene silencing approaches can also be carried out using genetically engineered viruses that can produce the desired dsRNA in the pest it targets.⁷⁴ The two later approaches involves transgenic organisms which is not widely acceptable over the globe and faces ethical issues. Moreover, development of transgenics requires high skill as well as the cost of development is too high. Table 1 elucidates some of the genes targeted in pests to induce the silencing effect, and the RNAi process involved therein.

Huvenne and Smagghe (2010)⁸² classified the RNAi process into two types based upon its silencing effect as – cell autonomous RNAi and non-cell autonomous RNAi (Fig. 5).

6. Limitations of RNAi based pest control

RNAi has become an important tool to silence targeted genes. It has several benefits when compared to other pest control agents available in the market. However, any novel inventions or products have certain levels of risk associated with them and so is the RNAi mediated pest control process (Table 2).

6.1. Impact on other beneficial insects

The RNAi process is although highly targeted and specific for a particular gene sequence, there remains a chance that it may sometimes affect a non-target insect that holds close genetic resemblance to the targeted pest. This may be a possible case if the non-targeted pest shares a common key gene with the pest and also has a close dsRNA sequence homology. The beneficial insect with respect to the pest if shares the common host and same feeding pattern, then could lead to its death.⁸³

6.2. Variation of susceptibility between species

Prediction or expecting a successful gene knockdown can be quite difficult as the susceptibility of pests to dsRNA is species dependent and may vary widely. When exposed to dsRNA, insect pests belonging to coleopteran species are mostly susceptible followed by the dipterans, and hymenopterans species. The

lepidopterans and hemipterans species are found to be very rarely susceptible.⁸⁴

6.3. Variation of susceptibility among species

Populations from within the same species show differential responses to external administration of dsRNA. Study conducted by Sugahara *et al.* (2017)⁸⁵ suggested that individuals belonging to the same laboratory strain can have different degrees of responses towards dsRNA. Similar results were also confirmed by other researchers.^{86,87}

6.4. Tissue dependent variation

Several inconsistencies have been observed in RNAi response within the same insects. The susceptibility of the insect pests to dsRNA mediated gene silencing can vary in its degree from tissues to tissues or cell to cell.⁸⁸ Telang *et al.* (2013)⁸⁹ reported lower RNAi efficiency in the ovarian and head tissues of *Aedes aegypti* as compared to other tissues upon external application of dsRNA. Tissue dependent RNAi susceptibility has also been observed in lepidopteran species.⁹⁰

6.5. High cost of production

The production/synthesis of target specific dsRNA for RNAi against crop pests is a costly affair and thus will increase the price of the final product available for pest control in crops like rice plants.¹

6.6. Extracellular degradation of dsRNA

Some insects on feeding upon dsRNA have almost no effect. The dsRNA upon reaching the digestive system of the insects get degraded by the nucleases present thus inhibiting the RNAi activity.⁹¹ A study by Wynant *et al.* (2014)⁹² demonstrated the dsRNA degrading ability of digestive solution obtained from the midgut of *Schistocerca gregaria*. The digestive enzyme solution was found to degrade almost 150 ng of the dsRNA within 5 minutes of exposure.

Table 2 Limitations of RNAi based approaches in pest control

Limitations of RNAi	Probable reasons	Reference
Lethal impact on beneficial insects	Non-targeted pest sharing common key genes/ similar dsRNA sequence homology with the target pest	83
Lack of uniform susceptibility between species	Enzymatic degradation of dsRNA, exposure dosage, impaired RNAi machinery, presence of virus in the target pest	84
Variation in RNAi responses within the same insect	Insufficient spread of RNAi response throughout the body, variation in pH among different organs	85 and 87
High production costs	High cost of chemicals, equipment, and maintenance conditions	1
Extracellular degradation of dsRNA	Degradation by nucleases present in the digestive system	91 and 92



7. Tools to overcome RNAi instability

The problems faced due to low RNAi sensitivity in the targeted pests needs to be looked upon seriously before applying it as a potential pesticidal agent. Facilitation of uptake of dsRNA and preventing its degradation can be carried out by means of efficient delivery systems. Some of the recent delivery systems include microorganisms like bacteria, and viruses, nanoparticle based carriers, liposomes, carrier proteins, and chemical modifications.⁹³

Microorganisms specifically genetically modified bacteria that lack the RNaseIII endonuclease have been used to deliver dsRNA within the insect cells.⁹⁴ The bacterial shell is thought of probably providing a protective effect to the dsRNA inside the digestive system. Pre-treatment of the bacterial cell by sonication have been found to improve the dsRNA release inside the insect most probably by weakening the bacterial cell wall.⁹⁵ The selection of bacteria for the purpose is an important factor and only symbiotic bacteria or yeasts must be selected to avoid any potential pathogenicity to other organisms.⁹⁶ Viruses can also be used as an effective tool for the

delivery of dsRNA into the intracellular environment in pests. Viruses have been found to be specific to certain particular hosts and thus can be carefully selected as a successful delivery agent in RNAi experiments.⁷⁴ Despite possessing several advantages as a delivery system for dsRNA, the application of the viral carriers *in vivo* has not yet been fully investigated due to many concerned safety issues. All the viruses are not host specific and may pose the chance of cross infecting several other beneficial insects which is a potential biosafety issue.

Nanoparticles can also be used as a delivery agent wherein dsRNA can be incorporated into these particles to enhance the stability and uptake efficiency. Chitosan derived nanoparticles have been found to efficiently deliver dsRNA through oral routes in *A. gambiae* and *A. aegyptia* resulting in knockdown of genes.⁹⁷ Synthetically modified polymer nanoparticles have also been used to a great extent. Uptake of dsRNA complexed with a fluorescent nanoparticle led to RNAi silencing of CHT10 gene in the larvae of Asian corn borer (*O. furnacalis*). The fluorescent nanoparticle not only helped to visualize the dsRNA after uptake but also prevented the aggregation of the same in




Fig. 6 Functionalized polymers and increased RNAi efficiency. (A) Schematic diagram showing steps in formation of cationic polymer based polyplexes with higher stability and RNAi efficiency. In a recent study several different combinations of polyplexes were designed by Christiaens *et al.* (2018)⁷⁹ and were characterized. Based upon the N/P ratios, charge, size, and degradation assay the polymer PAG87 was selected as the most suitable candidate for polyplex formation with the dsRNA. (B) A 100% complexation was achieved at N/P ratios of 2:1 and 4:1 as evident from the agarose gel data. (C) The zeta potential data also verified the same. (D) Results from an *ex vivo* assay revealed the protective nature of PAG87 due to the high content of guanidine against nucleolytic degradation of dsRNA. The polyplex when incubated in the gut juice obtained from *S. exigua* at pH 11 for different time periods followed by decomplexation of the dsRNA and analysis over a 1.5% agarose gel revealed protection of the dsRNA for as long as 30 hours. This supports the possible use of functionalized polymers like PAG87 in successful protection of the dsRNA at higher alkaline pH from degradation thus increasing the RNAi efficiency. (B–D) has been reproduced from Christiaens *et al.* (2018)⁷⁹ under a creative commons license (CC BY 4.0).



water.⁷⁸ Several nanoparticles have been designed from cationic polymer based derivatives functionalized with guanidine side groups. The functionalization helped protect the dsRNA from degradation under high alkaline pH which is a characteristic feature of the lepidopteran gut environment (Fig. 6). The pH stable nanoparticles when fed to larvae of *S. exigua* resulted in the knockdown of chitin synthase B gene thus leading to increased mortality.⁷⁹

Lipid based transfection agents referred to as 'liposomes' are also very instrumental in increasing RNAi efficiency. Formation of liposomes occurs naturally when the transfection agents are subjected to an aqueous environment. Bilayer lipid particles are formed when positively charged lipid molecules envelope the negatively charged dsRNA.⁹⁸ The liposome encapsulated dsRNA is facilitated entry into the cell through lipofection. Zhang *et al.* (2018)⁹⁹ had successfully performed liposome mediated uptake of dsRNA in *Rhipicephalus haemaphysaloides*.

Carrier proteins also called as cell penetrating peptides are also an excellent prospect for delivery of dsRNA into pest cells. These peptides are cationic in nature and comprises of short chains of amino acids (10–30) with a high occurrence of basic amino acid residues like lysine and arginine.¹⁰⁰ The cationic peptides facilitate the entry into the intracellular environment of the pest most possibly by endocytosis along with transporting the dsRNA. However, the exact cellular mechanism behind the carrier proteins mediated dsRNA uptake is still fuzzy.¹⁰¹ In a study by Gillet *et al.* (2017)¹¹⁵ to induce RNAi response in

Anthonomus grandis, a fusion protein comprising of a peptide transduction domain and dsRNA binding domain from human protein kinase R was designed. The protein transduction domain and dsRNA binding domain along with the dsRNA forms a ribonucleic protein particle which in turn facilitates the uptake into the gut of the insect. The ribonucleic protein particle was found to increase the knockdown of chitin synthase II gene in *A. grandis* as compared to the naked dsRNA.

Small RNA oligonucleotides like siRNA are not effective enough to initiate a RNAi response in pests, however chemically modifying these molecules has been found to improve their activity in terms of stability and uptake.¹⁰² Modified siRNAs targeting important genes in *Plutella xylostella* have been found to result in increased mortality.¹⁰³

8. Cationic polymers in RNAi based pest control

Cationic polymers are the positive charge bearing macromolecules. The charges may either be present in the backbone or in the side chains of the polymer. Most of these polymers contain functional amine groups that could be protonated.¹⁰⁴ Cationic polymers have been found to be quite instrumental in several fields which involves drug delivery, gene delivery, and as anti-microbial agents.

Cationic polymers act as an efficient non-viral agent for transfer of DNA material into the pests. Due to high positive




Fig. 7 Cationic nano-polymer based delivery of dsRNA/siRNA in pests. This figure has been reproduced from Yan *et al.* (2021)¹⁰⁹ under a creative commons license (CC BY-NC-ND 4.0).



charge densities, the cationic polymers interact with the negatively charged dsDNA thus forming stable complexes termed as 'polyplexes' or 'nanoplexes'.¹⁰⁵ The cationic polymers protect the dsDNA from degradation by enzymes and high alkaline pH present in the gut microenvironment of the pests. Moreover, the cationic polymers are biodegradable, less toxic, are structurally diverse, and pose high transfection efficiency.¹⁰⁶ These polymers facilitate internalization into cellular compartments and endosomal escape through a mechanism known as proton sponge¹⁰⁷ (Fig. 7). Cationic polymers also help in controlled release and the net positive charge of the complex facilitates binding to the anionic proteoglycans present on the cell surfaces.¹⁰⁸ The polyplex can be used as a formulation for spraying on leaves of the crops. The complex either is ingested or enters the pest through dermal penetration. The polyplex after binding to the cell membrane of the insect enters into the cell by endocytosis. Post the uptake by the cells, the polyplex generated endocytic vesicles travel through the microtubules to fuse with early endosomes which further mature into late endosomes at pH = 5.0–6.2. These polyplex fused endosomes finally enter into degradative lysosomes. The polyplex here needs to exit the endosome to prevent degradation by the lysosome which is carried out by a mechanism known as 'proton sponge effect'. Cationic polymers in general have a strong buffering capacity. The acidic environment inside the lysosome causes protonation of the amine groups

present in the polymer, leading to influx of water and lysis of the endosome thus releasing the dsDNA into the cytosol. The next step involves unbinding of the dsDNA from the polymer nanoparticle which occurs *via* competitive displacement of the polymer from the dsDNA by certain intracellular polyanions. The polymeric nanoparticles can also be designed to respond to intracellular stimuli like pH and certain reducers thus inducing the disassembly process.

Several types of cationic polymers based upon their structural variations have been used for control of pests in crops. The current section summarizes some of the important cationic polymers used in pest control which includes linear homopolymers (LP), branched polymers (BP), and guanylated nanoparticles (GNP)¹⁶ (Fig. 8). LPs (linear homopolymers) are the simplest type of cationic polymers used to impart RNAi effect. Some commonly used LPs are poly[2(dimethylamino)ethyl methacrylate] (pDMAEMA), polyethyleneimine (PEI), and poly-L-lysine (PLL).¹¹⁰ These type of polymers are made up of a single monomer that contains an amine group inside the polymeric chain backbone. The high pK_a of these cationic polymers (pDMAEMA: 7.4–7.8, PEI: 8.2–9.5, PLL: 9–11) favours the complexation with the phosphate backbone of the dsRNA through electrostatic forces. LPs till date have not been found to be efficient in the complexation and protection of the dsRNA besides the issue of non-specific cytotoxicity which may be




Fig. 8 Structural variation based different cationic polymers generally used in RNAi based plant pest control.



a major concern for other beneficial organisms. The cytotoxicity effect may arise due to interaction of the polymer with cell membranes in organisms, leading to formation of pores and eventually cell death. To overcome the drawbacks posed by LPs, several architectural and functional modifications have been designed to enhance dsRNA uptake, maintain stability, and reduce the cytotoxicity. The uses of star shaped polymers for dsRNA delivery in insects have been studied by several researchers.^{84,111} The issues have also been tried to be overcome by simple alteration in the chemistry of the polymers which includes but not limited to variation in the molecular weight, charge density, ionic strength, and pK_a .¹¹²

Branched polymers (BPs) were designed with the aim to improve the transfection and RNAi efficiency and reducing the levels of non-specific cytotoxicity as observed in case of LPs. Star cationic polymers with 3–5 arms shows reduced cytotoxicity as the branching increases as most of the nitrogen atoms involved in complexation remains within the dense core of the polymer. However, a higher amount of BPs is required to stabilize the DNA when compared with the LPs due to less availability of nitrogen complexing moieties.¹¹³

Several designs and developments have been made to functionalized polymers to increase their efficacy as a dsRNA vector. Certain pH responsive polymers have been prepared that releases the dsRNA within the endosome compartment of the cell by undergoing conformational changes due to a transition in pH.¹¹⁴ The pH transition range of the polymers however may not work for pest control strategies. Pests belonging to Lepidoptera family have a very alkaline intestinal gut pH and therefore may destabilize the dsRNA thus decreasing its efficacy. Polymers designed to protect the dsRNA under highly alkaline conditions need to seriously take into account the impact of pH on the complexation of polymers. Guanidine functionalized cationic polymers have been designed to protect the dsRNA over a high range of alkaline pH which is a characteristic feature of the lepidopteran gut microenvironment. These polymers facilitate in endocytic passage of the RNA through the cell membranes and escape from the endosomes.¹¹⁵ PGPMA (poly-[N-(3-guanidinopropyl)methacrylamide]) has been found to have a pK_a of 12.5 thus ensuring the protonation of guanidinium functional groups at high alkaline gut pH. Parsons *et al.* (2018),⁶⁰ complexed PGPMA with dsRNA to form a compact polyplex at pH 10. The polyplex brought about a 92% reduction in the CDC27 mRNA in Sf9 cell lines post 48 hours of incubation. Feeding assays on 2nd and 3rd instar larvae of *S. frugiperda* with a diet supplemented with the polyplex (PGPMA/CDC27) for a period of 7 days resulted in approximately 30% mortality after 29 days. In another study a series of copolymers of poly-N-2-aminoethylmethacrylate (PAEMA) and PDMAEMA were synthesized by free radical polymerization. The synthesized copolymers were guanidium functionalized by reacting 1H-pyrazole-1-carboxamide hydrochloride (HPC) with part of primary amine moieties of the PAEMA. The copolymer with highest guanidine content showed higher protection efficiency towards the dsRNA when incubated with the larval midgut juice of *S. exigua* (pH 7.5 and 11).⁷⁹ The guanylated copolymer when complexed with the dsRNA of the chitin synthase (ChSB) gene

protected the dsRNA for as long as 10 hours in the gut juice of *S. exigua* (pH = 11). *In vivo* feeding assays supported the use of guanidine functionalized polyplexes in enhancing the efficiency of RNAi in *S. exigua*. The polymer protected dsRNA was found to exhibit 53% mortality as compared to a low mortality rate of only 16% in case of the naked dsRNA. The association of guanidine functional groups with the copolymer has been proved to likely provide enhanced dsRNA protection and thus increase the efficacy of RNAi. A study conducted by Gurusamy *et al.* (2020)¹¹⁶ evaluated the role of chitosan–dsRNA polyplex in improving RNAi in *Spodoptera frugiperda*. The complex showed reduced accumulation in the endosomes of the Sf9 cells and in the larval tissues thus exhibiting the protective property of chitosan. Moreover, the polyplex when fed to the *S. frugiperda* larvae resulted in successful knockdown of the *iap* gene, thus leading to retardation of growth and mortality among the larvae. In a similar fashion, Wang *et al.* (2020)¹¹⁷ studied the role of cationic polymers chitosan and lipofectamine 2000 to specifically target the glyceraldehyde-3-phosphate dehydrogenase (G3PDH) gene in the rice pest *Chilo suppressalis*. The chitosan–G3PDH conjugate when fed to the 2nd instar larvae brought about a 45% reduction in the expression of the gene in the pest gut. Similarly lipofectamine 2000 reduced the expression of the targeted gene to 52% in the gut tissues.

The above discussions aptly support the role of polyplexes (cationic polymer–dsRNA complex) in maintaining the stability of the dsRNA in both *in vitro* and *in vivo*. Greater stability of the complex results in higher RNAi efficiency as observed in the several studies.

Cationic polymers have been known to provide several advantages over other pest control agents. The cationic polymers act as an excellent non-viral delivery agent thereby reducing the cost. They are quite easy to produce, preserve, and exhibit no potential pathogenicity unlike the viral carriers.¹⁷ Cationic polymers are important non-viral transfection agents. Cationic polymers due to their strong positive charges can attract and condense the dsDNA thus facilitating the entry through the cellular membranes. These polymers also enhance the endosomal escape ability.¹⁸ The cationic polymers when chemically modified behave as a more target specific unit, reduce toxicity, and improve the efficiency of transfection.¹¹⁸

The choice of cationic polymers for pest control is a crucial aspect to be considered while working on RNAi based pest control in crops. The cationic polymers to be used should not be toxic to the environment and its components. Moreover the selected polymers should be biodegradable in nature thereby avoiding environmental persistence. The physicochemical properties of the polymers can be used as a base for deciding on its usefulness as a delivery agent in pest control. A particle size of the polymer at nanoscale, higher numbers of positive surface charges, and a suitable spatial framework are favourable parameters for selection of the polymers.¹⁷

9. Alternatives to cationic polymers

The efficacy of RNAi mechanism purely depends upon the delivery or uptake of the intact dsRNA into the cells of the pests.



Table 3 Alternatives to cationic polymers for delivery of dsRNA

Delivery methods	Target species	Molecular impact	Effect	References
Cationic liposome	<i>D. melanogaster</i> , <i>D. sechellia</i> , <i>D. yakuba</i> , and <i>D. pseudoobscura</i> larvae	3' UTR of γ -tubulin gene	Mortality	119
Lipofectamine (liposome)	<i>Drosophila suzukii</i>	mRNA silencing	40–50% silencing	120
Chemical modification	—	Addition of methyl group to the 2' of ribosyl ring of 2nd base of siRNA	Increase in specificity of the dsRNA	121
Root drenching and trunk injection	<i>Diaphorina citri</i>	Silencing of arginine kinase (dsRNA-AK)	Increased mortality	122
	Brown plant hopper	Knockdown of carboxylesterase (Ces) and cytochrome P450 (Cyp18A1)	High mortality among BPH nymphs	123
	<i>Ostrinia furnacalis</i>	Silencing of Kunitz-type trypsin inhibitors (dsKTI)	High mortality rate	123
Application of bacteria and viruses	<i>Rhodnius prolixus</i>	Initiation of RNAi upon ingestion of recombinant bacteria	Knockdown of horizontally transmissible phenotypes	96
	<i>Bactericera cockerelli</i>	Recombinant TMV targeting actin and V-ATPase sequences	Decrease in mRNA abundance and progeny production	124
Engineered/transplastomic plants	<i>Leptinotarsa decemlineata</i>	—	100% larval mortality	97

There are several options other than cationic polymers to reduce the degradation of the dsRNA as well as increasing the cellular uptake efficiency. These includes liposomes, bringing chemical modifications, absorption into plants *via* roots, direct

injection into vessels, involvement of bacteria and viruses, and development of engineered or transplastomic plants (Table 3).

Liposomes being non-toxic and biodegradable in nature have been found to safely deliver exogenous RNA to the target




Fig. 9 Cationic liposomes as efficient transfection agents in RNAi. (A) The cationic lipids/liposomes are amphiphilic molecules with overall positive charge. They bind to the nucleic acid as well as cell membranes with electrostatic interactions. The lipid based complex enters the target cell through endocytosis and is subsequently released into the cytoplasm. If the exogenous nucleic acid to be supplied is a DNA, then it needs to be transported into the nucleus while in case of a mRNA molecule it stays within the cytoplasm.¹²⁵ (B) A study by Whyard *et al.* (2009)¹¹⁹ demonstrated the effect of different transfection agents and dsRNA concentration on the GUS gene in the gut of *D. melanogaster* larvae. The transfection agents when each coupled with a GUS-dsRNA concentration of 0.5 mg ml⁻¹ brought about an improved silencing effect on the GUS gene as compared to the dsRNA alone. Among all the combinations, Lipofectamine 2000 coupled dsRNA showed maximum GUS silencing activity (more than 50%) as evident from the graph. (Fig. 9B has been reproduced from Whyard *et al.* (2009)¹¹⁹ with permission from Elsevier, copyright 2009).



cells. Several researchers have demonstrated the effectiveness of liposomes in the RNAi process (Fig. 9). Lipofectamine, a liposome was successfully used as a transfection agent by Taning *et al.* (2016)¹²⁰ in *Drosophila suzukii*. Feeding naked dsRNA did not yield any result whereas lipofectamine complexed dsRNA led to a silencing efficiency of 40–50%. A similar study by Whyard *et al.* (2009)¹¹⁹ demonstrated successful RNAi silencing effect in four different species of *Drosophila* when fed with γ Tub23C-dsRNA encapsulated with different cationic liposomes.

Chemically modifying any one or both of the strands of the dsRNA can improve the stability of the molecule. It can also help in increasing the shelf life, bio-distribution, and specificity. However, the cost of production and safety concerns needs to be assessed prior to the modifications.

RNAi based silencing could also be initiated by supply of dsRNA *via* absorption in roots of the plants or injecting the same into the trunk or vessels. The sucking and chewing pests thereby acquire the dsRNA naturally.¹²⁶ Hunter *et al.* (2012)¹²² exposed citrus plant to dsRNA by means of root drenching and injection into the trunk. 2 g of dsRNA in 15 L of water was applied to the citrus plants and could be observed in the plant vessels until 7 weeks post the treatment. The experiment also demonstrated two hemipteran species and a leafhopper taking up the dsRNA feeding on the treated plant. Strategies like root absorption or trunk injection have some serious concerns associated. For the purpose, production of dsRNA in large mass is required thus making it a costly affair. Root application of dsRNA could be carried out on large fields through irrigation however the problem is with the short lived nature of the dsRNA in the soil. Trunk injection is more appropriate for sap sucking insects over the chewing insects. Treatment of crops like rice which are cultivated on a large scale by the method of trunk injection is almost impractical. Both trunk injection and root absorption requires repeated application at regular intervals and thus another drawback of the method.¹²⁷

Bacteria mediated delivery of dsRNA pose several advantages which includes low cost and possible large scale production. Continuous and large scale production of dsRNA is made possible by the bacterial species. Viruses can also act as an efficient vector for the production of dsRNA. Several plant viruses have been studied for triggering RNAi in plants.^{62,124} Plants react to infections caused by viruses through the siRNA pathway. Introduction of an insect specific RNAi inducer sequence into a plant virus will produce siRNAs specific to that insect. Insect feeding on the plant can uptake the siRNA thus leading to silencing effect and mortality.

Long dsRNA are required for an effective RNAi activity in insects. However, the dsRNA expressed in plants are mostly diced into siRNAs and then taken up by insects leading to a limited RNAi effect.¹²⁸ This problem can be overcome by engineering plants to express dsRNA in the organelles like chloroplasts which lack the RNAi processing ability. The chloroplasts are derived from cyanobacteria that lack RNAi pathway thus accumulating dsRNA.¹²⁹

9.1. Prospects and challenges

The world is facing an expansion of population and the soaring population has put tremendous pressure on the food chain. In order to meet with the food shortage the agricultural productions have become of paramount importance. The traditional problems of lower agricultural production such as pests *etc.* have been taken care of by the use of hormones and chemical fertilizers. These abusive use of pesticides has caused more harm than good in de-balancing the environment. The use of pesticide has not only contributed to environmental pollution but also deteriorated the quality of agricultural products.

With the growth in research tools in molecular biology, it is pertinent that scientists use such innovative tools to address the lacunae in the field. The field of RNAi offers some exciting solutions and potentials. There is tremendous scope with a lot of research to be done in the use of nucleic acid based techniques especially to improve their half-life. With a multidisciplinary approach and efforts of experts from diverse fields nucleic acids can be efficiently used for agricultural utility with solution to major problems of food, population, and pollution.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Mamoni Dash acknowledges the Ramalingaswami Fellowship 2016–17 (D.O.NO.BT/HRD/35/02/2006), Department of Biotechnology, Government of India. The funding from Biotechnology Industry Research Assistance Council (BIRAC) is highly acknowledged for the funding through the PACE grant BFD/RO/B.04/0350/2021-22, Institute of Life Sciences (ILS), Department of Biotechnology funded project BT/PR 36474/NNT/28/1701/2020, Government of India. We highly acknowledge Dr Amaresh C Panda for useful discussions regarding nucleic acids.

References

- 1 S. J. Fletcher, P. T. Reeves, B. T. Hoang and N. Mitter, A Perspective on RNAi-Based Biopesticides, *Front. Plant Sci.*, 2020, **11**, 51.
- 2 J. A. Heinemann and S. Walker, Environmentally applied nucleic acids and proteins for purposes of engineering changes to genes and other genetic material, *Biosafety and Health*, 2019, **1**(03), 113–123.
- 3 A. Székács, A. S. Ammour and M. L. Mendelsohn, *RNAi Based Pesticides*, Frontiers Media SA, 2021, p. 714116.
- 4 J. S. Whangbo and C. P. Hunter, Environmental RNA interference, *Trends Genet.*, 2008, **24**(6), 297–305.
- 5 S. Ivashuta, Y. Zhang, B. E. Wiggins, P. Ramaseshadri, G. C. Segers, S. Johnson, *et al.*, Environmental RNAi in herbivorous insects, *RNA*, 2015, **21**(5), 840–850.
- 6 N. Bensoussan, S. Dixit, M. Tabara, D. Letwin, M. Milojevic, M. Antonacci, *et al.*, Environmental RNA interference in



- two-spotted spider mite, *Tetranychus urticae*, reveals dsRNA processing requirements for efficient RNAi response, *Sci. Rep.*, 2020, **10**(1), 1–16.
- 7 N. Bensoussan, M. Milojevic, K. Bruinsma, S. Dixit, S. Pham, V. Singh, *et al.*, Localized efficacy of environmental RNAi in *Tetranychus urticae*, *Sci. Rep.*, 2022, **12**(1), 1–14.
 - 8 M. Mondal, J. Peter, O. Scarbrough and A. Flynt, Environmental RNAi pathways in the two-spotted spider mite, *BMC Genomics*, 2021, **22**(1), 1–11.
 - 9 A. Sharma, V. Kumar, R. Kumar, B. Shahzad, A. K. Thukral and R. Bhardwaj, Brassinosteroid-mediated pesticide detoxification in plants: A mini-review, *Cogent Food Agric.*, 2018, **4**(1), 1436212.
 - 10 K. C. Parsons, P. Mineau and R. B. Renfrew, Effects of pesticide use in rice fields on birds, *Waterbirds*, 2010, **33**(sp1), 193–218.
 - 11 J. A. Anderson, P. C. Ellsworth, J. C. Faria, G. P. Head, M. D. Owen, C. D. Pilcher, *et al.*, Genetically engineered crops: importance of diversified integrated pest management for agricultural sustainability, *Front. Bioeng. Biotechnol.*, 2019, **7**, 24.
 - 12 J. Niu, C. Taning, O. Christiaens, G. Smagghe and J. Wang, Rethink RNAi in Insect Pest Control: Challenges and Perspectives, *Advances in Insect Physiology*, 2018, pp. 1–17.
 - 13 D. Biedenkopf, T. Will, T. Knauer, L. Jelonek, A. Furch, T. Busche, *et al.*, Phloem-mediated spreading of SIGS-derived non-coding RNAs in *Hordeum vulgare*, *bioRxiv*, 2019, DOI: [10.1101/2019.12.30.891002](https://doi.org/10.1101/2019.12.30.891002).
 - 14 P. K. Das, C. Mohanty, G. K. Purohit, S. Mishra and S. Palo, Nanoparticle assisted environmental remediation: Applications, toxicological implications and recommendations for a sustainable environment, *Environ. Nanotechnol., Monit. Manage.*, 2022, **18**, 100679.
 - 15 J. Li and K. Kataoka, Chemo-physical strategies to advance the *in vivo* functionality of targeted nanomedicine: the next generation, *J. Am. Chem. Soc.*, 2020, **143**(2), 538–559.
 - 16 C. E. Pugsley, R. E. Isaac, N. J. Warren and O. J. Cayre, Recent Advances in Engineered Nanoparticles for RNAi-Mediated Crop Protection Against Insect Pests, *Front. Agron.*, 2021, **3**, 652981.
 - 17 N. Bono, F. Ponti, D. Mantovani and G. Candiani, Non-Viral *in Vitro* Gene Delivery: It is Now Time to Set the Bar, *Pharmaceutics*, 2020, **12**(2), 183.
 - 18 Q. Huang, S. Li, Y.-F. Ding, H. Yin, L.-H. Wang and R. Wang, Macrocyclic-wrapped polyethylenimine for gene delivery with reduced cytotoxicity, *Biomater. Sci.*, 2018, **6**(5), 1031–1039.
 - 19 A. H. A. Rashid and D. D. Lateef, Novel techniques for gene delivery into plants and its applications for disease resistance in crops, *Am. J. Plant Sci.*, 2016, **7**(1), 181–193.
 - 20 S. Shimizu-Sato, K. Tsuda, M. Nosaka-Takahashi, T. Suzuki, S. Ono, K. N. Ta, *et al.*, Agrobacterium-mediated genetic transformation of wild *Oryza* species using immature embryos, *Rice*, 2020, **13**(1), 1–13.
 - 21 W. Broothaerts, H. J. Mitchell, B. Weir, S. Kaines, L. Smith, W. Yang, *et al.*, Gene transfer to plants by diverse species of bacteria, *Nature*, 2005, **433**(7026), 629–633.
 - 22 D. S. Rathore and E. Mullins, Alternative Non-Agrobacterium based methods for plant transformation, *Annu. Plant Rev.*, 2018, 891–908.
 - 23 C. Darmawan, N. M. A. Wiendi, C. Utomo and T. Liwang, Electroporation-mediated genetic transformation of oil palm (*Elaeis guineensis*), *Biodiversitas*, 2020, **21**(8), 3720–3726.
 - 24 I. I. Ozyigit and K. Yucebilgili Kurtoglu, Particle bombardment technology and its applications in plants, *Mol. Biol. Rep.*, 2020, **47**(12), 9831–9847.
 - 25 H. Matsuoka, T. Komazaki, Y. Mukai, M. Shibusawa, H. Akane, A. Chaki, *et al.*, High throughput easy microinjection with a single-cell manipulation supporting robot, *J. Biotechnol.*, 2005, **116**(2), 185–194.
 - 26 P. Gurusaravanan, S. Vinoth and N. Jayabalan, An improved Agrobacterium-mediated transformation method for cotton (*Gossypium hirsutum* L. 'KC3') assisted by microinjection and sonication, *In Vitro Cell. Dev. Biol.: Plant*, 2020, **56**(1), 111–121.
 - 27 M. Tomizawa, F. Shinozaki, Y. Motoyoshi, T. Sugiyama, S. Yamamoto and M. Sueishi, Sonoporation: Gene transfer using ultrasound, *World J. Methodol.*, 2013, **3**(4), 39.
 - 28 S.-H. Cheon, K.-H. Lee, J.-Y. Kwon, S.-H. Choi, M.-N. Song and D.-I. Kim, Enhanced delivery of siRNA complexes by sonoporation in transgenic rice cell suspension cultures, *J. Microbiol. Biotechnol.*, 2009, **19**(8), 781–786.
 - 29 H. F. Kaeppeler, W. Gu, D. A. Somers, H. W. Rines and A. F. Cockburn, Silicon carbide fiber-mediated DNA delivery into plant cells, *Plant Cell Rep.*, 1990, **9**(8), 415–418.
 - 30 D. Songstad, D. Somers and R. Griesbach, Advances in alternative DNA delivery techniques, *Plant Cell, Tissue Organ Cult.*, 1995, **40**(1), 1–15.
 - 31 Z. Akram, S. Ali, G. M. Ali, Y. Zafar, Z. H. Shah and F. Alghabari, Whisker-mediated transformation of peanut with chitinase gene enhances resistance to leaf spot disease, *Crop Breed. Appl. Biotechnol.*, 2016, **16**, 108–114.
 - 32 V. Sidorov, D. Wang, E. D. Nagy, C. Armstrong, S. Beach, Y. Zhang, *et al.*, Heritable DNA-free genome editing of canola (*Brassica napus* L.) using PEG-mediated transfection of isolated protoplasts, *In Vitro Cell. Dev. Biol.: Plant*, 2022, **58**(3), 447–456.
 - 33 J. M. Dunwell, Transgenic approaches to crop improvement, *J. Exp. Bot.*, 2000, **51**(suppl_1), 487–496.
 - 34 R. E. Goodman, S. Vieths, H. A. Sampson, D. Hill, M. Ebisawa, S. L. Taylor, *et al.*, Allergenicity assessment of genetically modified crops—what makes sense?, *Nat. Biotechnol.*, 2008, **26**(1), 73–81.
 - 35 J. Gu and J. Yang, Nitrogen (N) transformation in paddy rice field: Its effect on N uptake and relation to improved N management, *Crop and Environment*, 2022, **1**(1), 7–14.
 - 36 K. Jantapoa, S. Pinita, L. Zhouc, W. Wangc and J. Chaiwanonb, Effects of propiconazole on rice growth and gene expression in response to nitrogen and phosphorus deficiencies, *ScienceAsia*, 2021, **47**, 19–27.



- 37 Z. Jia and N. von Wirén, Signaling pathways underlying nitrogen-dependent changes in root system architecture: from model to crop species, *J. Exp. Bot.*, 2020, **71**(15), 4393–4404.
- 38 Z.-x. Lu, X.-p. Yu, K.-l. Heong and C. Hu, Effect of Nitrogen Fertilizer on Herbivores and Its Stimulation to Major Insect Pests in Rice, *Rice Sci.*, 2007, **14**(1), 56–66.
- 39 G. Conway, *The doubly green revolution: food for all in the twenty-first century*, Cornell University Press, 2019.
- 40 E. Cardarelli and G. Bogliani, Effects of grass management intensity on ground beetle assemblages in rice field banks, *Agric., Ecosyst. Environ.*, 2014, **195**, 120–126.
- 41 N. Brzezina, B. Kopainsky and E. Mathijs, Can Organic Farming Reduce Vulnerabilities and Enhance the Resilience of the European Food System? A Critical Assessment Using System Dynamics Structural Thinking Tools, *Sustainability*, 2016, **8**(10), 971.
- 42 J. Mariyono, Green revolution- and wetland-linked technological change of rice agriculture in Indonesia, *Manag. Environ. Qual.*, 2015, **26**(5), 683–700.
- 43 I. K. A. Zwertvaegher, I. Van Daele, P. Verheesen, M. Peferoen and D. Nuyttens, Development and implementation of a laboratory spray device and rainfall simulator for retention research using small amounts of agroformulations, *Pest Manage. Sci.*, 2017, **73**(1), 123–129.
- 44 C. Thorburn, The Rise and Demise of Integrated Pest Management in Rice in Indonesia, *Insects*, 2015, **6**(2), 381–408.
- 45 C. Drum, *Soil chemistry of pesticides*, PPG Industries. Inc USA, 1980.
- 46 R. Pavela, History, presence and perspective of using plant extracts as commercial botanical insecticides and farm products for protection against insects—a review, *Plant Prot. Sci.*, 2016, **52**(4), 229–241.
- 47 D. R. George, R. D. Finn, K. M. Graham and O. A. E. Sparagano, Present and future potential of plant-derived products to control arthropods of veterinary and medical significance, *Parasites Vectors*, 2014, **7**(1), 28.
- 48 D. Gunnell, M. Eddleston, M. R. Phillips and F. Konradsen, The global distribution of fatal pesticide self-poisoning: Systematic review, *BMC Public Health*, 2007, **7**(1), 357.
- 49 P. N. Holmsgaard, S. Dealtry, V. Dunon, H. Heuer, L. H. Hansen, D. Springael, *et al.*, Response of the bacterial community in an on-farm biopurification system, to which diverse pesticides are introduced over an agricultural season, *Environ. Pollut.*, 2017, **229**, 854–862.
- 50 S. Fahad, S. Saud, A. Akhter, A. A. Bajwa, S. Hassan, M. Battaglia, *et al.*, Bio-based integrated pest management in rice: An agro-ecosystems friendly approach for agricultural sustainability, *J. Saudi Soc. Agric. Sci.*, 2021, **20**(2), 94–102.
- 51 S.-S. Liu, A. Rao and S. B. Vinson, Biological Control in China: Past, present and future—An introduction to this special issue, *Biol. Control*, 2014, **68**(1), 5.
- 52 K. Sorby, G. Fleischer and E. Pehu, *Integrated pest management in development: review of trends and implementation strategies*, 2003.
- 53 J. Holt, A. Cook, T. Perfect and G. Norton, Simulation analysis of Brown Planthopper population dynamics in tropical rice: a simulation analysis, *J. Appl. Ecol.*, 1987, **24**, 87–102.
- 54 K. Jabran, G. Mahajan, V. Sardana and B. S. Chauhan, Allelopathy for weed control in agricultural systems, *Crop Prot.*, 2015, **72**, 57–65.
- 55 S. Senthil-Nathan, M. Y. Choi, C. H. Paik, H. Y. Seo and K. Kalaivani, Toxicity and physiological effects of neem pesticides applied to rice on the *Nilaparvata lugens* Stål, the brown planthopper, *Ecotoxicol. Environ. Saf.*, 2009, **72**(6), 1707–1713.
- 56 J. Hu, C. Xiao and Y. He, Recent progress on the genetics and molecular breeding of brown planthopper resistance in rice, *Rice*, 2016, **9**(1), 30.
- 57 J. Cheng. Rice Planthoppers in the Past Half Century in China, in *Rice Planthoppers: Ecology, Management, Socio Economics and Policy*, ed. Heong K. L, Cheng J. and Escalada M. M., Springer, Netherlands, Dordrecht, 2015. pp. 1–32.
- 58 H. X. Xu, X. S. Zheng and Z. X. Lu, No. 34 Striped stem borer, *Chilo suppressalis*, in *Institute of Plant Protection, Chinese Academy of Agricultural Sciences & China Society of Plant Protection, Crop Diseases and Insect Pests in China*, China Agriculture Press, Beijing, 2015, pp. 124–130.
- 59 G. M. Gurr, Z. Lu, X. Zheng, H. Xu, P. Zhu, G. Chen, *et al.*, Multi-country evidence that crop diversification promotes ecological intensification of agriculture, *Nat. Plants*, 2016, **2**, 16014.
- 60 K. H. Parsons, M. H. Mondal, C. L. McCormick and A. S. Flynt, Guanidinium-Functionalized Interpolyelectrolyte Complexes Enabling RNAi in Resistant Insect Pests, *Biomacromolecules*, 2018, **19**(4), 1111–1117.
- 61 P. M. Bachman, R. Bolognesi, W. J. Moar, G. M. Mueller, M. S. Paradise, P. Ramaseshadri, *et al.*, Characterization of the spectrum of insecticidal activity of a double-stranded RNA with targeted activity against Western Corn Rootworm (*Diabrotica virgifera virgifera* LeConte), *Transgenic Res.*, 2013, **22**(6), 1207–1222.
- 62 R. S. Nandety, Y.-W. Kuo, S. Nouri and B. W. Falk, Emerging strategies for RNA interference (RNAi) applications in insects, *Bioengineered*, 2015, **6**(1), 8–19.
- 63 B. W. Han, W. Wang, C. Li, Z. Weng and P. D. Zamore, Noncoding RNA. piRNA-guided transposon cleavage initiates Zucchini-dependent, phased piRNA production, *Science*, 2015, **348**(6236), 817–821.
- 64 K. Y. Zhu and S. R. Palli, Mechanisms, Applications, and Challenges of Insect RNA Interference, *Annu. Rev. Entomol.*, 2020, **65**(1), 293–311.
- 65 I. Biryukova and T. Ye, Endogenous siRNAs and piRNAs derived from transposable elements and genes in the malaria vector mosquito *Anopheles gambiae*, *BMC Genomics*, 2015, **16**(1), 278.
- 66 L. Swevers, J. Liu and G. Smagghe, Defense Mechanisms against Viral Infection in *Drosophila*: RNAi and Non-RNAi, *Viruses*, 2018, **10**(5), 230.



- 67 K. J. Lucas, B. Zhao, S. Roy, A. L. Gervaise and A. S. Raikhel, Mosquito-specific microRNA-1890 targets the juvenile hormone-regulated serine protease JHA15 in the female mosquito gut, *RNA Biol.*, 2015, **12**(12), 1383–1390.
- 68 M. Ninova, S. Griffiths-Jones and M. Ronshaugen, Abundant expression of somatic transposon-derived piRNAs throughout *Tribolium castaneum* embryogenesis, *Genome Biol.*, 2017, **18**(1), 184.
- 69 K. Cappelle, C. F. R. de Oliveira, B. Van Eynde, O. Christiaens and G. Smagghe, The involvement of clathrin-mediated endocytosis and two Sid-1-like transmembrane proteins in double-stranded RNA uptake in the Colorado potato beetle midgut, *Insect Mol. Biol.*, 2016, **25**(3), 315–323.
- 70 N. Wynant, D. Santos, P. Van Wielendaele and J. Vanden Broeck, Scavenger receptor-mediated endocytosis facilitates RNA interference in the desert locust, *Schistocerca gregaria*, *Insect Mol. Biol.*, 2014, **23**(3), 320–329.
- 71 P. Baldrich, B. D. Rutter, H. Z. Karimi, R. Podicheti, B. C. Meyers and R. W. Innes, Plant Extracellular Vesicles Contain Diverse Small RNA Species and Are Enriched in 10- to 17-Nucleotide “Tiny” RNAs, *Plant Cell*, 2019, **31**(2), 315–324.
- 72 S. Liu, M. Jaouannet, D. A. Dempsey, J. Imani, C. Coustau and K. H. Kogel, RNA-based technologies for insect control in plant production, *Biotechnol. Adv.*, 2020, **39**, 107463.
- 73 O. Christiaens, S. Whyard, A. M. Vélez and G. Smagghe, Double-stranded RNA technology to control insect pests: Current status and challenges, *Front. Plant Sci.*, 2020, **11**, 451.
- 74 A. Kolliopoulou, C. N. T. Taning, G. Smagghe and L. Swevers, Viral Delivery of dsRNA for Control of Insect Agricultural Pests and Vectors of Human Disease: Prospects and Challenges, *Front. Physiol.*, 2017, **8**, 399.
- 75 M. Chen, A. Shelton and G.-y Ye, Insect-resistant genetically modified rice in China: from research to commercialization, *Annu. Rev. Entomol.*, 2011, **56**(1), 81–101.
- 76 Z. Li, Y. Xue, H. Zhou, Y. Li, B. Usman, X. Jiao, *et al.*, High-resolution mapping and breeding application of a novel brown planthopper resistance gene derived from wild rice (*Oryza. rufipogon* Griff), *Rice*, 2019, **12**(1), 41.
- 77 E. Fishilevich, A. J. Bowling, M. L. Frey, P.-H. Wang, W. Lo, M. Rangasamy, *et al.*, RNAi targeting of rootworm Troponin I transcripts confers root protection in maize, *Insect Biochem. Mol. Biol.*, 2019, **104**, 20–29.
- 78 B. He, Y. Chu, M. Yin, K. Müllen, C. An and J. Shen, Fluorescent Nanoparticle Delivered dsRNA Toward Genetic Control of Insect Pests, *Adv. Mater.*, 2013, **25**(33), 4580–4584.
- 79 O. Christiaens, M. G. Tardajos, Z. L. Martinez Reyna, M. Dash, P. Dubruel and G. Smagghe, Increased RNAi Efficacy in *Spodoptera exigua* via the Formulation of dsRNA With Guanlylated Polymers, *Front. Physiol.*, 2018, **9**, 316.
- 80 M. Zotti, E. A. Dos Santos, D. Cagliari, O. Christiaens, C. N. T. Taning and G. Smagghe, RNA interference technology in crop protection against arthropod pests, pathogens and nematodes, *Pest Manage. Sci.*, 2018, **74**(6), 1239–1250.
- 81 C. N. Taning, O. Christiaens, X. Li, L. Swevers, H. Casteels, M. Maes, *et al.*, Engineered flock house virus for targeted gene suppression through RNAi in fruit flies (*Drosophila melanogaster*) *in vitro* and *in vivo*, *Front. Physiol.*, 2018, **9**, 805.
- 82 H. Huvenne and G. Smagghe, Mechanisms of dsRNA uptake in insects and potential of RNAi for pest control: a review, *J. Insect Physiol.*, 2010, **56**(3), 227–235.
- 83 J. A. Baum, T. Bogaert, W. Clinton, G. R. Heck, P. Feldmann, O. Ilagan, *et al.*, Control of coleopteran insect pests through RNA interference, *Nat. Biotechnol.*, 2007, **25**(11), 1322–1326.
- 84 O. Christiaens, T. Dzhambazova, K. Kostov, S. Arpaia, M. R. Joga, I. Urru, *et al.*, Literature review of baseline information on RNAi to support the environmental risk assessment of RNAi-based GM plants, *EFSA Supporting Publ.*, 2018, **15**(5), 1424E.
- 85 R. Sugahara, S. Tanaka, A. Jouraku and T. Shiotsuki, Geographic variation in RNAi sensitivity in the migratory locust, *Gene*, 2017, **605**, 5–11.
- 86 H. M. Abd El Halim, B. M. H. Alshukri, M. S. Ahmad, E. Y. T. Nakasu, M. H. Awwad, E. M. Salama, *et al.*, RNAi-mediated knockdown of the voltage gated sodium ion channel TcNav causes mortality in *Tribolium castaneum*, *Sci. Rep.*, 2016, **6**(1), 29301.
- 87 J. Spit, A. Philips, N. Wynant, D. Santos, G. Plaetinck and J. Vanden Broeck, Knockdown of nuclease activity in the gut enhances RNAi efficiency in the Colorado potato beetle, *Leptinotarsa decemlineata*, but not in the desert locust, *Schistocerca gregaria*, *Insect Biochem. Mol. Biol.*, 2017, **81**, 103–116.
- 88 C. Lenaerts, D. Cools, R. Verdonck, L. Verbakel, J. Vanden Broeck and E. Marchal, The ecdysis triggering hormone system is essential for successful moulting of a major hemimetabolous pest insect, *Schistocerca gregaria*, *Sci. Rep.*, 2017, **7**(1), 46502.
- 89 A. Telang, J. A. Rechel, J. R. Brandt and D. M. Donnell, Analysis of ovary-specific genes in relation to egg maturation and female nutritional condition in the mosquitoes *Georgacraigius atropalpus* and *Aedes aegypti* (Diptera: Culicidae), *J. Insect Physiol.*, 2013, **59**(3), 283–294.
- 90 O. Terenius, A. Papanicolaou, J. S. Garbutt, I. Eleftherianos, H. Huvenne, S. Kanginakudru, *et al.*, RNA interference in Lepidoptera: an overview of successful and unsuccessful studies and implications for experimental design, *J. Insect Physiol.*, 2011, **57**(2), 231–245.
- 91 R. Katoch and N. Thakur, Insect gut nucleases: a challenge for RNA interference mediated insect control strategies, *Int. J. Biochem. Biotechnol.*, 2012, **1**, 198–203.
- 92 N. Wynant, D. Santos, R. Verdonck, J. Spit, P. Van Wielendaele and J. Vanden Broeck, Identification, functional characterization and phylogenetic analysis of double stranded RNA degrading enzymes present in the



- gut of the desert locust, *Schistocerca gregaria*, *Insect Biochem. Mol. Biol.*, 2014, **46**, 1–8.
- 93 E. Vogel, D. Santos, L. Mingels, T.-W. Verdonck and J. V. Broeck, RNA Interference in Insects: Protecting Beneficials and Controlling Pests, *Front. Physiol.*, 2019, **9**, 1912.
- 94 L. Timmons, D. L. Court and A. Fire, Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in *Caenorhabditis elegans*, *Gene*, 2001, **263**(1–2), 103–112.
- 95 M. Vatanparast and Y. Kim, Optimization of recombinant bacteria expressing dsRNA to enhance insecticidal activity against a lepidopteran insect, *Spodoptera exigua*, *PLoS One*, 2017, **12**(8), e0183054.
- 96 M. M. A. Whitten, P. D. Facey, R. Del Sol, L. T. Fernández-Martínez, M. C. Evans, J. J. Mitchell, *et al.*, Symbiont-mediated RNA interference in insects, *Proc. Biol. Sci.*, 2016, **283**(1825), 20160042.
- 97 X. Zhang, K. Mysore, E. Flannery, K. Michel, D. W. Severson, K. Y. Zhu, *et al.*, Chitosan/interfering RNA nanoparticle mediated gene silencing in disease vector mosquito larvae, *J. Visualized Exp.*, 2015, (97), e52523.
- 98 B. Dalby, S. Cates, A. Harris, E. C. Ohki, M. L. Tilkins, P. J. Price, *et al.*, Advanced transfection with Lipofectamine 2000 reagent: primary neurons, siRNA, and high-throughput applications, *Methods*, 2004, **33**(2), 95–103.
- 99 Y. Zhang, J. Cui, Y. Zhou, J. Cao, H. Gong, H. Zhang, *et al.*, Liposome mediated double-stranded RNA delivery to silence ribosomal protein P0 in the tick *Rhipicephalus haemaphysaloides*, *Ticks Tick-borne Dis.*, 2018, **9**(3), 638–644.
- 100 J. Durzyńska, Ł. Przysiecka, R. Nawrot, J. Barylski, G. Nowicki, A. Warowicka, *et al.*, Viral and Other Cell-Penetrating Peptides as Vectors of Therapeutic Agents in Medicine, *J. Pharmacol. Exp. Ther.*, 2015, **354**(1), 32.
- 101 S. Y. Choi and E. A. David, Cell Penetrating Peptides and the Mechanisms for Intracellular Entry, *Curr. Pharm. Biotechnol.*, 2014, **15**(3), 192–199.
- 102 M. R. Joga, M. J. Zotti, G. Smagghe and O. Christiaens, RNAi Efficiency, Systemic Properties, and Novel Delivery Methods for Pest Insect Control: What We Know So Far, *Front. Physiol.*, 2016, **7**, 553.
- 103 K. He, Y. Sun, H. Xiao, C. Ge, F. Li and Z. Han, Multiple miRNAs jointly regulate the biosynthesis of ecdysteroid in the holometabolous insects, *Chilo suppressalis*, *RNA*, 2017, **23**(12), 1817–1833.
- 104 I. Tabujew and K. Peneva, *Functionalization of cationic polymers for drug delivery applications*, 2014.
- 105 Y. Ho and H.-P. Too, CATIONIC POLYMER BASED GENE DELIVERY: UPTAKE AND INTRACELLULAR TRAFFICKING, *Cosmos*, 2014, **10**(01), 17–24.
- 106 Q. Liu, J. Xu, Y. Wang, C. Li, G. Han, J. Qi, *et al.*, Synergism of CmGV and *Bacillus thuringiensis* against larvae of *Cnaphalocrocis medinalis* Guénée, *J. Yangzhou Univ. Agric. Life Sci. Ed.*, 2013, **34**(4), 89–93.
- 107 A. Kichler, C. Leborgne, E. Coeytaux and O. Danos, Polyethylenimine-mediated gene delivery: a mechanistic study, *J. Gene Med.*, 2001, **3**(2), 135–144.
- 108 G. Creusat, A.-S. Rinaldi, E. Weiss, R. Elbaghdadi, J.-S. Remy, R. Mulherkar, *et al.*, Proton Sponge Trick for pH-Sensitive Disassembly of Polyethylenimine-Based siRNA Delivery Systems, *Bioconjugate Chem.*, 2010, **21**(5), 994–1002.
- 109 S. Yan, B.-Y. Ren and J. Shen, Nanoparticle-mediated double-stranded RNA delivery system: A promising approach for sustainable pest management, *Insect Sci.*, 2021, **28**(1), 21–34.
- 110 S. Agarwal, Y. Zhang, S. Maji and A. Greiner, PDMAEMA based gene delivery materials, *Mater. Today*, 2012, **15**(9), 388–393.
- 111 A. B. Cook, R. Peltier, M. Hartlieb, R. Whitfield, G. Moriceau, J. A. Burns, *et al.*, Cationic and hydrolysable branched polymers by RAFT for complexation and controlled release of dsRNA, *Polym. Chem.*, 2018, **9**(29), 4025–4035.
- 112 J. M. Layman, S. M. Ramirez, M. D. Green and T. E. Long, Influence of Polycation Molecular Weight on Poly(2-dimethylaminoethyl methacrylate)-Mediated DNA Delivery In Vitro, *Biomacromolecules*, 2009, **10**(5), 1244–1252.
- 113 C. V. Synatschke, A. Schallon, V. Jérôme, R. Freitag and A. H. Müller, Influence of polymer architecture and molecular weight of poly(2-(dimethylamino)ethyl methacrylate) polycations on transfection efficiency and cell viability in gene delivery, *Biomacromolecules*, 2011, **12**(12), 4247–4255.
- 114 Z. Cao, H. Xiao, L. Li, M. Liu, G. Lin, P. Zhai, *et al.*, The Codelivery of siRNA and QDs by pH-Responsive Micelle for Hepatoma Cancer Cells, *Front. Pharmacol.*, 2019, **10**, 1194.
- 115 F. X. Gillet, R. A. Garcia, L. L. P. Macedo, E. V. S. Albuquerque, M. C. M. Silva and M. F. Grossi-de-Sa, Investigating Engineered Ribonucleoprotein Particles to Improve Oral RNAi Delivery in Crop Insect Pests, *Front. Physiol.*, 2017, **8**, 256.
- 116 D. Gurusamy, K. Mogilicherla and S. R. Palli, Chitosan nanoparticles help double-stranded RNA escape from endosomes and improve RNA interference in the fall armyworm, *Spodoptera frugiperda*, *Arch. Insect Biochem. Physiol.*, 2020, **104**(4), e21677.
- 117 K. Wang, Y. Peng, J. Chen, Y. Peng, X. Wang, Z. Shen, *et al.*, Comparison of efficacy of RNAi mediated by various nanoparticles in the rice striped stem borer (*Chilo suppressalis*), *Pestic. Biochem. Physiol.*, 2020, **165**, 104467.
- 118 F. Haghirsadat, G. Amoabediny, S. Naderinezhad, T. Forouzanfar, M. N. Helder and B. Zandieh-Doulabi, Preparation of PEGylated cationic nanoliposome-siRNA complexes for cancer therapy, *Artif. Cells, Nanomed., Biotechnol.*, 2018, **46**(sup1), 684–692.
- 119 S. Whyard, A. D. Singh and S. Wong, Ingested double-stranded RNAs can act as species-specific insecticides, *Insect Biochem. Mol. Biol.*, 2009, **39**(11), 824–832.



Review

- 120 C. N. T. Taning, O. Christiaens, N. Berkvens, H. Casteels, M. Maes and G. Smaghe, Oral RNAi to control *Drosophila suzukii*: laboratory testing against larval and adult stages, *J. Pest Sci.*, 2016, **89**(3), 803–814.
- 121 A. L. Jackson, S. R. Bartz, J. Schelter, S. V. Kobayashi, J. Burchard, M. Mao, *et al.*, Expression profiling reveals off-target gene regulation by RNAi, *Nat. Biotechnol.*, 2003, **21**(6), 635–637.
- 122 W. B. Hunter, E. Glick, N. Paldi and B. R. Bextine, Advances in RNA interference: dsRNA treatment in trees and grapevines for insect pest suppression, *Southwest. Entomol.*, 2012, **37**(1), 85–87.
- 123 H. Li, R. Guan, H. Guo and X. Miao, New insights into an RNAi approach for plant defence against piercing-sucking and stem-borer insect pests, *Plant, Cell Environ.*, 2015, **38**(11), 2277–2285.
- 124 A. M. Khan, M. Ashfaq, Z. Kiss, A. A. Khan, S. Mansoor and B. W. Falk, Use of recombinant tobacco mosaic virus to achieve RNA interference in plants against the citrus mealybug, *Planococcus citri* (Hemiptera: Pseudococcidae), *PLoS One*, 2013, **8**(9), e73657.
- 125 A. Fus-Kujawa, P. Prus, K. Bajdak-Rusinek, P. Teper, K. Gawron, A. Kowalczyk, *et al.*, An overview of methods and tools for transfection of eukaryotic cells in vitro, *Front. Bioeng. Biotechnol.*, 2021, **634**, 701031.
- 126 E. C. de Andrade and W. B. Hunter, RNA interference–natural gene-based technology for highly specific pest control (HiSPeC), in, *RNA Interference*, I. Y. Abdurakhmonov, IntechOpen, London, 2016, DOI: [10.5772/61612](https://doi.org/10.5772/61612).
- 127 K. A. Murphy, C. A. Tabuloc, K. R. Cervantes and J. C. Chiu, Ingestion of genetically modified yeast symbiont reduces fitness of an insect pest *via* RNA interference, *Sci. Rep.*, 2016, **6**(1), 22587.
- 128 M. Kumar, G. P. Gupta and M. V. Rajam, Silencing of acetylcholinesterase gene of *Helicoverpa armigera* by siRNA affects larval growth and its life cycle, *J. Insect Physiol.*, 2009, **55**(3), 273–278.
- 129 J. Zhang, S. A. Khan, C. Hasse, S. Ruf, D. G. Heckel and R. Bock, Full crop protection from an insect pest by expression of long double-stranded RNAs in plastids, *Science*, 2015, **347**(6225), 991–994.

