



Cite this: *RSC Adv.*, 2022, 12, 8569

DOI: 10.1039/d2ra90024f

rsc.li/rsc-advances

Correction: Dipyrrolyl-bis-sulfonamide chromophore based probe for anion recognition

Namdev V. Ghule,^{ac} Sheshanath V. Bhosale^{*bd} and Sidhanath V. Bhosale^{*a}

Correction for 'Dipyrrolyl-bis-sulfonamide chromophore based probe for anion recognition' by Namdev V. Ghule *et al.*, *RSC Adv.*, 2014, 4, 27112–27115, DOI: 10.1039/C4RA04000G.

The authors regret that an incorrect version of Fig. 1 was included in the original article. The correct version of Fig. 1 is presented below.




Fig. 1 Color changes of receptor DPBS in chloroform upon addition of 5 equiv. of F^- , Cl^- , Br^- , I^- , $H_2PO_4^-$, HSO_4^- , ClO_4^- and AcO^- (tetra-butylammonium salts).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aPolymers and Functional Material Division, CSIR-Indian Institute of Chemical Technology, Hyderabad 50060, AP, India. E-mail: bhosale@iiict.re.in

^bSchool of Applied Sciences, RMIT University, GPO Box 2476V, Melbourne 3001, Victoria, Australia. E-mail: sheshanath.bhosale@rmit.edu.au; Tel: +61399252680

^cDepartment of Organic Chemistry, School of Chemical Sciences, North Maharashtra University, Jalgaon-425001, M.S., India

