


 Cite this: *RSC Adv.*, 2022, 12, 34568

## Correction: Palladium nanoparticles immobilized on polyethylenimine-derivatized gold surfaces for catalysis of Suzuki reactions: development and application in a lab-on-a-chip context

Prasad Anaspure, Subramanian Suriyanarayanan\* and Ian A. Nicholls

DOI: 10.1039/d2ra90120j

[rsc.li/rsc-advances](https://rsc.li/rsc-advances)

 Correction for 'Palladium nanoparticles immobilized on polyethylenimine-derivatized gold surfaces for catalysis of Suzuki reactions: development and application in a lab-on-a-chip context' by Prasad Anaspure *et al.*, *RSC Adv.*, 2021, 11, 35161–35164. <https://doi.org/10.1039/D1RA06851B>.


The authors regret that the turnover numbers (TONs) were not correctly given in the original article.

 In the abstract on page 35161, the corrected number should read  $3.4 \times 10^4$ .

The corrected versions of Table 1 and 2 are shown below.

Accordingly, Table 1-SI, Table 2-SI, Table 3-SI, and Table 4-SI in the original ESI have been revised; the ESI has been updated online.


An independent expert has viewed the corrected tables and has concluded that they are consistent with the discussions and conclusions presented.

**Table 1** Suzuki cross-coupling reactions of aryl halides with arylboronic acids using PEI/Pd as catalysts<sup>a</sup>


Entry	R <sub>1</sub>	X	R <sub>2</sub>	Amount of Pd, μg	Yield	TON
1	H	I	H	3.2	93%	$3.1 \times 10^4$
2	H	Br	H	2.8	95%	$3.4 \times 10^4$
3	H	I	2-CH <sub>3</sub>	3.9	82%	$2.2 \times 10^4$
4	H	I	3-OCH <sub>3</sub>	4.0	57%	$1.5 \times 10^4$
5	H	I	4-OCH <sub>3</sub>	3.7	84%	$2.4 \times 10^4$
6	H	I	2-CN	3.99	15%	$0.4 \times 10^4$
7	H	I	4-CN	3.6	95%	$2.8 \times 10^4$
8	4-CH <sub>3</sub>	Br	H	6.2	88%	$1.5 \times 10^4$
9	4-OCH <sub>3</sub>	Br	H	8.4	95%	$1.2 \times 10^4$
10	H	I	H3-NH <sub>2</sub>	3.5	n. r.	—
11	H	Cl	H	1.0	94%	$10.0 \times 10^4$
12	4-OCH <sub>3</sub>	Cl	H	1.62	80%	$5.3 \times 10^4$
13	4-CoCH <sub>3</sub>	Cl	H	1.5	n. r.	—

<sup>a</sup> General procedure: 1.0 mmol of aryl halide, 1.2 mmol of arylboronic acid, 2.0 mmol of K<sub>2</sub>CO<sub>3</sub> in H<sub>2</sub>O/EtOH. Turnover number TON = mol product/mol Pd. n. r. = no reaction.


## Correction

Table 2 Suzuki cross coupling reaction of iodobenzene and phenylboronic acid using PEI/Pd as catalyst<sup>a</sup>

Entry	Run	Conc. of Pd, $\mu$ g	Yield	TON
1	1 <sup>st</sup>	$\pm 2.3$	93%	$4.4 \times 10^4$
2	2 <sup>nd</sup>	$\pm 2.22$	89%	$4.3 \times 10^4$
3	3 <sup>rd</sup>	$\pm 2.22$	85%	$4.0 \times 10^4$
4	4 <sup>th</sup>	$\pm 2.15$	80%	$3.9 \times 10^4$

<sup>a</sup> General procedure: 1.0 mmol of aryl halide, 1.2 mmol of arylboronic acid, 2.0 mmol of K<sub>2</sub>CO<sub>3</sub> in H<sub>2</sub>O/EtOH. TON = mol product/mol.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

