

Cite this: *Chem. Sci.*, 2022, 13, 4909 All publication charges for this article have been paid for by the Royal Society of Chemistry

Photoinduced metal-free borylation of aryl halides catalysed by an *in situ* formed donor–acceptor complex†

Manhong Li,^{‡a} Siqi Liu,^{ID ‡b} Haoshi Bao,^a Qini Li,^a Yi-Hui Deng,^{bc} Tian-Yu Sun^{*b} and Leifeng Wang^{ID *a}

Organoboron compounds are very important building blocks which can be applied in medicinal, biological and industrial fields. However, direct borylation in a metal free manner has been very rarely reported. Herein, we described the successful direct borylation of haloarenes under mild, operationally simple, catalyst-free conditions, promoted by irradiation with visible light. Mechanistic experiments and computational investigations indicate the formation of an excited donor–acceptor complex with a -3.12 V reduction potential, which is a highly active reductant and can facilitate single-electron-transfer (SET) with aryl halides to produce aryl radical intermediates. A two-step one-pot method was developed for site selective aromatic C–H bond borylation. The protocol's good functional group tolerance enables the functionalization of a variety of biologically relevant compounds, representing a new application of aryl radicals merged with photochemistry.

Received 27th January 2022

Accepted 3rd April 2022

DOI: 10.1039/d2sc00552b

rsc.li/chemical-science

Introduction

Boronic esters are commonly encountered in pharmaceuticals, functional materials and agrochemicals.¹ Aryl halides and related arenes are the most commonly employed precursors to produce them.² Conventional routes for their synthesis include metalation (generally by Li or Mg reagents)³ followed by nucleophilic substitution and transition-metal catalysis,⁴ offering a broad substrate scope and good functional group compatibility if their disadvantages such as high cost, low reactivity and operational inconvenience could be ignored. Visible light catalysis⁵ has become a well-accepted and powerful method for the direct borylation of drug like molecules. Seminal studies by Larionov,⁶ Aggarwal,⁷ Wu,⁸ Schelter⁹ and others¹⁰ have established the viability and synthetic utility of this approach, circumventing the need for well recognized photocatalysts and metal additives. Studer¹¹ developed radical borylation of aryl iodides with bis(catecholato)diboron (B_2cat_2) as the boron source under mild conditions. Recently, Jiao¹² reported the

borylation of aryl halides using 4-phenylpyridine as an efficient catalyst and sodium methanolate as a strong base with bis(pinacolato)diboron (B_2pin_2). Different from Jiao's work, Li,¹³ Larionov¹⁴ and Lin¹⁵ independently described simple and efficient methods to convert aryl triflates and aryl halides into aryl radicals under UV irradiation or reductive electro-photocatalysis. There is a constant quest for more efficient clean borylation strategies (*e.g.*, metal-free visible light catalysis using an *in situ* formed photocatalyst) from simple and readily available starting materials.

In recent years, *in situ* formed donor–acceptor complexes have been successfully used in the single-electron-transfer (SET) process to produce radical species by hybridization and modulation of the relevant energy levels.^{16,17} As a result, visible light excitation can occur favoured by a red shift of the absorption band, although both donor and acceptor components are individually insensitive to visible light irradiation. This novel protocol can provide a new approach for aryl radical intermediates through C–X bond activation in a metal-free manner (Scheme 1).

In 2020, we reported a photoinduced dehalogenation of aryl halides.¹⁸ Spectroscopic, computational and chemical studies indicate that the formation of a twisted intramolecular charge-transfer (TICT) species enables the population of higher-energy doublet states. König¹⁹ developed a versatile photocatalytic strategy for the ipso-borylation of substituted arenes by using thiolate as a catalyst, forming a donor–acceptor complex between thiolate/ B_2pin_2 and boryl-anion-activated substrates. Inspired by our previous work and related reports,²⁰ we questioned whether borylation of aryl halides could be achieved with

^aSchool of Pharmaceutical Sciences (Shenzhen), Sun Yat-Sen University, No. 66, Gongchang Road, Shenzhen 518107, P. R. China. E-mail: wanglf33@mail.sysu.edu.cn


^bShenzhen Bay Laboratory, Shenzhen 518132, P. R. China. E-mail: Tian-Yu_Sun@pku.edu.cn

^cLab of Computational Chemistry and Drug Design, State Key Laboratory of Chemical Oncogenomics, Peking University Shenzhen Graduate School, Shenzhen 518055, P. R. China

† Electronic supplementary information (ESI) available. CCDC 2144922. For ESI and crystallographic data in CIF or other electronic format see <https://doi.org/10.1039/d2sc00552b>

‡ These authors contributed equally to this work.





Scheme 1 Synthetic approaches to aryl boronic esters.

the photocatalytic intermediate formed *in situ* employing simple available substrates (*e.g.*, N-containing heterocycles/organic base/boron source). Herein, we described the successful direct borylation of haloarenes under mild, operationally simple, catalyst-free conditions, promoted by irradiation with visible light.

Results and discussion

We made our initial discovery of the borylation reaction employing 4-bromoanisole as the model substrate (Table 1). First, under visible light photoirradiation, when the 2,2'-bipyridine (nitrogen containing heterocycle: NCH)/triethylamine


(NEt_3 ; final reductant)/ B_2pin_2 (boron source) system was evaluated in acetonitrile, the desired product can be detected in 27% yield (entry 1). In order to generate the optimal ate-intermediate to facilitate the single-electron-transfer (SET) process, a series of NCHs were evaluated (entries 2–5). Only trace desired product was detected using the 2-phenyl pyridine/ NEt_3 / B_2pin_2 system (entry 2) suggesting that the ate-intermediate formed with 2-phenyl pyridine couldn't deliver a single electron to the aromatic C–Br bond. To our delight, the yield was improved remarkably (49% yield) employing isoquinoline as the ate-intermediate precursor (entry 3). Neither acridine (entry 4) nor 4-trifluoromethyl pyridine (entry 5) could benefit the borylation reaction, indicating that steric resistance and electron-deficient factors of NCHs can decrease the reductive ability of the ate-intermediate. A brief investigation of the solvent effect demonstrates that DMSO (entry 6), DMF (entry 7) and DCE (entry 8) are not as efficient as acetonitrile. Gratifyingly, the desired product could be isolated with 81% yield *via* increasing the loading of B_2pin_2 / NEt_3 and prolonging the photoirradiation process (entry 9). Not surprisingly, control experiments without NCH (entry 10) and in the dark (entry 11) couldn't afford the desired product effectively.

With the optimized reaction conditions in hand, we first examined the substrate scope of aryl bromides (Table 2).

Table 1 Optimization of the reaction conditions^a

Entry	Solvent	B_2pin_2 (X eq.)	NEt_3 (Y eq.)	NCH	Time (h)	Yield (%) ^b
1	AcN	X = 2	Y = 3	A	12	27
2	AcN	X = 2	Y = 3	B	12	Trace
3	AcN	X = 2	Y = 3	C	12	49
4	AcN	X = 2	Y = 3	D	12	17
5	AcN	X = 2	Y = 3	E	12	Trace
6	DMSO	X = 2	Y = 3	C	12	Trace
7	DMF	X = 2	Y = 3	C	12	NR
8	DCE	X = 2	Y = 3	C	12	16
9	AcN	X = 4	Y = 5	C	36	87 (81) ^c
10	AcN	X = 4	Y = 5	—	36	14
11	AcN	X = 4	Y = 5	C	36	NR ^d

^a Reactions were performed with 0.2 mmol 4-bromoanisole, 1.0 mmol NEt_3 , 20 mol% NCH, 0.5 ml solvent and suitable B_2pin_2 for the indicated period of time under $2 \times 390 \text{ nm LED}$ (40 W) irradiation. ^b NMR yields using pyrazine or hexamethyl disiloxane as the internal standard. ^c Isolated yields. ^d Without irradiation.

Table 2 Substrate scope of aryl bromides^a

^a Reactions were run on a 0.2 mmol scale; isolated yield. ^b Irradiation for 48 h. ^c Using 1-bromo-4-chlorobenzene as the starting material.



Various *para*-substituted aryl bromides, including electron donating group substituted (**3a–3f**), unsubstituted (**3g**), unprotected $-OH$ (**3h**)/ $-NH_2$ (**3i**), electron withdrawing group substituted (**3j–3o**) and even simple alkene moieties (**3o**), could be well tolerated in this system, delivering the corresponding arylboronates in good to excellent yields. However, bromo-styrene was not tolerated. A dual borylation product (**3q**) could be synthesized using dihaloarene or haloaryl boronic ester. *Meta*-(**3r–3t**) and *ortho*-(**3u–3w**) substituted arylboronates can both be obtained in good yields (e.g., **3w**, 1-allyl-2-bromobenzene, 63% yield). Moreover, 2,5-(**3x**)/3,4-(**3y–3ac**)/3,5-disubstituted (**3ad–3ae**) aryl bromides produced borylation products in good yields (**3aa** and **3ac**, 69% and 64%, respectively). Heteroaromatic bromides, such as benzothiophene (**3ag**) and indole (**3ah**) derivatives, can be employed to access the desired borylation products with good efficiencies other than brominated benzofuran (**3af**, acceptable 23% yield). This methodology also worked well with a series of diboron reagents to produce arylboronates **4a–6a** with useful efficiencies (e.g., **4a**, 69% yield). In contrast, $B_2(OH)_4$ was not a good borylation candidate with a poor solubility in acetonitrile. Notably, except for aryl halides, **3a** could also be prepared through more challenging Ar–O (**7a**)/Ar–N (**8a**) bond cleavage under standard conditions. Surprisingly, reductive deoxygenation of fluorenone was accomplished with high efficiency (**9a**, 65%).

Next, we turned our attention to exploring the substrate scope of the borylation of unactivated aryl chlorides, since cleavage of the C–Cl bond in a metal-free manner remains quite challenging [e.g., for PhCl, $E_{red} = -3.04$ V vs. SCE].²¹ Our investigations revealed that both electron-rich and electron-deficient chlorobenzene derivatives could be employed to provide the corresponding arylboronates in good yields with a little more loading of NCH and longer irradiation time (Table 3; 68% NMR yield for 20 mol% NCH loading and 36 h irradiation vs. 82% NMR yield for 100 mol% NCH loading and


72 h irradiation). It's worth noting that the reaction was allowed for production of fluorinated boronic ester (**11h**). In addition, unprotected *para*, *meta*, and *ortho* $-OH/-NH_2$ substituted aryl chlorides (**10e–10f**; **10j**; **10l**) were well tolerated, illustrating the immediate utility of this approach in preparing value-added boronic ester in a protecting group-free manner. Similar to aryl bromide substrates, more electron-rich and sterically hindered aryl chlorides (**10m–10p**) could also be transformed into desired products with good yields using this new protocol (Table 4).

Site selective aromatic C–H functionalization provides rapid access to useful structural and functional molecular complexity.²² Based on the optimized reaction conditions, we developed a two-step one-pot method for simple aromatic C–H borylation. In the first step, almost equivalent regioselective bromination is accomplished using *N*-bromosuccinimide (NBS) as the bromination reagent, controlled by the electrophilicity and steric hindrance effect of the aromatic ring.²³ Subsequently, a series of aryl bromides without further purification were transformed into the corresponding boronic esters with high regioselectivities and efficiencies (e.g., **13a**, 73% yield; rr up to 40 : 1). Aromatic silyl-ethers (**13c–d**), unprotected phenol (**13e**), aniline (**13f**) and polysubstituted benzenes (**13g–h**) were well tolerated, delivering site selective products with moderate to good yields (Table 5).

The broad functional group tolerance and two-step one-pot method for site selective aromatic C–H borylation encouraged us to test this mild strategy for late stage functionalization of complex molecules, such as natural products and active pharmaceutical ingredients (APIs). Boronates derived from methyl dehydroabietate (**14**), amino acid tyrosine (**15**), and a variety of APIs (**16–18**) containing polar groups, sterically hindered substituents and polycyclic frameworks were still readily produced under the optimized conditions, highlighting the utility of this methodology in the late-stage functionalization of biologically relevant compounds.


To obtain insights into the reaction mechanism, an *N,N*-diboronate complex²⁴ (dimer **A**) was prepared through reductive coupling of isoquinoline and B_2pin_2 (Scheme 2a). Under photoirradiation, no reaction of 4-bromoanisole and dimer **A** was

Table 3 Substrate scope of aryl chlorides^a



^a Reactions were run on a 0.2 mmol scale; isolated yield. ^b 20 mol% NCH (isoquinoline) was used and irradiation for 36 h. ^c NMR yield.

Table 4 Two step, one-pot method for Ar C–H bond borylation^a



^a Reactions were run on a 0.2 mmol scale; isolated yield.




Table 5 Late stage functionalization of complex molecules^a

^a Reactions were run on a 0.2 mmol scale; isolated yield.

detected without addition of NEt_3 (see the ESI[†]). In contrast, the potential excited donor–acceptor complex was able to slowly reduce 4-bromoanisole into anisole in the presence of NEt_3 ; when additional B_2pin_2 (2.0 eq.) was applied as the trapping reagent without loading of NEt_3 , no dehalogenation product except boronic ester product was detected; furthermore, the reduction of 4-bromoanisole took place more efficiently in the presence of NEt_3 and B_2pin_2 , delivering the corresponding boronic ester as the major product (Scheme 2b). These observations indicated that the reduction ability of the coordination complex of acceptor dimer **A** and donor NEt_3 has indeed been enhanced after photoexcitation. To obtain a clearer picture of the complex of acceptor dimer **A** and donor NEt_3 , density functional theory (DFT) calculations were carried out. First, the computational UV-Vis spectrum was employed to compare with the experimental UV-Vis spectrum (Scheme 2c). Without NEt_3 , there exist two smooth peaks in the computational UV-Vis spectrum of dimer **A** and the peak at 227 nm is higher than that at 301 nm, which is in agreement with the experiments (Scheme 2c(I) and d(III)). When one NEt_3 molecule binds to dimer **A** *in silico*, the curve between the two peaks becomes sharp, which is also consistent with the experimental UV-Vis spectrum (Scheme 2c(II) and d(IV)). However, when two NEt_3 bind to dimer **A**, the computational UV-Vis spectrum reveals that the peak around 220 nm becomes lower than the peak around 300 nm, which is quite different from experiments (Scheme 2d(V)). Therefore, the computational results suggest that only one NEt_3 binds to dimer **A** under the optimized reaction conditions. In addition, the binding energy between dimer **A** and one NEt_3 is 11.6 kcal mol⁻¹ by calculation. Considering that the ratio of dimer **A** (20 mol% isoquinoline → 10 mol% dimer **A**) to NEt_3 (5.0 eq.) is 1 : 50 under the optimized reaction conditions, dimer **A** and complex **B** can equilibrate (Scheme 2d). After **B** is determined to be the complex combining acceptor dimer **A** and donor NEt_3 , time dependent TD-DFT calculations were conducted for the excited-state of **B**. The excited-state reduction potential of **B**^{*} is -3.12 V, indicating that **B**^{*} is a highly active reductant and can facilitate SET with aryl halides to produce aryl radical intermediates.


With this mechanistic insight, a plausible mechanism for the conversion of aryl halides to their corresponding boronic esters is proposed in Scheme 3. Reductive coupling of



Scheme 2 Mechanistic and computational experiments: (a) preparation of dimer **A**; (b) mechanistic experiments using dimer **A** (0.15 mmol)/4-bromoanisole (0.2 mmol)/ NEt_3 (1.0 mmol) with or without B_2pin_2 under 2×390 nm LEDs (40 W) irradiation for 12 h; (c) experimental UV-Vis spectra (I: dimer **A**/II: dimer **A** + NEt_3); (d) computational UV-Vis spectra (III: dimer **A**; IV: dimer **A** + NEt_3 ; V: dimer **A** + 2NEt_3); (e) binding between dimer **A** and one NEt_3 and then photoexcitation of **D**–**A** complex **B** by 390 nm light.

isoquinoline and B_2pin_2 generates dimer **A**, which can coordinate with NEt_3 *in situ* to afford intermediate **B**. Donor–acceptor complex **B** is then excited by 390 nm light, populating a highly reducing excited state **B**^{*}, which can undergo a single electron transfer process with an electronically matched aryl halide to generate a radical cation species **C**, aryl radical **D**, and regenerate isoquinoline/ B_2pin_2 . Radical cation **C** could be transformed into enamine **E** followed by hydrogen atom transfer and





Scheme 3 The plausible mechanism.

deprotonation processes. Finally, radical **D** is trapped by B_2pin_2 immediately, yielding the desired boronic ester.

Conclusions

In summary, a facile transition-metal-free conversion of aryl halides to the corresponding boronic esters through a radical process has been developed. A series of aryl boronic esters were synthesized with good functional group tolerance in moderate to excellent yields under mild conditions. Mechanistic experiments and computational investigations indicate the formation of an excited donor-acceptor complex, serving as the super single electron reductant. Furthermore, the key role of the donor-acceptor complex formed *in situ* was confirmed by control experiments and DFT calculations. The protocol's functional group tolerance and two-step one-pot method for site selective aromatic C-H bond borylation enabled the functionalization of a variety of biologically relevant compounds, representing a new application of aryl radical merged with photochemistry.

Data availability

Data for all compounds in this manuscript are available in the ESI,[†] which includes experimental details, characterisation data, computational data and copies of 1H , ^{13}C , ^{19}F , ^{11}B NMR spectra.

Author contributions

M.-H. L. performed the experiments with contributions from H.-S. B and Q.-N. L., S.-Q. L. and Y.-H. D. performed DFT calculations. T.-Y. S. and L.-F. W. supervised the project. All the authors analyzed the data, discussed results and contributed to the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We gratefully acknowledge the support of this research by the National Natural Science Foundation of China (No. 21933004), the Key-Area Research and Development Program of Guangdong Province (No. 2020B010188001), the Guangdong Basic and Applied Basic Research Foundation (No. 2022A1515011994), the Technology & Innovation Commission of Shenzhen Municipality (No. 75110-42100033), the Fundamental Research Funds for the Central Universities (No. 75110-30610023) and the Sun Yat-sen University Startup Fund (No. 75110-1841230 & 75110-18841290). The calculations were carried out at the SZBL supercomputing center.

Notes and references

- (a) D. G. Hall, *Boronic Acids: Preparation and Applications in Organic Synthesis Medicine and Materials*, Wiley-VCH, Weinheim, Germany, 2nd edn, 2011; (b) A. B. Cuenca, R. Shishido, H. Ito and E. Fernández, *Chem. Soc. Rev.*, 2017, **46**, 415–430; (c) Y.-M. Tian, X.-N. Guo, H. Braunschweig, U. Radius and T. B. Marder, *Chem. Rev.*, 2021, **121**, 3561–3597.
- (a) F. W. Friese and A. Studer, *Chem. Sci.*, 2019, **10**, 8503–8518; (b) Z.-H. Qiu and C.-J. Li, *Chem. Rev.*, 2020, **120**, 10454–10515; (c) S. K. Bose, L.-J. Mao, L. Kuehn, U. Radius, J. Nekvinda, W. L. Santos, S. A. Westcott, P. G. Steel and T. B. Marder, *Chem. Rev.*, 2021, **121**, 13238–13341; (d) E. M. Larin, J. Loup, I. Polishchuk, R. J. Ross, A. Whyte and M. Lautens, *Chem. Sci.*, 2020, **11**, 5716–5723; (e) T. Lim, J. Y. Ryoo and M. S. Han, *J. Org. Chem.*, 2020, **85**, 10966–10972; (f) G.-L. Gao, J.-X. Yan, K. Yang, F.-E. Chen and Q.-L. Song, *Green Chem.*, 2017, **19**, 3997–4001; (g) X.-H. Pu, J.-F. Hu, Y. Zhao and Z.-Z. Shi, *ACS Catal.*, 2016, **6**, 6692–6698; (h) A. Verma, K. Tomar and P. K. Bharadwaj, *Inorg. Chem.*, 2019, **58**, 1003–1006.
- (a) R. L. Letsinger and I. H. Skoog, *J. Org. Chem.*, 1953, **18**, 895–897; (b) M. Légaré, M. Rang, G. Bélanger-Chabot, J. I. Schweizer, I. Krummenacher, R. Bertermann, M. Arrowsmith, M. C. Holthausen and H. Braunschweig, *Science*, 2019, **363**, 1329–1332; (c) J. A. Knöller, G.-Y. Meng, X. Wang, D. Hall, A. Pershin, D. Beljonne, Y. Olivier, S. Laschat, E. Zysman-Colman and S. Wang, *Angew. Chem., Int. Ed.*, 2020, **59**, 3156–3160; (d) D.-T. Yang, J. Zheng, J.-B. Peng, X. Wang and S. Wang, *J. Org. Chem.*, 2021, **86**, 829–836.
- For reviews, see: (a) D. Wang and D. Astruc, *Chem. Soc. Rev.*, 2017, **46**, 816–854; (b) F. Song, T. Gou, B.-Q. Wang and Z.-J. Shi, *Chem. Soc. Rev.*, 2018, **47**, 7078–7115; (c) Y.-J. Quan and Z. W. Xie, *Chem. Soc. Rev.*, 2019, **48**, 3660–3673; (d) H.-M. Huang, P. Bellotti and F. Glorius, *Chem. Soc. Rev.*, 2020, **49**, 6186–6197; (e) H. Lu, T.-Y. Yu, P.-F. Xu and H. Wei, *Chem. Rev.*, 2021, **121**, 365–411; (f) U. B. Kim, D. J. Jung, H. J. Jeon, K. Rathwell and S. Lee, *Chem. Rev.*, 2020, **120**, 13382–13433; (g) J. Trouvé and R. Gramage-Doria, *Chem. Soc. Rev.*, 2021, **50**, 3565–3584; (h) A. Y. Chan, I. B. Perry, N. B. Bissonnette, B. F. Buksh, G. A. Edwards,



- L. I. Frye, O. L. Garry, M. N. Lavagnino, B.-X. Li, Y. Liang, E. Mao, A. Millet, J. V. Oakley, N. L. Reed, H. A. Sakai, C. P. Seath and D. W. C. MacMillan, *Chem. Rev.*, 2022, **122**, 1485–1542.
- 5 For reviews, see: (a) N. A. Romero and D. A. Nicewicz, *Chem. Rev.*, 2016, **116**, 10075–10166; (b) C.-S. Wang, P. H. Dixneuf and J.-F. Soulé, *Chem. Rev.*, 2018, **118**, 7532–7585; (c) L. Troian-Gautier, M. D. Turlington, S. A. M. Wehlin, A. B. Maurer, M. D. Brady, W. B. Swords and G. J. Meyer, *Chem. Rev.*, 2019, **119**, 4628–4683; (d) H.-M. Huang, P. Bellotti and F. Glorius, *Chem. Soc. Rev.*, 2020, **49**, 6186–6197; (e) X.-Y. Yu, J.-R. Chen and W.-J. Xiao, *Chem. Rev.*, 2021, **121**, 506–561; (f) N. L. Reed and T. P. Yoon, *Chem. Soc. Rev.*, 2021, **50**, 2954–2967; (g) J. D. Bell and J. A. Murphy, *Chem. Soc. Rev.*, 2021, **50**, 9540–9685; (h) M. S. Galliher, B. J. Roldan and C. R. J. Stephenson, *Chem. Soc. Rev.*, 2021, **50**, 10044–10057; (i) F.-D. Lu, J. Chen, X. Jiang, J.-R. Chen, L.-Q. Lu and W.-J. Xiao, *Chem. Soc. Rev.*, 2021, **50**, 12808–12827; (j) M. A. Bryden and E. Zysman-Colman, *Chem. Soc. Rev.*, 2021, **50**, 7587–7680; (k) N. Holmberg-Douglas and D. A. Nicewicz, *Chem. Rev.*, 2022, **122**, 1925–2016; (l) S. P. Pitre and L. E. Overman, *Chem. Rev.*, 2022, **122**, 1717–1751; (m) K. Kwon, R. T. Simons, M. Nandakumar and J. L. Roizen, *Chem. Rev.*, 2022, **122**, 2353–2428; (n) L. Chang, Q. An, L. Duan, K. Feng and Z. Zuo, *Chem. Rev.*, 2022, **122**, 2429–2486.
- 6 S. Jin, H. T. Dang, G. C. Haug, R. He, V. D. Nguyen, V. T. Nguyen, H. D. Arman, K. S. Schanze and O. V. Larionov, *J. Am. Chem. Soc.*, 2020, **142**, 1603–1613.
- 7 A. Noble, R. S. Mega, D. Pflästerer, E. L. Myers and V. K. Aggarwal, *Angew. Chem., Int. Ed.*, 2018, **57**, 2155–2159.
- 8 J.-H. Xu, J.-L. Cao, X.-Y. Wu, H. Wang, X.-N. Yang, X.-X. Tang, R.-W. Toh, R. Zhou, E. K. L. Yeow and J. Wu, *J. Am. Chem. Soc.*, 2021, **143**, 13266–13273.
- 9 Y.-S. Qiao, Q.-M. Yang and E. J. Schelter, *Angew. Chem., Int. Ed.*, 2018, **57**, 10999–11003.
- 10 (a) Y.-M. Tian, X.-N. Guo, I. Krummenacher, Z. Wu, J. Nitsch, H. Braunschweig, U. Radius and T. B. Marder, *J. Am. Chem. Soc.*, 2020, **142**, 18231–18242; (b) J. H. Docherty, K. Nicholson, A. P. Dominey and S. P. A. Thomas, *ACS Catal.*, 2020, **10**, 4686–4691; (c) Z.-J. Kuang, H.-H. Chen, J. Qiu, Z.-L. Ou, Y. Lan and Q.-L. Song, *Chem*, 2020, **6**, 2347–2363; (d) G. Zhang, M.-Y. Li, W.-B. Ye, Z.-T. He, C.-G. Feng and G.-Q. Lin, *Org. Lett.*, 2021, **23**, 2948–2953.
- 11 Y. Cheng, C. Mück-Lichtenfeld and A. Studer, *Angew. Chem., Int. Ed.*, 2018, **57**, 16832–16836.
- 12 (a) L. Zhang and L. Jiao, *J. Am. Chem. Soc.*, 2017, **139**, 607–610; (b) L. Zhang and L. Jiao, *J. Am. Chem. Soc.*, 2019, **141**, 9124–9128; (c) H. Yang, L. Zhang, F.-Y. Zhou and L. Jiao, *Chem. Sci.*, 2020, **11**, 742–747.
- 13 (a) W.-B. Liu, X.-B. Yang, Y. Gao and C.-J. Li, *J. Am. Chem. Soc.*, 2017, **139**, 8621–8627; (b) W. Liu, J. Li, C.-Y. Huang and C.-J. Li, *Angew. Chem., Int. Ed.*, 2020, **59**, 1786–1796.
- 14 (a) A. M. Mfuh, J. D. Doyle, B. Chhetri, H. D. Arman and O. V. Larionov, *J. Am. Chem. Soc.*, 2016, **139**, 2985–2988; (b) S.-F. Jin, H. T. Dang, G. C. Haug, R. He, V. D. Nguyen, V. T. Nguyen, H. D. Arman, K. S. Schanze and O. V. Larionov, *J. Am. Chem. Soc.*, 2020, **142**, 1603–1613.
- 15 H. Kim, H. Kim, T. H. Lambert and S. Lin, *J. Am. Chem. Soc.*, 2020, **142**, 2087–2092.
- 16 (a) S. Takeuchi, T. Nakagawa and Y. Yokoyama, *Chem. Commun.*, 2020, **56**, 6492–6494; (b) M. Madhu, R. Ramakrishnan, V. Vijay and M. Hariharan, *Chem. Rev.*, 2021, **121**, 8234–8284; (c) K. Kohara, A. Trowbridge, M. A. Smith and M. J. Gaunt, *J. Am. Chem. Soc.*, 2021, **143**, 19268–19274.
- 17 (a) Y. Wang, W. Zhu, W. Du, X. Liu, X. Zhang, H. Dong and W. Hu, *Angew. Chem., Int. Ed.*, 2018, **57**, 3963–3967; (b) T. J. Whitemore, A. Millet, H. J. Sayre, C. Xue, B. S. Dolinar, E. G. White, K. R. Dunbar and C. Turro, *J. Am. Chem. Soc.*, 2018, **140**, 5161–5170.
- 18 I. A. MacKenzie, L.-F. Wang, N. P. R. Onuska, O. F. Williams, K. Begam, A. M. Moran, B. D. Dunietz and D. A. Nicewicz, *Nature*, 2020, **580**, 76–80.
- 19 S. Wang, H. Wang and B. König, *Chem*, 2021, **7**, 1653–1665.
- 20 (a) A. B. Grommet, L. M. Lee and R. Klajn, *Acc. Chem. Res.*, 2020, **53**, 2600–2610; (b) M. Baroncini, S. Silvi and A. Credi, *Chem. Rev.*, 2020, **120**, 200–268; (c) Si. Witzel, A. S. K. Hashmi and J. Xie, *Chem. Rev.*, 2021, **121**, 8868–8925; (d) W.-S. Guo, Q. Wang and J.-P. Zhu, *Chem. Soc. Rev.*, 2021, **50**, 7359–7377; (e) J. P. Barham and B. König, *Angew. Chem., Int. Ed.*, 2020, **59**, 11732–11747; (f) L. M. Moreau, E. Lapsheva, J. I. Amaro-Estrada, M. R. Gau, P. J. Carroll, B. C. Manor, Y. Qiao, Q. Yang, W. W. Lukens, D. Sokaras, E. J. Schelter, L. Maron and C. H. Booth, *Chem. Sci.*, 2022, **13**, 1759; (g) H.-Y. Li, X.-X. Tang, J.-H. Pang, X.-Y. Wu, E. K. L. Yeow, J. Wu and S. Chiba, *J. Am. Chem. Soc.*, 2021, **143**, 481–487; (h) L. F. T. Novaes, J.-J. Liu, Y.-F. Shen, L. X. Lu, J. M. Meinhardt and S. Lin, *Chem. Soc. Rev.*, 2021, **50**, 7941–8002.
- 21 L. Pause, M. Robert and J.-M. Savéant, *J. Am. Chem. Soc.*, 1999, **121**, 7158–7159.
- 22 (a) C. B. Kelly and R. Padilla-Salinas, *Chem. Sci.*, 2020, **11**, 10047–10060; (b) L. Zhang and T. Ritter, *J. Am. Chem. Soc.*, 2022, **144**, 2399–2414; (c) F. Berger and T. Ritter, *Synlett*, 2022, **33**, 339–345; (d) R. Cannalire, S. Pelliccia, L. Sancineto, E. Novellino, G. C. Tron and M. Giustiniano, *Chem. Soc. Rev.*, 2021, **50**, 766–897; (e) B. Prabagar, Y.-Q. Yang and Z.-Z. Shi, *Chem. Soc. Rev.*, 2021, **50**, 11249–11269.
- 23 (a) X. Xiong, F. Tan and Y.-Y. Yeung, *Org. Lett.*, 2017, **19**, 4243–4246; (b) D. R. Motati, D. Uredi and E. B. Watkins, *Chem. Sci.*, 2018, **9**, 1782–1788; (c) R.-J. Tang, T. Milcent and B. Crousse, *J. Org. Chem.*, 2018, **83**, 930–938; (d) W.-J. Wang, X.-Y. Li, X.-X. Yang, L.-S. Ai, Z.-W. Gong, N. Jiao and S. Song, *Nat. Commun.*, 2021, **12**, 3873–3882.
- 24 D.-P. Chen, G.-Q. Xu, Q.-H. Zhou, L. W. Chung and W.-J. Tang, *J. Am. Chem. Soc.*, 2017, **139**, 9767–9770.

