

Cite this: *Chem. Sci.*, 2022, 13, 9265

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 29th May 2022

Accepted 14th July 2022

DOI: 10.1039/d2sc02985e

rsc.li/chemical-science

# Asymmetric higher-order [10 + *n*] cycloadditions of palladium-containing 10 $\pi$ -cycloaddends†

Ao Li,<sup>‡a</sup> Yang Gao,<sup>‡a</sup> Jian-Bin Lu,<sup>a</sup> Zhi-Chao Chen,<sup>a</sup> Wei Du<sup>ID \*a</sup> and Ying-Chun Chen<sup>ID \*ab</sup>


We uncovered an asymmetric higher-order [10 + 2] cycloaddition reaction between diverse activated alkenes and a new type of  $\pi$ -allylpalladium complex-containing dipole-type 10 $\pi$ -cycloaddend, which was generated *in situ* from 2-methylene-1-indanols via a dehydrative insertion and deprotonation strategy under double activation of Pd(0) and phosphoric acid. A similar strategy was applied to an asymmetric higher-order [10 + 8] cycloaddition reaction or [10 + 4] cycloaddition reaction by using a heptafulvene derivative or a cyclic enone, respectively, as the acceptor. A variety of polycyclic frameworks imbedding an indene core were generally furnished in moderate to excellent yields with high levels of enantioselectivity by employing a newly designed chiral phosphoramidite ligand.

## Introduction

Higher-order cycloaddition involving conjugated systems with more than 6 $\pi$ -electrons, which enables rapid construction of complicated cyclic frameworks,<sup>1</sup> has received considerable attention since its discovery in the 1960s.<sup>2</sup> In spite of its charm in organic synthesis, higher-order cycloaddition generally suffers from low reactivity, and poor periselectivity and stereoselectivity; thus a variety of pre-prepared cyclic polyenes (>4 $\pi$ ) have been commonly utilised, including fulvenes,<sup>3</sup> Cr(0)-cycloheptatriene complexes,<sup>4</sup> tropone and its analogues,<sup>5</sup> and 3*H*-pyrrolizines,<sup>6</sup> as well as amino-stabilised isobenzofulvenes,<sup>7</sup> in combination with diverse 2 $\pi$ - or 4 $\pi$ -systems under different conditions (Scheme 1a). Recently, significant progress in the field of higher-order cycloaddition reactions has been made with some well-designed carbonyl substrates, which could be activated by a suitable organocatalyst to generate several types of 6 $\pi$ , 8 $\pi$ , 10 $\pi$  or even 12 $\pi$ -cycloaddends *in situ*, typically featuring HOMO-raised polyenamine<sup>8</sup> or polyenolate species<sup>9</sup> (Scheme 1b). However, the development of relevant cycloaddends embedding a reactive metal-complexed motif

catalytically, which can be successfully applied in higher-order cycloaddition reactions, has not been disclosed yet.

Over the past few decades, the zwitterionic reagents bearing a  $\pi$ -allylmetal moiety, usually generated *in situ* from allylic alcohol derivatives under palladium or iridium catalysis, have been extensively utilised as valuable 1,*n*-dipoles for various asymmetric formal cycloaddition reactions.<sup>10</sup> However, such a reaction strategy, through designing suitable conjugated  $\pi$ -systems, has not been envisioned for potentially developing higher-order cycloadditions.<sup>11</sup> As a result, it would be particularly intriguing to uncover a new type of metal-embedding



**Scheme 1** Summary of typical cycloaddends (>4 $\pi$ ) used in higher-order cycloadditions and our design. (a) Diverse pre-prepared cycloaddends. (b) Diverse *in situ* formed cycloaddends via organocatalysis. (c) This work: *in situ* formed  $\pi$ -allylpalladium-containing 10 $\pi$ -cycloaddends.

<sup>a</sup>Key Laboratory of Drug-Targeting and Drug Delivery System of the Education Ministry and Sichuan Province, Sichuan Research Center for Drug Precision Industrial Technology, West China School of Pharmacy, Sichuan University, Chengdu 610041, China. E-mail: duweiyb@scu.edu.cn; ycchen@scu.edu.cn; Fax: +86 28 85502609

<sup>b</sup>College of Pharmacy, Third Military Medical University, Shapingba, Chongqing 400038, China

† Electronic supplementary information (ESI) available: Experimental procedures, spectroscopic data for new compounds, NMR and HRMS spectra and HPLC chromatograms, and the CIF file of enantiopure product 3a. CCDC 2174555. For ESI and crystallographic data in CIF or other electronic format see <https://doi.org/10.1039/d2sc02985e>

‡ A. Li and Y. Gao contributed equally to this work.


dipole, which could participate in asymmetric higher-order cycloadditions as cycloaddends with more than  $6\pi$ -electrons. With these considerations, we envisaged that allylic alcohol or its derivative **1**, readily available from 1-indanone, would undergo oxidative addition under Pd(0) catalysis. The resultant  $\pi$ -allylpalladium complex **I**, having an indene-type structure,<sup>12</sup> would feasibly increase the acidity of the benzylic C–H.<sup>13</sup> Thus, zwitterionic intermediate **II** would be generated after deprotonation, as outlined in Scheme 1c, which might perform as a unique metal-containing  $10\pi$ -cycloaddend to undertake higher-order  $[10 + n]$  cycloaddition reactions with suitable electrophilic counterparts.

## Results and discussion

### Condition optimisation

The initial attempt with allylic carbonate **1a** and activated alkene  $\alpha$ -cyano chalcone **2a** proved to be unsuccessful in

**Table 1** Optimisation of reaction conditions of the asymmetric  $[10 + 2]$  cycloaddition reaction<sup>a</sup>



Entry	<b>1</b>	<b>L</b>	<b>A</b>	Yield <sup>b</sup> (%)	ee <sup>c</sup> (%)
1 <sup>d</sup>	<b>1a</b>	—	—	—	—
2 <sup>d</sup>	<b>1b</b>	—	—	65	—
3	<b>1b</b>	<b>L1</b>	—	45	–65
4	<b>1c</b>	PPh <sub>3</sub>	—	<5%	—
5	<b>1c</b>	PPh <sub>3</sub>	( <i>R</i> )- <b>A1</b>	65	–12
6	<b>1c</b>	<b>L1</b>	( <i>R</i> )- <b>A1</b>	40	–5
7	<b>1c</b>	<b>L2</b>	( <i>R</i> )- <b>A1</b>	65	–25
8	<b>1c</b>	<b>L3</b>	( <i>R</i> )- <b>A1</b>	65	–30
9	<b>1c</b>	<b>L4</b>	( <i>R</i> )- <b>A1</b>	60	86
10	<b>1c</b>	<b>L4</b>	( <i>S</i> )- <b>A1</b>	68	91
11	<b>1c</b>	<b>L5</b>	( <i>S</i> )- <b>A1</b>	75	76
12 <sup>e</sup>	<b>1c</b>	<b>L4</b>	( <i>S</i> )- <b>A1</b>	72	91
13 <sup>e</sup>	<b>1c</b>	<b>L4</b>	<b>A2</b>	99	91


<sup>a</sup> Unless noted otherwise, reactions were performed with substrate **1** (0.1 mmol),  $\alpha$ -cyano chalcone **2a** (0.12 mmol), Pd<sub>2</sub>(dba)<sub>3</sub> (5 mol%), **L** (20 mol%), additive (20 mol%) and 4 Å MS (100 mg) in toluene (1 mL) at 50 °C under Ar for 24 h. After completion, Et<sub>3</sub>N (20 mol%) was added, and the mixture was stirred at rt for 2 h. <sup>b</sup> Yield of isolated product **3a**. <sup>c</sup> Determined by HPLC analysis on a chiral stationary phase, and dr > 19 : 1 by <sup>1</sup>H NMR analysis. <sup>d</sup> Pd(PPh<sub>3</sub>)<sub>4</sub> (10 mol%) was used. <sup>e</sup> At 30 °C.

toluene at 50 °C under the catalysis of Pd(PPh<sub>3</sub>)<sub>4</sub> (Table 1, entry 1). It was speculated that enhancing the acidity of the benzylic C–H, by introducing an electron-withdrawing group at the indane ring, would be helpful for the formation of the desired dipole intermediate. Pleasingly, 6-nitro-substituted substrate **1b** showed good reactivity under the identical catalytic conditions. The expected formal  $[10 + 2]$  cycloadduct **3a'** was detected, whereas product **3a**, after isomerisation of the double bond, was found to be the more thermally stable one. Moreover, by simple treatment with catalytic amounts of Et<sub>3</sub>N in one pot, pure **3a** was efficiently isolated in a moderate yield with excellent diastereoselectivity (entry 2). We next explored the asymmetric version by utilising Pd<sub>2</sub>(dba)<sub>3</sub> and chiral ligands. Unfortunately, only moderate enantioselectivity was attained with phosphoramidite **L1** after extensively screening diverse ligands (entry 3).<sup>14</sup> We turned to explore the reaction by using a new substrate and a new catalytic system. Compared to allylic carbonates or esters, free allylic alcohols represent more atom-economic and environmentally benign precursors for generating the corresponding  $\pi$ -allyl species under transition metal catalysis, as water is the sole side product.<sup>15</sup> Nevertheless, alcohol **1c** suffered from low reactivity under the catalysis of Pd(0) due to the poor leaving ability of the hydroxyl group (entry 4). Subsequently, we tried to add a phosphoric acid as a co-catalyst, as it has been demonstrated to be beneficial for the oxidative addition of Pd(0) towards allylic alcohols and also might be beneficial for enantiocontrol.<sup>16</sup> To our gratification, the cycloaddition of **1c** and **2a** proceeded smoothly by using BINOL-derived (*R*)-**A1** as an additive and PPh<sub>3</sub> as the ligand, and **3a** was afforded in a moderate yield albeit with low enantioselectivity (entry 5). It should be noted that such a dehydration strategy has not been previously utilised in Pd-based dipole chemistry.<sup>10</sup> Moreover, it also suggested that acid (*R*)-**A1** would play dual roles in sequential activation of both substrates **1c** and **2a**. Consequently, some chiral ligands were investigated in combination with (*R*)-**A1**. Chiral phosphoramidites **L1** and **L2** showed moderate reactivity but still with low enantiocontrol (entries 6 and 7). While the newly designed phosphoramidite **L3** gave fair enantioselectivity (entry 8), (*R*)-BINOL derived **L4**, a diastereomer of **L3**, exhibited much higher enantioselectivity (entry 9). Moreover, an improved yield with a better ee value was obtained by using the combination of acid (*S*)-**A1** and **L4** (entry 10). Nevertheless, ligand **L5** with a smaller TES group delivered reduced enantioselectivity (entry 11). A slightly higher yield was attained at 30 °C (entry 12). It was further found that the chiral acid was not necessary, and even higher catalytic activity and exclusive regioselectivity with retained enantioselectivity was achieved with simple phosphoric acid **A2** (entry 13).

### Substrate investigation of asymmetric $[10 + 2]$ cycloadditions

Subsequently, we explored the substrate scope and limitations of the asymmetric formal  $[10 + 2]$  cycloaddition reaction under the cooperative catalysis of Pd/**L4** and phosphoric acid **A2**. As summarised in Table 2, an array of  $\alpha$ -cyano chalcones **2** were first tested in the reactions with benzocyclopentenol **1c**. Acceptors **2** with different aromatic substituents at the  $\beta$ -



**Table 2** Substrate scope of asymmetric [10 + 2] cycloadditions between benzocyclopentenols **1** and  $\alpha$ -cyano chalcones **2**<sup>a</sup>


Entry	EWG	R <sup>1</sup> , R <sup>2</sup>	Yield <sup>b</sup> (%)	ee <sup>c</sup> (%)
1	6-NO <sub>2</sub>	Ph, Ph	<b>3a</b> , 99	91 <sup>d</sup>
2	6-NO <sub>2</sub>	Ph, 2-ClC <sub>6</sub> H <sub>4</sub>	<b>3b</b> , 96	93
3 <sup>e</sup>	6-NO <sub>2</sub>	Ph, 3-ClC <sub>6</sub> H <sub>4</sub>	<b>3c</b> , 89 (80)	82 (92)
4	6-NO <sub>2</sub>	Ph, 4-BrC <sub>6</sub> H <sub>4</sub>	<b>3d</b> , 67	89
5 <sup>e</sup>	6-NO <sub>2</sub>	Ph, 4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	<b>3e</b> , — (70)	— (90)
6	6-NO <sub>2</sub>	Ph, 2-MeOC <sub>6</sub> H <sub>4</sub>	<b>3f</b> , 99	95
7 <sup>e</sup>	6-NO <sub>2</sub>	Ph, 3-MeOC <sub>6</sub> H <sub>4</sub>	<b>3g</b> , 99 (75)	87 (93)
8	6-NO <sub>2</sub>	Ph, 4-MeOC <sub>6</sub> H <sub>4</sub>	<b>3h</b> , 99	88
9	6-NO <sub>2</sub>	Ph, 2-naphthyl	<b>3i</b> , 99	91
10	6-NO <sub>2</sub>	Ph, 2-thienyl	<b>3j</b> , 70	88
11 <sup>e</sup>	6-NO <sub>2</sub>	2-BrC <sub>6</sub> H <sub>4</sub> , Ph	<b>3k</b> , — (73)	— (91)
12 <sup>e</sup>	6-NO <sub>2</sub>	3-BrC <sub>6</sub> H <sub>4</sub> , Ph	<b>3l</b> , — (75)	— (82)
13	6-NO <sub>2</sub>	4-ClC <sub>6</sub> H <sub>4</sub> , Ph	<b>3m</b> , 99	93
14	6-NO <sub>2</sub>	4-BrC <sub>6</sub> H <sub>4</sub> , Ph	<b>3n</b> , 76	94
15 <sup>e</sup>	6-NO <sub>2</sub>	2-MeC <sub>6</sub> H <sub>4</sub> , Ph	<b>3o</b> , 86 (90)	83 (89)
16 <sup>e</sup>	6-NO <sub>2</sub>	3-MeC <sub>6</sub> H <sub>4</sub> , Ph	<b>3p</b> , 88 (72)	82 (91)
17	6-NO <sub>2</sub>	4-MeC <sub>6</sub> H <sub>4</sub> , Ph	<b>3q</b> , 99	92
18	6-NO <sub>2</sub>	2-Naphthyl, Ph	<b>3r</b> , 91	90
19	4-NO <sub>2</sub>	Ph, Ph	<b>3s</b> , 99	89
20	5-NO <sub>2</sub>	Ph, Ph	<b>3a</b> , 61	88
21	6-CN	Ph, Ph	<b>3t</b> , 99	89
22 <sup>f</sup>	6-NO <sub>2</sub>	Ph, Ph	<b>3a</b> , 91	88

<sup>a</sup> Unless noted otherwise, reactions were performed with allylic alcohol **1** (0.1 mmol), activated alkene **2** (0.12 mmol), Pd<sub>2</sub>(dba)<sub>3</sub> (5 mol%), **L4** (20 mol%), acid **A2** (20 mol%) and 4 Å MS (100 mg) in toluene (1 mL) at 30 °C under Ar for 2–24 h. After completion, Et<sub>3</sub>N (20 mol%) was added, and the mixture was stirred at rt for 0.5–2 h. <sup>b</sup> Yield of the isolated product. <sup>c</sup> Determined by HPLC analysis on a chiral stationary phase; dr > 19 : 1 by <sup>1</sup>H NMR analysis. <sup>d</sup> The absolute configuration of enantiopure **3a** was determined by X-ray analysis. The other products were assigned by analogy. <sup>e</sup> Data in parentheses were obtained with acid (*S*)-**A1** (20 mol%). <sup>f</sup> On a 1.0 mmol scale.


position, including electron-deficient and -rich ones, smoothly gave corresponding products **3a–3j** in moderate to good yields with high stereoselectivity (Table 2, entries 1–10). The one with a *p*-nitrophenyl group showed low reactivity under the standard conditions, but desired product **3e** was obtained in a good yield with excellent enantioselectivity by using (*S*)-**A1** as the additive (entry 5). Notably, the halogen-substituted ones were compatible with the reactions (entries 2–4). Similarly good results were generally produced for substrates **2** with diverse  $\alpha'$ -aroyl groups (entries 11–18), whereas (*S*)-**A1** was found to be helpful in some cases. Unfortunately, the activated alkenes with aliphatic substituents failed to afford the cycloadducts, probably because of the undesired acidic vinylogous C–H of these substrates.<sup>14</sup> On the other hand, benzocyclopentenols **1** with a nitro group at different positions on the phenyl ring were applied in the reactions with **2a**. When the one with a 4-nitro substituent was utilised, cycloadduct **3s** was afforded in good yield with good enantioselectivity (entry 19).<sup>14</sup> Nevertheless, the alcohol with a 5-nitro group also delivered product **3a**, same as that from **1c** (6-

nitro), indicating an isomerisation process of the 10 $\pi$ -intermediate would be involved (entry 20).<sup>14</sup> In addition, a 6-cyano-substituted alcohol showed comparable reactivity, and the expected cycloadduct **3t** was attained quantitatively with high enantioselectivity (entry 21). We also conducted the reaction on a larger scale, and similar good results were afforded (entry 22).

### More substrate exploration

Apart from  $\alpha$ -cyano chalcones, we successfully extended the asymmetric [10 + 2] cycloaddition reactions to other types of activated alkenes for constructing polycyclic frameworks with more structural diversity. It was found that barbiturate-derived alkenes **4** could be well assembled with alcohol **1c** under the standard catalytic conditions. As summarised in Table 3, alkenes **4** with a different aryl or heteroaryl substituent underwent the cycloaddition reaction smoothly, furnishing corresponding spirocyclic architectures **5a–5e** in moderate to good yields with high stereoselectivity (entries 1–7). Besides, the one with a 2-styryl group also worked well, and product **5h** was obtained in moderate yield and enantioselectivity (entry 8). (*S*)-**A1** was further tested when the reactions did not work well (entries 2 and 7).


Moreover, the formal [10 + 2] cycloaddition reaction could be extended to benzylidene Meldrum's acid **6** by using the combination of Pd/**L4** and (*S*)-**A1**, delivering product **7** in a moderate yield with high enantioselectivity, whereas a higher yield with lower enantiocontrol was observed with acid **A2** (Scheme 2). Interestingly, when barbiturate-heptafulvene **8** was employed,<sup>17</sup> an asymmetric [10 + 8] higher-order cycloaddition reaction was applicable, and polycyclic **9** was constructed in a moderate yield with excellent enantiocontrol. Moreover, the assembly of carbonate **1g** and 2-benzylidenbenzo[*b*]thiophen-

**Table 3** Substrate scope of asymmetric [10 + 2] cycloadditions between benzocyclopentenol **1c** and barbiturate-derived alkenes **4**<sup>a</sup>


Entry	R	Yield <sup>b</sup> (%)	ee <sup>c</sup> (%)
1	Ph	<b>5a</b> , 99	93
2 <sup>d</sup>	4-BrC <sub>6</sub> H <sub>4</sub>	<b>5b</b> , — (60)	— (94)
3	4-MeC <sub>6</sub> H <sub>4</sub>	<b>5c</b> , 99	93
4	4-MeOC <sub>6</sub> H <sub>4</sub>	<b>5d</b> , 99	80
5	2-Naphthyl	<b>5e</b> , 75	93
6	2-Furyl	<b>5f</b> , 89	96
7 <sup>d</sup>	2-Thienyl	<b>5g</b> , 76 (64)	80 (87)
8	2-Styryl	<b>5h</b> , 91	80

<sup>a</sup> Unless noted otherwise, reactions were performed with allylic alcohol **1c** (0.1 mmol), alkene **4** (0.12 mmol), Pd<sub>2</sub>(dba)<sub>3</sub> (0.005 mmol, 5 mol%), **L4** (20 mol%), **A2** (15 mol%) and 4 Å MS (100 mg) in toluene (1 mL) at 10–15 °C for 2–12 h under Ar. After completion, Et<sub>3</sub>N (20 mol%) was added, and the mixture was stirred at rt for 2 h. <sup>b</sup> Yield of the isolated product. <sup>c</sup> Determined by HPLC analysis on a chiral stationary phase. <sup>d</sup> Data in parentheses were obtained with acid (*S*)-**A1** (20 mol%).





Scheme 2 More exploration of higher-order cycloadditions.

3(2*H*)-one **10** was successful under the cooperative catalysis of Pd/L4 and benzoic acid, and [10 + 4] cycloadduct **11** embedding an oxepine motif was isolated in a moderate yield with high enantiocontrol, along with the observation of minor [10 + 2] product **12**. Interestingly, using *o*-fluorobenzoic acid as the additive, cycloadduct **12** with similar enantiocontrol was delivered as the major one after the treatment with Et<sub>3</sub>N, albeit in a slightly lower yield.<sup>14</sup>

### Mechanistic proposal

To gain more insight into the mechanism, a control experiment with the analogous acyclic alcohol **13** was conducted. As shown in Scheme 3a, no reaction occurred in combination with acceptor **2a** in the presence of Pd/L4 and acid **A2**, which suggested that

expected zwitterionic intermediate **III** might not be generated *via* a similar oxidative addition/deprotonation process under the standard conditions.<sup>18</sup> These results indicated that the indene-based structure was crucial for the vinylogous activation of the benzylic C–H group to generate the active dipole species. Therefore, as outlined in Scheme 3b, it was proposed that allylic alcohol **1c** would be partially protonated in the presence of phosphoric acid **A2** and undergo oxidative addition with complex Pd(0)/L4 along with the release of H<sub>2</sub>O. The  $\pi$ -allylpalladium complex moiety of the resultant intermediate **I** would further enhance the acidity of the benzylic C–H group, and a deprotonation process would occur to give dipole **II**, which would more reasonably exist as a polyconjugated 10 $\pi$ -type cycloaddend. Subsequently, acid **A2** would act as a Brønsted acid to activate  $\alpha$ -cyano chalcone **2a**, rendering the assembly with dipole **II** to deliver adduct **IV**. An intramolecular allylic alkylation would be followed to afford **3a'** together with the regeneration of Pd(0). Finally, an isomerisation process took place with the assistance of Et<sub>3</sub>N to furnish the thermally more stable product **3a**.

## Conclusions

A new type of metal-containing 10 $\pi$ -cycloaddend could be generated from benzocyclopentenols with an electron-withdrawing group *via* a deprotonation strategy under the cooperative catalysis of Pd(0) and phosphoric acid. The subsequent asymmetric higher-order [10 + 2] cycloaddition reactions with diverse activated alkenes were efficiently accomplished by employing a newly designed phosphoramidite ligand, producing polycyclic frameworks embedding an indene core in moderate to excellent yields with good to excellent enantioselectivity. In addition, [10 + 8] and [10 + 4] higher-order cycloaddition reactions were similarly realised with a heptafulvene derivative or a cyclic enone substrate, respectively. The newly designed metal-containing 10 $\pi$ -synthons would have more potential in asymmetric reactions, and the results will be reported in due course.

## Data availability

The data that support the findings of this study are available in the ESI† or on request from the corresponding author.

## Author contributions


The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We are grateful for the financial support from NSFC (21971166, 21961132004, 21931006 and 21921002) and 111 Project (B18035).



Scheme 3 Mechanistic proposal. (a) Control experiment. (b) Proposed catalytic mechanism.





## Notes and references

- (a) D. McLeod, M. K. Thøgersen, N. I. Jessen, K. A. Jørgensen, C. S. Jamieson, X.-S. Xue, K. N. Houk, F. Liu and R. Hoffmann, *Acc. Chem. Res.*, 2019, **52**, 3488; (b) S. Frankowski, M. Romaniszyn, A. Skrzyńska and Ł. Albrecht, *Chem.-Eur. J.*, 2020, **26**, 2120; (c) N. I. Jessen, D. McLeod and K. A. Jørgensen, *Chem*, 2022, **8**, 20; (d) M.-M. Zhang, B.-L. Qu, B. Shi, W.-J. Xiao and L.-Q. Lu, *Chem. Soc. Rev.*, 2022, **51**, 4146.
- (a) K. N. Houk and R. B. Woodward, *J. Am. Chem. Soc.*, 1970, **92**, 4143; (b) K. N. Houk and R. B. Woodward, *J. Am. Chem. Soc.*, 1970, **92**, 4145 For related initial contributions, see: (c) R. C. Cookson, B. V. Drake, J. Hudec and A. Morrison, *Chem. Commun.*, 1966, 15; (d) S. Ito, Y. Fujise, T. Okuda and Y. Inoue, *Bull. Chem. Soc. Jpn.*, 1966, **39**, 1351; (e) L. A. Paquette and J. H. Barrett, *J. Am. Chem. Soc.*, 1966, **88**, 2590.
- (a) K. N. Houk, L. J. Luskus and N. S. Bhacca, *J. Am. Chem. Soc.*, 1970, **92**, 6392; (b) T. C. Wu and K. N. Houk, *J. Am. Chem. Soc.*, 1985, **107**, 5308; (c) W. Friedrichsen, W. Seidel and T. Debaerdemaeker, *J. Heterocycl. Chem.*, 1983, **20**, 1621; (d) D. McLeod, A. CherubiniCelli, N. Sivasothirajah, C. H. McCulley, M. L. Christensen and K. A. Jørgensen, *Chem.-Eur. J.*, 2020, **26**, 11417; (e) Y. Hayashi, H. Gotoh, M. Honma, K. Sankar, I. Kumar, H. Ishikawa, K. Konno, H. Yui, S. Tsuzuki and T. Uchimar, *J. Am. Chem. Soc.*, 2011, **133**, 20175.
- For a review, see: (a) J. H. Rigby, *Acc. Chem. Res.*, 1993, **26**, 579 For selected examples, see: (b) J. H. Rigby and H. S. Ateeq, *J. Am. Chem. Soc.*, 1990, **112**, 6442; (c) K. Chaffee, P. Huo, J. B. Sheridan, A. Barbieri, A. Aistars, R. A. Lalancette, R. L. Ostrander and A. L. Rheingold, *J. Am. Chem. Soc.*, 1995, **117**, 1900; (d) J. H. Rigby, K. M. Short, H. S. Ateeq and J. A. Henshilwood, *J. Org. Chem.*, 1992, **57**, 5290.
- For selected examples, see: (a) G. Bertuzzi, D. McLeod, L.-M. Mohr and K. A. Jørgensen, *Chem.-Eur. J.*, 2020, **26**, 15491; (b) M. Xie, X. Liu, X. Wu, Y. Cai, L. Lin and X. Feng, *Angew. Chem., Int. Ed.*, 2013, **52**, 5604; (c) B. M. Trost, P. J. McDougall, O. Hartmann and P. T. Wathen, *J. Am. Chem. Soc.*, 2008, **130**, 14960; (d) P. Yu, C. Q. He, A. Simon, W. Li, R. Mose, M. K. Thøgersen, K. A. Jørgensen and K. N. Houk, *J. Am. Chem. Soc.*, 2018, **140**, 13726; (e) M. Romaniszyn, K. Gronowska and Ł. Albrecht, *J. Org. Chem.*, 2019, **84**, 9929; (f) S. Frankowski, A. Skrzyńska and Ł. Albrecht, *Chem. Commun.*, 2019, **55**, 11675; (g) R. Mose, G. Preegel, J. Larsen, S. Jakobsen, E. H. Iversen and K. A. Jørgensen, *Nat. Chem.*, 2017, **9**, 487; (h) H. Liu, Y. Wu, Y. Zhao, Z. Li, L. Zhang, W. Yang, H. Jiang, C. Jing, H. Yu, B. Wang, Y. Xiao and H. Guo, *J. Am. Chem. Soc.*, 2014, **136**, 2625.
- (a) D. Johnson and G. Jones, *J. Chem. Soc., Perkin Trans. 1*, 1972, 840; (b) N. I. Jessen, M. Bura, G. Bertuzzi and K. A. Jørgensen, *Angew. Chem., Int. Ed.*, 2021, **60**, 18527.
- K. Hafner and W. Bauer, *Angew. Chem., Int. Ed. Engl.*, 1968, **7**, 297.
- (a) Z. Zhou, Z.-X. Wang, Y.-C. Zhou, W. Xiao, Q. Ouyang, W. Du and Y.-C. Chen, *Nat. Chem.*, 2017, **9**, 590; (b) G. Bertuzzi, M. K. Thøgersen, M. Giardinetti, A. Vidal-Albalat, A. Simon, K. N. Houk and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2019, **141**, 3288; (c) B. S. Donslund, A. Monleón, T. A. Palazzo, M. L. Christensen, A. Dahlgaard, J. D. Erickson and K. A. Jørgensen, *Angew. Chem., Int. Ed.*, 2018, **57**, 1246; (d) B. S. Donslund, N. I. Jessen, G. Bertuzzi, M. Giardinetti, T. A. Palazzo, M. L. Christensen and K. A. Jørgensen, *Angew. Chem., Int. Ed.*, 2018, **57**, 13182; (e) D. McLeod, J. A. Izzo, D. K. B. Jørgensen, R. F. Lauridsen and K. A. Jørgensen, *ACS Catal.*, 2020, **10**, 10784; (f) J. Zhao, X. Zheng, Y.-S. Gao, J. Mao, S.-X. Wu, W.-L. Yang, X. Luo and W.-P. Deng, *Chin. J. Chem.*, 2021, **39**, 3219; (g) N. I. Jessen, G. Bertuzzi, M. Bura, M. L. Skipper and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2021, **143**, 6140; (h) G. Bertuzzi, V. Corti, J. A. Izzo, S. Ričko, N. I. Jessen and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2022, **144**, 1056.
- (a) Q. Peng, S.-J. Li, B. Zhang, D. Guo, Y. Lan and J. Wang, *Commun. Chem.*, 2020, **3**, 177; (b) Y. Yang, Y. Jiang, W. Du and Y.-C. Chen, *Chem.-Eur. J.*, 2020, **26**, 1754.
- (a) T. Hashimoto and K. Maruoka, *Chem. Rev.*, 2015, **115**, 5366; (b) J.-J. Feng and J. Zhang, *ACS Catal.*, 2016, **6**, 6651; (c) B. D. W. Allen, C. P. Lakeland and J. P. A. Harrity, *Chem.-Eur. J.*, 2017, **23**, 13830; (d) N. De and E. J. Yoo, *ACS Catal.*, 2018, **8**, 48; (e) T.-R. Li, Y.-N. Wang, W.-J. Xiao and L.-Q. Lu, *Tetrahedron Lett.*, 2018, **59**, 1521.
- For selected examples with non-conjugated dipole species, see: (a) L.-C. Yang, Y.-N. Wang, R. Liu, Y. Luo, X. Q. Ng, B. Yang, Z.-Q. Rong, Y. Lan, Z. Shao and Y. Zhao, *Nat. Chem.*, 2020, **12**, 860; (b) Y.-N. Wang, L.-C. Yang, Z.-Q. Rong, T.-L. Liu, R. Liu and Y. Zhao, *Angew. Chem., Int. Ed.*, 2018, **57**, 1596; (c) Q.-L. Zhang, Q. Xiong, M.-M. Li, W. Xiong, B. Shi, Y. Lan, L.-Q. Lu and W.-J. Xiao, *Angew. Chem., Int. Ed.*, 2020, **59**, 14096.
- (a) T. Hayashi, T. Suzuka, A. Okada and M. Kawatsura, *Tetrahedron: Asymmetry*, 2004, **15**, 545; (b) H.-L. Cui, X.-H. Sun, L. Jiang, L. Dong and Y.-C. Chen, *Eur. J. Org. Chem.*, 2011, 7366; (c) J. Zhang, H.-H. Wu and J. Zhang, *Org. Lett.*, 2017, **19**, 6080.
- Z. Wang, Z. He, L. Zhang and Y. Huang, *J. Am. Chem. Soc.*, 2018, **140**, 735.
- Please see the ESI† for more details.
- (a) B. Sundararaju, M. Achard and C. Bruneau, *Chem. Soc. Rev.*, 2012, **41**, 4467; (b) A. Baeza and C. Nájera, *Synthesis*, 2014, **46**, 25; (c) N. A. Butt and W. Zhang, *Chem. Soc. Rev.*, 2015, **44**, 7929.
- (a) D. Parmar, E. Sugiono, S. Raja and M. Rueping, *Chem. Rev.*, 2014, **114**, 9047; (b) G. Jiang and B. List, *Angew. Chem., Int. Ed.*, 2011, **50**, 9471; (c) Z.-L. Tao, W.-Q. Zhang, D.-F. Chen, A. Adele and L.-Z. Gong, *J. Am. Chem. Soc.*, 2013, **135**, 9255; (d) C.-C. Tsai, C. Sandford, T. Wu, B. Chen, M. S. Sigman and F. D. Toste, *Angew. Chem., Int. Ed.*, 2020, **59**, 14647.
- X. Chen, M. K. Thøgersen, L. Yang, R. F. Lauridsen, X.-S. Xue, K. A. Jørgensen and K. N. Houk, *J. Am. Chem. Soc.*, 2021, **143**, 934.



18 The corresponding 1,3-dipole could be formed from the allylic carbonate under the catalysis of Pd(0) and special ligands, see: (a) B. M. Trost and Z. Zuo, *Angew. Chem., Int. Ed.*, 2020, **59**, 1243; (b) B. M. Trost, Z. Jiao and C.-I. Hung, *Angew. Chem., Int. Ed.*, 2019, **58**, 15154.

*Ed.*, 2020, **59**, 1243; (b) B. M. Trost, Z. Jiao and C.-I. Hung, *Angew. Chem., Int. Ed.*, 2019, **58**, 15154.

