Volume 59 Number 5 16 January 2023 Pages 493-648

# ChemComm

Chemical Communications



ISSN 1359-7345



**COMMUNICATION** Glenn M. Sammis, Nicholas D. Ball *et al.* Facile synthesis of sulfonyl fluorides from sulfonic acids

## ChemComm

Check for updates

Cite this: Chem. Commun., 2023,



**View Article Online** 

View Journal | View Issue

### COMMUNICATION

#### Facile synthesis of sulfonyl fluorides from sulfonic acids<sup>†</sup>

Brodie J. Thomson,<sup>a</sup> Samuel R. Khasnavis,<sup>b</sup> Emma C. Grigorian,<sup>‡b</sup> Rohun Krishnan,<sup>‡<sup>b</sup></sup> Theodore D. Yassa,<sup>‡<sup>b</sup></sup> Kelvin Lee,<sup>b</sup> Glenn M. Sammis <sup>(D)</sup> \*<sup>a</sup> and Nicholas D. Ball 🕩 \*b

DOI: 10.1039/d2cc05781f

Received 25th October 2022 Accepted 6th December 2022

rsc.li/chemcomm

**59**, 555

Herein, we demonstrate two complementary strategies for the syntheses of sulfonyl fluorides using sulfonic acids and their salts. One strategy involves the conversion of sulfonic acid sodium salts to sulfonyl fluorides using thionyl fluoride in 90-99% yields in one hour. Lessons learned from the mechanism of this reaction also have enabled a complementary deoxyfluorination of sulfonic acids using Xtalfluor-E<sup>®</sup> – a bench stable solid – allowing for the conversion of both aryl and alkyl sulfonic acids and salts to sulfonyl fluorides in 41-94% yields. Notably, using Xtalfluor-E<sup>®</sup> enabled milder conditions and the use of both sulfonic acids and their sodium salts.

Sulfonyl fluorides have been widely adopted as a novel set of click reagents since their introduction by Sharpless in 2014.<sup>1</sup> The functional group is valuable due to its widespread applications in polymer chemistry, surface modifications, catalysis,<sup>2</sup> and as precursors in the formation of additional sulfur(vi) species.<sup>1,3</sup> In addition to its impact on synthesis, sulfonyl fluorides have found important applications as biological probes and protein inhibitors.<sup>4</sup> Their high desirability across many scientific fields has instigated a robust development of synthetic strategies to access sulfonyl fluorides, including from sulfonamides,<sup>5</sup> thiols and disulfides,<sup>6</sup> sulfonyl hydrazides,<sup>7</sup> sulfinic acids,<sup>7</sup> Grignard reagents,<sup>8</sup> arynes,<sup>9</sup> aryl halides,<sup>10</sup> carboxylic acids,<sup>11</sup> and diazonium salts.<sup>12</sup> Despite these innovations, an underexplored synthetic approach to sulfonyl fluorides has been to employ sulfonic acids. This strategy enables access to sulfonyl fluorides from stable, readily accessible S(vi) compounds without involving oxidation, reactive intermediates, or specialized compounds like sulfonamides or hydrazides.5,7 Current strategies toward the conversion of sulfonic acids to sulfonyl fluorides require two- step processes involving the in situ formation of sulfonyl chlorides

(Scheme 1). Recent advances using this approach include a onepot, two-step protocol, forming the sulfonyl chloride using trichloroacetonitrile13 or cyanuric chloride14 before Cl-/F- halogen exchange. While an improvement over the two-pot procedures, these strategies either are restricted to aryl sulfonic acids or require long reaction times (24 h). Furthermore, neither method was successful when using with electron poor substrates.

An alternative deoxyfluorination approach that uses a reagent to both activate the sulfonic acid and provide a fluoride source would enable the direct access of sulfonyl fluorides from sulfonic acids and salts in a single step (Scheme 1). Efforts toward this direct deoxyfluorination strategy have been limited. Recently, 1,1,2,2-tetrafluoroethyl-*N*,*N*-dimethylamine<sup>15</sup> has been employed with only a single substrate, and a method that uses carbonyl fluoride<sup>16</sup> as the reactive intermediate was only demonstrated with five substrates. Liskamp and coworkers have also demonstrated the ability of amino sulfur(IV) fluorides, including DAST, Xtalfluor-E<sup>®</sup>, and Xtalfluor-M<sup>®</sup> to access a few





Ar/Alk

`OH/Na

<sup>&</sup>lt;sup>a</sup> Department of Chemistry, University of British Columbia, 2036 Main Mall,

Vancouver, British Columbia V6T 1Z1, Canada. E-mail: gsammis@chem.ubc.ca

<sup>&</sup>lt;sup>b</sup> Department of Chemistry, Pomona College, 645 North College Avenue, Claremont, California 91711, USA. E-mail: nicholas.ball@pomona.edu

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: https://doi.org/ 10.1039/d2cc05781f

<sup>‡</sup> These authors contributed equally.

#### Communication

sulfonyl fluoride inhibitors. However, yields were limited, and the reactions required heating reaction solutions at reflux containing HF and sulfonic acids over long reaction times.<sup>17</sup> Collectively, these examples highlight the challenges of using sulfonic acids to make sulfonyl fluorides *via* deoxyfluorination, and the need for new methodologies. Herein, we demonstrate two complementary strategies toward sulfonyl fluorides directly from sulfonic acids and their salts using mild reaction conditions.

We began our investigation by exploring the use of thionyl fluoride –  $SOF_2$  – in the deoxyfluorination of sulfonate salts. Thionyl fluoride has recently been reported in a rapid deoxyfluorination of carboxylic acids to form the corresponding acyl fluorides.<sup>18</sup> Notably, thionyl fluoride is believed to activate carboxylic acids *via* acyl fluorosulfinate intermediates, facilitating rapid conversion to the corresponding acyl fluoride. Importantly, thionyl fluoride gas can be generated *in situ* in the amounts needed using readily available bench-stable reagents. We hypothesized that thionyl fluoride could also be used to activate more challenging sulfonic acid derivatives toward sulfonyl fluorides.

Our studies began by treating pyridinium *p*-toluenesulfonic acid with a solution of thionyl fluoride in acetonitrile due to the solvent's previously reported effectiveness in the deoxyfluorination of carboxylic acids (Table 1, entry 1).<sup>18</sup> No product was detected, and 4-methylbenzenesulfonic anhydride was observed as the exclusive product. Changing the reaction solvent to DMF in order to access similar intermediates reported in the COF2-mediated synthesis of sulfonic acids<sup>16</sup> led to no reaction at 80 °C (entry 2), but the desired product was formed in 48% yield when the reaction was heated to 115 °C (entry 3). Elevating the temperature further to 130 °C increased the formation of 3b to 92% yield. Increasing the amount of thionyl fluoride to 3 equivalents afforded the desired sulfonyl fluoride (3a) in 98% yield in only 1 hour (entry 5). Sodium salt derivatives, which represent the most accessible source of sulfonic acids, generally afforded low yields (entry 6). To increase the yield, BF3 OEt2 was added as it has been shown to increase the electrophilicity of sulfur(IV) reagents.<sup>19</sup> Gratifyingly, 3a was formed in quantitative yields by <sup>19</sup>F NMR in 1 hour (entry 7).<sup>20</sup>

			,			
		0,0 S,0 <sup>⊖</sup> x <sup>⊕</sup> 4a	SOF <sub>2</sub> , a	dditive ∽ nt, 1 h	S F 3a	
Entry	Х	Additive	Solvent	$T(^{\circ}C)$	SOF <sub>2</sub> equiv. <sup>a</sup>	Yield (%)
1	Pyridinium	_	MeCN	80	2	$0^b$
2	Pyridinium		DMF	80	2	0
3	Pyridinium		DMF	115	2	48
4	Pyridinium		DMF	130	2	92
5	Pyridinium		DMF	130	3	98
6	Na		DMF	130	3	25
7	Na	BF <sub>3</sub> ·OEt <sub>2</sub>	DMF	130	3	99

Table 1 Optimization of thionyl fluoride mediated deoxyfluorination

Yields determined by <sup>19</sup>F NMR spectroscopy at 0.25 mmol. <sup>*a*</sup> Equivalents of thionyl fluoride gas dissolved in DMF. <sup>*b*</sup> Only 4-methylbenzenesulfonic anhydride was detected by <sup>1</sup>H NMR spectroscopy.

With our optimized protocol in hand, the substrate scope was investigated with a range of aromatic and aliphatic acids (Table 2). Alkyl substitution had no effect on the reaction, with **3a–3c** all proceeding in near-quantitative yield. Similarly, the reaction was largely insensitive to electron rich and electron poor phenyl derivatives (**3d–3g**) Notably, this is the highest reported yield for 4-nitro derivative (**3g**) by any method without direct halogen exchange from sulfonyl chloride.<sup>21</sup> Naphthylsulfonyl fluoride **3h** was also synthesized in quantitative yield. Pyridine sulfuryl fluoride **3i** was only synthesized in 14% <sup>19</sup>F NMR yield, and significant quantities of the respective sulfonic acid anhydride were observed. Similarly, sulfuryl fluoride derivative **3j** could not be synthesized by this method.

Significantly, we found our optimized protocol could also be applied to the synthesis of aliphatic substrates with products 3k-3l produced in  $\geq 92\%$  yield, as determined by <sup>19</sup>F NMR spectroscopy. There are also few methods to make short chain sulfonyl fluorides with ease without going from a sulfonyl chloride. This method shows a higher yield for pentyl derivative 3k compared to the only existing method for this substrate.<sup>14</sup>

We next investigated the role of DMF in our optimized reaction. Heating a solution of thionyl fluoride dissolved in DMF to 130 °C resulted in the formation of  $FSO_2^{-}$ , indicated by a signal at +36 ppm by <sup>19</sup>F NMR spectroscopy.<sup>22</sup> The presence of



Isolated yields of sulfonyl fluorides are reported using a 0.5 mmol of sulfonic acid sodium salt. <sup>19</sup>F NMR yields are reported in brackets. <sup>*a*</sup> Isolated yields are reported using 0.5 mmol of pyridinium sulfonic acid. <sup>*b*</sup> Yield adjusted for 91% due to residual diethyl ether content.



 $FSO_2^-$  confirmed that deoxygenation of DMF was occurring within our reaction system forming either compound 5 or 7 (Scheme 2a).

Conversely, heating a solution of thionyl fluoride dissolved in chlorobenzene – a solvent with sufficiently high polarity and a boiling point comparable to DMF – to 130 °C in the presence of pyridinium *p*-toluene sulfonic acid afforded no detectable product by <sup>19</sup>F NMR spectroscopy, and only anhydride **9** was detected (Scheme 2b). The similarly poor product yield using acetonitrile in the initial reaction optimisation (Table 1, entry 1), in which **9** was the exclusive product, suggest that intermediate **8** (Scheme 2b) is selectively formed in the absence of DMF. Another equivalent of a sulfonate salt can then react with the S(IV) center of intermediate **8**, reforming **8**, or it reacts with the S(VI) center to form sulfonic anhydride **9**.

When DMF is used as a solvent, it reacts with  $SOF_2$  to presumingly form intermediate 5 (see ESI<sup>†</sup>). Intermediate 5, or Vilsmeier-like intermediate 7 may then react with the sulfonic acid in solution to form a DMF-activated sulfonic acid before subsequently reacting with free fluoride to afford the desired sulfonyl fluoride (Scheme 2a).

We were intrigued by intermediate 7 given its similarity in structure to the non-gaseous, deoxyfluorinating agents Xtalfluor-E<sup>®</sup> (10) and Xtalfluor-M<sup>®</sup> (a morpholine derivative of Xtalfluor-E<sup>®</sup>).<sup>23</sup> We hypothesized that a sulfonate could add to Xtalfluor-E<sup>®</sup> (10) generating a species akin to 6 (intermediate 11) that could undergo addition of fluoride to give a sulfonyl fluoride 3. (Scheme 2c). Although the reagents were previously reported to form sulfonyl fluorides using harsh reaction conditions, we were optimistic that using the Xtalfluor salts under our optimized protocol may afford a more practical method.

Using our previous protocol with Xtalfluor- $E^{(R)}$ , sulfonyl fluoride (4a) was successfully formed in 80% yield using pyridinium

Table 3 XtalFluor-E/M<sup>®</sup> acid sources

		Xtalfluor salt, additive				
Entry	Х	Additive	Solvent	$T(^{\circ}C)$	Xtalfluor salt	Yield <sup>a</sup> (%)
1	Pyridinium	_	DMF	130	Е	80
2	Pyridinium		MeCN	rt	E	92
3	$\dot{H}^{b}$	Pyridine	MeCN	50	Е	30
4	Na	NaF	MeCN	50	E	91
5	Na	NaF	MeCN	50	Μ	92
-		10				1.

 $^a$  Yields determined by  $^{19}{\rm F}$  NMR spectroscopy at 0.25 mmol.  $^b$  The monohydrate of the acid was used.

acid and DMF as a solvent (Table 3, entry 1). Notably, the yields improved to 92% when the reaction was performed at room temperature in acetonitrile (entry 2), which offers a milder approach to previously reported methods.<sup>17</sup> Similar to our reaction trials with thionyl fluoride, the yields of **4a** were significantly reduced when sulfonic acid sodium salts were tested (entry 3); however, the addition of sodium fluoride and an increased temperature circumvented this issue and led to the formation of **3a** in 92% yield in under 1 hour (entry 4). The reaction was equally efficient when Xtalfluor-M<sup>®</sup> was utilized (entry 5). However, we continued our study using Xtalfluor-E<sup>®</sup> due to its lower cost.<sup>24</sup>

Xtalfluor-E<sup>®</sup> was successful with all the tested aromatic sulfonic acids, affording similar yields to thionyl fluoride (between 42 and 100%) by <sup>19</sup>F NMR spectroscopy (Table 4). The reaction was more sensitive to electronic effects compared to the thionyl fluoride-mediated approach, where electronically rich systems (e.g. 3a, f) afforded up to quantitative yields by <sup>19</sup>F NMR spectroscopy. Conversely, electronically poor substrates, such as 3g and 3n, required longer reaction times. The comparatively lower isolated yield of 3g is due to its poor solubility in MeCN. Additionally, most alkyl and some benzyl substrates required extended reaction times. Unlike thionyl fluoride, heteroatomic substrate 30 was tolerated well under the optimized conditions, affording 71% isolated yield. We next tested aliphatic substrate pentanesulfonic acid (3k). However, pentene was identified as the major product.<sup>25</sup> Longer-chain aliphatic substrates, such as 3l, prevented the formation of the discovered alkenes, affording excellent isolated yields in 94% with extended reaction times.

Compared to existing methods, the Xtalfluor-E<sup>®</sup> deoxyfluorination of aromatic sulfonic acids under our optimized conditions displayed milder reaction conditions in expedited timeframes. Furthermore, the salts are easy to access, store and handle. This method provides an excellent alternative route to access versatile sulfonyl fluoride moieties.

Overall, we have demonstrated two separate methods for accessing sulfonyl fluorides utilizing the sulfur (v) oxidation in sulfur fluoride reagents thionyl fluoride and Xtalfluor-E<sup>®</sup>. Thionyl fluoride was found to be highly effective in the deoxy-fluorination of both aliphatic and aromatic substrates, where further investigations found the successful reaction was a result

Table 4 Xtalfluor-E  $^{(\!R\!)}$  -mediated deoxyfluorination of sulfonic acids and sulfonate salts



Isolated yields of sulfonyl fluorides are reported using a 0.250 mmol of sulfonic acid or sulfonate salt. Pyridine was used with all sulfonic acids and NaF was used with all sulfonate salts (see ESI). <sup>19</sup>F NMR yields are reported in brackets. Some reactions were run for extended periods, either <sup>*a*</sup> 16 hours, <sup>*b*</sup> 3 hours, or <sup>*c*</sup> 8 hours.

of a novel DMF-activated intermediate. The use of Xtalfluor-E<sup>(R)</sup> was subsequently demonstrated to offer competitively high yields in the conversion of aromatic sulfonic acids across a diverse array of electronically rich and poor substrates. This alternative method offers an additional approach to access sulfonyl fluorides using a solid, bench-stable reagent with significantly milder conditions and ease-of-handling compared to thionyl fluoride and current reported methods.

N. D. B. thanks the National Institutes of General Medical Sciences of the National Institutes of Health for their funding (NIH-R15-GM134457-01A1). G. M. S. thank the University of British Columbia (UBC), and the Natural Sciences and Engineering Research Council of Canada (2020-RGPIN-04653) for funding. Student support was provided in the form of a MITACS Accelerate fellowships (B. J. T.).

#### Conflicts of interest

There are no conflicts to declare.

#### Notes and references

1 J. Dong, L. Krasnova, M. G. Finn and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2014, **53**, 9430–9448.

- (a) P. K. Chinthakindi and P. I. Arvidsson, Eur. J. Org. Chem., 2018, 3648–3666; (b) A. S. Barrow, C. S. Smedley, Q. Zheng, S. Li, J. Dong and J. E. Moses, Chem. Soc. Rev., 2019, 48, 4731–4758; (c) Z. Liu, J. Li, S. Li, G. Li, K. B. Sharpless and P. Wu, J. Am. Chem. Soc., 2018, 140, 2919–2925; (d) R. Lekkala, R. Lekkala, B. Moku, K. P. Rakesh and H.-L. Qin, Org. Chem. Front., 2019, 6, 3490–3516; (e) C. Lee, A. J. Cook, J. E. Elisabeth, N. C. Friede, G. M. Sammis and N. D. Ball, ACS Catal., 2021, 11, 6578–6589; (f) T. S.-B. Lou and M. C. Willis, Nat. Rev. Chem., 2022, 6, 146–162.
- 3 P. Mukherjee, C. P. Woroch, L. Cleary, M. Rusznak, R. W. Franzese, M. R. Reese, J. W. Tucker, J. M. Humphrey, S. M. Etuk, S. C. Kwan, C. W. Am Ende and N. D. Ball, *Org. Lett.*, 2018, **20**, 3943–3947.
- 4 A. Narayanan and L. H. Jones, Chem. Sci., 2015, 6, 2650-2659.
- 5 M. Pérez-Palau and J. Cornella, Eur. J. Org. Chem., 2020, 17, 2497–2500.
- 6 For representative examples, see: (a) G. Laudadio, A. A. Bartolomeu, L. M. H. M. Verwijlen, Y. Cao, K. T. de Oliveira and T. Noël, J. Am. Chem. Soc., 2019, 141, 11832–11836; (b) S. W. Wright and K. N. Hallstrom, J. Org. Chem., 2006, 71, 1080–1084.
- 7 L. Tang, Y. Yang, L. Wen, X. Yang and Z. Wang, Green Chem., 2016, 18, 1224–1228.
- 8 C. Lee, N. D. Ball and G. S. Sammis, Chem. Commun., 2019, 55, 14753-14756.
- 9 J. Kwon and B. Moon Kim, Org. Lett., 2019, 21, 428-433.
- 10 (a) A. T. Davies, J. M. Curto, S. W. Bagley and M. C. Davies, *Chem. Sci.*, 2017, 8, 1233–1237; (b) A. L. Tribby, I. Rodríguez, S. Sharffudin and N. D. Ball, *J. Org. Chem.*, 2017, 82, 2294–2299.
- 11 Z. Ma, Y. Liu, X. Ma, X. Hu, Y. Guo, Q.-Y. Chen and C. Liu, Org. Chem. Front., 2022, 9, 1115–1120.
- 12 For representative examples, see: (a) Z. Ma, L. Shan, X. Ma, X. Hu, Y. Guo, Q.-Y. Chen and C. Liu, *J. Fluorine Chem.*, 2022, 254, 109948; (b) T. Zhong, M.-K. Pang, Z.-D. Chen, B. Zhang, J. Weng and G. Lu, *Org. Lett.*, 2020, 22, 3072–3078; (c) Y. Liu, D. Yu, Y. Gup, J.-C. Xiao, Q.-Y. Chen and C. Liu, *Org. Lett.*, 2020, 22, 2281–2286.
- 13 J.-G. Kim and D. O. Jang, Synlett, 2010, 20, 3049.
- 14 Y. Jiang, N. S. Alharbi, B. Sun and H.-L. Qin, RSC Adv., 2019, 9, 13863–13867.
- 15 V. A. Petrov, S. Swearingen, W. Hong and W. C. Petersen, *J. Fluorine Chem.*, 2001, **109**, 25–31.
- 16 S. Zhao, Y. Guo, Z. Su, C. Wu, W. Chen and Q.-Y. Chen, *Chin. J. Chem.*, 2021, **39**, 1225–12320.
- 17 For representative examples, see: (a) R. Artschwager, D. J. Ward, S. Gannon, A. J. Brouwer, H. van de Langemheen, H. Kowalski and R. M. J. Liskamp, J. Med. Chem., 2018, 61, 5396–5411; (b) A. J. Brouwer, N. H. Álvarez, A. Ciaffoni, H. van de Langemheen and R. M. J. Liskamp, Bioorg. Med. Chem., 2016, 24, 3429–3435; (c) A. J. Brouwer, T. Ceylan, A. M. Jonker, T. van der Linden and R. M. J. Liskamp, Bioorg. Med. Chem., 2011, 19, 2397–2406.
- 18 C. Lee, B. J. Thomson and G. M. Sammis, Chem. Sci., 2022, 13, 188–194.
- 19 I. Ruppert, J. Fluorine Chem., 1982, 20, 79-84.
- 20 Sulfuryl fluoride also failed in yielding any desired product under similar conditions, suggesting the sulfur(w) oxidation was required for the success of the reaction and highlighted the unique reactivity of thionyl fluoride.
- 21 For representative examples, see: (a) L. Tang, X. Zhao, X. Yang, Y. Zhou and G. Zhou, *CN Pat.*, 105198683, 2015; (b) T. T. Bui, V. H. Tran and H.-K. Kim, *Adv. Synth. Catal.*, 2022, **364**, 341–347; (c) D. Louvel, A. Chelagha, J. Rouillon, P.-A. Payard, L. Khrouz, C. Monnereau and A. Tlili, *Chem. Eur. J.*, 2021, **27**, 8704–8707.
- 22 K. Gulbe, J. Luginina, E. Jansons, A. Kinens and M. Turks, *Beilstein J. Org. Chem.*, 2021, 17, 964–976.
- 23 F. Beaulieu, L.-P. Beuregard, G. Courchesne, M. Couturier, F. LaFlamme and A. L'Heureux, Org. Lett., 2009, 11, 5050–5053.
- 24 Xtlalfluor-E<sup>®</sup> is currently priced \$196.00/25g USD: Sigma-Aldrich, https://www.sigmaaldrich.com/CA/en/product/aldrich/719439, accessed October 2022. Xtalfluor-M<sup>®</sup> is currently priced at \$302.00/25g USD: Sigma-Aldrich, https://www.sigmaaldrich.com/CA/en/product/aldrich/ 719447?gclsrc=aw.ds&gclsrc=ds&gclsrc=aw.ds, accessed October 2022.
- 25 For a discussion on the formation of alkene side-products, see the ESI<sup>†</sup>.