


Showcasing research from Professor Tianlei Zhang's laboratory, School of Chemistry & Environment Science, Shaanxi University of Technology, Hanzhong, China.

A possible atmospheric source of  $\text{HNO}_3$ : the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  in the presence of water monomer, water dimer, and sulfuric acid

The effect of  $\text{H}_2\text{O}$ ,  $(\text{H}_2\text{O})_2$  and  $\text{H}_2\text{SO}_4$  on the ammonolysis of  $t\text{-N}_2\text{O}_4$  to form  $\text{HNO}_3$  was studied by a quantum chemical method and Master equation rate calculations. Results reveal that the ammonolysis of  $t\text{-N}_2\text{O}_4$  with  $(\text{H}_2\text{O})_2$  and  $\text{H}_2\text{SO}_4$  are barrierless or nearly barrierless reactions. Considering the effective rate constant,  $(\text{H}_2\text{O})_2$  outperforms the other catalysts in the range of 280-320 K (0 km). Moreover, the effect of  $\text{H}_2\text{SO}_4$  is obvious at higher altitudes of 5-30 km. In general, this work will give new insights into how the neutral and acidic catalysts affect the formation of  $\text{HNO}_3$ .

As featured in:



See Tianlei Zhang *et al.*, *Environ. Sci.: Atmos.*, 2023, 3, 1407.



Cite this: *Environ. Sci.: Atmos.*, 2023, 3, 1407

## A possible atmospheric source of HNO<sub>3</sub>: the ammonolysis reaction of *t*-N<sub>2</sub>O<sub>4</sub> in the presence of water monomer, water dimer, and sulfuric acid†

Ruxue Mu,‡<sup>a</sup> Weixin Zhou,‡<sup>a</sup> Zhaozhao Hong,<sup>a</sup> Rui Wang,<sup>a</sup> Quan Liu,<sup>a</sup> Qiang Zhang,<sup>a</sup> Min Jiang,<sup>a</sup> Balaganesh Muthiah<sup>b</sup> and Tianlei Zhang<sup>a\*</sup>

Although the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> is one of the potential sources of HNO<sub>3</sub> formation, the available studies have only focused on its naked reaction. Herein, the effect of important neutral and acidic trace gases, water monomer, water dimer, and sulfuric acid, on the formation of HNO<sub>3</sub> from the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> was studied by the quantum chemical method of CCSD(T)/aug-cc-pVTZ//B3LYP-D3/6-311++G(3df,2pd) and the Master equation/Rice–Ramsperger–Kassel–Marcus (ME/RRKM) rate calculations. The quantum chemical results revealed that the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> with (H<sub>2</sub>O)<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub> are barrierless or nearly barrierless reactions, potentially lowering the energy barrier to 3.4–4.1 kcal mol<sup>-1</sup>. The calculated effective rate constant illustrates that (H<sub>2</sub>O)<sub>2</sub> (100% RH) dominates over H<sub>2</sub>O and H<sub>2</sub>SO<sub>4</sub> within the range of 280–320 K (0 km), with an effective rate constant that is 1–3 orders of larger magnitude, whereas H<sub>2</sub>SO<sub>4</sub> (10<sup>8</sup> mol cm<sup>-3</sup>) is the most favorable catalyst within the troposphere between 5 and 30 km. However, the contributions of H<sub>2</sub>O, (H<sub>2</sub>O)<sub>2</sub>, and H<sub>2</sub>SO<sub>4</sub> are not apparent in the gas-phase ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> within the range of 213–320 K and 0–30 km because their effective rate constants were at least 4 orders of magnitude lower than the corresponding rate constant of the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub>. In general, the current findings shed fresh light on neutral (H<sub>2</sub>O and (H<sub>2</sub>O)<sub>2</sub>) and acidic (H<sub>2</sub>SO<sub>4</sub>) catalysts that not only affect energy barriers but also have an impact on the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> in neutral and acidic conditions.

Received 24th June 2023  
Accepted 3rd August 2023

DOI: 10.1039/d3ea00095h

rsc.li/esatmospheres

### Environmental significance

Nitrogen tetroxide (N<sub>2</sub>O<sub>4</sub>) is considered to be a dimer of nitrogen dioxide (NO<sub>2</sub>) and plays an important role in the formation of acid rain. The fact is that the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> is one of the potential sources of HNO<sub>3</sub> formation; thus, the effect of water monomer, water dimer, and sulfuric acid on the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> was studied by quantum chemical method and Master equation rate calculations. The quantum chemical results reveal that the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> with (H<sub>2</sub>O)<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub> are barrierless or nearly barrierless reactions. In terms of the effective rate constant, (H<sub>2</sub>O)<sub>2</sub> outperforms the other catalysts in the temperature range 280–320 K (0 km). Moreover, the effect of H<sub>2</sub>SO<sub>4</sub> on the ammonolysis reaction of *t*-N<sub>2</sub>O<sub>4</sub> is obvious at higher altitudes of 5–30 km. In general, this work will give a new insights into how the neutral and acidic catalysts affect the formation of HNO<sub>3</sub>.

## 1. Introduction

Nitrogen dioxide (NO<sub>2</sub>),<sup>1</sup> as one of the most significant NO<sub>x</sub>, is not only a precursor to the photochemical formation of ozone in the troposphere,<sup>2</sup> but it can also contribute to the formation of photochemical smog and cause significant health and

environmental hazards.<sup>3</sup> Nitrogen tetroxide (N<sub>2</sub>O<sub>4</sub>), a dimer of nitrogen dioxide (NO<sub>2</sub>),<sup>4</sup> can be used for nitration, nitrosation, and oxidation.<sup>5</sup> In addition, N<sub>2</sub>O<sub>4</sub> is an oxidizing agent for auto-igniting fuels and plays an important role in the formation of acid rain.<sup>6</sup> Due to the fact that N<sub>2</sub>O<sub>4</sub> is a highly toxic chemical species that hinders experimental studies,<sup>7</sup> quantum calculations are a trend in current research to probe the N<sub>2</sub>O<sub>4</sub>-related reaction mechanisms. As the most major loss route of N<sub>2</sub>O<sub>4</sub> in the atmosphere, the hydrolysis of N<sub>2</sub>O<sub>4</sub> is potentially important in the lower atmosphere and plays an important role in the formation of HONO, a major source of OH pollution in the urban atmosphere.<sup>8,9</sup>

Several investigations have shown that the less stable *t*-N<sub>2</sub>O<sub>4</sub> (*trans*-N<sub>2</sub>O<sub>4</sub>) is substantially more reactive than *s*-ONONO<sub>2</sub> (symmetric N<sub>2</sub>O<sub>4</sub>); hence, it was selected as a starting point for

<sup>a</sup>Institute of Theoretical and Computational Chemistry, Shaanxi Key Laboratory of Catalysis, School of Chemical & Environment Science, Shaanxi University of Technology, Hanzhong, Shaanxi 723000, P. R. China. E-mail: ztianlei88@163.com; Fax: +86-916-2641083; Tel: +86-916-2641083

<sup>b</sup>Department of Chemical Engineering, National Taiwan University, Taipei 106, Taiwan



† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3ea00095h>

‡ Ruxue Mu and Weixin Zhou have contributed equally to this work.



studying the hydrolysis of NO<sub>2</sub> dimers.<sup>9–12</sup> The reaction barrier for the hydrolysis of *t*-N<sub>2</sub>O<sub>4</sub> to form HONO was in the range of 10.8–11.9 kcal mol<sup>-1</sup> at different theoretical levels.<sup>9,11,12</sup> Several groups reported that using H<sub>2</sub>O and (H<sub>2</sub>O)<sub>2</sub> as catalysts stabilized the reactant complexes by 3.5–6.9 kcal mol<sup>-1</sup> and reduced the energy barrier to 6.2 kcal mol<sup>-1</sup>. Consequently, Zhang *et al.*<sup>9</sup> revealed that, for the hydrolysis of *t*-N<sub>2</sub>O<sub>4</sub>, the H<sub>2</sub>SO<sub>4</sub> catalyst is more effective than H<sub>2</sub>O and (H<sub>2</sub>O)<sub>2</sub> catalysts, resulting in not only a higher binding energy of 15.0 kcal mol<sup>-1</sup> for the reactant complex but also a lower energy barrier of 3.8 kcal mol<sup>-1</sup>.

In the atmosphere, the hydrolysis of *t*-N<sub>2</sub>O<sub>4</sub> to produce HONO is the most major loss route of *t*-N<sub>2</sub>O<sub>4</sub>.<sup>10,11</sup> As a complement to the loss of *t*-N<sub>2</sub>O<sub>4</sub>, the ammonolysis reaction of *t*-N<sub>2</sub>O<sub>4</sub> can form HNO<sub>3</sub>,<sup>13</sup> which could be competitive with the main source of HNO<sub>3</sub>, the gas-phase reaction of NO<sub>2</sub> with the hydroxyl (OH) radical<sup>18,14</sup> during the day and the hydrolysis reaction of N<sub>2</sub>O<sub>5</sub><sup>15–17</sup> at night, in polluted areas of NH<sub>3</sub>. The ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> (shown in eqn (1) and (2)) investigated by Lin *et al.*<sup>13</sup> reveals that the energy barrier of the *t*-N<sub>2</sub>O<sub>4</sub> + NH<sub>3</sub> reaction was determined to be 5.3 kcal mol<sup>-1</sup> and the corresponding rate constant at low temperature was not pressure-dependent. As far as we know, the effect of neutral and acidic gases on the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub>, which plays a significant catalytic role in hydrogen abstraction reactions,<sup>9,15,18–37</sup> has not been explored.



As in the previous studies on N<sub>2</sub>O<sub>4</sub> + H<sub>2</sub>O,<sup>11,12</sup> HO<sub>2</sub> + NO<sub>2</sub>,<sup>38</sup> H<sub>2</sub>CO + NH<sub>3</sub>,<sup>39</sup> and SO<sub>2</sub> + NO<sub>2</sub><sup>40</sup> reactions, water molecules were found to play an essential role in enhancing the stability of pre-reactive complexes and lowering the apparent activation energies of the transition states. In addition to water monomer, some recent works addressed the potential role of water dimer,<sup>18,41</sup> which may play a significant catalytic role in hydrogen abstraction reactions because its concentrations can reach  $9 \times 10^{14}$  mol cm<sup>-3</sup>.<sup>42,43</sup> Aside from water monomer and water dimer, acidic<sup>9,15,30–37</sup> gas species in the atmosphere may also be effective in lowering the energy barriers for hydrogen transfer reactions<sup>30–34</sup> and atmospheric hydrolysis reactions<sup>9,15,35–37</sup> in the gas phase. The presence of H<sub>2</sub>SO<sub>4</sub><sup>37,44</sup> in the atmosphere was considered to be a more effective catalyst than neutral catalysts,<sup>25,28–30,33,45</sup> which not only greatly reduces the energy barriers<sup>27,28,33</sup> but also facilitates the transfer of hydrogen,<sup>30–33,46</sup> and thus, H<sub>2</sub>SO<sub>4</sub> was regarded as either good acceptors or good donors of H in the catalytic gas reactions. These situations stimulated our interest in studying the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> by neutral (H<sub>2</sub>O and (H<sub>2</sub>O)<sub>2</sub>) and acidic (H<sub>2</sub>SO<sub>4</sub>) gases.

In this work, using global minimum searching combined with quantum chemical methods, we first obtained the stable structures of the reactant complexes of *t*-N<sub>2</sub>O<sub>4</sub>⋯NH<sub>3</sub>⋯X (X = H<sub>2</sub>O, (H<sub>2</sub>O)<sub>2</sub>, and H<sub>2</sub>SO<sub>4</sub>). The ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> in the presence of X was then studied using the stable molecular clusters *t*-N<sub>2</sub>O<sub>4</sub>⋯NH<sub>3</sub>⋯X. Finally, the effective rate constant for

the ammonolysis of *t*-N<sub>2</sub>O<sub>4</sub> with X was estimated at temperatures ranging from 213 to 320 K and altitudes ranging from 0 to 30 km.

## 2. Calculation details

### 2.1 Electronic structure calculations

The molecular geometries of the isolated reactants, pre-reactive complexes, transition states, post-reactive complexes, and products of the ammonolysis reaction of *t*-N<sub>2</sub>O<sub>4</sub> without and with X were optimized using the B3LYP-D3 method<sup>47–53</sup> with 6-311++G(3df,2pd) basis set in Gaussian 09 suites.<sup>54</sup> The D3 method has been reliably utilized to describe the noncovalent interaction and the equilibrium structure of atmospheric clusters and reactions.<sup>51,55</sup> Notably, the keyword “stable = opt” is added in the calculations at the B3LYP-D3/6-311++G(3df,2pd) level to ensure that all the geometries are stable. Frequency calculations were calculated at the same level for all stationary points to check that all transition states have the same character as a first-order saddle point with a single imaginary frequency and that other stationary points correspond to minima on potential energy surfaces (PESs). The scaling factor employed to adjust the ZPEs was 0.969.<sup>56,57</sup> To ensure that the optimized transition state connects the desired pre- and post-reactive complexes, intrinsic reaction coordinate (IRC) calculations<sup>58–60</sup> were performed at the same level. To improve the accuracy of the relative energies, single-point energy calculation was performed at the CCSD(T)/aug-cc-pVTZ<sup>61–63</sup> level by using the Gaussian 09 software.<sup>54</sup> It was noted that the *T*<sub>1</sub> diagnostic values for closed-shell in Table S3† were 0.02 less than the standard value,<sup>64,65</sup> showing the multireference calculations for recovering non-dynamical correlation were not a problem, and the single reference method of CCSD(T)/aug-cc-pVTZ is reliable to single point energy calculation.

Global minimum searching combined with quantum chemical methods was employed to obtain the most stable structures of the reactant complexes of *t*-N<sub>2</sub>O<sub>4</sub>⋯NH<sub>3</sub>⋯X. Initially, 500 structures with low energies were auto-produced by ABCluster software<sup>66,67</sup> with TIP4P<sup>68,69</sup> model for H<sub>2</sub>O, (H<sub>2</sub>O)<sub>2</sub>, and CHARMM<sup>70</sup> force field for *t*-N<sub>2</sub>O<sub>4</sub>, NH<sub>3</sub>, and H<sub>2</sub>SO<sub>4</sub>. Then, pre-optimized by the semi-empirical method of PM7<sup>71</sup> in MOPAC 2016.<sup>72</sup> Next, the structures with the N(*t*-N<sub>2</sub>O<sub>4</sub>)⋯N(NH<sub>3</sub>) interaction of electron donor–acceptor (EDA) and facilitating the transfer of hydrogen atom from *t*-N<sub>2</sub>O<sub>4</sub> to NH<sub>3</sub> were selected to optimize at the B3LYP-D3/6-311+G(d,p) level. Subsequently, 50 isomers with an order of electronic energies were chosen to optimize at the level of B3LYP-D3/6-311+G(2d,2p). Finally, the global minimum isomers within 5.0 kcal mol<sup>-1</sup> (the electric energy) were re-optimized at the B3LYP-D3/6-311++G(3df,2pd) level.

### 2.2 Rate constant calculations

The rate constants for the ammonolysis reaction of *t*-N<sub>2</sub>O<sub>4</sub> without and with X were calculated in two steps. First, the high-pressure-limit (HPL) rate constants were calculated by using the VRC-VTST calculations in Polyrate 2017 software.<sup>73</sup> The details





of VRC-VTST can be seen in Table S7 of the ESI.† Then, based on the HPL rate constants, the rate constants for the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  without and with X at different temperatures and pressures were calculated using the Master Equation Solver for Multi-Energy Well Reactions (MESMER) program.<sup>74</sup> The Inverse Laplace Transform (ILT) approach was used to analyze the barrierless step from distinct reactants to the pre-reactive complex.<sup>75,76</sup> Meanwhile, the transition step from the pre-reactive complex to the post-reactive complex occurring through the transition state was applied to the RRKM theory.<sup>77,78</sup> ILT methods and RRKM theory can be, respectively, expressed in eqn (3) and (4).

$$k(E) = \frac{W(E - E_0)}{h\rho(E)} \quad (3)$$

$$k^\infty(\beta) = \frac{1}{Q(\beta)} \int_0^\infty k(E)\rho(E)\exp(-\beta E)dE \quad (4)$$

In eqn (3) and (4), where  $W(E - E_0)$  is the rovibrational sum of states (SOS) at the optimized transition state (TS) geometry,  $E_0$  is the reaction threshold energy,  $h$  is Planck's constant,  $\rho(E)$  is the density of rovibrational states of the reactant, and  $Q(\beta)$  is the corresponding canonical partition function. Moreover, the electronic geometries, vibrational frequencies, and rotational constants were calculated at the B3LYP-D3/6-311++G(3df,2pd) level, and single-point energy calculations were refined at the CCSD(T)/aug-cc-pVTZ level for the modeling. The one-dimensional asymmetric Eckart potential was used to treat the tunneling effect in the RRKM calculation. In addition, the Lennard-Jones (L-J) parameters  $\varepsilon/k_B = 71.4$  K and  $\sigma = 3.798$  Å were used for  $\text{N}_2$ ,<sup>79</sup>  $\varepsilon/k_B = 200.0$  K and  $\sigma = 3.50$  Å were used for  $t\text{-N}_2\text{O}_4$ ,<sup>80</sup> while  $\varepsilon/k_B = 481.0$  K and  $\sigma = 2.92$  Å were estimated for  $\text{NH}_3$ .<sup>80</sup>

### 3. Results and discussions

The pre-reactive complex in each reaction channel was denoted by “IM” followed by a number, whereas the transition state and post-reactive complexes were denoted by “TS” and “IMF”, respectively. Species in the presence of  $\text{H}_2\text{O}$ ,  $(\text{H}_2\text{O})_2$ , and  $\text{H}_2\text{SO}_4$  were denoted by the suffixes “WM”, “WD”, and “SA”.

#### 3.1 Mechanism and rate constants for the ammonolysis reaction of $t\text{-N}_2\text{O}_4$

The ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  has been extensively investigated from the theoretical viewpoint.<sup>13</sup> Here, this reaction has been reinvestigated at the CCSD(T)/aug-cc-pVTZ//B3LYP-D3/6-311++G(3df,2pd) level to check the catalytic effect of X. Our results shown in Fig. 1 were found to be very mechanistically and energetically similar to the work reported by Lin *et al.*<sup>13</sup> All the relative energy values qualitatively matched (see Table S2†). As seen in Fig. 1, the  $t\text{-N}_2\text{O}_4 + \text{NH}_3$  reaction occurred through a ring formation mechanism, resulting in the formation of a six-membered ring complex  $t\text{-N}_2\text{O}_4 \cdots \text{NH}_3$  with a binding energy of  $0.5 \text{ kcal mol}^{-1}$ . Then, the terminal O1 atom of  $t\text{-N}_2\text{O}_4$  abstracts the H atom of  $\text{NH}_3$  along with the N-N bond formation to form the product complex  $\text{HNO}_3 \cdots \text{NH}_2\text{NO}$ . From an energy point of view, the barrier height of the  $t\text{-N}_2\text{O}_4 + \text{NH}_3$  reaction was  $3.7 \text{ kcal mol}^{-1}$ , revealing that  $t\text{-N}_2\text{O}_4$  can easily react with  $\text{NH}_3$  in the gas phase.

The calculated rate constants for the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  are listed in Table 1. In the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$ , the hindered internal rotation (HIR)<sup>77,81–86</sup> correction at 760 Torr has a moderate effect, increasing the rate constants by a factor of 1.25 to 1.28. Meanwhile, the almost unchanged rate constants for the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  at different atmospheric pressures revealed that the pressure (10–760 Torr)<sup>87</sup> has little effect on the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  within the temperature range of 280–320 K.<sup>88,89</sup>

#### 3.2 Mechanism and rate constants for the ammonolysis reaction of $t\text{-N}_2\text{O}_4$ with $\text{H}_2\text{O}$ and $(\text{H}_2\text{O})_2$

Fig. 2a and b show the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  assisted by  $\text{H}_2\text{O}$  (Channel WM) and  $(\text{H}_2\text{O})_2$  (Channel WD), where both  $\text{H}_2\text{O}$  and  $(\text{H}_2\text{O})_2$  served as a “bridge” to promote hydrogen atom transfer from the N3 atom of  $\text{NH}_3$  to the terminal O1 atom of  $t\text{-N}_2\text{O}_4$ . In the case of Channel WM, the reaction can occur either (a) between  $\text{NH}_3$  and monohydrated  $t\text{-N}_2\text{O}_4$  ( $t\text{-N}_2\text{O}_4 \cdots \text{H}_2\text{O}$ ) or (b) between hydrated  $\text{NH}_3$  ( $\text{NH}_3 \cdots \text{H}_2\text{O}$ ) and  $t\text{-N}_2\text{O}_4$ . The binding energy of  $t\text{-N}_2\text{O}_4 \cdots \text{H}_2\text{O}$  was  $2.9 \text{ kcal mol}^{-1}$ , which was consistent with the previously calculated value of  $3.5 \text{ kcal mol}^{-1}$  at the CCSD(T)-F12a/cc-pVDZ-F12//M06-2X/6-311++G(3df,2pd) level.<sup>9</sup>





Fig. 1 Potential energy profiles for the  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \rightarrow \text{HNO}_3 + \text{NH}_2\text{NO}$  reaction at the CCSD(T)/aug-cc-pVTZ//B3LYP-D3/6-311++G(3df,2pd) level.



**Table 1** The rate constants ( $k_R$ ) ( $\text{cm}^3 \text{mol}^{-1} \text{s}^{-1}$ ) for the  $t\text{-N}_2\text{O}_4 + \text{NH}_3$  reaction without and with HIR treatments within the temperature range of 280–320 K and pressure range of 10–760 Torr

$T$ (K)	HIR impact <sup>a</sup>			Pressure impact				
	With HIR treatments	Without HIR treatments	Factor <sup>b</sup>	10 Torr	50 Torr	100 Torr	300 Torr	760 Torr
280	$6.67 \times 10^{-17}$	$5.20 \times 10^{-17}$	1.28	$5.03 \times 10^{-17}$	$5.14 \times 10^{-17}$	$5.16 \times 10^{-17}$	$5.18 \times 10^{-17}$	$5.20 \times 10^{-17}$
290	$7.81 \times 10^{-17}$	$6.13 \times 10^{-17}$	1.27	$5.90 \times 10^{-17}$	$6.07 \times 10^{-17}$	$6.10 \times 10^{-17}$	$6.12 \times 10^{-17}$	$6.13 \times 10^{-17}$
298	$8.80 \times 10^{-17}$	$6.95 \times 10^{-17}$	1.27	$6.65 \times 10^{-17}$	$6.88 \times 10^{-17}$	$6.92 \times 10^{-17}$	$6.94 \times 10^{-17}$	$6.95 \times 10^{-17}$
300	$9.06 \times 10^{-17}$	$7.17 \times 10^{-17}$	1.26	$6.84 \times 10^{-17}$	$7.09 \times 10^{-17}$	$7.13 \times 10^{-17}$	$7.16 \times 10^{-17}$	$7.17 \times 10^{-17}$
310	$1.04 \times 10^{-16}$	$8.32 \times 10^{-17}$	1.26	$7.84 \times 10^{-17}$	$8.20 \times 10^{-17}$	$8.26 \times 10^{-17}$	$8.30 \times 10^{-17}$	$8.32 \times 10^{-17}$
320	$1.19 \times 10^{-16}$	$9.57 \times 10^{-17}$	1.25	$9.40 \times 10^{-17}$	$9.40 \times 10^{-17}$	$9.55 \times 10^{-17}$	$9.55 \times 10^{-17}$	$9.57 \times 10^{-17}$

<sup>a</sup> HIR impact represents the hindered internal rotations treatment. <sup>b</sup> Factor denotes the rate ratio between with HIR treatments and without HIR treatments.



**Fig. 2** Potential energy profiles for the  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \rightarrow \text{HNO}_3 + \text{NH}_2\text{NO}$  reaction catalyzed by  $\text{H}_2\text{O}$  and  $(\text{H}_2\text{O})_2$  at the CCSD(T)/aug-cc-pVTZ//B3LYP-D3/6-311++G(3df,2pd) level (a–c) denotes the values respectively reported from ref. 9, 90, and 91.

The binding energy of  $\text{NH}_3 \cdots \text{H}_2\text{O}$  was  $4.4 \text{ kcal mol}^{-1}$ , which agreed well with the calculated values of  $4.4\text{--}4.6 \text{ kcal mol}^{-1}$ .<sup>90,91</sup> The stability of  $\text{NH}_3 \cdots \text{H}_2\text{O}$  was  $1.5 \text{ kcal mol}^{-1}$  higher than that of  $t\text{-N}_2\text{O}_4 \cdots \text{H}_2\text{O}$ . So, the ammonolysis of  $t\text{-N}_2\text{O}_4$  in the presence of  $\text{H}_2\text{O}$  mainly takes place *via* the collision of  $\text{NH}_3 \cdots \text{H}_2\text{O}$  with  $t\text{-N}_2\text{O}_4$  to form the quasi-planar eight-membered ring reactant complex IM\_WM1. The energy of the reactant complex IM\_WM1 was  $4.7 \text{ kcal mol}^{-1}$  lower than that of the isolated reactants  $t\text{-N}_2\text{O}_4 + \text{H}_2\text{O} + \text{NH}_3$ . In the complex IM\_WM1,  $\text{H}_2\text{O}$  played the roles of a single acceptor and donor of hydrogen

bonds. After the formation of the complex IM\_WM1, the reaction proceeded to form a hydrogen-bonded complex,  $\text{HNO}_3 \cdots \text{H}_2\text{O} \cdots \text{NH}_2\text{NO}$  (denoted IMF\_WM), through the transition state TS\_WM, with an energy barrier of  $4.2 \text{ kcal mol}^{-1}$ .

The reaction  $t\text{-N}_2\text{O}_4 + \text{NH}_3 + (\text{H}_2\text{O})_2$  can be initiated by the reactant complex IM\_WD1, which can be formed from  $t\text{-N}_2\text{O}_4 \cdots (\text{H}_2\text{O})_2 + \text{NH}_3$  or  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots (\text{H}_2\text{O})_2$ . It is clear from Fig. 2b that  $\text{NH}_3$  was most likely bound to  $(\text{H}_2\text{O})_2$  prior to  $t\text{-N}_2\text{O}_4$ . The reactant complex IM\_WD1 has a quasi-planar structure similar to that of complex IM\_WM1 and can be regarded as  $\text{H}_2\text{O}$  in



IM\_WM1, which was replaced by  $(\text{H}_2\text{O})_2$ . The binding energy of IM\_WD1 was  $1.7 \text{ kcal mol}^{-1}$  from  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots (\text{H}_2\text{O})_2$ . After complex IM\_WD1, the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  with  $(\text{H}_2\text{O})_2$  can form the product complex  $\text{HNO}_3 \cdots (\text{H}_2\text{O})_2 \cdots \text{NH}_2\text{NO}$  (labeled as IMF\_WD) through the transition state TS\_WD with an energy barrier of  $0.3 \text{ kcal mol}^{-1}$ . Three hydrogen atom transfer mechanisms occurred at TS\_WD, as well as the simultaneous formation of the N(2)–N(3) bond. In comparison to  $\text{H}_2\text{O}$  in Channel WM,  $(\text{H}_2\text{O})_2$  in Channel WD played a more obvious catalytic role in promoting the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$ . When  $(\text{H}_2\text{O})_2$  was used as a catalyst in Channel WD, it stabilized the reactant complex by a further  $4.9 \text{ kcal mol}^{-1}$  and decreased the reaction barrier by  $3.9 \text{ kcal mol}^{-1}$ . The more pronounced catalytic effect of  $(\text{H}_2\text{O})_2$  could be attributed to two factors. On the one hand,  $(\text{H}_2\text{O})_2$  may improve the  $\text{N}(t\text{-N}_2\text{O}_4) \cdots \text{N}(\text{NH}_3)$  interaction compared to  $\text{H}_2\text{O}$ . For example, in the reactant complex IM\_WD1, the strengthening of the  $\text{N}(t\text{-N}_2\text{O}_4) \cdots \text{N}(\text{NH}_3)$  interaction was shown by the shortening of the bond distance  $\text{N}(2) \cdots \text{N}(3)$  ( $2.18 \text{ \AA}$ , shown in Fig. 2b), which is less than the corresponding value in the reactant complex IM\_WM1 ( $2.21 \text{ \AA}$ , shown in Fig. 2a). On the other hand, when  $\text{H}_2\text{O}$  was replaced by  $(\text{H}_2\text{O})_2$ , however, the transition state extended from an eight-member ring (TS\_WM) to a ten-member ring (TS\_WD). This structural change reduces the ring tension of the transition state to a certain extent, lowering the reaction energy barrier.

As shown in Table 2, within the range of 213–320 K and 0–30 km,<sup>88,89</sup> the rate constant for the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  assisted by  $\text{H}_2\text{O}$  ( $k_{\text{WM}}$ ) was predicted to be  $3.94 \times 10^{-20}$  to  $1.93 \times 10^{-19} \text{ cm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ , which was 2–3 orders of magnitude lower than that of the naked ammonolysis reaction of  $t\text{-N}_2\text{O}_4$ . The calculated rate constant for the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  assisted by  $(\text{H}_2\text{O})_2$  ( $k_{\text{WD}}$ ) was  $3.66 \times 10^{-16}$  to  $1.98 \times 10^{-15} \text{ cm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ , which was 1–2 orders of magnitude greater than the naked ammonolysis reaction of  $t\text{-N}_2\text{O}_4$ . The ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  assisted by  $(\text{H}_2\text{O})_2$  was more kinetically favorable than the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  with  $\text{H}_2\text{O}$ , with a rate constant that was 3–5 orders of magnitude greater.

### 3.3 Mechanism and rate constants for the ammonolysis reaction of $t\text{-N}_2\text{O}_4$ with $\text{H}_2\text{SO}_4$

As shown in Fig. S3,<sup>†</sup> nine geometrical isomers of the reactant complex  $t\text{-N}_2\text{O}_4 \cdots \text{NH}_3 \cdots \text{H}_2\text{SO}_4$  (labeled as IM\_SAn,  $n = 1-9$ ) were found at the B3LYP-D3/6-311++G(3df,2pd) level, with complex IM\_SA1 being the most stable. Based on complex IM\_SA1, Fig. 3 presents the potential energy surface (PES) for the ammonolysis of  $t\text{-N}_2\text{O}_4$  in the presence of  $\text{H}_2\text{SO}_4$  (Channel SA). In the case of Channel SA, the reaction can occur (a) between  $\text{NH}_3$  and  $t\text{-N}_2\text{O}_4 \cdots \text{H}_2\text{SO}_4$  or (b) between  $t\text{-N}_2\text{O}_4$  and  $\text{NH}_3 \cdots \text{H}_2\text{SO}_4$ . The binding energy of  $\text{NH}_3 \cdots \text{H}_2\text{SO}_4$  was  $14.7 \text{ kcal mol}^{-1}$ , which was in good agreement with the previously reported value.<sup>92</sup> The large binding energy of  $\text{NH}_3 \cdots \text{H}_2\text{SO}_4$  indicates that in the reaction of  $t\text{-N}_2\text{O}_4 + \text{NH}_3 + \text{H}_2\text{SO}_4$ ,  $\text{NH}_3$  is easily bound to the isolated  $\text{H}_2\text{SO}_4$ . In this sense, the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  with  $\text{H}_2\text{SO}_4$  mainly takes place *via* the collision of  $t\text{-N}_2\text{O}_4$  with  $\text{NH}_3 \cdots \text{H}_2\text{SO}_4$ , resulting in the reactant complex IM\_SA1.

The energy of the reactant complex IM\_SA1 was  $15.0 \text{ kcal mol}^{-1}$  lower than that of the separate reactants  $t\text{-N}_2\text{O}_4$ ,  $\text{NH}_3$ , and  $\text{H}_2\text{SO}_4$ . In the complex IM\_SA1,  $\text{H}_2\text{SO}_4$  served as a single donor and acceptor of hydrogen bonds to form a ring-like structure with the binary complex of  $t\text{-N}_2\text{O}_4 \cdots \text{NH}_3$ . The stability of complex IM\_SA1 increased by  $5.4-10.3 \text{ kcal mol}^{-1}$  when compared to the reactant complexes IM\_WM1 and IM\_WD1, with the distance of the  $\text{N}(t\text{-N}_2\text{O}_4) \cdots \text{N}(\text{NH}_3)$  bond reduced by  $0.01-0.04 \text{ \AA}$ . After the formation of the complex IM\_SA1, Channel SA proceeded through the transition state TS\_SA to form a ten-membered ring hydrogen-bonded complex  $\text{HNO}_3 \cdots \text{H}_2\text{SO}_4 \cdots \text{NH}_2\text{NO}$  (labeled as IMF\_SA). Similar to the transition state TS\_WM described above, TS\_SA was in the middle of a double hydrogen atom transfer process, with the  $\text{H}_2\text{SO}_4$  moiety serving as a bridge for the hydrogen transfer. From the viewpoint of the energy barrier height, Channel SA was a barrierless process. In comparison to  $\text{H}_2\text{O}$  and  $(\text{H}_2\text{O})_2$ ,  $\text{H}_2\text{SO}_4$  could lower the energy barrier, at least by  $0.7-4.6 \text{ kcal mol}^{-1}$ . Complex IMF\_SA showed a ten-membered ring structure. It had a binding energy of  $28.0 \text{ kcal mol}^{-1}$  to the separate reactants  $t\text{-N}_2\text{O}_4$ ,  $\text{NH}_3$ , and  $\text{H}_2\text{SO}_4$ , which was  $13.0 \text{ kcal mol}^{-1}$  lower than the

**Table 2** Calculated rate constants ( $k$ ,  $\text{cm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ ) for the  $t\text{-N}_2\text{O}_4 + \text{NH}_3$  reaction with  $\text{H}_2\text{O}$ ,  $(\text{H}_2\text{O})_2$ , and  $\text{H}_2\text{SO}_4$  calculated by master equation within the range of 213–320 K and 0–30 km<sup>a,b</sup>

Altitude	$T$ (K)	$k_{\text{R}}$	$k_{\text{WM}}$	$k_{\text{WD}}$	$k_{\text{SA}}$
0 km	280	$5.20 \times 10^{-17}$	$1.16 \times 10^{-19}$	$6.05 \times 10^{-16}$	$1.49 \times 10^{-16}$
	290	$6.13 \times 10^{-17}$	$1.33 \times 10^{-19}$	$5.29 \times 10^{-16}$	$1.50 \times 10^{-16}$
	298	$6.95 \times 10^{-17}$	$1.48 \times 10^{-19}$	$4.76 \times 10^{-16}$	$1.50 \times 10^{-16}$
	300	$7.17 \times 10^{-17}$	$1.52 \times 10^{-19}$	$4.64 \times 10^{-16}$	$1.51 \times 10^{-16}$
	310	$8.32 \times 10^{-17}$	$1.72 \times 10^{-19}$	$4.11 \times 10^{-16}$	$1.52 \times 10^{-16}$
	320	$9.57 \times 10^{-17}$	$1.93 \times 10^{-19}$	$3.66 \times 10^{-16}$	$1.54 \times 10^{-16}$
5 km	259.3	$3.59 \times 10^{-17}$	$8.59 \times 10^{-20}$	$8.30 \times 10^{-16}$	$1.48 \times 10^{-16}$
10 km	229.7	$1.92 \times 10^{-17}$	$5.32 \times 10^{-20}$	$1.40 \times 10^{-15}$	$1.51 \times 10^{-16}$
15 km	212.6	$1.24 \times 10^{-17}$	$3.94 \times 10^{-20}$	$1.98 \times 10^{-15}$	$1.56 \times 10^{-16}$
20 km	215.5	$1.28 \times 10^{-17}$	$4.15 \times 10^{-20}$	$1.86 \times 10^{-15}$	$1.55 \times 10^{-16}$
25 km	218.6	$1.37 \times 10^{-17}$	$4.38 \times 10^{-20}$	$1.74 \times 10^{-15}$	$1.54 \times 10^{-16}$
30 km	223.7	$1.55 \times 10^{-17}$	$4.79 \times 10^{-20}$	$1.57 \times 10^{-15}$	$1.53 \times 10^{-16}$

<sup>a</sup>  $k_{\text{R}}$ ,  $k_{\text{WM}}$ ,  $k_{\text{WD}}$ , and  $k_{\text{SA}}$  were respectively denoted the rate constants for the  $t\text{-N}_2\text{O}_4 + \text{NH}_3$ ,  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots \text{H}_2\text{O}$ ,  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots (\text{H}_2\text{O})_2$ , and  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots \text{H}_2\text{SO}_4$  reactions. <sup>b</sup> The 0–30 km data were reported from ref. 88 and 89.






Fig. 3 Potential energy profiles for the  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \rightarrow \text{HNO}_3 + \text{NH}_2\text{NO}$  reaction catalyzed by  $\text{H}_2\text{SO}_4$  at the CCSD(T)/aug-cc-pVTZ//B3LYP-D3/6-311+G(3df,2pd) level (a) The values were reported from ref. 92.

reactant complex IM\_SA1. The rate constant of the  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots \text{H}_2\text{SO}_4$  reaction is one order of magnitude more than that of the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  without a catalyst, as shown in Table 2. This result revealed that  $\text{H}_2\text{SO}_4$  in the  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots \text{H}_2\text{SO}_4$  reaction plays a favorable catalytic role in promoting the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$ . Thus, the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  assisted by  $(\text{H}_2\text{O})_2$  and  $\text{H}_2\text{SO}_4$  was more kinetically favorable than the reaction with  $\text{H}_2\text{O}$ .

### 3.4 Kinetics and implication in atmospheric chemistry

According to previous reports,<sup>93–96</sup> it is clear that the rate constant listed in Table 2 is insufficient to predict the atmospheric importance of the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  assisted by  $(\text{H}_2\text{O})_2$  and  $\text{H}_2\text{SO}_4$ . To understand the atmospheric effect of  $(\text{H}_2\text{O})_2$  and  $\text{H}_2\text{SO}_4$  on the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$ , we introduced the effective rate constants ( $k'$ ) to calculate the relative efficiency of neutral and acid trace gases affecting the atmospheric reaction<sup>97–100</sup> and to compare the rate constant for the naked reaction.

On the basis of the rate constant for Channels WM, WD, and SA, the equilibrium constant for the bimolecular formation ( $\text{NH}_3 \cdots \text{H}_2\text{O}$ ,  $\text{NH}_3 \cdots (\text{H}_2\text{O})_2$ , and  $\text{NH}_3 \cdots \text{H}_2\text{SO}_4$ ) and the

concentrations of  $\text{H}_2\text{O}$ ,  $(\text{H}_2\text{O})_2$ , and  $\text{H}_2\text{SO}_4$  are listed in Tables S4–6,<sup>†</sup> as stated in eqn (5)–(7).

$$k'_{\text{WM}} = k_{\text{WM}} \times K_{\text{eq}}(\text{NH}_3 \cdots \text{H}_2\text{O}) \times [\text{H}_2\text{O}] \quad (5)$$

$$k'_{\text{WD}} = k_{\text{WD}} \times K_{\text{eq}}(\text{NH}_3 \cdots (\text{H}_2\text{O})_2) \times [(\text{H}_2\text{O})_2] \quad (6)$$

$$k'_{\text{SA}} = k_{\text{SA}} \times K_{\text{eq}}(\text{NH}_3 \cdots \text{H}_2\text{SO}_4) \times [\text{H}_2\text{SO}_4] \quad (7)$$

where  $k_{\text{WM}}$ ,  $k_{\text{WD}}$ , and  $k_{\text{SA}}$  are the bimolecular rate constants for Channel WM, Channels WD, and SA, respectively;  $K_{\text{eq}}(\text{NH}_3 \cdots \text{H}_2\text{O})$ ,  $K_{\text{eq}}(\text{NH}_3 \cdots (\text{H}_2\text{O})_2)$ , and  $K_{\text{eq}}(\text{NH}_3 \cdots \text{H}_2\text{SO}_4)$  are the equilibrium constants for the formation of complexes  $\text{NH}_3 \cdots \text{H}_2\text{O}$ ,  $\text{NH}_3 \cdots (\text{H}_2\text{O})_2$ , and  $\text{NH}_3 \cdots \text{H}_2\text{SO}_4$ .  $[\text{H}_2\text{O}]$ ,  $[(\text{H}_2\text{O})_2]$ , and  $[\text{H}_2\text{SO}_4]$  represent the concentrations of  $\text{H}_2\text{O}$ ,  $(\text{H}_2\text{O})_2$ , and  $\text{H}_2\text{SO}_4$  taken from previous reports.<sup>44,87</sup> The  $k'$  for the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  with X at 0 km altitude and at different altitudes (5–30 km) in the troposphere was calculated.

**3.4.1 Zero kilometer altitude.** As seen in Table 3, the calculated value of  $k'_{\text{SA}}$  can compete with  $k'_{\text{WD}}$  at 280 K. With increasing temperature, the calculated  $k'_{\text{SA}}$  was  $\sim 1\text{--}3$  orders of magnitude smaller than the values of  $k'_{\text{WD}}$ , showing that the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  was superior to that of an acidic

Table 3 The effective rate constants ( $k'$ ,  $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ ) for the  $t\text{-N}_2\text{O}_4 + \text{NH}_3$  reaction assisted by  $\text{H}_2\text{O}$ ,  $(\text{H}_2\text{O})_2$ , and  $\text{H}_2\text{SO}_4$  within the temperature range of 280–320 K<sup>a,b</sup> (0 km)

Catalysts	T/K	280 K	290 K	298 K	300 K	310 K	320 K
$k'_{\text{WM}}$	100% RH <sup>c</sup>	$4.29 \times 10^{-22}$	$6.96 \times 10^{-22}$	$1.02 \times 10^{-21}$	$1.11 \times 10^{-21}$	$1.70 \times 10^{-21}$	$2.49 \times 10^{-21}$
$k'_{\text{WD}}$	100% RH <sup>c</sup>	$2.45 \times 10^{-21}$	$3.68 \times 10^{-21}$	$5.16 \times 10^{-21}$	$5.46 \times 10^{-21}$	$7.70 \times 10^{-21}$	$1.02 \times 10^{-20}$
$k'_{\text{SA}}$	$[\text{H}_2\text{SO}_4]^c = 10^8 \text{ mol cm}^3$	$1.39 \times 10^{-21}$	$5.49 \times 10^{-22}$	$2.72 \times 10^{-22}$	$2.32 \times 10^{-22}$	$1.03 \times 10^{-22}$	$4.71 \times 10^{-23}$

<sup>a</sup>  $k'_{\text{WM}}$ ,  $k'_{\text{WD}}$ , and  $k'_{\text{SA}}$  were respectively denoted the effective rate constants for the  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots \text{H}_2\text{O}$ ,  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots (\text{H}_2\text{O})_2$ , and  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots \text{H}_2\text{SO}_4$  reactions. <sup>b</sup> The values of temperature were reported from ref. 88 and 89. <sup>c</sup> The values of concentrations were reported from ref. 44 and 87.





**Table 4** The effective rate constants ( $k'$ ,  $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ ) for the  $t\text{-N}_2\text{O}_4 + \text{NH}_3$  reaction with  $\text{H}_2\text{O}$ ,  $(\text{H}_2\text{O})_2$ , and  $\text{H}_2\text{SO}_4$  at different altitudes in troposphere<sup>a</sup>

Altitude (km) <sup>b</sup>	$T$ (K) <sup>b</sup>	$P$ (torr) <sup>b</sup>	$k'_{\text{WM}}$	$k'_{\text{WD}}$	$k'_{\text{SA}}$	$k'_{\text{SA}}/k'_{\text{tot}}$
5	259.3	406.75	$5.62 \times 10^{-23}$	$1.50 \times 10^{-22}$	$1.81 \times 10^{-21}$	$5.05 \times 10^{-5}$
10	229.7	202.16	$2.12 \times 10^{-23}$	$1.74 \times 10^{-22}$	$9.47 \times 10^{-21}$	$4.93 \times 10^{-4}$
15	212.6	91.20	$1.38 \times 10^{-25}$	$2.95 \times 10^{-26}$	$3.74 \times 10^{-21}$	$3.02 \times 10^{-4}$
20	215.5	41.04	$6.12 \times 10^{-26}$	$4.52 \times 10^{-27}$	$4.00 \times 10^{-22}$	$3.11 \times 10^{-5}$
25	218.6	19.00	$3.05 \times 10^{-26}$	$8.56 \times 10^{-28}$	$2.71 \times 10^{-21}$	$1.97 \times 10^{-4}$
30	223.7	8.36	$1.33 \times 10^{-26}$	$1.12 \times 10^{-28}$	$7.88 \times 10^{-21}$	$5.06 \times 10^{-4}$

<sup>a</sup>  $k'_{\text{WM}}$ ,  $k'_{\text{WD}}$ , and  $k'_{\text{SA}}$  were respectively denoted the effective rate constants for the  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots \text{H}_2\text{O}$ ,  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots (\text{H}_2\text{O})_2$ , and  $t\text{-N}_2\text{O}_4 + \text{NH}_3 \cdots \text{H}_2\text{SO}_4$  reactions;  $k'_{\text{tot}} = k_{\text{R}} + k'_{\text{WD}} + k'_{\text{SA}}$ . <sup>b</sup> The values of altitude, temperature and pressure were reported from ref. 87, 88 and 89.

( $\text{H}_2\text{SO}_4$ ) catalyst in the presence of neutral ( $\text{H}_2\text{O}$  and  $(\text{H}_2\text{O})_2$ ) catalysts. However, the  $\text{H}_2\text{SO}_4$ -catalyzed reaction can be neglected because its calculated  $k'_{\text{SA}}$  was at least 4 orders of magnitude lower than the corresponding value of  $k_{\text{R}}$  in the naked reaction of the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$ . This indicated that the contributions of  $\text{H}_2\text{SO}_4$  to the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  in atmospheric chemistry are not obvious within the temperature range of 280–320 K (at 0 km altitude).

**3.4.2 Higher altitudes.** The  $k'$  for the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  with  $\text{H}_2\text{O}$ ,  $(\text{H}_2\text{O})_2$ , and  $\text{H}_2\text{SO}_4$  were calculated within the 5–30 km altitude range, and the calculated  $k'$  is listed in Table 4. It can be seen in Table 4 that the contribution of  $\text{H}_2\text{SO}_4$  was most obvious in the catalysts of  $\text{H}_2\text{O}$ ,  $(\text{H}_2\text{O})_2$ , and  $\text{H}_2\text{SO}_4$  within the altitude range of 5–30 km, since the value of  $k'_{\text{SA}}$  was larger by 1–8 orders of magnitude than that of  $k'_{\text{WD}}$ . In order to quantitatively assess the impact of  $\text{H}_2\text{SO}_4$  on the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$ , the total rate constant  $k'_{\text{tot}}$  can be calculated using eqn (8). The branching ratio for  $k'_{\text{SA}}/k'_{\text{tot}}$  in Table 4 was calculated to be  $3.11 \times 10^{-5}$ – $5.06 \times 10^{-4}$  at 5–30 km. This indicates that the contribution of the  $\text{H}_2\text{SO}_4$ -assisted ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  can be negligible in atmospheric gas-phase chemistry.

$$k'_{\text{tot}} = k_{\text{R}} + k'_{\text{WD}} + k'_{\text{SA}} \quad (8)$$

## 4. Summary and conclusions

In this work, the favorable route for the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  in the presence of neutral ( $\text{H}_2\text{O}$  and  $(\text{H}_2\text{O})_2$ ) and acidic ( $\text{H}_2\text{SO}_4$ ) catalysts was investigated using the quantum chemical method of CCSD(T)/aug-cc-pVTZ//B3LYP-D3/6-311++G(3df,2pd) and the master equation. The calculated results show that the energy barrier for the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  increased when  $\text{H}_2\text{O}$  was present, but when  $(\text{H}_2\text{O})_2$  was present, the reaction energy barrier decreased to  $0.3 \text{ kcal mol}^{-1}$ , which was  $3.4 \text{ kcal mol}^{-1}$  lower than the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  without the catalyst, especially when  $\text{H}_2\text{SO}_4$  was directly involved in the reaction that is a barrierless process. In terms of the effective rate constant,  $(\text{H}_2\text{O})_2$  outperforms the other catalysts in the temperature range of 280–320 K (0 km). Moreover, the effect of  $\text{H}_2\text{SO}_4$  on the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$  is obvious at higher altitudes of 5–30 km. Overall, this work will

give a new insight into how the neutral and acidic catalysts affect the formation of  $\text{HNO}_3$  from the ammonolysis reaction of  $t\text{-N}_2\text{O}_4$ . As  $\text{HNO}_3$  is an important source of acid rain, the present work will provide a potential formation pathway for  $\text{HNO}_3$ , which plays a crucial role in the formation of acid rain.

## Author contributions

Ruxue Mu: investigation, data curation, visualization, writing – original draft. Weixin Zhou: supervision, writing – review and editing, project administration. Zhaozhao Hong: visualization, data curation. Rui Wang: writing – review and editing, project administration. Quan Liu: visualization, data curation. Qiang Zhang: formal analysis. Min Jiang: formal analysis. Balaganesh Muthiah: supervision, writing – review and editing. Tianlei Zhang: writing – review and editing, project administration.

## Conflicts of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

This work was supported by the National Natural Science Foundation of China (no. 22073059, 22203052); the Natural Science Foundation of Shaanxi Province (no. 2023-YBGY-486, 2022JM-133); the Funds of Graduate Innovation of Shaanxi University of Technology (no. SLGYCX2304); the authors also thank Prof. Mark A. Blitz (from University of Leeds) for providing the use of the MESMER program.

## References

- M. T. Martins-Costa, J. M. Anglada, J. S. Francisco and M. F. Ruiz-López, Theoretical investigation of the photoexcited  $\text{NO}_2 + \text{H}_2\text{O}$  reaction at the air-water interface and its atmospheric implications, *Chem. – Eur. J.*, 2019, 25, 13899–13904.
- A. Bacak, M. Cooke, M. Bardwell, M. McGillen, A. Archibald, L. Huey, D. Tanner, S. Utembe, M. Jenkin and R. Derwent, Kinetics of the  $\text{HO}_2 + \text{NO}_2$  Reaction: On the impact of new





- gas-phase kinetic data for the formation of HO<sub>2</sub>NO<sub>2</sub> on HO<sub>x</sub>, NO<sub>x</sub> and HO<sub>2</sub>NO<sub>2</sub> levels in the troposphere, *Atmos. Environ.*, 2011, **45**, 6414–6422.
- 3 J. H. Seinfeld and S. N. Pandis, *Atmospheric Chemistry and Physics: from Air Pollution to Climate Change*, John Wiley & Sons, 2016.
  - 4 W.-G. Liu and W. A. Goddard III, First-principles study of the role of interconversion between NO<sub>2</sub>, N<sub>2</sub>O<sub>4</sub>, *cis*-ONO-NO<sub>2</sub>, and *trans*-ONO-NO<sub>2</sub> in chemical processes, *J. Am. Chem. Soc.*, 2012, **134**, 12970–12978.
  - 5 C. Nonnenberg, I. Frank and T. M. Klapötke, Ultrafast cold reactions in the bipropellant monomethylhydrazine/nitrogen tetroxide: CPMD simulations, *Angew. Chem., Int. Ed.*, 2004, **43**, 4586–4589.
  - 6 L. Dejun, T. Gan, Y. Zhengwei, J. Guofeng, W. ZHANG, W. Yin and W. Huali, Stress corrosion behavior of 2195-T8 Al-Li alloy with an artificial pit exposed to a 30vol% HNO<sub>3</sub> solution, *Chin. J. Aeronaut.*, 2023, **36**, 304–315.
  - 7 Y. Guo, Z. Huang, G. Tian, W. Wu, J. Lin and X. Chang, Isomerization and reaction process of N<sub>2</sub>O<sub>4</sub>(H<sub>2</sub>O)<sub>n</sub>, *RSC Adv.*, 2023, **13**, 12469–12475.
  - 8 B. J. Finlayson-Pitts and J. N. Pitts Jr, *Chemistry of the Upper and Lower Atmosphere: Theory, Experiments, and Applications*, Elsevier, 1999.
  - 9 T. Zhang, M. Wen, X. Cao, Y. Zhang, Z. Zeng, X. Guo, C. Zhao, M. Lily and R. Wang, The hydrolysis of NO<sub>2</sub> dimer in small clusters of sulfuric acid: A potential source of nitrous acid in troposphere, *Atmos. Environ.*, 2020, **243**, 117876.
  - 10 L. Li, Z. Duan, H. Li, C. Zhu, G. Henkelman, J. S. Francisco and X. C. Zeng, Formation of HONO from the NH<sub>3</sub>-promoted hydrolysis of NO<sub>2</sub> dimers in the atmosphere, *Proc. Natl. Acad. Sci.*, 2018, **115**, 7236–7241.
  - 11 R. Zhu, K.-Y. Lai and M.-C. Lin, Ab initio chemical kinetics for the hydrolysis of N<sub>2</sub>O<sub>4</sub> isomers in the gas phase, *J. Phys. Chem. A*, 2012, **116**, 4466–4472.
  - 12 R. Putikam and M. C. Lin, A novel mechanism for the isomerization of N<sub>2</sub>O<sub>4</sub> and its implication for the reaction with H<sub>2</sub>O and acid rain formation, *Int. J. Quantum Chem.*, 2018, **118**, e25560.
  - 13 H. P. Trac, T. Le Huyen and M.-C. Lin, A Computational Study on the Redox Reactions of Ammonia and Methylamine with Nitrogen Tetroxide, *J. Phys. Chem. A*, 2020, **124**, 9923–9932.
  - 14 K. Mondal, S. Biswas, A. Chattopadhyay, P. Chatterjee and T. Chakraborty, Gas-phase oxidation of NO<sub>2</sub> to HNO<sub>3</sub> by phenol: atmospheric implications, *ACS Earth Space Chem.*, 2021, **5**, 2131–2141.
  - 15 M. Wen, R. Li, T. Zhang, C. Ding, Y. Hu, R. Mu, M. Liang, T. Ou and B. Long, A potential source of tropospheric secondary organic aerosol precursors: The hydrolysis of N<sub>2</sub>O<sub>5</sub> in water dimer and small clusters of sulfuric acid, *Atmos. Environ.*, 2022, **287**, 119245.
  - 16 J. G. Calvert, A. Lazrus, G. L. Kok, B. G. Heikes, J. G. Walega, J. Lind and C. A. Cantrell, Chemical mechanisms of acid generation in the troposphere, *Nature*, 1985, **317**, 27–35.
  - 17 J. I. Steinfeld, Atmospheric chemistry and physics: from air pollution to climate change, *Environ Sci Policy*, 1998, **40**, 26.
  - 18 L. P. Viegas and A. n. J. Varandas, Role of (H<sub>2</sub>O)<sub>n</sub> (n = 2-3) Clusters on the HO<sub>2</sub> + O<sub>3</sub> Reaction: A Theoretical Study, *J. Phys. Chem. B*, 2016, **120**, 1560–1568.
  - 19 B. Long, J. L. Bao and D. G. Truhlar, Unimolecular reaction of acetone oxide and its reaction with water in the atmosphere, *Proc. Natl. Acad. Sci. U.S.A.*, 2018, **115**, 6135–6140.
  - 20 X. Chen, C. Tao, L. Zhong, Y. Gao, W. Yao and S. Li, Theoretical study on the atmospheric reaction of SO<sub>2</sub> with the HO<sub>2</sub> and HO<sub>2</sub>···H<sub>2</sub>O complex formation HSO<sub>4</sub> and H<sub>2</sub>SO<sub>3</sub>, *Chem. Phys. Lett.*, 2014, **608**, 272–276.
  - 21 R. J. Buszek, J. S. Francisco and J. M. Anglada, Water effects on atmospheric reactions, *Int. Rev. Phys. Chem.*, 2011, **30**, 335–369.
  - 22 C. Iuga, J. R. Alvarez-Idaboy, L. Reyes and A. Vivier-Bunge, Can a single water molecule really catalyze the acetaldehyde + OH reaction in tropospheric conditions?, *J. Phys. Chem. Lett.*, 2010, **1**, 3112–3115.
  - 23 B. Du and W. Zhang, Theoretical study on the water-assisted reaction of NCO with HCHO, *J. Phys. Chem. A*, 2013, **117**, 6883–6892.
  - 24 P. S. Peters, D. Duflo, A. Faure, C. Kahane, C. Ceccarelli, L. Wiesenfeld and C. Toubin, Theoretical investigation of the isomerization of *trans*-HCOH to H<sub>2</sub>CO: an example of a water-catalyzed reaction, *J. Phys. Chem. A*, 2011, **115**, 8983–8989.
  - 25 B. Long, J. L. Bao and D. G. Truhlar, Atmospheric chemistry of Criegee intermediates: Unimolecular reactions and reactions with water, *J. Am. Chem. Soc.*, 2016, **138**, 14409–14422.
  - 26 F. Xu, X.-F. Tan, Z.-G. Dong, D.-S. Ren and B. Long, Hydrolysis of ketene catalysed by nitric acid and water in the atmosphere, *Environ. Chem.*, 2020, **17**, 457–467.
  - 27 K. Li, X. Song, T. Zhu, C. Wang, X. Sun, P. Ning and L. Tang, Mechanistic and kinetic study on the catalytic hydrolysis of COS in small clusters of sulfuric acid, *Environ. Pollut.*, 2018, **232**, 615–623.
  - 28 B. Long, X.-F. Tan, C.-R. Chang, W.-X. Zhao, Z.-W. Long, D.-S. Ren and W.-J. Zhang, Theoretical studies on gas-phase reactions of sulfuric acid catalyzed hydrolysis of formaldehyde and formaldehyde with sulfuric acid and H<sub>2</sub>SO<sub>4</sub>···H<sub>2</sub>O complex, *J. Phys. Chem. A*, 2013, **117**, 5106–5116.
  - 29 X.-F. Tan, B. Long, D.-S. Ren, W.-J. Zhang, Z.-W. Long and E. Mitchell, Atmospheric chemistry of CH<sub>3</sub>CHO: the hydrolysis of CH<sub>3</sub>CHO catalyzed by H<sub>2</sub>SO<sub>4</sub>, *Phys. Chem. Chem. Phys.*, 2018, **20**, 7701–7709.
  - 30 R. J. Buszek, A. Sinha and J. S. Francisco, The isomerization of methoxy radical: intramolecular hydrogen atom transfer mediated through acid catalysis, *J. Am. Chem. Soc.*, 2011, **133**, 2013–2015.
  - 31 J. Elm, M. Bilde and K. V. Mikkelsen, Influence of nucleation precursors on the reaction kinetics of methanol with the OH radical, *J. Phys. Chem. A*, 2013, **117**, 6695–6701.



- 32 A. Karton, Inorganic acid-catalyzed tautomerization of vinyl alcohol to acetaldehyde, *Chem. Phys. Lett.*, 2014, **592**, 330–333.
- 33 M. Torrent-Sucarrat, J. S. Francisco and J. M. Anglada, Sulfuric acid as autocatalyst in the formation of sulfuric acid, *J. Am. Chem. Soc.*, 2012, **134**, 20632–20644.
- 34 Y. Zhang, Y. Cheng, T. Zhang, R. Wang, J. Ji, Y. Xia, M. Lily, Z. Wang and B. Muthiah, A computational study of the  $\text{HO}_2 + \text{SO}_3 \rightarrow \text{HOSO}_2 + {}^3\text{O}_2$  reaction catalyzed by a water monomer, a water dimer and small clusters of sulfuric acid: kinetics and atmospheric implications, *Phys. Chem. Chem. Phys.*, 2022, **24**, 18205–18216.
- 35 T. Zhang, Y. Zhang, S. Tian, M. Zhou, D. Liu, L. Lin, Q. Zhang, R. Wang and B. Muthiah, Possible atmospheric source of  $\text{NH}_2\text{SO}_3\text{H}$ : the hydrolysis of  $\text{HNSO}_2$  in the presence of neutral, basic, and acidic catalysts, *Phys. Chem. Chem. Phys.*, 2022, **24**, 4966–4977.
- 36 Y. Cheng, R. Wang, Y. Chen, S. Tian, N. Gao, Z. Zhang and T. Zhang, Hydrolysis of  $\text{SO}_3$  in Small Clusters of Sulfuric Acid: Mechanistic and Kinetic Study, *ACS Earth Space Chem.*, 2022, **6**, 3078–3089.
- 37 J. Liu, S. Fang, Z. Wang, W. Yi, F.-M. Tao and J.-y. Liu, Hydrolysis of sulfur dioxide in small clusters of sulfuric acid: Mechanistic and kinetic study, *Environ. Sci. Technol.*, 2015, **49**, 13112–13120.
- 38 T. Zhang, R. Wang, H. Chen, S. Min, Z. Wang, C. Zhao, Q. Xu, L. Jin, W. Wang and Z. Wang, Can a single water molecule really affect the  $\text{HO}_2 + \text{NO}_2$  hydrogen abstraction reaction under tropospheric conditions?, *Phys. Chem. Chem. Phys.*, 2015, **17**, 15046–15055.
- 39 S. Sarkar and B. Bandyopadhyay, Aldehyde as a potential source of aminol in troposphere: Influence of water and formic acid catalysis on ammonolysis of formaldehyde, *Atmos. Environ.*, 2019, **213**, 223–230.
- 40 Z. Wang, C. Zhang, G. Lv, X. Sun, N. Wang and Z. Li, Synergistic reaction of  $\text{SO}_2$  with  $\text{NO}_2$  in presence of  $\text{H}_2\text{O}$  and  $\text{NH}_3$ : A potential source of sulfate aerosol, *Int. J. Mol. Sci.*, 2019, **20**, 3746.
- 41 B. Du and W. Zhang, The effect of  $(\text{H}_2\text{O})_n$  ( $n = 1-2$ ) or  $\text{H}_2\text{S}$  on the hydrogen abstraction reaction of  $\text{H}_2\text{S}$  by OH radicals in the atmosphere, *Comput. Theor. Chem.*, 2015, **1069**, 77–85.
- 42 T. Zhang, W. Wang, P. Zhang, J. Lü and Y. Zhang, Water-catalyzed gas-phase hydrogen abstraction reactions of  $\text{CH}_3\text{O}_2$  and  $\text{HO}_2$  with  $\text{HO}_2$ : a computational investigation, *Phys. Chem. Chem. Phys.*, 2011, **13**, 20794–20805.
- 43 M. E. Dunn, E. K. Pokon and G. C. Shields, Thermodynamics of forming water clusters at various temperatures and pressures by Gaussian-2, Gaussian-3, complete basis set-QB3, and complete basis set-APNO model chemistries; implications for atmospheric chemistry, *J. Am. Chem. Soc.*, 2004, **126**, 2647–2653.
- 44 H. Zhang, O. Kupiainen-Määttä, X. Zhang, V. Molinero, Y. Zhang and Z. Li, The enhancement mechanism of glycolic acid on the formation of atmospheric sulfuric acid-ammonia molecular clusters, *J. Chem. Phys.*, 2017, **146**, 184308.
- 45 L. Zhang and B. Long, Hydrolysis of formyl fluoride catalyzed by sulfuric acid and formic acid in the atmosphere, *ACS Omega*, 2019, **4**, 18996–19004.
- 46 R. Asatryan and E. Ruckenstein, Dihydrogen catalysis: a remarkable avenue in the reactivity of molecular hydrogen, *Catal. Rev.*, 2014, **56**, 403–475.
- 47 S. Grimme, S. Ehrlich and L. Goerigk, Effect of the damping function in dispersion corrected density functional theory, *J. Comput. Chem.*, 2011, **32**, 1456–1465.
- 48 J. Witte, M. Goldey, J. B. Neaton and M. Head-Gordon, Beyond energies: Geometries of nonbonded molecular complexes as metrics for assessing electronic structure approaches, *J. Chem. Theory Comput.*, 2015, **11**, 1481–1492.
- 49 J. Rezac, Non-covalent interactions atlas benchmark data sets 2: Hydrogen bonding in an extended chemical space, *J. Chem. Theory Comput.*, 2020, **16**, 6305–6316.
- 50 J. Rezáč, An integrative framework for computational chemistry, *J. Comput. Chem.*, 2016, **37**, 1230–1237.
- 51 J. Elm, M. Bilde and K. V. Mikkelsen, Assessment of density functional theory in predicting structures and free energies of reaction of atmospheric prenucleation clusters, *J. Chem. Theory Comput.*, 2012, **8**, 2071–2077.
- 52 N. Mardirossian and M. Head-Gordon, How accurate are the Minnesota density functionals for noncovalent interactions, isomerization energies, thermochemistry, and barrier heights involving molecules composed of main-group elements?, *J. Chem. Theory Comput.*, 2016, **12**, 4303–4325.
- 53 A. T. Pereira, A. J. Ribeiro, P. A. Fernandes and M. J. Ramos, Benchmarking of density functionals for the kinetics and thermodynamics of the hydrolysis of glycosidic bonds catalyzed by glycosidases, *Int. J. Quantum Chem.*, 2017, **117**, e25409.
- 54 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson and *et al.*, *Gaussian09*, 1, Gaussian, Inc., Wallingford CT, 2009, 121, pp. 150–166.
- 55 J. Elm, M. Bilde and K. V. Mikkelsen, Assessment of binding energies of atmospherically relevant clusters, *Phys. Chem. Chem. Phys.*, 2013, **15**, 16442–16445.
- 56 A. F. Rodrigues-Oliveira, F. W. M. Ribeiro, G. Cervi and T. C. Correra, Evaluation of common theoretical methods for predicting infrared multiphotonic dissociation vibrational spectra of intramolecular hydrogen-bonded ions, *ACS Omega*, 2018, **3**, 9075–9085.
- 57 M. Kieninger and O. N. Ventura, Calculations of the infrared and Raman spectra of simple thiols and thiol-water complexes, *Int. J. Quantum Inf.*, 2011, **111**, 1843–1857.
- 58 H. Hratchian and H. Schlegel, Using Hessian updating to increase the efficiency of a Hessian based predictor-corrector reaction path following method, *J. Chem. Theory Comput.*, 2005, **1**, 61–69.
- 59 H. P. Hratchian and H. B. Schlegel, Accurate reaction paths using a Hessian based predictor-corrector integrator, *J. Chem. Phys.*, 2004, **120**, 9918–9924.
- 60 S. Maeda, Y. Harabuchi, Y. Ono, T. Taketsugu and K. Morokuma, Intrinsic reaction coordinate: Calculation,



- bifurcation, and automated search, *Int. J. Quantum Chem.*, 2015, **115**, 258–269.
- 61 J. Pople, R. Krishnan, H. Schlegel and J. Binkley, Electron correlation theories and their application to the study of simple reaction potential surfaces, *Int. J. Quantum Chem.*, 1978, **14**, 545–560.
- 62 Y. S. Lee, S. A. Kucharski and R. J. Bartlett, A coupled cluster approach with triple excitations, *J. Chem. Phys.*, 1984, **81**, 5906–5912.
- 63 R. A. Kendall, T. H. Dunning Jr and R. J. Harrison, Electron affinities of the first-row atoms revisited. Systematic basis sets and wave functions, *J. Chem. Phys.*, 1992, **96**, 6796–6806.
- 64 T. J. Lee and P. R. Taylor, A diagnostic for determining the quality of single-reference electron correlation methods, *Int. J. Quantum Chem.*, 1989, **36**, 199–207.
- 65 T. J. Lee, Comparison of the  $T_1$  and  $D_1$  diagnostics for electronic structure theory: a new definition for the open-shell  $D_1$  diagnostic, *Chem. Phys. Lett.*, 2003, **372**, 362–367.
- 66 J. Zhang and M. Dolg, ABCluster: the artificial bee colony algorithm for cluster global optimization, *Phys. Chem. Chem. Phys.*, 2015, **17**, 24173–24181.
- 67 J. Zhang and M. Dolg, Global optimization of clusters of rigid molecules using the artificial bee colony algorithm, *Phys. Chem. Chem. Phys.*, 2016, **18**, 3003–3010.
- 68 W. L. Jorgensen, J. Chandrasekhar, J. D. Madura, R. W. Impey and M. L. Klein, Comparison of simple potential functions for simulating liquid water, *J. Chem. Phys.*, 1983, **79**, 926–935.
- 69 D. J. Wales and M. P. Hodges, Global minima of water clusters  $(\text{H}_2\text{O})_n$ ,  $n \leq 21$ , described by an empirical potential, *Chem. Phys. Lett.*, 1998, **286**, 65–72.
- 70 A. D. MacKerell Jr, D. Bashford, M. Bellott, R. L. Dunbrack Jr, J. D. Evanseck, M. J. Field, S. Fischer, J. Gao, H. Guo and S. Ha, All-atom empirical potential for molecular modeling and dynamics studies of proteins, *J. Phys. Chem. B*, 1998, **102**, 3586–3616.
- 71 J. J. Stewart, Optimization of parameters for semiempirical methods VI: more modifications to the NDDO approximations and re-optimization of parameters, *J. Mol. Model.*, 2013, **19**, 1–32.
- 72 J. Stewart, *MOPAC2016 Stewart Computational Chemistry*, Colorado Springs, CO, OpenMOPAC, 2016.
- 73 Y. Chuang, J. Corchado, P. Fast, J. Villa, E. Coitino, W. Hu, Y. Liu, G. Lynch, K. Nguyen and C. Jackels, *Polyrate-version 8.2*, University of Minnesota, Minneapolis, 1999.
- 74 D. R. Glowacki, C.-H. Liang, C. Morley, M. J. Pilling and S. H. Robertson, MESMER: an open-source master equation solver for multi-energy well reactions, *J. Phys. Chem. A*, 2012, **116**, 9545–9560.
- 75 G. Horváth, I. Horváth, S. A.-D. Almousa and M. Telek, Numerical inverse Laplace transformation using concentrated matrix exponential distributions, *Perform. Evaluation*, 2020, **137**, 102067.
- 76 A. Kumar, S. Mallick and P. Kumar, Oxidation of HOSO' by Cl': a new source of SO<sub>2</sub> in the atmosphere?, *Phys. Chem. Chem. Phys.*, 2021, **23**, 18707–18711.
- 77 T. V.-T. Mai, M. V. Duong, H. T. Nguyen and L. K. Huynh, Ab initio kinetics of the HOSO<sub>2</sub> + <sup>3</sup>O<sub>2</sub> → SO<sub>3</sub> + HO<sub>2</sub> reaction, *Phys. Chem. Chem. Phys.*, 2018, **20**, 6677–6687.
- 78 S. Mallick, A. Kumar and P. Kumar, Oxidation of HOSO by NH<sub>2</sub>: A new path for the formation of an acid rain precursor, *Chem. Phys. Lett.*, 2021, **773**, 138536.
- 79 H. Hippler, J. Troe and H. Wendelken, Collisional deactivation of vibrationally highly excited polyatomic molecules. II. Direct observations for excited toluene, *J. Chem. Phys.*, 1983, **78**, 6709–6717.
- 80 R. J. Kee, G. Dixon-Lewis, J. Warnatz, M. E. Coltrin and J. A. Miller, A Fortran computer code package for the evaluation of gas-phase multicomponent transport properties, *Sandia National Laboratories Report SAND86-8246*, 1986, vol. 13, pp. 80401–81887.
- 81 T. V.-T. Mai, H. T. Nguyen and L. K. Huynh, Ab initio dynamics of hydrogen abstraction from N<sub>2</sub>H<sub>4</sub> by OH radicals: an RRKM-based master equation study, *Phys. Chem. Chem. Phys.*, 2019, **21**, 23733–23741.
- 82 T. V.-T. Mai, H. T. Nguyen and L. K. Huynh, Kinetics of hydrogen abstraction from CH<sub>3</sub>SH by OH radicals: An *ab initio* RRKM-based master equation study, *Atmos. Environ.*, 2020, **242**, 117833.
- 83 X. T. Le, T. V.-T. Mai, M. v. Duong and L. K. Huynh, Kinetics of hydrogen abstraction from desflurane by OH and Cl radicals—A theoretical study, *Chem. Phys. Lett.*, 2019, **728**, 142–147.
- 84 T. V.-T. Mai, P. Raghunath, X. T. Le, L. K. Huynh, P.-C. Nam and M.-C. Lin, Ab initio chemical kinetics for the HCCO + OH reaction, *Chem. Phys. Lett.*, 2014, **592**, 175–181.
- 85 T. V.-T. Mai, M. v. Duong, H. T. Nguyen, K. C. Lin and L. K. Huynh, Ab initio chemical kinetics of the CH<sub>2</sub>OO + C<sub>2</sub>F<sub>4</sub> reaction, *Chem. Phys. Lett.*, 2018, **706**, 280–284.
- 86 T. V.-T. Mai and L. K. Huynh, Ab initio kinetics of the C<sub>2</sub>H<sub>2</sub> + NH<sub>2</sub> reaction: a revisited study, *Phys. Chem. Chem. Phys.*, 2019, **21**, 17232–17239.
- 87 J. M. Anglada, G. J. Hoffman, L. V. Slipchenko, M. M. Costa, M. F. Ruiz-Lopez and J. S. Francisco, Atmospheric significance of water clusters and ozone-water complexes, *J. Phys. Chem. A*, 2013, **117**, 10381–10396.
- 88 B. Bandyopadhyay, P. Kumar and P. Biswas, Ammonia catalyzed formation of sulfuric acid in troposphere: the curious case of a base promoting acid rain, *J. Phys. Chem. A*, 2017, **121**, 3101–3108.
- 89 S. Mallick, S. Sarkar, B. Bandyopadhyay and P. Kumar, Effect of ammonia and formic acid on the OH' + HCl reaction in the troposphere: Competition between single and double hydrogen atom transfer pathways, *J. Phys. Chem. A*, 2018, **122**, 350–363.
- 90 T. Zhang, K. Zhai, Y. Zhang, L. Geng, Z. Geng, M. Zhou, Y. Lu, X. Shao and M. Lily, Effect of water and ammonia on the HO + NH<sub>3</sub> → NH<sub>2</sub> + H<sub>2</sub>O reaction in troposphere: Competition between single and double hydrogen atom transfer pathways, *Comput. Theor. Chem.*, 2020, **1176**, 112747.
- 91 S. Sarkar and B. Bandyopadhyay, Theoretical investigation of the relative impacts of water and ammonia on the



- tropospheric conversion of  $\text{N}_2\text{O}_5$  to  $\text{HNO}_3$ , *Phys. Chem. Chem. Phys.*, 2021, **23**, 6651–6664.
- 92 B. Long, X. F. Tan, Y. B. Wang, J. Li, D. S. Ren and W. J. Zhang, Theoretical Studies on Reactions of OH with  $\text{H}_2\text{SO}_4 \cdots \text{NH}_3$  Complex and  $\text{NH}_2$  with  $\text{H}_2\text{SO}_4$  in the Presence of Water, *ChemistrySelect*, 2016, **1**, 1421–1430.
- 93 S. Sarkar, S. Mallick, P. Kumar and B. Bandyopadhyay, Isomerization of methoxy radical in the troposphere: competition between acidic, neutral and basic catalysts, *Phys. Chem. Chem. Phys.*, 2017, **19**, 27848–27858.
- 94 R. J. Buszek, M. Torrent-Sucarrat, J. M. Anglada and J. S. Francisco, Effects of a single water molecule on the OH +  $\text{H}_2\text{O}_2$  reaction, *J. Phys. Chem. A*, 2012, **116**, 5821–5829.
- 95 J. Gonzalez, J. M. Anglada, R. J. Buszek and J. S. Francisco, Impact of water on the OH + HOCl reaction, *J. Am. Chem. Soc.*, 2011, **133**, 3345–3353.
- 96 A. Parandaman, J. E. Perez and A. Sinha, Atmospheric decomposition of trifluoromethanol catalyzed by formic acid, *J. Phys. Chem. A*, 2018, **122**, 9553–9562.
- 97 G. Shen, M. Suto and L. Lee, Reaction rate constant of  $\text{SO}_3 + \text{CH}_3\text{OH}$  in the gas phase, *Int. J. Chem. Kinet.*, 1990, **22**, 633–639.
- 98 L. Liu, J. Zhong, H. Vehkamäki, T. Kurtén, L. Du, X. Zhang, J. S. Francisco and X. C. Zeng, Unexpected quenching effect on new particle formation from the atmospheric reaction of methanol with  $\text{SO}_3$ , *Proc. Natl. Acad. Sci.*, 2019, **116**, 24966–24971.
- 99 J. Gonzalez, M. Torrent-Sucarrat and J. M. Anglada, The reactions of  $\text{SO}_3$  with  $\text{HO}_2$  radical and  $\text{H}_2\text{O} \cdots \text{HO}_2$  radical complex. Theoretical study on the atmospheric formation of  $\text{HSO}_5$  and  $\text{H}_2\text{SO}_4$ , *Phys. Chem. Chem. Phys.*, 2010, **12**, 2116–2125.
- 100 Y.-Q. Sun, X. Wang, F.-Y. Bai and X.-M. Pan, Theoretical study of the hydrolysis of HOSO +  $\text{NO}_2$  as a source of atmospheric HONO: effects of  $\text{H}_2\text{O}$  or  $\text{NH}_3$ , *Environ. Chem.*, 2016, **14**, 19–30.

