



Cite this: *Mater. Adv.*, 2023,
4, 6449

Correction: DFT investigation of the oxygen reduction reaction over nitrogen (N) doped graphdiyne as an electrocatalyst: the importance of pre-adsorbed OH* and the solvation effect

Yuelin Wang,^a Thanh Ngoc Pham,^a Harry H. Halim,^a Likai Yan^c and Yoshitada Morikawa^{*ab}

DOI: 10.1039/d3ma90095a

rsc.li/materials-advances

Correction for 'DFT investigation of the oxygen reduction reaction over nitrogen (N) doped graphdiyne as an electrocatalyst: the importance of pre-adsorbed OH* and the solvation effect' by Yuelin Wang *et al.*, *Mater. Adv.*, 2023, <https://doi.org/10.1039/d3ma00502j>.

The authors regret that on the 5th page of the article, in the three paragraphs from “Consequently, upon applying a limiting potential of 0.22 V...” to “...cannot be used as an ORR electrocatalyst.”, all instances of ‘sp-N1GDY(OH)/GDY’ and ‘sp-N2GDY(OH)/GDY’ were mislabelled. They should be read as ‘sp-N1GDY(OH)/G’ and ‘sp-N2GDY(OH)/G’, respectively.

In Fig. 4, section (a), ‘sp-N1GDY/G (OH)’ and ‘sp-N2GDY/G (OH)’ were mislabelled. They should be read as ‘sp-N1GDY(OH)/G’ and ‘sp-N2GDY(OH)/G’, respectively. The correctly labelled Fig. 4 is given below.

In the section titled “The electronic structure of the active site relates to O₂ activation”, the O₂ dissociation barrier values for sp-N1GDY(OH)/G and Pyri-NGDY/G are given incorrectly. The correct text should be “. . .and O₂ dissociation barrier is high (1.86 eV and 1.52 eV)”.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Department of Precision Engineering, Graduate School of Engineering, Osaka University, 2-1, Yamada-oka, Osaka 565-0871, Suita, Japan.

E-mail: morikawa@prec.eng.osaka-u.ac.jp

^b Research Center for Precision Engineering, Graduate School of Engineering, Osaka University, 2-1 Yamada-oka, Suita 565-0871, Osaka, Japan

^c Institute of Functional Material Chemistry, Key Laboratory of Polyoxometalate Science of Ministry of Education, Faculty of Chemistry, Northeast Normal University, Changchun, 130024, P. R. China






Fig. 4 (a) PDOS of C 2p of the active site in sp-N1GDY/G, sp-N1GDY(OH)/G, sp-N2GDY/G, sp-N2GDY(OH)/G, and Pyri-NGDY/G. (b) The relationship between PDOS height value at the Fermi level and the adsorption energy of O₂.

