


 Cite this: *New J. Chem.*, 2023, 47, 20358

 Received 17th September 2023,
 Accepted 19th October 2023

DOI: 10.1039/d3nj04371a

rsc.li/njc

Catalytic prenylation of natural polyphenols†

 Yi Du,^a Iman Korchi,^a Aleksandr E. Rubtsov^{ab} and Andrei V. Malkov^{id}*^a

Prenylated polyphenols occur naturally and exhibit biological activity superior to the ubiquitous parent polyphenols. However, their low abundance limits the wider application of these derivatives. In this work, we present an expedient, single-step catalytic protocol for introducing terpene fragments into the aromatic rings of the widely available natural stilbenoids and flavonoids to upgrade their therapeutic potential.

Introduction

Flavonoids and stilbenoids are ubiquitous secondary metabolites in plants, leaves, flowers, fruits and seeds.^{1,2} Bioactive flavonoids acquired a proven track record of beneficial effects on human health;³ the pharmaceutical, medicinal, nutraceutical and cosmetics industries actively use flavonoids in their development.^{3,4} However, the most common polyphenols do not reach therapeutic plasma concentration levels due to the low solubility and poor absorption of the compounds. This low bioavailability can be improved through chemical modifications that alter the polarity and lipophilicity of the polyphenols, as well as the biological functions that depend on the structure. Among the chemical modification strategies, prenylation is often considered an essential way to boost the potency of the parent flavonoid.^{5–8} Prenylated analogues have been detected for most flavonoids, such as chalcones, stilbenes, flavanones, flavones, flavonols and isoflavones, with some examples shown in Fig. 1. While flavonoids are usually present in reasonable quantities in nature, the levels of prenylated flavonoids are much lower. The low abundance in nature limits their application as medicines and nutraceuticals, therefore it is important to develop facile synthetic approaches.

Chemical synthesis of prenylated flavonoids usually relies on multistep sequences involving the protection of hydroxyls, *O*-prenylation followed by the Claisen rearrangement of the *O*- to *C*-isomer and deprotection.^{8–12} Alternative approaches are




Fig. 1 Selected examples of natural prenylated polyphenols.

based on Suzuki–Miyaura cross-coupling, which requires prior synthesis of the respective halides or boronic acid derivatives,^{13,14} or the use of chemoenzymatic methods.^{15,16}

Herein, we present the development of a simple, single-step catalytic Friedel–Crafts alkylation of unprotected natural polyphenols using prenyl alcohol and its analogues.

Results and discussion

Optimisation studies


Catalytic C–H allylation of aromatic compounds is a widely used synthetic transformation for the formation of C–C bonds.¹⁷ Of the plethora of methods for the modification of electron-rich arenes, Friedel–Crafts alkylation looks appealing from the practical viewpoint because it can employ unprotected allylic alcohols as the

^a Department of Chemistry, Loughborough University, Loughborough, LE11 3TU, UK. E-mail: A.Malkov@lboro.ac.uk

^b Department of Chemistry, Perm State University, Bukireva 15, Perm 614990, Russia

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3nj04371a>



Table 1 Optimisation of prenylation of resveratrol **1a**^a

Entry	Catalyst	Additive	Solvent	Yield ^b (%)	Ratio 3/4
1	(<i>acac</i>) ₂ MoCl ₂	AgOTf	CH ₃ CN	13	40/60
2	Fe(OTf) ₃	—	CH ₃ CN	48	1/99
3	In(OAc) ₃	—	CH ₃ CN	nr	—
4	InCl ₃	—	CH ₃ CN	nr	—
5	InCl ₃	AgOTf	CH ₃ CN	37	1/99
6	AlCl ₃	—	CH ₃ CN	42	25/75
7	AlCl ₃	AgOTf	CH ₃ CN	65	26/74
8	AlCl ₃	AgOTf	Et ₂ O	Trace	—
9	AlCl ₃	AgOTf	MeOH	nr	—
10	AlCl ₃	AgOTf	DCM	Trace	—
11	AlCl ₃	AgOTf	Acetone	85	21/79
12	AlCl ₃	AgOTf	EtOAc	81	31/69
13 ^c	AlCl ₃	AgOTf	Acetone	59	24/76
14 ^d	AlCl ₃	AgOTf	Acetone	42	24/76


^a Conditions: **1a** (1 mmol), **2a** (2 mmol), catalyst (10 mol%) and additive (10 mol%); in 24 mL solvent under N₂ at RT overnight. ^b Combined isolated yield; nr: no reaction. ^c Catalyst: 10 mol%, additive: 20 mol%. ^d Catalyst: 10 mol%; additive: 30 mol%.

electrophiles.^{18,19} The reaction can be performed using catalytic quantities of Lewis or Brønsted acids.^{18,20,21}

However, despite the apparent simplicity, there are very few reports of the Friedel–Crafts prenylation of polyphenolic compounds, which mainly focus on the allylation of related resorcinol analogues.^{22–24} The reason for this may stem from the inactivation of the Lewis catalysts by binding to the phenolic groups. Therefore, successful examples^{22,24,25} usually employ the excess of promoters.

With the aim to develop a catalytic system for the modification of natural polyphenols, prenylation of resveratrol **1a** was selected for optimisation studies (Table 1).

First, a catalytic system developed by us previously for the allylation of electron-rich arenes was examined.²¹ It employed the Lewis acidic complex (*acac*)₂MoCl₂ activated by silver triflate by replacing the chloride with a non-nucleophilic anion (entry 1), however, the yield and regioselectivity were low. Next, a series of other metal salts were examined. With Fe(OTf)₃ (entry 2), 2-prenylresveratrol **4aa** was obtained in a modest yield with only trace quantities of the 4-isomer **3aa**. Indium(III) acetate and chloride were inactive (entries 3 and 4), but InCl₃ activated by AgOTf worked similarly to Fe(OTf)₃, with a slightly reduced yield (entry 5). The increased yield was obtained with the catalytic system based on the equimolar quantities of AlCl₃ and AgOTf (entry 7). AlCl₃ worked even without silver triflate (entry 6) but with lower product yield. Notably, AlCl₃ produced a 3 : 1 mixture of regioisomers **4aa** and **3aa**. This could be viewed



Scheme 1 Prenylation of protected resveratrol.

as a disadvantage, however, taking into account that the isomers are readily separable by chromatography and that synthesis of isomer **3aa**, a naturally occurring arachidin-2, requires multistep sequences,^{26,27} this reaction can serve as a viable alternative for obtaining practical quantities of **3aa**, in a single step. In this context, the molybdenum catalyst (entry 1) exhibited the highest proportion of the symmetrical **3aa** in the product mixture. To boost the reaction yield, a protection strategy was explored, where the hydroxy groups were protected as silyl ethers (Scheme 1). A nearly equimolar mixture of **3aa** and **4aa** has been achieved, though, despite some improvement in the ratio towards **3aa**, the overall yield remained modest.

Next, the reaction solvents were briefly assessed (entries 8–12) with a focus on greener solvents with lower environmental impact.^{28,29} Ether and methanol were found unsuitable (entries 8 and 9); as was non-green dichloromethane (entry 10). The highest yields were obtained in acetone and ethyl acetate (entries 11 and 12). An increase in the number of equivalents of AgOTf to AlCl₃ proved to be detrimental to the reaction yield but the product ratio remained the same (entries 13 and 14). Thus, the conditions of entry 11 were selected as optimal.


The reaction scope

With the optimal protocol in hand, the reaction scope has been investigated using naturally occurring polyphenols. The results are presented according to the polyphenol classes.

Following the optimisation experiments, the allylation of resveratrol was assessed first. Chemical yields of the isolated pure compounds are shown in Table 2. Two entries are shown for the reaction of resveratrol with prenyl. Thus, under the optimal conditions (entry 1), the isolated yields of both regioisomers **3aa** and **4aa** are presented. At the same time, this reaction can occur regioselectively, furnishing only **4aa** (entry 2). In this instance, Fe(OTf)₃ had been used as a catalyst.

The reaction of resveratrol with geraniol mirrored that of the prenyl: a 3 : 1 mixture of products **4ab** and **3ab** was produced in a high overall yield of 92% (entry 3). The two compounds were readily separated by chromatography. The more abundant isomer **4ab** is a bioactive macatrichocarpin F isolated from *Macaranga* plants.³⁰ This compound previously was synthesised through a sequence involving *O*-alkylation followed by the *O*- to *C*-isomerisation.^{31,32} The allylation with cinnamyl alcohol produced an unknown symmetrical isomer **3ac** as a single product in a 60% yield (entry 4). The reason for such a switch in the



Table 2 Allylation of resveratrol **1a**^a

Entry	Alcohol 2, R	3, Yield (%)	4, Yield (%)
1	Prenyl alcohol, 2a	3aa , 18	4aa , 67
2	Prenyl alcohol, 2a ^b	—	4aa , 48
3	Geraniol, 2b	3ab , 22	4ab , 70
4	Cinnamyl alcohol, 2c	3ac , 60	—

^a Conditions: **1a** (1 mmol), **2** (2 mmol), AlCl₃ (10 mol%) and AgOTf (10 mol%); in 24 mL solvent under N₂ at RT overnight, unless stated otherwise. The regioisomers were separated by flash chromatography, yields are given for isolated analytically pure compounds. ^b The reaction was performed in CH₃CN using Fe(OTf)₃ (10 mol%) as a catalyst.

regioselectivity is unknown but may relate to the formation of a more stabilised cation intermediate.

Prenylation of flavones, chrysin **1b** and luteolin **1c** proved to be more challenging (Fig. 2). Despite good overall yields, mixtures of compounds were produced. Thus, the reaction of chrysin with prenyl gave a mixture of mono and di-prenylation products, **3ba** and **5ba**, respectively. The initially formed 6-isomer **3ba** likely undergoes the second alkylation to furnish **5ba**. Nonetheless, the compounds can be readily separated by chromatography. In contrast, luteolin gave a 3:1 mixture of





Fig. 2 Allylation of flavones, chrysin **1b**, and luteolin **1c**. (For conditions, see Table 1, entry 11, yields are given for isolated, analytically pure compounds. ^a Separated chromatographically from the mixture of **3ba** and **5ba**. ^b Separated chromatographically from the mixture of **3ca** and **4ca**).

monosubstituted isomers **3ca** and **4ca**, which had similar chromatographic mobility and were difficult to separate; only isomer **3ca** was isolated in pure form. Other isomer **4ca** was characterised in the mixture with **3ca**.

Flavonols quercetin **1d** and myricetin **1e** are even more densely substituted with hydroxyl groups and, therefore potentially represented even more challenging targets for the catalytic prenylation. However, they turned out to be good substrates (Scheme 2). Thus, alkylation of quercetin with prenyl, geraniol and cinnamyl alcohol occurred uneventfully furnishing good yields of the respective monosubstituted derivatives **3da**, **3db** and **3dc** as single regioisomers. Note that compound **3da** is a natural polyphenol gancaonin P found in liquorice.³³ The reaction of myricetin with geraniol followed the trend to afford **3ab** in a 73% yield. However, the prenylation of myricetin,



Scheme 2 Allylation of flavonols, (A) quercetin **1d** and (B) myricetin **1e**. (For conditions, see Table 1, entry 11, yields are given for isolated, analytically pure compounds. ^a Separated chromatographically from the mixture of **3ea** and **5ea**).



along with the expected **3ea** (24%), showed the formation of dialkylated product **5ea** (62%). So far, the attempts at reversing the ratio in favour of monoalkylation have been unsuccessful, but the products are readily separable by simple flash chromatography.

Experimental

General procedure for prenylation of polyphenols

A polyphenol (1 mmol) was added to the solution of AlCl₃ (10 mol%) and AgOTf (10 mol%) in acetone (20 mL). The mixture was stirred for 20 min. Then, a solution of prenyl or other allylic alcohol (2 mmol) in acetone (4 mL) was added dropwise to this mixture under an inter atmosphere. The resulting solution was stirred at room temperature overnight. The reaction progress was monitored by TLC. The work-up involved dilution with water and extraction with EtOAc. The crude product was purified by chromatography on silica to afford the target compound. For details, see ESI.†

Conclusions

In conclusion, we developed an expedient catalytic method for the introduction of allyl groups, such as prenyl, geranyl and cinnamyl, into the aromatic rings of natural polyphenols under mild reaction conditions. Resveratrol, flavones and flavonols proved to be competent substrates for this reaction furnishing practical yields of the alkylation products, some of the synthesised analogues are occurring in nature. Chemical prenylation can modify the structure of polyphenols, potentially leading to enhanced biological activity and improved pharmacological properties as evidenced by a superior bioactivity of some naturally occurring prenylated analogues compared to the parent polyphenols. Therefore, the developed single-step method may serve as a convenient tool for upgrading the bioactivity of the abundant natural polyphenols.

Author contributions

Conceptualization, A. V. M.; funding acquisition, A. V. M. methodology, Y. D., I. K. and A. V. M. formal analysis, Y. D., I. K. and A. E. R.; investigation, Y. D. and I. K.; writing – original draft, A. V. M. and A. E. R.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Y. D. and A. V. M. thank H2020 BBI JU for the PHENOLEXA Grant 101023225. A. E. R. thanks Perm Research and Education Centre “Rational Subsoil Use”, 2023.

Notes and references

- 1 N. Shen, T. Wang, Q. Gan, S. Liu, L. Wang and B. Jin, Plant flavonoids: Classification, distribution, biosynthesis, and antioxidant activity, *Food Chem.*, 2022, **383**, 132531.
- 2 P. Gupta, J. Kour, M. Bakshi and R. Kalsi, in *Nutraceuticals and Health Care*, ed. J. Kour and G. A. Nayik, Academic Press, 2022, ch. 5, pp. 105–113.
- 3 S. Chibisov, E. Kharlitskaya, R. B. Singh, A. Itharat, E. On-Saard, H.-R. Park, J. Chaudhury, S. Chakravorty, O. K. Gupta and M. M. A. Smail, in *Functional Foods and Nutraceuticals in Metabolic and Non-Communicable Diseases*, ed. R. B. Singh, Elsevier, 2022, ch. 44, pp. 671–689.
- 4 A. Ullah, S. Munir, S. L. Badshah, N. Khan, L. Ghani, B. G. Poulson, A. H. Emwas and M. Jaremko, Important Flavonoids and Their Role as a Therapeutic Agent, *Molecules*, 2020, **25**, 5243.
- 5 B. Botta, A. Vitali, P. Menendez, D. Misiiti and G. Delle Monache, Prenylated Flavonoids: Pharmacology and Biotechnology, *Curr. Med. Chem.*, 2005, **12**, 713–739.
- 6 S. Shi, J. Li, X. Zhao, Q. Liu and S.-J. Song, A comprehensive review: Biological activity, modification and synthetic methodologies of prenylated flavonoids, *Phytochemistry*, 2021, **191**, 112895.
- 7 L. M. Dominika, J. Stefano, D. Karel and S. Dallacqua, C-prenylated flavonoids with potential cytotoxic activity against solid tumor cell lines, *Phytochem. Rev.*, 2019, **18**, 1051–1100.
- 8 Y. Zhang, X. Zhang, Z. Xiao, X. Zhang and H. Sun, Hypoglycemic and hypolipidemic dual activities of extracts and flavonoids from *Desmodium caudatum* and an efficient synthesis of the most potent 8-prenylquercetin, *Fitoterapia*, 2022, **156**, 105083.
- 9 W. Li, L. Shu, K. Liu and Q. Wang, Regioselective synthesis of C-prenylated flavonoids via intramolecular [1,3] or [1,5] shift reaction catalyzed by acidic clays, *Tetrahedron Lett.*, 2019, **60**, 151138.
- 10 T. Kawamura, M. Hayashi, R. Mukai, J. Terao and H. Nemoto, An Efficient Method for C8-Prenylation of Flavonols and Flavanones, *Synthesis*, 2012, 1308–1314.
- 11 H. Sun, Y. Li, X. Zhang, Y. Lei, W. Ding, X. Zhao, H. Wang, X. Song, Q. Yao, Y. Zhang, Y. Ma, R. Wang, T. Zhu and P. Yu, Synthesis, α -glucosidase inhibitory and molecular docking studies of prenylated and geranylated flavones, isoflavones and chalcones, *Bioorg. Med. Chem. Lett.*, 2015, **25**, 4567–4571.
- 12 S. Tischer and P. Metz, Selective C-6 prenylation of flavonoids via europium(III)-catalyzed Claisen rearrangement and cross-metathesis, *Adv. Synth. Catal.*, 2007, **349**, 147–151.
- 13 M. Hurtová, D. Biedermann, Z. Osifová, J. Cvačka, K. Valentová and V. Křen, Preparation of Synthetic and Natural Derivatives of Flavonoids Using Suzuki–Miyaura Cross-Coupling Reaction, *Molecules*, 2022, **27**, 967.
- 14 G. Kwesiga, J. Greese, A. Kelling, E. Sperlich and B. Schmidt, The Suzuki–Miyaura Cross-Coupling–Claisen Rearrangement–Cross-Metathesis Approach to Prenylated Isoflavones, *J. Org. Chem.*, 2023, **88**, 1649–1664.



- 15 X. Yang, J. Yang, Y. Jiang, H. Yang, Z. Yun, W. Rong and B. Yang, Regiospecific synthesis of prenylated flavonoids by a prenyltransferase cloned from *Fusarium oxysporum*, *Sci. Rep.*, 2016, **6**, 24819.
- 16 T. Kumano, S. B. Richard, J. P. Noel, M. Nishiyama and T. Kuzuyama, Chemoenzymatic syntheses of prenylated aromatic small molecules using *Streptomyces* prenyltransferases with relaxed substrate specificities, *Bioorg. Med. Chem.*, 2008, **16**, 8117–8126.
- 17 S. Dutta, T. Bhattacharya, D. B. Werz and D. Maiti, Transition-metal-catalyzed C-H allylation reactions, *Chemistry*, 2021, **7**, 555–605.
- 18 R. Ortiz and R. P. Herrera, *Molecules*, 2017, **22**, 574.
- 19 M. Rueping and B. J. Nachtsheim, *Beilstein J. Org. Chem.*, 2010, **6**(No. 6), DOI: [10.3762/bjoc.6.6](https://doi.org/10.3762/bjoc.6.6).
- 20 W. Bonrath, C. Dittel, L. Giraudi, T. Netscher and T. Pabst, Rare earth triflate catalysts in the synthesis of Vitamin E and its derivatives, *Catal. Today*, 2007, **121**, 65–70.
- 21 A. V. Malkov, P. Spoor, V. Vinader and P. Kočovský, Molybdenum(IV) Complexes as Efficient, Lewis Acidic Catalysts for Allylic Substitution. Formation of C–C and C–N Bonds, *J. Org. Chem.*, 1999, **64**, 5308–5311.
- 22 N. G. Jentsch, X. Zhang and J. Magolan, Efficient synthesis of cannabigerol, grifolin, and piperogalin via alumina-promoted allylation, *J. Nat. Prod.*, 2020, **83**, 2587–2591.
- 23 J. Zhang, W. Xiong, Y. Wen, X. Fu, X. Lu, G. Zhang and C. Wang, Magnesium dicarboxylates promote the prenylation of phenolics that is extended to the total synthesis of icaritin, *Org. Biomol. Chem.*, 2022, **20**, 1117–1124.
- 24 Y. W. Tang, C. J. Shi, H. L. Yang, P. Cai, Q. H. Liu, X. L. Yang, L. Y. Kong and X. B. Wang, Synthesis and evaluation of isoprenylation-resveratrol dimer derivatives against Alzheimer's disease, *Eur. J. Med. Chem.*, 2019, **163**, 307–319.
- 25 M. Osorio, M. Carvajal, A. Vergara, E. Butassi, S. Zacchino, C. Mascayano, M. Montoya, S. Mejias, M. C. S. Martín and Y. Vásquez-Martínez, Prenylated flavonoids with potential antimicrobial activity: Synthesis, biological activity, and in silico study, *Int. J. Mol. Sci.*, 2021, **22**, 5472.
- 26 B. H. Park, H. J. Lee and Y. R. Lee, Total synthesis of chiricanine A, arahypin-1, trans -arachidin-2, trans -arachidin-3, and arahypin-5 from peanut seeds, *J. Nat. Prod.*, 2011, **74**, 644–649.
- 27 A. M. Hartung, J. A. Beutler, H. A. Navarro, D. F. Wiemer and J. D. Neighbors, Stilbenes as κ -selective, non-nitrogenous opioid receptor antagonists, *J. Nat. Prod.*, 2014, **77**, 311–319.
- 28 D. Prat, A. Wells, J. Hayler, H. Sneddon, C. R. McElroy, S. Abou-Shehadeh and P. J. Dunn, CHEM21 selection guide of classical- and less classical-solvents, *Green Chem.*, 2015, **18**, 288–296.
- 29 F. P. Byrne, S. Jin, G. Paggiola, T. H. M. Petchey, J. H. Clark, T. J. Farmer, A. J. Hunt, C. Robert McElroy and J. Sherwood, Tools and techniques for solvent selection: green solvent selection guides, *Sustain. Chem. Process.*, 2016, **4**, 7.
- 30 M. Tanjung, L. D. Juliawaty, E. H. Hakim and Y. M. Syah, Flavonoid and stilbene derivatives from *Macaranga trichocarpa*, *Fitoterapia*, 2018, **126**, 74–77.
- 31 T. Zhou, Y. Jiang, B. Zeng and B. Yang, The cancer preventive activity and mechanisms of prenylated resveratrol and derivatives, *Curr. Res. Toxicol.*, 2023, **5**, 100113.
- 32 T. Puksasook, S. Kimura, S. Tadtong, J. Jiaranaikulwanitch, J. Pratuangdejkul, W. Kitphati, K. Suwanborirux, N. Saito and V. Nukoolkarn, Semisynthesis and biological evaluation of prenylated resveratrol derivatives as multi-targeted agents for Alzheimer's disease, *J. Nat. Med.*, 2017, **71**, 665–682.
- 33 T. Nomura, T. Fukai and T. Akiyama, Chemistry of phenolic compounds of licorice (*Glycyrrhiza* species) and their estrogenic and cytotoxic activities, *Pure Appl. Chem.*, 2002, **74**, 1199–1206.

