# **RSC** Advances



View Article Online

View Journal | View Issue

## PAPER

Check for updates

Cite this: RSC Adv., 2023, 13, 3173

Received 2nd November 2022 Accepted 19th December 2022

DOI: 10.1039/d2ra06930j

rsc.li/rsc-advances

#### 1 Introduction

Black and odorous water (BOW) is an extreme example of organic pollution in water.<sup>1-3</sup> BOW may have a pungent smell, which some people may find unpleasant or disgusting.

There are two main mechanisms of water-blackening: one is insoluble substances in water in the form of solid/adsorbed particles; the other is colored organic compounds dissolved in water.<sup>4</sup>

There are four types of odorants: methane ( $CH_4$ ), hydrogen sulfide ( $H_2S$ ), ammonia ( $NH_3$ ) and other small-molecule gases. If the water body is seriously polluted by organic matter, the aerobic decomposition of organic matter makes the oxygen consumption rate in the water body higher than the reoxygenation rate, thereby resulting in water hypoxia. In anoxic water, odor production is synchronized with blackening, and organic matter is decomposed anaerobically to produce  $CH_4$ ,  $H_2S$ ,  $NH_3$ and other odorous and volatile small molecular compounds, which overflow the water surface and enter the atmosphere,

## Removal of ammonia nitrogen from black and odorous water by macrophytes based on laboratory microcosm experiments

Shangmin Huangfu,<sup>a</sup> Fulai Zhou,<sup>b</sup> Xianyun Zheng,<sup>c</sup> Xiaoping Zhang<sup>\*a</sup> and Lifang Hu<sup>1</sup>

In recent years, the removal mechanism of ammonia nitrogen in black and odorous water (BOW), especially in the process of phytoremediation, has been a research "hotspot". Here, the migration process of ammonia nitrogen in macrophytes (*Acorus calamus, Canna indica* and *Eichhornia crassipes*) was detected by Fourier transform infrared (FT-IR) spectroscopy. Experiments revealed that the concentration of ammonia nitrogen (NH<sub>4</sub><sup>+</sup>-N) was reduced significantly. Maximum reduction in the NH<sub>4</sub><sup>+</sup>-N concentration was obtained in 75% BOW: the absorption of NH<sub>4</sub><sup>+</sup>-N was >90% in *A. calamus* and *C. indica*, and >80% in *E. crassipes*. After two 10 days cultivations, in the culture dishes of *A. calamus* and *C. indica*, absorption of NH<sub>4</sub><sup>+</sup>-N was >90% whereas, in the culture dishes of *E. crassipes*, absorption of NH<sub>4</sub><sup>+</sup>-N was ~50% and ~60%. FT-IR spectroscopy showed that NH<sub>4</sub><sup>+</sup>-N, NO<sub>2</sub><sup>-</sup>-N and NO<sub>3</sub><sup>-</sup>-N could be absorbed by the root and migrate to the stem and leaf of macrophytes. NH<sub>4</sub><sup>+</sup>-N and NO<sub>2</sub><sup>-</sup>-N were transformed, and the direction was NH<sub>4</sub><sup>+</sup>-N  $\rightarrow$  NO<sub>2</sub><sup>-</sup>-N  $\rightarrow$  NO<sub>3</sub><sup>-</sup>-N. The migration rate of NH<sub>4</sub><sup>+</sup>-N in *C. indica* was faster because of its regular and smooth capillaries according to scanning electron microscopy. Our study on the removal and transformation mechanism of ammonia nitrogen in BOW could be an important reference for other bodies of water.

thereby emitting odor. Ding *et al.*<sup>5</sup> found that the odor of the water body was caused mainly by the release of H<sub>2</sub>S and NH<sub>3</sub> during the decomposition of organic matter containing sulfur or nitrogen.

Aquatic plants (also known as "macrophytes") can take-up nitrogen compounds in water through absorption and adsorption,<sup>6</sup> which can reduce the nitrogen content in eutrophic water effectively. William<sup>7</sup> pointed out that the effect of a macrophytebased ecological-treatment system is as good as that of typical chemical-treatment system. Macrophytes increase the concentration of dissolved oxygen, provide oxygen for aquatic animals and aerobic microorganisms in the water, restore the damaged aquatic ecosystem, and promote aerobic microorganisms to effectively degrade organic nutrients in the water through respiration; the growth of macrophytes consumes nitrogen in the water, thereby avoiding eutrophication effectively.

Macrophytes mainly absorb nitrogen compounds in the sediment through the roots, then distribute them to the branches and, finally, release them into the water through the living plant or the decay of a dead plant.<sup>8</sup> In natural waters, large-root plants (*e.g.*, submerged macrophytes) have more competitive advantages than phytoplankton because they can absorb nutrients directly from sediment.<sup>9</sup>

Macrophytes can absorb nitrogen compounds directly from water or sediment.<sup>10</sup> The storage of nitrogen compounds in submerged plants is more stable than that in algae.<sup>11</sup> Ammonia

<sup>&</sup>quot;College of Life Science, Anhui Normal University, Wuhu, 241000, China. E-mail: 634732130@qq.com

<sup>&</sup>lt;sup>b</sup>School of Chemical and Environmental Engineering, Anhui Polytechnic University, Wuhu, 241000, China

School of Chemical Engineering, Anhui University of Science and Technology, Huainan, 232001, China. E-mail: hulf@aust.edu.cn

nitrogen  $(NH_4^+-N)$  can be removed by plant uptake, but nitrification and denitrification remain the main mechanisms of removal. The physiological and metabolic activities of submerged plants are directly related to the migration and transformation of nutrients.<sup>12</sup> Submerged macrophytes can control the growth of phytoplankton, provide shelter for zooplankton-feeding animals, repair the aquatic ecosystem as well as maintain the balance of the aquatic ecosystem by biological manipulation.<sup>13</sup>

Yao *et al.*<sup>14</sup> studied the purification effect of six macrophytes (*Ceratophyllum demersum*, *Vallisneria* species, *Vallisneria macrocephala*, *Hydrilla verticillata*, *Sagittaria humilis*, and crown grass) in simulated wastewater. With an increase in treatment time, the proportion of ammonia nitrogen decreased and the proportion of nitrate nitrogen increased. *Vallisneria* species had a good purification effect on nitrogen compounds.

In the present study, three types of macrophytes, *Acorus calamus, Canna indica* and *Eichhornia crassipes*, were used to remove ammonia from BOW. The migration and transformation of  $NH_4^+$ -N,  $NO_2^-$ -N and  $NO_3^-$ -N were investigated by Fourier transform infrared (FT-IR) spectroscopy.

#### 2 Experimental section

#### 2.1 Absorption experiment

The macrophytes were *A. calamus*, *C. indica* and *E. crassipes* in their mature period. The water quality was simulated to be black and odorous. According to the average concentration of the water body on site and plant tolerance, the initial concentration of ammonia nitrogen was set at 24 mg  $L^{-1}$  for the first 10 days and 16 mg  $L^{-1}$  for the second 10 days.

The experiment was simulated in the laboratory. Each plant was placed into five BOW concentrations: 0%, 25%, 50%, 75% and 100% (BOW/pure water, vol%), respectively. Each treatment was repeated thrice. All treatments were carried out in a bucket with a volume of 20 L and duration of 20 days.



Fig. 1 Standard curve for testing the concentration of ammonia nitrogen.

At 8 am on the days 1, 3, 5, 10, 14, 16 and 20 of the experiment, samples were taken to determine the amount of ammonia nitrogen in treated water samples, and BOW of identical concentration was replenished on day-10.

The NH<sub>4</sub><sup>+</sup>-N concentration was tested using Nessler's reagent spectrophotometry method. The standard curve is shown in Fig. 1. The spectral intensity *vs.* concentration of ammonia nitrogen yielded the equation  $y = 0.1881 \times -0.003$  and the coefficient of association was 0.9993.

#### 2.2 Characterization method

The transformation of nitrogen-containing compounds was characterized by FT-IR spectroscopy carried out on a vector 33 FT-IR spectrophotometer (Bruker, USA) with a DTGS detector at a resolution of 4 cm<sup>-1</sup>. The microcapillary structure of aquatic plants was visualized on a Flex-SEM1000 electron microscope (Hitachi, Japan).

#### 3 Results and discussion

The absorption performances of *A. calamus, C. indica* and *E. crassipes* for ammonia nitrogen in five types of BOW are shown in Fig. 2 and Fig. 3.

After 10 days of cultivation in BOW with an initial concentration of ammonia nitrogen of 24 mg L<sup>-1</sup>, the concentration of ammonia nitrogen in the utensils employed for cultivating *A. calamus* and *C. indica* was <2 mg L<sup>-1</sup> (Fig. 2). After 10 days, the concentration of ammonia nitrogen in the utensils employed for cultivating *A. calamus* and *C. indica* remained <2 mg L<sup>-1</sup>. These results showed that *A. calamus* and *C. indica* had a good absorption effect on ammonia nitrogen. For *E. crassipes*, the concentration of ammonia nitrogen in the culture dish was ~8 mg L<sup>-1</sup> after two 10 days cultivations, which indicated that the absorption effect of *E. crassipes* was inferior to that of *A. calamus* and *C. indica*.

Fig. 3 shows that, after two 10 days cultivations, in the culture dishes of *A. calamus* and *C. indica*, absorption of ammonia nitrogen was >90% whereas, in the culture dishes of *E. crassipes*, absorption of ammonia nitrogen was ~50% and ~60%. Among the five types of BOW, 75% BOW had the highest absorption efficiency of ammonia nitrogen, irrespective of whether *A. calamus*, *C. indica* or *E. crassipes* was the macrophyte.

FT-IR spectroscopy is an important means to study the absorption of ammonia nitrogen by plants and its transformation within plants. The characterization results are shown in Fig. 4–8. We focused on analyses of the changes in the nitrogen-containing compounds  $NH_4^+$ -N,  $NO_2^-$ -N and  $NO_3^-$ -N. According to the literature,<sup>15,16</sup> the characteristic peaks (in cm<sup>-1</sup>) of  $NH_4^+$ -N are 1738 and 1459, whereas they are 1511 and 1332 for  $NO_2^-$ -N, and 1414 cm<sup>-1</sup> for  $NO_3^-$ -N.

The FT-IR spectra of the root, stem and leaf of *A. calamus* cultivated in BOW for 10 days are shown in Fig. 4. Fig. 4A reveals the FT-IR spectrum of the root of *A. calamus* cultivated in BOW for 10 days. Irrespective of the BOW concentration (0–100%), the characteristic peaks of  $NH_4^+$ -N,  $NO_2^-$ -N and  $NO_3^-$ -N appeared in the spectrum, thereby indicating that the root







Fig. 3 Absorption efficiency of ammonia nitrogen in three macrophytes. (A) Acorus calamus; (B) Canna indica; (C) Eichhornia crassipes.

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.







Fig. 5 FT-IR spectra of the root (A), stem (B) and leaf (C) of A. calamus cultivated in black and odorous water for 20 days.

absorbed NH<sub>4</sub><sup>+</sup>-N, NO<sub>2</sub><sup>-</sup>-N and NO<sub>3</sub><sup>-</sup>-N in BOW. Fig. 4B reveals the FT-IR spectrum of the stem of A. calamus cultivated in BOW for 10 days. Irrespective of the BOW concentration (0-100%), the characteristic peaks of NH4+-N, NO2--N and NO3--N appeared in the spectrum, thereby indicating that NH<sub>4</sub><sup>+</sup>-N, NO<sub>2</sub><sup>-</sup>-N and NO<sub>3</sub><sup>-</sup>-N in the root migrated to the stem. However, compared with the root, the intensity of the characteristic peak of  $NH_4^+$ -N at 1738 cm<sup>-1</sup> and 1459 cm<sup>-1</sup> in the FT-IR spectrum of the stem was weakened significantly, and the relative intensity of the peaks at 1511  $\text{cm}^{-1}$  and 1414  $\text{cm}^{-1}$  was enhanced. These findings indicated that the concentration of NO<sub>2</sub><sup>-</sup>-N and NO<sub>3</sub><sup>-</sup>-N in the stem was greater than that in the root. This phenomenon may have been caused by the transformation of NH<sub>4</sub><sup>+</sup>-N to NO<sub>2</sub><sup>-</sup>-N, and a small amount of NO<sub>2</sub><sup>-</sup>-N being transformed to NO<sub>3</sub><sup>-</sup>-N. Fig. 4C denotes the FT-IR spectrum of a leaf of A. calamus after 10 days of cultivation in BOW. Irrespective of the BOW concentration (0–100%), the characteristic peaks of  $NO_3^{-1}$ -N were the strongest, whereas the characteristic peaks of NH<sub>4</sub><sup>+</sup>-N and NO<sub>2</sub><sup>-</sup>-N were very weak. These observations suggested that NH<sub>4</sub><sup>+</sup>-N, NO<sub>2</sub><sup>-</sup>-N and NO<sub>3</sub><sup>-</sup>-N in the root migrated to the leaf through the stem, NH<sub>4</sub><sup>+</sup>-N and NO<sub>2</sub><sup>-</sup>-N were transformed, and the likely equation is:  $NH_4^+ N \rightarrow NO_2^- N \rightarrow NO_3^- N$ .

The FT-IR spectra of the root, stem and leaf of *A. calamus* cultivated in BOW for 20 days are shown in Fig. 5. Irrespective of the BOW concentration (0–100%), the characteristic peaks of  $NH_4^+$ -N,  $NO_2^-$ -N and  $NO_3^-$ -N were observed (Fig. 5A), thereby indicating that the root absorbed  $NH_4^+$ -N,  $NO_2^-$ -N and  $NO_3^-$ -N in BOW, but the  $NH_4^+$ -N concentration was stronger compared

with that at 10 days. Compared with Fig. 5B, the relative intensity of the peak at 1738 cm<sup>-1</sup> in Fig. 4B was reduced significantly. This observation indicated that, compared with 10 days previously, the NH<sub>4</sub><sup>+</sup>-N level in the stem had decreased significantly. In Fig. 5C, the peak at 1738 cm<sup>-1</sup> is present, but the peaks at 1511 cm<sup>-1</sup> and 1332 cm<sup>-1</sup> are absent. A possible reason is that the conversion of NH<sub>4</sub><sup>+</sup>-N to NO<sub>2</sub><sup>-</sup>-N was slower than the conversion of NO<sub>2</sub><sup>-</sup>-N to NO<sub>3</sub><sup>-</sup>-N in the leaf.

The FT-IR spectra of the root, stem and leaf of C. indica cultivated in BOW for 10 days are shown in Fig. 6. Fig. 6A reveals that NH<sub>4</sub><sup>+</sup>-N, NO<sub>2</sub><sup>-</sup>-N and NO<sub>3</sub><sup>-</sup>-N were absorbed in the root irrespective of the BOW concentration. At 100% BOW, the NH4+-N concentration in the root was the highest whereas, at the other BOW percentages, the NH4<sup>+</sup>-N concentration was much lower than that observed in A. calamus. In the stem of C. indica, residual NH<sub>4</sub><sup>+</sup>-N was present, but very little NO<sub>2</sub><sup>-</sup>-N was present in BOW (except at 100% BOW). These data indicated that the speed of migration and transformation of NO<sub>2</sub><sup>-</sup>-N was rapid. In Fig. 6C, the peak at 1332 cm<sup>-1</sup> (characteristic peak of NO<sub>2</sub><sup>-N</sup>) was present, but peaks at 1738, 1459 and 1511 cm<sup>-1</sup> (characteristic peaks of NH4<sup>+</sup>-N and NO2<sup>-</sup>-N) were absent, which indicated that a small amount of NO2--N remained in the leaf during migration and transformation. Compared with A. *calamus*, residual amounts of NH<sub>4</sub><sup>+</sup>-N and NO<sub>2</sub><sup>-</sup>-N were present. However, inorganic nitrogen in C. indica can be transformed rapidly to  $NO_3^{-}$ -N due to the straight-channel capillaries in C. indica.



Fig. 6 FT-IR spectra of the root (A), stem (B) and leaf (C) of Canna indica cultivated in black and odorous water for 10 days.







Fig. 8 FT-IR spectra of the root (A) and leaf (B) of *Eichhornia crassipes* cultivated in black and odorous water for 10 days and the leaf (C) cultivated for 20 days.



ig. 9 Scanning electron micrographs of dehydrated Canna indica.

After 20 days of cultivation in BOW,  $NH_4^+$ -N,  $NO_2^-$ -N and  $NO_3^-$ -N appeared in the root of *C. indica* (Fig. 7A). However, only  $NO_3^-$ -N was present in the stem and leaf:  $NH_4^+$ -N and  $NO_2^-$ -N were absent (Fig. 7B and C). A possible reason was that because the cultivation time was longer, the absorption capacity was weakened, and  $NO_2^-$ -N was transformed completely into  $NO_3^-$ -N, and the transformation rate of  $NH_4^+$ -N and  $NO_2^-$ -N was greater than their migration rate.

The FT-IR spectra of root and leaf of *E. crassipes* cultivated in BOW for 10 days, and that of the leaf cultivated in BOW for 20 days, are shown in Fig. 8. The migration and transformation performance of  $NH_4^+$ -N,  $NO_2^-$ -N and  $NO_3^-$ -N in *E. crassipes* were analogous to those of *A. calamus* and *C. indica*. The difference was that the concentrations of  $NH_4^+$ -N and  $NO_2^-$ -N were very low, which may have been due to the weak absorption of *E. crassipes*.

The capillary structure of aquatic plants is important for the migration of ammonia nitrogen, nitrite and nitrate. Scanning electron micrographs (Fig. 9) revealed many capillary structures in *C. indica*, and the tube walls were smooth. Three types of inorganic nitrogen compounds migrate in macrophytes and are transformed in the order  $NH_4^+$ ·N  $\rightarrow NO_2^-$ ·N  $\rightarrow NO_3^-$ ·N. There are many ammonia- and nitrite-oxidizing bacteria in macrophytes. They are responsible for the oxidation of ammonia (*i.e.*, ammonia nitrogen is oxidized to nitrite if sufficient oxygen is present) and nitrite (*i.e.*, nitrite is oxidized to nitrate by nitrite oxidoreductase). Both processes use oxygen as the electronacceptor.

Our study is based on biological nitrification, and the conversion mechanism of ammonia nitrogen is shown in eqn (1) and (2).

$$2NH_4^+ + 3O_2 \rightarrow 2NO_2^- + 2H_2O + 4H^+$$
(1)

$$2\mathrm{NO}_2^- + \mathrm{O}_2 \to 2\mathrm{NO}_3^- \tag{2}$$

The absorption performances of *A. calamus* and *C. indica* for  $NH_4^+$ -N were more robust than those of *E. crassipes* because *C. indica* and *A. calamus* grew rapidly after they were moved into wastewater, their height increased significantly, and the root system extended swiftly. At the end of our experiment, some

roots extended into the bottom of the water tank. Also, the biomass accumulation of *C. indica* and *A. calamus* was greater than that of *E. crassipes*.

#### 4 Conclusions

The migration process of NH<sub>4</sub><sup>+</sup>-N in A. calamus, C. indica and E. crassipes was detected via FT-IR spectroscopy. NH4+-N and  $NO_2^{-}N$  were transformed in the direction  $NH_4^{+}N \rightarrow NO_2^{-}N$  $\rightarrow$  NO<sub>3</sub><sup>-</sup>-N from the root to stem and leaf. A multitude of ammonia- and nitrite-oxidizing bacteria are present in macrophytes, and they are responsible for ammonia oxidation and nitrite oxidation. The migration rate of NH4+-N in C. indica was faster because of its regular and smooth capillaries. A. calamus and *C. indica* could absorb more NH<sub>4</sub><sup>+</sup>-N than *E. crassipes*. This was because C. indica and A. calamus grew rapidly upon transfer to wastewater, their height increased significantly, and the root system extended swiftly. The efficiency and quantity of transformation will be the focus of further research. Our study on the removal and transformation mechanism of ammonia nitrogen in BOW could be an important reference for other bodies of water.

### Data availability

The data used to support the findings of this study are available from the corresponding author upon request.

### Conflicts of interest

There are no conflicts of interest to declare.

#### Acknowledgements

This work was supported by Research Foundation of the Institute of Environment-friendly Materials and Occupational Health of Anhui University of Science and Technology (Wuhu) (ALW2020YF09).

### References

- 1 S. Sarigai, J. Yang, A. Zhou, *et al.*, Monitoring urban blackodorous water by using hyperspectral data and machine learning, *Environ. Pollut.*, 2021, **269**(10), 116166.
- 2 H. He, Q. Yu, C. Lai, *et al.*, The treatment of black-odorous water using tower bipolar electro-flocculation including the removal of phosphorus, turbidity, sulfion, and oxygen enrichment, *Front. Environ. Sci. Eng.*, 2021, **15**(2), 1–13.
- 3 L. Ye, S. Yang, P. Wu, *et al.*, Design and Application of Monitoring Station of Black and Odorous Water Body Based on Industrial Computer and Remote Computer, *IOP Conf. Ser.: Earth Environ. Sci.*, 2020, **508**(1), 012100.
- 4 P. Wang, Y. Weng, P. Zhuang, *et al.*, Analysis of Formation and Mechanisms of Black and Smelly River Water in Island Cities, *Meteorol. Environ. Res.*, 2018, **9**(6), 42–48.
- 5 Q. Ding, L.-H. Tang and D. Xie, Forming mechanism of black-odor of a campus lake, *Ind. Water Wastewater*, 2012, **43**(3), 28–30.
- 6 L. Marchand, M. Mench, D. L. Jacob, *et al.*, Metal and metalloid removal in constructed wetlands, with emphasis on the importance of plants and standardized measurements: a review, *Environ. Pollut.*, 2010, **158**(12), 3447–3461.
- 7 J. William, The role of water plant in water treatment, *Agric. Eng.*, 1986, 57(6), 9–10.
- 8 H. Qin, Z. Zhang, M. Liu, *et al.*, Site test of phytoremediation of an open pond contaminated with domestic sewage using

water hyacinth and water lettuce, *Ecol. Eng.*, 2016, **15**, 753–762.

- 9 Z. Y. Wu, D. R. Qiu, F. He, *et al.*, Effects of rehabilitation of submerged macrophytes on nutrient level of a eutrophic lake, *Chin. J. Appl. Ecol.*, 2003, **14**(8), 1351–1353.
- 10 S. Elisa, B. Marco, M. Marco, *et al.*, An ounce of prevention is worth a pound of cure: Managing macrophytes for nitrate mitigation in irrigated agricultural watersheds, *Sci. Total Environ.*, 2018, **647**, 301–312.
- 11 L. Huang, N. C. Wu, T. Tang, *et al.*, Enrichment and removal of nutrients in eutrophic water by aquatic macrophytes, *China Environ. Sci.*, 2010, **30**, 1–6.
- 12 Y. X. Zhong, H. Y. Hu and Y. Qian, Advances in utilization of macrophytes in water pollution control, *Tech. Equip. Environ. Pollut. Control*, 2003, 4(2), 36–40.
- 13 D. B. Sylvia, T. Samuel and T. Ludwig, Stabilizing the clearwater state ineutrophic ponds after biomanipulation: Submerged vegetation *versus* recolonization, *Hydrobiologia*, 2012, 689(1), 61–176.
- 14 Y. Yao, L. Z. Huang, S. Y. Chen, *et al.*, Purification effect of different submerged plants on nitrogen and phosphorus in water, *J. Zhejiang Agric. Sci.*, 2011, **4**, 789–792.
- 15 E. Ito, Y. J. Mergler, B. E. Nieuwenhuys, *et al.*, IR mechanistic studies on NO reduction with NH<sub>3</sub> in the presence of oxygen over cerium-exchanged mordenite, *J. Chem. Soc., Faraday Trans.*, 1996, **92**(10), 1799–1806.
- 16 H. Wang, L. Lv, D. H. Wang, et al., NO catalytic oxidation over copper manganese oxide at room temperature, J. Beijing Univ. Chem. Technol. (Nat. Sci. Ed.), 2010, 37(4), 24.