RSC Advances



View Article Online

View Journal | View Issue

PAPER

Received 21st April 2023

Accepted 28th April 2023

DOI: 10.1039/d3ra02637j

rsc.li/rsc-advances

Check for updates

Cite this: RSC Adv., 2023, 13, 14958

A sustainable metal and base-free direct amidation of esters using water as a green solvent[†]

Nanthini Rajendran,‡^a Kiruthigadevi Kamaraj,‡^a Saranya Janakiraman,^a Mary Saral,^a Pierre H. Dixneuf^{b*b} and Charles Beromeo Bheeter^b*^a

Herein, we report a simple and efficient synthetic approach for direct amidation of esters *via* C(acyl)–O bond cleavage without any additional reagents or catalysts, using only water as a green solvent. Subsequently, the reaction byproduct is recovered and utilized for the next phase of ester synthesis. This method emphasized metal-free, additive-free, and base-free characteristics making it a new, sustainable, and eco-friendly way to realize direct amide bond formation. In addition, the synthesis of the drug molecule diethyltoluamide and the Gram-scale synthesis of a representative amide are demonstrated.

Amide bond formation is an essential step in pharmaceutical industries for drug designing, synthesis of polymers and material products since amide bonds are an indispensable constituent in peptide derivatives, natural products, and drugs.¹ Ester derivatives can be powerful alternatives for amidation reactions rather than the commonly used organic acid halides due to their environment-friendly nature and ready availability. Oils, fatty acids, triesters of glycerol, and fruits are good sources of esters. Thus the methods of conversion of esters into amides are becoming significant.² Traditionally, most pharmaceutical industries follow the acid-amine coupling method with an overstoichiometric amount of coupling reagents and bases for amide bond formation.³

Wu and Beller *et al.* successfully developed a new catalytic method for the synthesis of amides using CO gas by amino carbonylation methodology.⁴ Another well-known approach in amide bond formation is the C(acyl)–O bond cleavage of esters *via* the catalytic pathway, using metal catalysts,⁵ (Scheme 1, Part A, B) strong base^{5,6} (Scheme 1, Part C), various additives,⁷ and often non-greener solvents.⁸ This method requires suitable substituents for activating esters⁹ and for effective conversion. In addition, aggressive organometallic reagents such as BuLi, AlMe₃, Al(O-*t*Bu) along with Ni(cod)^{5g} catalyst or strong bases like LiHMDS^{5a} and NaH¹⁰ are required for activating some esters. When highly activated esters are used, amidation is achieved using K₂CO₃ as a base with amines.¹¹

While the cleavage of esters yields carbonylative and decarbonylative¹² products, the selectivity is controlled intermittently

^aDepartment of Chemistry, School of Advanced Sciences, Vellore of Institute of Technology, Vellore-632014, TamilNadu, India. E-mail: charlesbheeter@vit.ac.in ^bUniversity of Rennes, ISCR, UMR CNRS 6226, 35000 Rennes, France

† Electronic supplementary information (ESI) available. See DOI: https://doi.org/10.1039/d3ra02637j

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence. Open Access Article. Published on 16 May 2023. Downloaded on 7/24/2025 8:04:47 AM.

by the nature of the ligands.^{5b,12c} For instance, Ni-(*N*-heterocyclic carbene) ligand gives carbonylative product¹³ while [Ni-bis-phosphine] ligand yields decarbonylative product.^{12e,14} Other non-noble metals also cleaved the C(acyl)–O bond of esters with hybrid ligands such as NHC with nitrogen donors in Mn(pincer)¹⁵ (Scheme 1, Part B) and La(OTf)₃ as lewis acid catalyst.¹⁶ Besides, the amidation process is achieved through transition metal complexes such as Pd–NHC,¹⁷*a,b* Ni–NHC, Pd-Phosphine ligands, Mn pincer complex,¹⁵ and heterobimetallic lanthanide sodium alkaloids.¹⁸ In all these methods, the need



Scheme 1 Different approaches towards C(acyl)–O bond cleavage of esters to produce amides. (A) Precious metal approach for amide bond formation from esters (i)^{5a}, (ii)^{7a} (B) base metal approach for amide bond¹⁸ (C) metal-free with a strong base approach for amide bond formation^{8a} (D) this work.

[‡] Contributed equally to this work.

for a catalyst, reagents, additives, and/or base, non-green solvents are prevalent.

Hence, an environmentally friendly methodology is necessary for direct amidation reactions, especially for the industrial synthesis of amides in large quantities. Herein, we are reporting amide bond formation via the coupling of inactivated esters with amines in the presence of water in an oil bath at 110 °C with the recovery of side product phenol, which is further used in the synthesis of ester moieties. Overall, the conversion of simple esters into amides under metal and base-free conditions using water as a green solvent leading to an atom-economical with reduced cost is expected to be excellent progress in the area of amide synthesis. To the best of our knowledge, a sustainable direct amidation of esters under metal-free, additive-free, and base-free conditions using water as a solvent has not been explored.

At the outset, we have outlined the model scheme for the amidation of esters using phenyl benzoate (Table 1, 1a) and benzylamine (Table 1, 2a) as the ester and amine sources, respectively. Initially, we preferred using heterogeneous catalysts over traditional ones as the former can be recycled and reused several times. Accordingly, the reaction was performed using silica material, namely KIT-6 (mesoporous silica nanoparticles 10-100 µm) and KIT-6 with metal precursor (Fe-KIT-6), which furnished an amidation product (Table 1, 3a) with 78% and 82% yield respectively (Table 1, entries 1-2). Then we moved to another silica material, SBA-15 (mesoporous silica nanoparticles $< 150 \mu m$), which yielded 60% of the product (3a). Subsequently, we tried the reaction with normal mesoporous silica 200-400 mesh (25 mg), which yielded 80% of the product (3a) using THF at 110 °C (Table 1, entries 3-4).

Interestingly, standard mesoporous silica produced a very good yield, better than SBA-15, KIT-6, and comparable to Fe-KIT-6. As the reaction proceeded well with ordinary mesoporous silica, we wanted to determine the catalyst's role in the reaction. Subsequently, we tested for the catalyst-free

Table 1	Reaction optimization ^{<i>a</i>} $\downarrow \qquad \qquad$			
	Entry	Catalyst	Solvent Yie	ld (%)
Entry	Catalyst		Solvent	Yield (%)
1	Fe KIT-6		THF	82
2	KIT-6		THF	78
3	SBA-15		THF	60
4	Silica		THF	80
5	Catalyst-free		THF	80
6	Catalyst-free		CH ₃ CN	78
7	Catalyst-free		Water	95
8	Catalyst-free		Ethanol	45
9	Catalyst-free		Methanol	32

^{*a*} Conditions: 0.2525mmol. phenyl benzoate benzylamine 0.5050 mmol, catalyst 25 mg, solvent 3 mL, 110 °C in oil bath, 12 h, isolated yields.

performance of the reaction with different solvents. Surprisingly, the reaction proceeded well without a catalyst. So, to study the catalyst-free reaction further, we selected two different types of solvents according to their nature. The polar aprotic solvents like tetrahydrofuran and acetonitrile yielded 80% and 78% products (Table 1, entries 5-6). On the other hand, polar protic solvents, ethanol, and methanol showed low yields of 45% and 32%, respectively (Table 1, entries 8-9). Interestingly, when we tested with a universal solvent, water, it was observed to form the desired product with an excellent yield of 95% after 12 h (Table 1, entry 7). Hence, we concluded that the amidation process of esters proceeds well without a catalyst using the universal solvent water. This approach is best as it is greener and more atom economical than all previous approaches known so far. After optimization, we screened different esters such as methyl benzoate, ethyl benzoate, iso-propyl-benzoate, benzyl benzoate, and phenyl benzoate. Under our standard reaction condition methyl and ethyl benzoate gave a yield of <5%, iso-propyl- and benzyl benzoate gave 9% and 13% yield respectively, whereas phenyl benzoate gave the desired product with a 95% yield (Scheme 2).

From this, we hypothesized that phenol could act as a good leaving group in phenylesters compared to others esters without coupling reagent and base. Also, the inherent nature of the phenoxy group in phenyl benzoate could be related to its ability to involve in hydrogen bonding interaction with water to facilitate the amidation reaction of esters in water.¹⁹ Accordingly, the affinity of the phenolic moiety of the phenyl esters in water plays an important role than in organic solvents.

In the first part of the substrate scope, we explored the activity of aromatic esters with aliphatic amines employing the novel methodology. Benzylamine-bearing electron-donating and electron-withdrawing substituents in the para position, such as 4-methyl benzylamine with phenyl ester, give 76% yield, 4-methoxy benzylamine afforded 89% yield and 4-chloro benzylamine with 84% yield (Table 2, 3b-3d). Then we introduced substituents in ester moieties such as 4-chloro-, 2-methyl-, 3methyl- and 2-methoxy-phenyl benzoate, which afforded corresponding amides with 94%, 94%, 96%, and 73% of yields respectively (Table 2, 3e-3h). Later, we introduced benzylaminebearing electron-withdrawing substituents in ortho positions, such as 2-fluoro benzylamine, which converted into respective amide with a moderate yield of 39% (Table 2, 3i). Next, we introduced the 2-methoxy (Table 2, 3j-3k), and 3-methoxy (Table 2, 31-3m) substituents in ester moieties, and we correlated it with different substituents in benzyl amines such as 2-



Scheme 2 Optimization of different esters.

 Table 2
 Substrate scope for amidation of aromatic esters^a



^{*a*} Reaction conditions: 1 esters (1 eq), 2 amines (1.2 eq), H₂O (3 mL), 110 °C, 12 h. ^{*b*} 1v (2.2 eq), 2v (1.0 eq).

chloro-, 2-fluoro-, 4-fluoro- and 4-chloro-benzylamine which afforded 72% to 96% yield (Table 2, 3**j**-3**m**).

Also, 4-methoxy benzylamine with 2-methyl phenyl benzoate, 1-phenyl naphtholate, and 4-chloro phenyl benzoate as ester moieties yielded 55% to 57% of respective amides (Table 2, 3n-**3p**). In the case of 4-chloro phenyl benzoate with 2-methoxy benzylamine and 4-nitro phenyl benzoate with 4-fluoro benzylamine afforded yields of 49% and 73%, respectively (Table 2, 3q-3r). Likewise, 4-bromo phenyl benzoate afforded 94% yield with benzylamine (Table 2, 3s), and 4-bromo phenyl benzoate afforded 89% yield with morpholine (Table 2, 3t). The reaction between phenyl 2,2 diphenyl acetate with benzylamine afforded a 48% yield (Table 2, 3u). But under the same optimized conditions, n-Boc piperazine was completely inactive towards phenyl esters (Table 2, 3v). Aliphatic cyclic amines were also utilized under the optimized reaction conditions. Pyrrolidine underwent the amidation process with 4-nitro-, 2-fluoro-, and 3methyl-phenyl benzoate affording 78%, 91% and 94% yield, respectively (Table 2, 3w-3y). The reaction of piperidine with phenyl benzoate, 4-fluoro-, and 4-methyl-phenyl benzoate provided 48%, 59%, and 98% of corresponding amides (Table 2, 3z, 3aa, 3ab). Morpholine reacted with phenyl benzoate to give a 48% yield while the reaction with 4-nitro phenyl benzoate afforded only a 27% yield (Table 2, 3ac, 3ad).

This may be due to the influence of the electronic nature of ester moieties in the amidation process. Particularly, when the alkyl chain of the amine was lengthened as in the reaction of phenethylamine with ester moieties such as 4-fluoro phenyl benzoate, 2 phenyl naphtholate and 3-fluoro phenyl benzoate giving corresponding amides, which gave a good yield of 63%, 63%, and 80% (Table 2, **3ae**, **3af**, **3ag**). Phenyl benzoate, 2-fluoro- and 2-methoxy-phenyl benzoate coupled with *n*-butylamine provided the corresponding amides with moderate to good yields of 38%, 85%, and 95% (Table 2, **3ah**, **3ai**, **3aj**).

In the second part of the substrate scope, we have explored the activity of heteroaromatic esters with aliphatic and aromatic amines *via* new methodology. When benzylamine is reacted with phenyl thiophene-2-carboxylate and phenyl furan-2carboxylate, 87%, and 88% yield respectively (Table 3, **3ak**, **3al**). Phenyl thiophene-2-carboxylate, when coupled with piperidine and morpholine gives 44%, and 56% yield (Table 3, **3am**, **3an**). To evaluate the scope of aromatic amines, a diverse range of heterocyclic esters was explored. Accordingly, phenyl furan-2-carboxylate coupled successfully with aniline, 4-toluidine, 4-bromo aniline, and 4-anisidine affording 54%, 30%, 40%, and 51% yield respectively (Table 3, **3ao**, **3ap**, **3aq**, **3ar**).

In the third part of the substrate scope, we explored the activity of aliphatic esters, amino acid ester with aliphatic and aromatic amines. A selection of fused aliphatic esters was employed for the reaction as they are vital components of bioactive compounds and useful intermediates in drug synthesis. Phenyl cyclopropane carboxylate coupled with benzylamine yields 96% of the product (Table 4, **3as**). Phenyl cyclopropane carboxylate reacts with aniline provide 42% to 81% of yield (Table 4, **3at**, **3au**, and **3av**). Phenyl cyclobutane carboxylate reacts with aniline affording a corresponding yield of 74% (Table 4, **3aw**), and the reaction with morpholine yields a trace amount of product (Table 4, **3ax**).

Also, this reaction was influential in the context of the direct synthesis of biologically relevant compounds like *tert*-butyl(S)-2-(phenyl carbamoyl)pyrrolidine-1-carboxylate (Table 4, **3ay**) from n-boc L-proline an amino acid with 52% yield. Significantly, we prepared the drug, diethyltoluamide (Table 4, **3az**), an insect repellent,²⁰ by aminolysis of *meta*-methyl phenyl benzoate with diethylamine to give a moderate yield of 45%.

To prove the proposed methodology's environment-friendly nature, the side product phenol (Scheme 3) was recovered as it could be used further in the preparation of ester derivatives (see the ESI† for more details).

Having demonstrated that water is an effective solvent for a range of direct amidation products employing the present



 a Reaction conditions: 1 esters (1 eq), 2 amines (1.2 eq), H_2O (3 mL), 110 °C, 12 h.

 Table 4
 Substrate scope for amidation of aromatic amine^a



^a Conditions: ester (1 eq), amine (1.2 eq), solvent 3 mL, 110 °C, 12 h.



Scheme 3 Recycling of side product.



methodology, we sought to exemplify the method on a Gram scale. We assessed the Gram scale synthesis (Scheme 4) with a mixture of phenyl benzoate **1a** (**1g**, one equiv), benzylamine **2a** (810 mg, 1.5 equiv), which was stirred in the presence of H_2O at 110 °C for 12 h to afford the formation of 953 mg of benzyl benzamide **3a** with 90% yield.

We have found a metal-free, base/additive-free amidation protocol under aqueous conditions. Based on the substrate scope and Gram scale experiment, we show that the aromatic, aliphatic, and amino acid ester derivatives with aliphatic and aromatic amines exhibit good activity to give the corresponding amidation products. Furthermore, successful recovery of side product for further use in ester derivative synthesis is done for making the discussed approach sustainable. This is the first report on ester acyl(C–O) bond cleavage for amide formation with water as an eco-friendly benign solvent.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

NR and KK acknowledge VIT for providing a fellowship. Authors acknowledge VIT for seed grants (SG20210184), infrastructure

and instrumentation facilities, and SERB (CRG/2021/07286) for research funding.

Notes and references

- (a) M. T. Sabatini, L. T. Boulton, H. F. Sneddon and T. D. Sheppard, *Nat. Catal.*, 2019, 2, 10–17; (b) C. L. Allen and J. M. J. Williams, *Chem. Soc. Rev.*, 2011, 40, 3405–3415; (c) E. Valeur and M. Bradley, *Chem. Soc. Rev.*, 2009, 38, 606–631; (d) G. A. Filonenko, R. van Putten, E. J. M. Hensen and E. A. Pidko, *Chem. Soc. Rev.*, 2018, 47, 1459–1483; (e) D. G. Brown and J. Boström, *J. Med. Chem.*, 2016, 59, 4443–4458.
- 2 (a) R. Takise, K. Muto and J. Yamaguchi, Chem. Soc. Rev., 2017, 46, 5864–5888; (b) M. Ishikawa, Y. Warita, E. Takahisa and Y. Ohkubo, in Comprehensive Natural Products II, ed. H.-W. Liu and L. Mander, Elsevier, Oxford, 2010, pp. 595–629, DOI: DOI: 10.1016/B978-008045382-8.00107-6; (c) A. J. Meléndez-Martínez, P. Mapelli-Brahm, D. Hornero-Méndez and I. M. Vicario, in Carotenoid Esters in Foods: Physical, Chemical and Biological Properties, The Royal Society of Chemistry, 2019, pp. 1–50, DOI: 10.1039/9781788015851-00001.
- 3 (a) N. G. Anderson, in *Practical Process Research and Development*, ed. N. G. Anderson, Academic Press, Oxford, 2nd edn, 2012, pp. 89–120, DOI: DOI: 10.1016/B978-0-12-386537-3.00004-6; (b) R. M. de Figueiredo, J.-S. Suppo and J.-M. Campagne, *Chem. Rev.*, 2016, 116, 12029–12122; (c) F. Albericio and A. El-Faham, *Org. Process Res. Dev.*, 2018, 22, 760–772.
- 4 L. Wu, X. Fang, Q. Liu, R. Jackstell, M. Beller and X.-F. Wu, *ACS Catal.*, 2014, 4, 2977–2989.
- 5 (a) T. Zhou, G. Li, S. P. Nolan and M. Szostak, Org. Lett., 2019, 21, 3304-3309; (b) T. Ben Halima, J. K. Vandavasi, M. Shkoor and S. G. Newman, ACS Catal., 2017, 7, 2176-2180; (c) Mondal, M. Subaramanian, A. Nandakumar and A. Balaraman, Org. Lett., 2018, 20, 3381-3384; (d) E. M. B. Kurosawa, R. Isshiki, K. Muto and J. Yamaguchi, J. Am. Chem. Soc., 2020, 142, 7386-7392; (e) A. H. Dardir, P. R. Melvin, R. M. Davis, N. Hazari and M. Mohadjer Beromi, J. Org. Chem., 2018, 83, 469-477; (f) K. Kim and S. H. Hong, J. Org. Chem., 2015, 80, 4152-4156; (g) L. Hie, N. F. Fine Nathel, X. Hong, Y.-F. Yang, K. N. Houk and N. K. Garg, Angew. Chem., Int. Ed., 2016, 55, 2810-2814; (h) Y.-L. Zheng and S. Newman, ACS Catal., 2019, 9, 4426-4433; (i) W. B. Wang and E. J. Roskamp, J. Org. Chem., 1992, 57, 6101-6103; (j) G. Li and M. Szostak, Nat. Commun., 2018, 9, 4165; (k) J. L. Vrijdag, F. Delgado, N. Alonso, W. M. De Borggraeve, N. Pérez-Macias and J. Alcázar, Chem. Commun., 2014, 50, 15094-15097.
- 6 (a) L. H.-G. Kim BoRam, K. Seung-Beom, G. H. Sung, J.-J. Kim, J. K. Park, L. Sang-Gyeong and Y. Y. Jin, Synthesis, 2012, 44, 42–50; (b) R. Zhang, W.-Z. Yao, L. Qian, W. Sang, Y. Yuan, M.-C. Du, H. Cheng, C. Chen and X. Qin, Green Chem., 2021, 23, 3972–3982; (c) W. I. Nicholson, F. Barreteau, J. A. Leitch, R. Payne, I. Priestley, E. Godineau, C. Battilocchio and D. L. Browne, Angew.

Chem., Int. Ed., 2021, **60**, 21868–21874; (*d*) T. Ohshima, Y. Hayashi, K. Agura, Y. Fujii, A. Yoshiyama and K. Mashima, *Chem. Commun.*, 2012, **48**, 5434–5436.

- 7 (a) G. Li and M. Szostak, Synthesis, 2020, 52, 2579-2599; (b) N. Caldwell, C. Jamieson, I. Simpson and A. J. B. Watson, Chem. Commun., 2015, 51, 9495-9498; (c) A. Ramirez, B. Mudryk, L. Rossano and S. Tummala, J. Org. Chem., 2012, 77, 775-779; (d) C. G. McPherson, N. Caldwell, C. Jamieson, I. Simpson and A. J. B. Watson, Org. Biomol. Chem., 2017, 15, 3507-3518; (e) D. A. Rankic, C. M. Stiff, C. W. am Ende and J. M. Humphrey, J. Org. Chem., 2017, 82, 12791-12797; (f) K. E. Price, C. Larrivée-Aboussafy, Lillie, R. W. McLaughlin, B. M. I. Mustakis, K. W. Hettenbach, J. M. Hawkins and R. Vaidyanathan, Org. Lett., 2009, 11, 2003-2006.
- 8 (a) Y. Liu, S. Shi, M. Achtenhagen, R. Liu and M. Szostak, *Org. Lett.*, 2017, **19**, 1614–1617; (b) M. M. Rahman, G. Li and M. Szostak, *J. Org. Chem.*, 2019, **84**, 12091–12100; (c) P. V. Ramachandran, H. J. Hamann and S. Choudhary, *Org. Lett.*, 2020, **22**, 8593–8597.
- 9 (a) L. Ling, C. Chen, M. Luo and X. Zeng, Org. Lett., 2019, 21, 1912–1916; (b) C. W. Cheung, N. Shen, S.-P. Wang, A. Ullah, X. Hu and J.-A. Ma, Org. Chem. Front., 2019, 6, 756–761; (c) L.-Y. Chen and M.-F. Wu, Synthesis, 2019, 51, 1595–1602; (d) T. Ooi, E. Tayama, M. Yamada and K. J. S. Maruoka, Synlett, 1999, 6, 729–730; (e) X. Yang and V. B. Birman, Org. Lett., 2009, 11, 1499–1502; (f) Y. Dongqiang and X. Jiaxi, Curr. Microw. Chem., 2020, 7, 74–82.
- S. A. Rzhevskiy, A. A. Ageshina, G. A. Chesnokov,
 P. S. Gribanov, M. A. Topchiy, M. S. Nechaev and
 A. F. Asachenko, *RSC Adv.*, 2019, 9, 1536–1540.

- 11 S. Karthik, R. Sreedharan and T. Gandhi, *ChemistrySelect*, 2019, 4, 175–180.
- 12 (a) K. Amaike, K. Muto, J. Yamaguchi and K. Itami, J. Am. Chem. Soc., 2012, 134, 13573-13576; (b) K. Muto, J. Yamaguchi, D. G. Musaev and K. Itami, Nat. Commun., 2015, 6, 7508; (c) A. Chatupheeraphat, H.-H. Liao, W. Srimontree, L. Guo, Y. Minenkov, A. Poater, L. Cavallo and M. Rueping, J. Am. Chem. Soc., 2018, 140, 3724-3735; (d) T. B. Boit, A. S. Bulger, J. E. Dander and N. K. Garg, ACS Catal., 2020, 10, 12109-12126; (e) L. Guo and M. Rueping, Acc. Chem. Res., 2018, 51, 1185-1195.
- 13 T. Ben Halima, J. Masson-Makdissi and S. G. Newman, *Angew. Chem., Int. Ed.*, 2018, 57, 12925–12929.
- 14 C. A. Malapit, M. Borrell, M. W. Milbauer, C. E. Brigham and M. S. Sanford, J. Am. Chem. Soc., 2020, 142, 5918–5923.
- 15 Z. Fu, X. Wang, S. Tao, Q. Bu, D. Wei and N. Liu, J. Org. Chem., 2021, 86, 2339–2358.
- 16 H. Morimoto, R. Fujiwara, Y. Shimizu, K. Morisaki and T. Ohshima, Org. Lett., 2014, 16, 2018–2021.
- 17 S. Shi and M. Szostak, ChemComm, 2017, 53, 10584-10587.
- 18 Z. Li, C. Wang, Y. Wang, D. Yuan and Y. Yao, Asian J. Org. Chem., 2018, 7, 810–814.
- 19 R. Parthasarathi, V. Subramanian and N. Sathyamurthy, J. Phys. Chem., 2005, 109, 843–850.
- 20 D. L. Sudakin and T. Osimitz, in *Hayes' Handbook of Pesticide Toxicology*, ed. R. Krieger, Academic Press, New York, 3rd edn, 2010, pp. 2111–2125, DOI: DOI: 10.1016/B978-0-12-374367-1.00098-7.