RSC Advances

PAPER

Cite this: RSC Adv., 2023, 13, 21327

Comparative antisickling and antioxidant activities of Pseudobombax ellipticum cultivars in relation to their metabolite profiling using LC/MS

Ahmed S. Mohamed[,](http://orcid.org/0009-0009-5728-9187) \mathbb{D}^{*a} Omnia Y. Abd El Dayem, ^b Ali M. El Shamy,^a Fatma S. El Sakhawy^a and Rania A. El Gedaily^a

Pseudobombax ellipticum is native to South America and is cultivated worldwide mostly for its medicinal benefits. The plant is used traditionally in respiratory disorders such as dry cough, in the treatment of fever and stomach pain, and as an antimicrobial and analgesic. The antisickling and antioxidant effects of the flowers of P. ellipticum (Kunth) Dugand (red) and P. ellipticum cultivar alba (white) were compared using an in vitro assay in 2% sodium metabisulfite sickling induction model, DPPH, and metal chelation assays. Both red and white flowers exhibited antioxidant and antisickling activities. In DPPH assay, lower IC₅₀ (34.89 \pm 0.98 and 53.28 \pm 1.14 μ g mL⁻¹) in red and white flowers respectively were detected relative to Trolox as a positive control (56.82 \pm 0.87 µg mL^{−1}). Comparable metal chelation activity (81.4 and 77.8 µM EDTA equivalent/mg) was detected in red and white flowers of both cultivars respectively. The average readings of the "reversal of sickling test "revealed a decrease in sickling percent from 49% to 15% in red flowers and to 18% in white flowers. Also, polymerization inhibition rate was increased from 0.34 to 1 and to 0.92 in red and white flowers respectively. Total phenolics, flavonoids and anthocyanins were quantified in red and white flowers as (163.9, 43.13 mg gallic acid equivalent/g extract), (71.92, 34.5 mg rutin equivalent/g extract) and (127.0, 85.9 mg pelargonidine-3-mono glucoside equivalent/kg extract), respectively. Liquid chromatography mass spectrometry (LC-MS) analysis was further employed for detection and identification of anthocyanins in flower extracts. Eight new anthocyanins were identified for the first time in genus Pseudobombax. These results reveal the potential role for both red and white flower extracts as possible antisickling agents in sickle cell anemia management. **PAPER**
 (a) Cheek for updates

Comparative antisickling and antioxidant activities

Comparative of *Pseudobombax* ellipticum cultivars in relation

Series *nsc Anche* 2003. Ahmed S. Mohamed, O^{ne} Omnia Y. Abd El Dayem

Received 18th May 2023 Accepted 6th July 2023

DOI: 10.1039/d3ra03312k

rsc.li/rsc-advances

1. Introduction

The family Bombacaceae is a small family of flowering plants termed the Kapok family, referring to the cottony white fibers (bombax or kapok) surrounding their seeds.^{1,2} They are characterized by their unique brush-like flowers in late winter.¹ Members of this family are used traditionally for the management of respiratory illnesses, gastrointestinal ailments and diabetes.^{3,4} The genus Pseudobombax comprises 30 species; P. ellipticum (Kunth) Dugand also known as Bombax ellipticum is cultivated in Egypt.⁵ Its flowers show beautiful 5-inch long red (P. ellipticum (Kunth) Dugand) or white (P. ellipticum cultivar alba Hort.) stamens, looking very much like a shaving brush.³ Few reports were found concerning the biological activities and chemical composition of P. ellipticum. Aqueous and ethanol extracts of the cortex exhibited significant antioxidant activity. 6 From its stem bark, β -lupeol was

isolated and showed potent gastro-protective effect.⁷ Cyanidin-3,5-diglucoside was isolated from the flowers.² Sickle cell anemia (SCA) is the most common genetic blood disorder worldwide.⁸ According to estimates, Africa accounts for 75– 85% of children born with SCA. Hemoglobin S (HbS) is produced instead of normal adult haemoglobin due to a mutation in the haemoglobin beta gene (HBB).⁹ Normal RBCs have a 120 day lifespan and a flexible biconcave disk-like shape that facilitates easy passage through the microvasculature¹⁰ while under hypoxic conditions, HbS polymerizes, resulting in "sickle cells," hard and deformed RBCs with shortened life span (10–20 days) that cause decreased microcirculation and hemolysis.¹¹ Hence, antioxidants are very important in SCA management.^{12,13} Pharmaceuticals that are now available for treating SCA are either not sufficiently effective, not entirely safe, or too expensive warranting for developing effective, safe, and affordable drugs from natural sources.¹⁴ Interestingly, Ceiba pentandra (L.) Gaertn (Bombacaceae) is used traditionally in Congo to treat SCA and it's aqueous bark extract exhibited significant antisickling and antithrombing activities.¹⁵ Also, previous reports attributed the antisickling effect of several plants to their high

a Department of Pharmacognosy, Faculty of Pharmacy, Cairo University, Kasr Al-Aini St., Cairo 11562, Egypt. E-mail: ahmed.abobakr@pharma.cu.edu.eg

b Clinical and Chemical Pathology Department, Faculty of Medicine, Cairo University Al-Saray St., El Manial Cairo 11956, Egypt

RSC Advances **Paper** Paper **Paper Paper Paper**

anthocyanin content, present mainly in the flowers.^{16,17} As little data was available concerning P. ellipticum flowers phytochemical makeup especially anthocyanins, and further prompted by the well reported anthocyanins antisickling potential, this study aims to identify phytoconstituents especially anthocyanins of potential effects in SCA using LC/MS. The antioxidant and antisickling activities of both P. ellipti cum red flowers (PR) as well as *P. ellipticum* cultivar alba white flowers (PW) were compared in relation to their metabolite fingerprint.

2. Material and methods

2.1. Blood samples

The blood utilized in this study was drawn from patients known to have sickle cell anemia (Hemoglobin SS) from the hematology clinic at Cairo University Children Hospital "Abu L-Reesh Mounira". Informed consents were obtained from human participants of this study and all experiments were approved by the national ethic committee (No. MP3122) at Cairo University. Blood samples withdrawn from the selected patients were confirmed to contain homozygous (SS) red blood cells by haemoglobin electrophoresis carried out on cellulose acetate at pH 8.5. Samples were kept at 4 °C once they were determined that they included SS red blood cells.¹⁶

2.2. Plant material

P. ellipticum red flowers (PR) and P. ellipticum cultivar alba white flowers.

(PW) were collected from Mostafa El Abed botanical garden, during May 2020. Dr Mohamed El Gibali, Senior Botanist & Consultant at Orman Botanic Garden, Giza, Egypt, kindly carried out the plant identification. A vouchered specimen of the plant, with serial number (20.09.2022) was kept at the Pharmacognosy Department, Faculty of Pharmacy, Cairo University.

2.3. Preparation of flower extracts for antisickling assay

For preparation of extracts, fresh flowers $(50 g)$ from each plant were dissected by scissors to small pieces and separately extracted by 200 mL 100% methanol in 1 : 4 solid/solvent ratio using ultrasonic for 2 h. Extracts were then filtered via cotton pad and evaporated using rotavapor at temperature of 30 °C that yielded 3 g residue for red flowers and 2.8 g residue for white flowers which were kept till further anaylsis.¹⁸

2.4. Preparation of anthocyanins enriched extracts (AEE)

For another (50 g) of fresh flowers, 200 mL acidified methanol (0.1% formic acid in methanol) was used in a ratio of 1 : 4 solid/ solvent and extraction was done in the dark, in amber colored bottles kept in the refrigerator to avoid against anthocyanins degradation. Extraction was done for 3 days followed by sonication for 10 minutes prior to filtration. Extract was evaporated using rotavapor at temperature of 30 °C to yield 2.13 g residue for red flowers and 1.75 g residue for white flowers and kept at -20 °C till further analysis.^{18,19}

2.5. Total phenolics assay

Total phenolics were measured in flower samples using Folin-Cioacalteu assay.²⁰ A stock solution of 1 mg mL⁻¹ of gallic acid in methanol was prepared as the standard, and seven serial dilutions were made at concentrations of 1000, 800, 600, 400, 200, 100, and 50 μg mL⁻¹. Samples from the AEE of white and red flowers were prepared in a concentration of 4 mg mL^{-1} in methanol. Multimode plate reader (Fluostar Omega, BMG LABTECH, Germany) was used to record the outcomes. Six replicates of the 7 standards and 2 samples were pipetted into the wells of the plate measured at 630 nm. The average of the absorbance readings of the 6 replicates for gallic acid standard was recorded and calibration curve was established. From the established calibration curve, the linear regression equation was extracted for gallic acid standard (Fig. 1A). By substitution in the linear regression equation, the total phenolics were detected. RSC Advances

anthose on each the contentry P. effective former 2.5, Total phenolies assay

the right can voo a contentry contentry P. effective former 2.5, Total phenolies were measured in flower samples as

the right pr

2.6. Total flavonoids assay

Total flavonoids were determined using colorimetric determination of flavonoids – AlCl₃ complex.²¹ A stock solution of 1 mg mL^{-1} of rutin in 100% methanol was prepared as the standard, and seven serial dilutions were made at concentrations of 1000, 600, 400, 200, 100, 50 and 10 µg mL⁻¹. Multimode plate reader (Fluostar Omega, BMG LABTECH, Germany) was used to record the outcomes. Six replicates of the 7 standards and 2 samples were pipetted into the wells of the plate measured at 420 nm. The average of the absorbance readings of the 6 replicates for rutin standard was recorded and calibration curve was established. From the established calibration curve, the linear regression equation was extracted for rutin standard (Fig. 1B). By substitution in the linear regression equation, the total flavonoids were detected.

2.7. Total anthocyanins assay

Using potassium chloride (0.025 M, pH 1.0) and sodium acetate (0.4 M, pH 4.5) buffer systems, the total anthocyanins content was measured using the pH-differential method.²² Each buffer solution (9 mL) was combined with 1 mL of AEE of red and white flowers. The absorbance (Abs) was determined using UV/ Vis spectrophotometer, Jenway, England set at 520 and 700 nm, respectively.

2.8. Liquid chromatography-mass spectrometry (LC-MS) profiling

An Agilent LC-MS system composed of an Agilent 1290 Infinity II UHPLC coupled to an Agilent 6545 ESI-Q-TOF-MS in positive mode was used to obtain Ultra-high-performance liquid chromatograms. Aliquots (1 μ L) of methanolic extracts (5 mg mL⁻¹ in MeOH) were analyzed on a Kinetex phenyl-hexyl $(1.7 \mu m, 2.1)$ \times 50 mm) column. Isocratic elution of 90% A (A: 100% H₂O + 0.1% formic acid) for 1 min followed by 6 min linear gradient elution to 100% B (95% MeCN + 5% H₂O + 0.1% formic acid) with a flow rate of 0.4 mL min^{-1} . ESI conditions were set with the capillary temperature at 320 °C, source voltage at 3.5 kV and

Fig. 1 The established calibration curves with the extracted linear regression equations for gallic acid (A), rutin (B) and EDTA (C) standards.

a sheath gas flow rate of 11 L min⁻¹. Ions detected in the full scan at an intensity above 1000 counts at 6 scans per s, with an isolation width of 1.3 \sim m/z, a maximum of 9 selected precursors per cycle and using ramped collision energy $(5 \times m/z/100 +$ 10 eV). Purine C₅H₄N₄ [M + H]⁺ ion (*m*/z 121.050873) and hexakis $(1H, 1H, 3H$ -tetrafluoropropoxy)-phosphazene $C_{18}H_{18}F_{24}N_3O_6P_3$ [M + H]⁺ ion (*m*/z 922.009798) were used as internal lock masses.²³ Metabolites were identified based on their retention times, molecular formulae and their fragmentation patterns, compared to earlier reported data aided with GNPS spectral library search and Sirius.

2.9. Antioxidant activity

2.9.1. 2,2-Diphenyl-1-picrylhydrazyl (DPPH) method. One of the most popular techniques for evaluating antioxidant activity is DPPH based in color change in DPPH solution to turn from deep violet color to light yellow color during the reaction. The EC50 of the compounds is used to display the assay's results.²⁴ In order to determine a range of concentrations within which the inhibitory concentration 50 (IC_{50}) lies, initial solutions of red and white flowers AEE were prepared at concentration spanning from 1000 and 100 µg mL⁻¹ in methanol. Extracts that exceeded 50% inhibition in any of the initial screening step,

concentrations were serially diluted to provide 5 concentrations. Briefly, 100 μ L of freshly prepared DPPH reagent $(0.1\%$ in methanol) were added to 100 μ L of the samples (dissolved in methanol) in 96 wells plate $(n = 6)$, the reaction was incubated for 30 min in dark at room temperature. Decrease in DPPH colour intensity was assessed at 540 nm using a Multimode plate reader (Fluostar Omega, BMG LABTECH, Germany). Trolox was used as standard prepared at 100, 80, 60, 50, 40, 20 and 10 µg mL^{-1} . The results of six separate trials were combined to produce a mean and standard deviation (SD) for each sample. By converting the concentrations to their logarithmic value and choosing the nonlinear inhibitor regression equation log(inhibitor) vs. normalised response (variable slope equation), the data were analyzed using Microsoft Excel®, and the IC_{50} value was determined using GraphPad Prism $5\degree$.²⁵ The IC₅₀ was determined from preliminary screening to be less than 100 µg mL⁻¹. For each sample, the following serial dilutions were created: 100, 80, 60, 50, 40, 20 and 10 μ g mL⁻¹.

2.9.2. Metal chelation assay. Metal chelation assay is based on that ferrozine (3-(2-pyridyl)-5,6-diphenyl-1,2,4-triazine-p,p0 disulfonic acid monosodium salt hydrate) can bind to $Fe²⁺$ ions to produce a blue color. The iron-ferrozine complex's blue color is attenuated upon binding of compounds and/or extracts to iron $(Fe²⁺)$. Using EDTA as a positive control, the colour reduction is used to gauge the compound's iron chelating action.²⁶ Serial dilutions (11) of 0.1 mM EDTA stock solution in water were created at concentrations of 5, 10, 15, 20, 25, 30, 35, 40, 50, 60, and 70 μ M. Samples of enriched extracts from red and white flowers were prepared at a concentration of 0.625 mg mL^{-1} in methanol. The assay was conducted using the Santos et al. method,²⁶ with a few minor adjustments for microplates. In a 96well plate $(n = 6)$, 20 µL of freshly made ferrous sulphate (0.3 mM) was combined with 50 μ L of the samples. Then, each well received 30 µL of ferrozine (0.8 mM). For 10 minutes, the reaction mixture was incubated at room temperature. The reduction in the intensity of the generated colour was measured at 562 nm and percentage inhibition was calculated using this formula $=$ (average absorbance of blank − average absorbance of the test/ average absorbance of blank) \times 100. The multimode plate reader (Fluostar Omega, BMG LABTECH, Germany) was used to record the results. From the established calibration curve plotted from % inhibition of different concentrations of EDTA standard, the linear regression equation was extracted (Fig. 1C) and the results of the samples were presented as μ M EDTA equivalent/mg sample. After subtraction from the blank (reagent $+$ solvent), the average of the readings of the 6 replicates was taken.

2.10. Biological assay

2.10.1. Sickling reversal test. Using previously established methods, the plants' capacity to reverse the sickling status of the RBCs was determined.16,27 Sickle cell blood was washed twice using five times the volume of phosphate-buffered saline (PBS) with a pH of 7.4 by centrifugation at 4000 rpm for three minutes and the flower's total extracts were resuspended in absolute ethanol in a concentration of (0.25, 0.5 and 1 mg mL⁻¹). For the control test, 50 µL of washed blood (WB) were added to 950 μ L PBS into a clean Eppendorf tube and to 450 μ L PBS and mixed with 500 μ L of freshly prepared 2% sodium metabisulfite into another clean one. For the inhibition test, 50 μ L of WB were added to 400 μ L PBS followed by 50 μ L from the flower's total extracts and mixed with 500 μ L of freshly prepared 2% sodium metabisulfite into clean Eppendorf tubes. Control Tubes represent the initial percentage of sickling and sickling percentage after induction with sodium metabisulfite respectively. Ten µL from each mixture were spotted on a microscope slide covered with a cover slip. To seal the edges of the cover slip, paraffin wax was carefully applied and thus completely preventing air entry. By counting normal and sickled red cells under a $40\times$ light microscope, the ability of the flowers extracts to reverse the sickling of RBCs was observed and documented after incubation of all slides for 2 h at 37 \degree C. Morphologically, the red blood cells were classified as normal or sickled by observing their shapes. The elongated, C-like, or wrinkled shapes were considered sickled while biconcave or rounded shapes were taken to be normal. Using the following formula, the percentage sickled red blood cells was calculated: Red blood cells percent of sickling $(\%)$ = (number of counted sickled red blood cells divided by the total number of counted red blood cells) \times 100. For each concentration, the experiment was conducted three times.

2.10.2. Polymerization inhibition test. Following a previous procedure, the polymerization inhibition test was conducted.¹¹ In this procedure, the turbidity of the polymerizing solution of RBCs at a wavelength of 700 nm at a temperature of 26 °C was measured. After adding 0.88 mL of freshly made 2% sodium metabisulfite, into a clean cuvette, 0.1 mL of PBS and 0.02 mL of sickled blood, which had been washed twice in five volumes of PBS (1 mL of blood in 5 mL of PBS) and then resuspended in PBS at a ratio of 1 : 1 was added. This serves as control test. For the inhibition test, 0.1 mL of PBS was substituted with 0.1 mL of the flower's total extracts that also were resuspended in absolute ethanol in a concentration of (0.25, 0.5 and 1 mg mL^{-1}). For 60 minutes, the absorbance was measured at 700 nm immediately and at periodic intervals. Using the following formula, the rate of polymerization in percentage was computed: rate of polymerization $(R_p) = |(\text{final})|$ absorbance – initial absorbance)/60] \times 100. For each concentration, the experiment was repeated three times. At time zero $(i.e.,$ immediately after addition of sodium metabisulfite), the initial absorbance of the polymerizing solution of sodium metabisulfite was measured and subtracted from the final absorbance taken at the end of 60 min. Dividing the resulting value by 60 gives the rate of polymerization inhibition. The statistical significance of difference was calculated by GraphPad Prism 9.5.1 using analysis of variance where value of $p < 0.05$ was considered significantly different. RSC Advances Article on Experimental control and the method on 14 AM. This article is limited to the control of the

3. Results

3.1. Total phenolics, flavonoids and anthocyanins determination

Total phenolics expressed as gallic acid/equivalent were detected at 163.9 \pm 6.05 and 43.13 \pm 4.2 mg g⁻¹ in red and white flower extracts, respectively. Likewise, higher flavonoids level expressed as rutin/equivalent was detected in red flowers at 71.92 \pm 4.0 versus 34.5 \pm 3.8 mg g⁻¹ extract in white flowers. Considering the abundance of anthocyanins in flowers, their levels were estimated as mg pelargonidine-3-mono glucoside. The total anthocyanins content was detected at 127.0 and 85.9 mg kg⁻¹ in red and white flowers, respectively.

3.2. Liquid chromatography-mass spectrometry (LC-MS) profiling of flower extracts

LC-MS was further used to identify metabolites, mainly anthocyanins as depicted in (Fig. 2). A total of 10 peaks were identified in both flowers (Table 1). Of the 9 anthocyanins detected (Fig. 3), only cyanidin-3,5-di-O-glucoside was previously identi fied.² Cyanidin-3,5-di-O-glucoside (peak 2, 611 MS1 m/z), cyaniding O-hexoside (peak 5, 449 MS1 m/z), cyaniding Orutinoside (peak 6, 595 MS1 m/z) and cyanidin O-deoxyhexoside (peak 7, 433 MS1 m/z) produced a characteristic fragment ion at m/z 287 after the loss of a diglucose moiety, a hexose moiety, (a deoxyhexose and hexose) moiety and a deoxyhexose moiety, respectively.^{28,29} The 100% abundance of m/z at 287 indicated it as cyanidin aglycone in these glycosidic conjugates. Another aglycone was detected in (peak 1, 579 MS1 m/z) with a fragment

Fig. 2 Base peak chromatograms of LC/MS analysis of P. ellipticum (Kunth) Dugand (A) and P. ellipticum cultivar alba Hort. (B) flower's anthocyanins enriched extracts (AEE) in positive mode on an Agilent LC-ESI-Q-TOF-MS instrument.

ion at m/z 271 after the loss of a (deoxyhexose and hexose) moiety. The m/z at 271 with 100% intensity indicates pelargonidin aglycone leading to peak 1 identification as pelargonidin O-deoxyhexose hexoside.²⁸ Delphinidin O-hexoside (peak 3, 465 MS1 m/z) yielded a fragment ion at m/z 303 resulting from the loss of hexose moiety.³⁰ The m/z at 303 with 100% intensity indicates delphinidin aglycone. Petonidin O-hexoside (peak 8, 479 MS1 m/z) and petonidin O-deoxyhexose hexoside (peak 9, 625 MS1 m/z) produced a fragment ion at m/z 317 after the loss of a hexose moiety and (a deoxyhexose and hexose) moiety, respectively.²⁸ The m/z at 317 with 100% intensity indicates petunidin aglycone. Finally, peonidin (5-methoxycyanidin) (peak 10, 301 MS1 m/z) yielded a fragment ion at m/z 286 resulting from the loss of a methyl group.³⁰

3.3. Antioxidant activities

Both red and white flowers showed a potential DPPH radical scavenging activities with IC₅₀ value of $(34.89 \pm 0.98$ and 53.28

 \pm 1.14 $\rm \mu g\,mL^{-1})$ respectively relative to Trolox with IC $_{50}$ (56.82 \pm 0.87 µg mL⁻¹). Red flowers exhibited higher activity than white ones in case of DPPH assay. Likewise, red flowers showed stronger metal chelation effect as another antioxidant action as manifested by 81.45 μ M EDTA equivalent/mg sample versus 77.8 in case of white flowers.

3.4. SCA in vitro assays

3.4.1. Sickling reversal test. Microscopical examination of sickle cells showed elongated or C-like shapes, whilst normal RBCs appeared rounded or biconcave (Fig. 4 and 5). By using sodium metabisulfite, sickled cell percentage increased from 8 to 49%. In contrast, red flowers extracts $(0.25, 0.5$ and 1 mg mL⁻¹) showed decrease in proportion of sickling to 20, 18 and 15% respectively, where white flowers extracts showed 24, 21 and 18% inhibition at a dose of 0.25, 0.5 and 1 mg mL^{-1} , respectively (Fig. 6).

Fig. 3 Structures of identified anthocyanins in P. ellipticum (Kunth) Dugand and P. ellipticum cultivar alba Hort. flower's anthocyanins enriched extract (AEE) in positive mode on an Agilent LC-ESI-Q-TOF-MS instrument.

3.4.2. Polymerization inhibition test. PR and PW showed significant increasing in the rate of polymerization inhibition (P value = 0.0008). Red flower extracts (0.25, 0.5 and 1 $\text{mg }\text{mL}^{-1}\text{)}$

increased the rate of polymerization inhibition to (0.68, 0.92 and 1) respectively, whereas white flower showed 0.54, 0.68 and 0.92 increase in rate at same dose of 0.25, 0.5 and 1 mg mL^{-1}

$$
\frac{1}{2}
$$

Fig. 4 Morphology of sickled RBCs in absence of flowers extracts with approximately 60% elongated C shaped sickled cells.

Fig. 5 Morphology of recovered RBCs in the presence of PR extract (1 mg mL⁻¹) with approximately 90% normal rounded cells.

Fig. 6 Sickling reversal activity of (PR) and (PW) at different extract concentrations.

(Fig. 7). The rate of polymerization inhibition of the control was 0.34. Both assays revealed higher efficacy for red flower extract compared to white flower and are in agreement with antioxidant effect.

4. Discussion

SCA is a common blood disorder, affecting millions of people worldwide.⁸ Ethno medicinal plants have a great importance in the treatment of various ailments including SCA where, phenolic compounds like flavonoids and anthocyanins appeared to exert potential antisickling activity.^{27,31,32} Anthocyanins especially exhibited promising antioxidant and antisickling activities.^{16,17} Anthocyanins ability to attach to proteins may account for their antisickling effect and to further inhibition of

Fig. 7 Rate of polymerization inhibition (RPI%) of (PR) and (PW) at different extract concentrations with significant difference relative to control.

the polymerization of deoxy HbS inside the RBCs.¹⁶ Anthocyanins also exhibit a stabilizing effect on RBCs membrane and enhance the solubility of deoxy HbS, thereby inhibiting gelification.¹⁶ In this study, the flowers anthocyanins content was quantified and LC-MS revealed that nine anthocyanins were found enriched in both flowers. Antioxidants are potential management strategies in SCA ^{13,27} Interestingly, the flowers extracts exhibited potential antioxidant activity by DPPH and metal chelation assays. Also, both flower extracts exhibited a potential antisickling and hemoglobin polymerization inhibition activities by preventing or reversing the sickling of RBCs. In accordance with previous reports higher content of active constituents mainly anthocyanins attributed to the higher antioxidant and antisickling activities in PR. As illustrated, PR and PW were rich in phenolics, flavonoids and anthocyanins with a higher content in case of PR bringing them (especially PR) as a promising trial in treatment of SCA.

5. Conclusion

The phytoconstituents of PR and PW especially anthocyanins were investigated using LC-MS. Their ethanolic extracts were able to stop or reverse sickling and haemoglobin polymerization of human sickled RBCs, thereby lowering the percentage of sickled cells and the rate of haemoglobin polymerization. Beside the antisickling activity, the extracts exhibited also potential antioxidant activity. These findings may represent a primary screening step highlighting the antisickling potential of the flowers of both cultivars. Further research is needed to detect the compounds responsible for the activity. Also, clinical studies need to be conducted to measure the activity in vivo and at the gene level. **PSC Advances**
 SC Article on 1990
 SC Articles Articles

Author contributions

Ahmed S. Mohamed: methodology, software, investigation, writing - original draft Omnia Y. Abd El Dayem: methodology, investigation Ali M. El Shamy: conceptualization, supervision, writing – reviewing and editing Fatma S. El Sakhawy: conceptualization, supervision, writing – reviewing and editing Rania A. El Gedaily: conceptualization, validation, supervision, writing – reviewing and editing.

Conflicts of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- 1 B. F. Mohamed, G. Das, H. Shin, S. S. Ningthoujam and A. Das Talukdar, Systematics, Phytochemistry, Biological Activities and Health Promoting Effects of the Plants from the Subfamily Bombacoideae (Family Malvaceae), Plants, 2021, 10, 1–38.
- 2 J. Refaat, S. Y. Desoky, M. A. Ramadan and M. S. Kamel, Bombacaceae: a phytochemical review, Plants, 2013, 51(1), 100–130.
- 3 N. Navarrette-Tindall and M. A. Orellana Núñez, Pseudobombax ellipticum (Kunth) Dugand, in Tropical Tree Seed Manual, 2003, pp. 664–665.
- 4 A. Ankli, M. Heinrich, P. Bork, L. Wolfram, P. Bauerfeind, R. Brun, C. Schmid, C. Weiss, R. Bruggisser, J. Gertsch,

M. Wasescha and O. Sticher, J. Ethnopharmacol., 2002, 79, 43–52.

- 5 C. C. Walker, A survey of succulents of the mallow family (Malvaceae), CactusWorld, 2021, vol. 39(2).
- 6 R. C. Díaz, G. Espinosa, C. A. Ilizaliturri, D. González and V. G. Cilia, Antioxidant Activity Assessment of Plants Used in Huastec Traditional Medicine, Mexico, J. Pharm., Chem. Biol. Sci., 2015, 3, 488–498.
- 7 A. Navarrete, Gastroprotective effect of β -LupeoL: roLe of prostaGLandins, suLfhydryLs and nitric oxide, Rev. Latinoam. Quim., 2009, 37, 133–143.
- 8 E. Wastnedge, D. Waters, S. Patel, K. Morrison, M. Y. Goh, D. Adeloye and I. Rudan, The global burden of sickle cell disease in children under five years of age: a systematic review and meta-analysis, J. Commun. Healthcare, 2018, 8, $1 - 9$
- 9 B. V. M. Ingram, Gene mutations in human haemoglobin: the chemical difference between normal and sickle cell hjemoglobin, Nature, 1957, 180, 326–328.
- 10 A. K. Geller and M. K. O'Connor, The sickle cell crisis: a dilemma in pain relief, Mayo Clin. Proc., 2008, 83, 320–323.
- 11 I. O. Nurain, C. O. Bewaji, J. S. Johnson, R. D. Davenport and Y. Zhang, Potential of Three Ethnomedicinal Plants as Antisickling Agents, Mol. Pharm., 2017, 14, 172–182.
- 12 E. A. Elbostany, E. A. Elghoroury, E. H. Thabet, A. A. Rashad, E. A. Rasheed, G. S. M. El-Saeed, D. A. Abdelhalim, S. N. Abdelfattah, I. I. Salama and N. Salama, Oxidative stress and hepcidin expression in pediatric sickle cell anemia with iron overload, Hematol./Oncol. Stem Cell Ther., 2021, 11, DOI: [10.1016/j.hemonc.2021.11.003](https://doi.org/10.1016/j.hemonc.2021.11.003).
- 13 L. M. Al-Naama, M. K. Hassan and J. K. Mehdi, Association of erythrocytes antioxidant enzymes and their cofactors with markers of oxidative stress in patients with sickle cell anemia, Qatar Med. J., 2016, DOI: [10.5339/qmj.2015.14](https://doi.org/10.5339/qmj.2015.14).
- 14 N. L. Yembeau, P. C. Biapa Nya, C. A. Pieme, K. D. Tchouane, C. B. Kengne Fotsing, P. J. Nya Nkwikeu, A. F. Feudjio and P. B. Telefo, Ethnopharmacological Study of the Medicinal Plants Used in the Treatment of Sickle Cell Anemia in the West Region of Cameroon, J. Evidence-Based Complementary Altern. Med., 2022, 1–10.
- 15 M. M. Nsimba, J. N. Lami, Y. Hayakawa, C. Yamamoto and T. Kaji, Decreased thrombin activity by a Congolese herbal medicine used in sickle cell anemia, J. Ethnopharmacol., 2013, 148, 895–900.
- 16 P. T. Mpiana, K. T. N. N. Ngbolua, M. T. Bokota, T. K. Kasonga, E. K. Atibu, D. S. T. Tshibangu and V. Mudogo, In vitro effects of anthocyanin extracts from Justicia secunda Vahl on the solubility of haemoglobin S and membrane stability of sickle erythrocytes, J. Blood Transfus., 2010, 8, 248–254.
- 17 D. D. Tshilanda, D. V. Onyamboko, D. T. Mwanangombo, P. V. Tsalu, N. K. Misengabu, D. S. T. Tshibangu, K. N. Ngbolua and P. T. Mpiana, In vitro Antisickling Activity of Anthocyanins from Ocimum canun (Lamiacea), Journal of Advancement in Medical and Life Sciences, 2015, $3(2), 1-5.$
- 18 P. S. Vankar and J. Srivastava, Evaluation of Anthocyanin Content in Red and Blue Flowers, Int. J. Food Eng., 2010, 6(4), 7, DOI: [10.2202/1556-3758.1907](https://doi.org/10.2202/1556-3758.1907).
- 19 S. Silva, E. Costa, C. Calhau, R. Morais and M. Pintado, Anthocyanin Extraction from Plant Tissues: A Review, Crit. Rev. Food Sci. Nutr., 2017, 3072–3083.
- 20 E. Attard, A rapid microtitre plate Folin-Ciocalteu method for the assessment of polyphenols, J. Biol. Chem., 2013, 8, 48–53.
- 21 M. Kiranmai, C. B. M. Kumar and M. Ibrahim, Comparison of total avanoid content of Azadirachta indica root bark extracts prepared by different methods of extraction, Res. J. Pharm., Biol. Chem. Sci., 2011, 2(3), 254–261.
- 22 J. Lee, R. W. Durst, R. E. Wrolstad, K. W. Barnes, T. Eisele, M. M. Giusti, J. Haché, H. Hofsommer, S. Koswig, D. A. Krueger, S. Kupina, S. K. Martin, B. K. Martinsen, T. C. Miller, F. Paquette, A. Ryabkova, G. Skrede, U. Trenn and J. D. Wightman, Determination of Total Monomeric Anthocyanin Pigment Content of Fruit Juices, Beverages, Natural Colorants, and Wines by the pH Differential Method: Collaborative Study, J. AOAC Int., 2005, 88(5), 1269–1278. **Paper Solution** (Solution Commons Articles), 10 M. The entry one is commons and the entry of the entr
	- 23 M. R. Morgan, M. Rogers-Carter and J. P. Christianson, Simultaneous LC-MS quantification of anthocyanins and non-anthocyanin phenolics from blueberries with widely divergent profiles and biological activities, *Physiol. Behav.*, 2017, 176, 139–148.
	- 24 Z. Chen, R. Bertin and G. Froldi, EC50 estimation of antioxidant activity in DPPH? assay using several statistical programs, Food Chem., 2013, 138, 414–420.
	- 25 C. Manach, A. Scalbert, C. Morand, C. Rémésy and L. Jiménez, Polyphenols: food sources and bioavailability, Am. J. Clin. Nutr., 2004, 79, 727–747.
- 26 M. Fidelis, J. S. Santos, A. L. K. Coelho, O. Y. Rodionova, A. Pomerantsev and D. Granato, Authentication of juices from antioxidant and chemical perspectives: a feasibility quality control study using chemometrics, Food Control, 2017, 73, 796–805.
- 27 N. Pauline, B. N. P. Cabral, P. C. Anatole, A. M. V. Jocelyne, M. Bruno and N. Y. Jeanne, The in vitro antisickling and antioxidant effects of aqueous extracts Zanthoxyllum heitzii on sickle cell disorder, BMC Complementary Altern. Med., 2013, 13, 162, DOI: [10.1186/1472-6882-13-162](https://doi.org/10.1186/1472-6882-13-162).
- 28 I. Kim and J. Lee, Variations in Anthocyanin Profiles and Antioxidant Activity of 12 Genotypes of Mulberry (Morus spp.) Fruits and Their Changes during Processing, Antioxidants, 2020, 9, 242, DOI: [10.3390/antiox9030242](https://doi.org/10.3390/antiox9030242).
- 29 K. Schütz, M. Persike, R. Carle and A. Schieber, Characterization and quantification of anthocyanins in selected artichoke (Cynara scolymus L.) cultivars by HPLC-DAD-ESI-MSn, Anal. Bioanal. Chem., 2006, 384, 1511–1517, DOI: [10.1007/s00216-006-0316-6](https://doi.org/10.1007/s00216-006-0316-6).
- 30 E. S. Rebecca, C. B. John, C. Casey and W. Steven, LC-MS/MS and UPLC-UV Evaluation of Anthocyanins and Anthocyanidins during Rabbiteye Blueberry Juice Processing, Beverages, 2017, 3, 56, DOI: [10.3390/](https://doi.org/10.3390/beverages3040056) [beverages3040056](https://doi.org/10.3390/beverages3040056).
- 31 P. T. Mpiana, D. S. T. Tshibangu, O. M. Shetonde and K. N. Ngbolua, In vitro antidrepanocytary actvity (antisickle cell anemia) of some congolese plants, Phytomedicine, 2007, 14, 192–195.
- 32 M. Oyenike, H. Akpan, O. Otulana, A. Adefule, K. Adedokun, W. Oluogun, M. Muhibi and H. Ojokuku, In-vitro antisickling and membrane stability potentials of Mishenland polyherbal extract on sickle red blood cells, Egypt. J. Haematol., 2019, 44, 65.