RSC Advances



View Article Online

View Journal | View Issue

PAPER



Cite this: RSC Adv., 2023, 13, 35292

Received 21st July 2023 Accepted 23rd November 2023

DOI: 10.1039/d3ra04931k

rsc.li/rsc-advances

1. Introduction

The N-heterocyclic carbene (NHC) complexes were first synthesized from imidazolium salts.¹ NHCs are derivatives of persistent carbene compounds having divalent carbons. These heterocyclic species have two nitrogen atoms attached with a carbene carbon within the ring arrangement.² The NHC complexes have been considered for numerous applications in various areas such as organometallic drugs,³ in the industry as catalysts for organic alterations and polymerization reactions, in dye-sensitized thin-film solar cells and in thousands of antimicrobials and anti-tumor medicines.⁴

Since then, the synthesis of novel NH-free carbene ligands and their corresponding metallic compounds such as with silver and selenium have become an area of extensive research.⁵

Characterization of three different benzimidazolium ligands and their organoselenium complexes by using density functional theory and Raman spectroscopy[†]

Sobia Younas,‡^a Aqdas Riaz,‡^a Haq Nawaz, ^b*^a Muhammad Irfan Majeed, ^s*^a Muhammad Adnan Iqbal,^a Nosheen Rashid,^b Areeba Altaf,^a Umar Sohail Shoukat,^a Faisal Jamil,^a Aafia Sehar,^a Sania Munir,^a Mahrosh Javed^a and Muhammad Imran ^c

In the present study, Raman spectroscopy (RS) along with density functional theory (DFT) calculations have been performed for the successful characterization and confirmation of the formation of three different selenium-based N-heterocyclic carbene (NHC) complexes from their respective salts. For this purpose, mean RS features and DFT calculations of different ligands and their respective selenium NHC complexes are compared. The identified characteristic RS and DFT features, of each of these ligands and their selenium complexes, show that the polarizability of benzimidazolium rings increases after complex formation with selenium. This has been shown by the enhanced intensity of the associated Raman peaks, therefore, confirming the formation of newly formed bonds. The complex formation is also confirmed by the identification of several new peaks in the spectra of complexes and these Raman bands were absent in the spectra of the ligands. Moreover, Raman spectral data sets are analyzed using a multivariate data analysis technique of Principal Component Analysis (PCA) to observe the efficiency of the RS analysis. The results presented in this study have proved the RS technique, along with DFT, an undoubtedly fast approach for the confirmation of synthesis of selenium based NHC-complexes.

> Various researches have shown that protein-rich foods such as eggs, fish and meat also contain huge amounts of selenium, with its antioxidant activity through enzymes such as glutathione peroxidase.⁶ Many of the applications regarding selenium-containing NHC complexes have also been studied in effective cancer treatment, anti-septic agents, antibiotics, antioxidants and stress relieving medicines.^{7,8}

> The various organoselenium complexes are being synthesized in several ways and for their exact characterization, different physical methods such as melting point and spectroscopic analytical approaches, for instance, Electrospray Ionization Mass Spectrometry (ESI-MS), Fourier-transform infrared spectroscopy (FT-IR), Nuclear Magnetic Resonance (NMR) spectroscopy and X-ray crystallography are being used.9,10 However, available literature reveals that these above-illustrated approaches do not provide a direct detection pathway within one certain analytic cycle as these techniques have many limitations like consumption, require more sample, expensive, often require complex sample preparation and various intermediate steps which are expensive and laborious.11,12 Raman spectroscopy has been used to minimize the limitations of the above illustrated techniques as it is inexpensive, based on nondestructive interaction between light with chemical bonds within a material, requires no complex sample preparation and

^aDepartment of Chemistry, University of Agriculture Faisalabad, Faisalabad 38000, Pakistan. E-mail: haqchemist@yahoo.com; irfan.majeed@uaf.edu.pk

^bDepartment of Chemistry, University of Education, Faisalabad Campus, Faisalabad 38000, Pakistan

^cDepartment of Chemistry, Faculty of Science, King Khalid University, P.O. Box 9004, Abha 61413, Saudi Arabia

[†] Electronic supplementary information (ESI) available. See DOI: https://doi.org/10.1039/d3ra04931k

[‡] First two authors have equal contribution.

Paper

requires sample only in a small amount. Raman spectroscopy has also been used to study crystallinity, chemical structure, polymorphism, phase, molecular interactions and to minimize the limitations in the characterizations of organometallic compounds such as dimethylselone, seleno urea, organo selenols, organo selenium carbene, organo selenium halides and transition organo selenium compounds.^{11,13,14}

This paper unravels a significant indication of the successful formation of selenium complexes from their particular NHC salts through Raman spectral (RS) analysis along with the support of DFT analysis with great precision. Raman spectroscopy is a molecular vibrational spectroscopic technique that provides vastly valuable evidence about the structures of the molecules based on the Raman scattering effect.¹⁵ This spectroscopic technique provides sensitive information which is specific to the distinct fingerprint of constituent particles in a sample under observation,¹⁶ thus enabling the detection of analytes.¹⁷ This broadened scope of the Raman spectral technique over the other traditional techniques include its good spatial resolution in spectroscopic information with nondestructive identification of target components under ordinary conditions.¹⁸

In the current study, the successful structural elucidation of three different selenium-carbene coordination complexes (with respect to nitrogen attached side chains) has been accomplished through the RS technique along with the DFT method to further explore Raman spectroscopy for such applications. Raman spectroscopy along with a multivariate data analysis technique, Principal Component Analysis (PCA), has been employed for the characterization of three structurally different (differing in substituent at nitrogen) ligands/benzimidazolium salts (ligand-I, ligand-II and ligand-III) and their respective organometallic compounds mono-NHC-selenium complexes (complex-I, complex-II and complex-III). This investigation has led to the advancement in Raman-based approaches for differentiation between a variety of ligands and their particular complexes and also provides an understanding of confirmation of the formation of coordinate covalent bonds in these complexes.

2. Material and methods

2.1 Synthesis of organometallic compounds

Three different salts are named as ligand-I (L1), ligand-II (L2) and ligand-III (L3) and their corresponding selenium adducts are named as complex-I (C1), complex-II (C2) and complex-III (C3) have been synthesized. The details of synthesis and other characterizations will be reported separately and are beyond the scope of the current study.

2.2 DFT calculations

Computational calculations of benzimidazolium ligands and their organo-selenium complexes to get the structural parameters were performed by using Gaussian 09 (ref. 19) and Gauss-View 06.²⁰ DFT measurements are based at B3LYP (Becke's three parameter exchange functional B3 combined with LeeYang–Parr correlation functional LYP).²¹

2.3 Raman spectral acquisition

Raman spectral acquisition of three different ligands and their respective selenium complexes as described in Fig. 1 was characterized using Raman spectrometer (Peak Seeker Pro-785; Agiltron, USA) using a near-infrared (IR) diode laser with a wavelength of 785 nm. To acquire Raman spectral data, a small quantity, 25 mg of each sample was placed on a small grove within the clean aluminum slide and fifteen spectra of each sample were recorded in the 200–1800 cm⁻¹ wavenumber range with the spectral resolution of $\approx 9 \text{ cm}^{-1}$ and 17 points spectral window at 150 mW laser power and 30 seconds of integration time. The monochromatic laser light was passed through the optics and the attached optical microscope with the incorporated lens of $40 \times /0.60$ and a spot size of 2 μ m. To acquire better Raman scattering, the laser was focused just on the surface of the sample on a slide under analysis. Throughout getting Raman spectra, the same focusing level was ensured every time during each measurement using a camera fixed with the microscope. The laser power was also optimized with great care to avoid photodegradation. During the whole process of Raman spectra recording, it was made certain that all the samples, both the ligands as well as their selenium complexes were provided with the persistent conditions. Because of the rapidity of this Raman instrumental technique, it was possible to record measurements of all the samples in a single day to overcome any kind of relevant issues.

2.4 Raman spectral data preprocessing

Before the analysis, the application of preprocessing on raw data was of great significance to achieve robust information and for the enhancement of distinct and prominent features. For this purpose, pre-processing of RS data for three pairs of ligands with their subsequent selenium complex compounds was accomplished using Matlab 7.8 with established protocols. This preprocessing included range selection, baseline correction, normalization and smoothing and substrate removal. A suitable and applicable "The Savitzky–Golay Smoothing method" (order 1, 17 point window) was processed to get all the spectra smoothed and a rubber band algorithm was carried out as well to get baseline corrected.

2.5 Raman spectral data analysis

The RS data was examined by making comparisons of the mean Raman spectra of each ligand with that of its corresponding metal complex and shown in Fig. 2–4. A vast literature has been studied to ascribe the Raman peaks and their obtained DFT results have been mentioned in Table 1. The interpretation has been made for the final results. Additional analysis has also been performed by using principal component analysis (PCA), an unsupervised multivariate data analysis technique that involves the transformation of correlated variables into uncorrelated variables called principal components (PCS). A linear transformation has been performed on the data to capture most



Fig. 1 Structural formulas of benzimidazolium salts (ligand-I, ligand-II and ligand-III) and their respective selenium-N-heterocyclic carbene complexes (complex-I, complex-II and complex-III).



Fig. 2 Molecular electrostatic potential structures of ligands and complexes

of the variance by the first few principal components. The first principal component (PC-1) captures the most variance, which has been followed by the second principal component (PC-2).

3. Results and discussion

3.1 Molecular electrosttic potential (MEP)

The Molecular Electrostatic Potential (MEP) is a useful tool for predicting the polarity and polarizability of the compounds. In Fig. 2, different colors represent different electrostatic potential values. In MEP calculations, red, orange and yellow colors represent high polarity while blue and green represent lower polarity. In the structures of ligands L1, L2 and L3, the presence of benizimidazole's nitrogens and ionizable bromine make it highly polar and shown as red and orange areas. While in the cases of complexes C1, C2 and C3, the polarity is still present (due to benzimidazole's nitrogens) but reduced because of the removal of bromide. Less polar areas are found around the benzimidazole ring. The surrounding regions having alkyl chains are shown as light blue and green indicating the surrounding of the benzimidazole as less polar and high polarizable as it is more exposed to external electric field. However, with the cases of L2 and C2, -NO₂ is present as substituent that contains strongly electronegative atoms that's why shown as red indicting the surrounding of the benzimidazole ring as highly polar site (Fig. 2).

3.2 Mean Raman spectral analysis

Fig. 3–5 represent the mean Raman spectra of ligands and their respective selenium complexes. Common mean Raman spectral features which are represented by dashed lines as well as highly differentiating features which are represented by solid lines have been manifested drawing vertical lines in spectra and are collectively summarized in Table 1 along with the appropriate assignments and relevant citations are made for corresponding literature. The highly common features which are present in both mean spectra of each metal-ligand pair are the indication of the chemical structure's similarity between the ligand and its metallic counterpart, while the obvious differences can be specifically associated with the attachment of the selenium metal with the complex. Two types of differences are noticeable demonstrating the attachment of selenium metal within a complex including enhancement in intensities and shifts in Raman values of peak positions. The intensity differences and shifts in peaks in these mean spectral features signal the confirmation of stable carbon–selenium dative bond formation.

As selenium forms a double bond with the carbene carbon of benzimidazole ring, vanishing of charge on ligand-I which is salt of bromide ion, transforms polarity of the ring into polarizability, as a result enhancing Raman spectral features at 331 cm^{-1} (imidazole ring wagging accompanied with the attached chains deformation), 713 cm⁻¹ (C-N out of plane stretching vibrations of imidazole rings), 750 cm⁻¹ (out-ofplane bending of benzene in benzimidazole), 736 cm^{-1} (wagging of C-H in benzene ring). The most significant differentiating features have been manifested through dashed lines at 1416 cm^{-1} in C1, probably due to C=Se vibrations while a strong peak appearance at 1186 cm^{-1} is the indication of activation of C-H in-plane bending of benzimidazole ring and a shoulder peak at 1342 cm⁻¹ is due to C-H deformation in benzene ring after complex formation. Notably, these vibrations/spectral features are absent in the L1 structure. Selenium complex formation has resulted in some of the spectral features to decrease in intensities or diminish completely in mean Raman spectra of complexes such as a band in ligand at 1313 cm^{-1} (C–H deformation of CH₂ in aliphatic side chains) is vague in complex and a large peak at 1610 cm^{-1}



Fig. 3 Comparison of mean Raman spectra of ligand-I (L1) with complex-I (C1).

in ligand has become a relatively small peak in C1 which indicates the reduced C=C stretching vibrations of benzimidazole ring after C1 formation. In the same way, relatively intense

bands at 779 $\rm cm^{-1}$ and 878 $\rm cm^{-1}$ (C-H out of plane bending in benzimidazole) in the spectrum of C1 also confirm selenium complex formation.



Fig. 4 Comparison of mean Raman spectra of ligand-II (L2) with the mean spectrum of complex-II (C2).

(cc) BY-NC

Table 1 Observed Raman bands in mean Raman spectra of benzimidazolium salts (L1, L2 and L3) and their respective selenium-N-heterocyclic carbene complexes (C1, C2 and C3) along with their DFT calculations^{α}

Observed L1 (cm ⁻¹)	DFT b3lyp L1 (cm^{-1})	Observed C1 (cm^{-1})	DFT b3lyp C1 (cm^{-1})	Observed L2 (cm^{-1})	DFT b3lyp L2 (cm^{-1})	Observed C2 (cm^{-1})	DFT b3lyp C2 (cm ⁻¹)	Observed L3 (cm^{-1})	DFT b3lyp L3 (cm ⁻¹)	Observed C3 (cm ⁻¹)	DFT b3lyp C3 (cm ⁻¹)	Peak assignment	References
267	270	267	268	267	267	267	265	266	266	_	_	(C–C–C) o-p bending	22
286	283	286	287	281	281	281	281	280	280	280	280	CH ₂ rocking	23
	—	—	—	—	—			293	—	293	291	CH ₂ rocking	23
_	—	_	—	311	308	_	—	309	310	_	—	Def vibrations in C– C–C	24
331	340	331	328	331	334	331	326	331	330	331	330	Ring wag + chain def	25
367	369	367	372	370	366	370	369	_	_	—	—	C–C def	26
_	_	_	_	385	376	385	390	_	_	_	_	Bz ring bending modes	27
_	—	—	—	—	—	—	—	421	421	421	421	Ring rotation 5- member ring	25 and 28
_	—	_	_	434	434	434	441	434	434	434	432	C–C def of aliphatic chain	23 and 26
458	458	458	457	_	_	_	_	457	459	457	462	C-C-C o-p bend	29
476	476	476	476	474	475	474	475	_		_		C-C skeleton def	30
_	_	_	_	_	_	_		499	498	499	498	C-N o-p bend	25
539	546	539	539	_	_	_	_	532	_	532	_	benzimidazole C-N-C def aliphatic	31
005	010	000	005					002		001		chain	01
_	_	_	_	548	546	548	547	_	_	_	_	Wag mode of NO ₂	32
_	_	_	_	577	577	577	579	572	582	572	582	Ring def	23
												benzimidazole	
_	—	_	—	_	—	_	—	609	609	609	609	C-C-C def i-p	33
617	615	617	610	_	_	_	_	_	_	_	_	C-C-C symmetric	34
017	015	017	019									bend phenyl ring	34
	_	_	_	627	626	_		_	_	_	_	C-C-C def i-p	33 and 34
				027	020							benzimidazole	oo unu o i
_	_	—	_	—	_	651	651	_	_	—	_	Phenyl ring	35
												vibrations	
713	713	713	713	713	_	713	713	713	713	713	713	C–N o-p str of 5- membered ring	23
736	738	736	737	_	_	_	_	_	_	_	_	Bz C-H sym wag	36
750	752	750	750	_	_	750	750	_	_	_	_	Bz o-p bend	23 and 37
779	782	779	781	780	784	780	787	782	780	782	780	C-H o-p bend	22 23 and
,,,,	/02	,,,,	/01	700	/01	700	, 0,	702	700	702	700	benzimidazole	37
822	825	822	820	826	827	826	826			822	822	C–C str	26
840	_	840	840	_	_	_	_	843	846	843	845	C–C str	38
_	_	_	_	863	863	_	_	_	_	_	_	NO ₂ sci	32
878	—	878	—	_	—	_	—	_	—	_	—	C–H o-p bend	23 and 39
				~~~					~~~			benzimidazole	
	_	_	_	887	876			890	895	_	_	C-H o-p bend	22
897	898	_	_			895	896	_	—		_	CH ₂ SCI	40
 067		067		942	938	942			 050		067	CH war	37 26 and 40
907 1010	903 1015	1010	909 1010	1014	1000	1014	1018	903 1016	939 1023	1023	907 1023	$C_{113}$ wag $C_{-}C_{-}C_{-}C_{-}$ trigonal bend	20 and 40
1019	1015	1019	1019	1014	1009	1014	1018	1010	1023	1023	1023	benzimidazole	39
	_	_	_	1049	1045		_			_	_	Ring def 5-member	25
_	_	_	_	_	_	_	_	10/1	10/1	_	_	c-c str mode of aliphatic	23 and 41
_		—	—	1098	1089	1098	1089	_	_	—	_	C–N str	18
1138	1139	1130	1135	_	—	1130	1130	1137	1143	1137	1137	C–C str aliphatic vibration	41
_	_	_	_	1144	1141	1156	1154	_	_	_	_	C–H sym bend	42
		1100	4462	4475	44.60	1102	4462	4400	1400	44.02	44.62	benzimidazole	
_	—	1186	1182	11/5	1169	1183	1182	1183	1183	1183	1183	1-p C–H bend benzimidazole	3/
1215	1213	1215	1212		_	_		_	_	_	_		23 and 26

Table 1 (Contd.)

$\begin{array}{cccccccccccccccccccccccccccccccccccc$	nt References
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	liphatic
-       -       -       -       -       -       -       1225       1225       1214       1218       C-N str benzimi         -       -       1246       1244       -       -       -       -       1225       1214       1218       C-N str benzimi         -       -       1246       1244       -       -       -       -       -       -       C-C str chain         -       -       -       -       -       -       -       -       -       C-C str chain         -       -       1297       1298       1283       1290       1293       1298       1294       1294       1294       1294       CH ₂ def chain         1313       1313       -       -       1318       1318       -       -       1313       1313       -       -       C-H def         1351       1351       1351       1354       1358       1369       1367       1354       1352       1354       1362       Ring str         -       -       1416       1417       -       -       1411       1411       -       -       1412       1410       C==Se         1434	•
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	43 and 44
-       1246       1244        -       -       -       -       -       C-C strate chain          -       -       -       1258       1256       1269       1260       1256       -       -       ip bz rin chain          -       1297       1298       1283       1290       1293       1298       1294       1294       1294       1294       1294       CH2 def chain         1313       1313       -       -       1318       1318       -       -       1313       1313       -       -       C-H def chain         1351       1351       1351       1351       1354       1358       1369       1367       1354       1352       1354       1362       Ring str         -       -       1416       1417       -       -       1411       1411       -       -       1412       1410       C=Se         1434       1432       1445       1443       1432       1448       1443       1433       1427       1446       1438       CH def	azole
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	liphatic side 23 and 26
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
-       -       1297       1298       1283       1290       1293       1298       1294       1294       1294       1294       1294       CH2 def chain         1313       1313       -       -       1318       1318       -       -       1313       1313       -       -       CH2 def chain         1313       1313       -       -       1318       1318       -       -       1313       1313       -       -       C-H def         1351       1351       1351       1351       1354       1358       1369       1367       1352       1354       1362       Ring str         -       -       1416       1417       -       -       1411       1411       -       -       1412       1410       C=Se         1434       1432       1445       1443       1432       1448       1433       1427       1446       1438       CH def	g def 27
1313         1318         1313         C-H def         1342       1340       1342       1342       1342       1342       1340       C-H def         1351       1351       1351       1351       1354       1358       1369       1367       1354       1352       1354       1362       Ring str           1416       1417         1411       1411        1412       1410       C=Se         1434       1432       1443       1432       1448       1433       1427       1446       1438       CH def	liphatic 23 and 26
1342       1340       1342       1342       1342       1340       C-H def         1351       1351       1351       1351       1354       1358       1369       1367       1354       1352       1354       1362       Ring str         -       -       1416       1417       -       -       1411       1411       -       1412       1410       C=Se         1434       1432       1443       1432       1448       1433       1427       1446       1438       CH def	of CH ₂ 45
1351       1351       1351       1354       1358       1369       1367       1354       1352       1354       1362       Ring str         -       -       1416       1417       -       -       1411       1411       -       -       1412       1410       C=Se         1434       1432       1445       1443       1432       1448       1433       1427       1446       1438       CH def	n bz ring 27 and 37
1416 1417 1411 1411 1412 1410 C=Se 1434 1432 1445 1443 1434 1432 1448 1448 1433 1427 1446 1438 CH def	midazole 27 and 37
1434         1432         1443         1434         1432         1448         1433         1427         1446         1438         CH def	46
	41 and 47
1590 1589 1590 1588 C-C-C s	r mode of 33
vibration	in phenyl
ring	
1610 — 1610 — 1613 1618 1613 1617 1611 — 1611 — C=C str	mode 18, 22, 37
benzimi	azole and 48

Comparatively enhanced bands associated with aliphatic side chains at 267 cm⁻¹ and 458 cm⁻¹ (C-C-C out-of-plane bending), 822 cm⁻¹, 840 cm⁻¹, 1215 cm⁻¹, 1246 cm⁻¹ and shift of 1138 cm⁻¹ to 1130 cm⁻¹ (C-C stretching associated with aliphatic chains on both sides of imidazole rings) and at 967  $\text{cm}^{-1}$  (CH₃ wagging) are most likely due to the reason that lone pairs at nitrogen atoms become delocalized after complex making, causing less (-I) inductive effect resulting in a lesser amount of polarity and more polarizability of alkyl chains hence enhancing Raman spectral signals in the spectrum of the complex. The bands in the L1 spectrum at 1434  $cm^{-1}$  and 1313 cm⁻¹ are associated with C-H deformations of CH₂ which are characteristic of hydrocarbon chains. Similarly, these Raman spectral features are in correspondence with the Raman spectral features calculated from density functional theory (DFT) as in ESI as Fig. S1.[†]

The distinct features in mean RS analysis have been observed, in which features related to benzimidazole ring structure at 331  $\text{cm}^{-1}$  (ring wag with chain twisting), 385  $\text{cm}^{-1}$ (benzene ring bending modes), 577 cm⁻¹ (ring deformation), 627 cm⁻¹ (in-plane C-C-C deformation), 750 cm⁻¹ (Bz out-ofplane bending), 780 cm⁻¹ (C-H out of plane bending), 826 cm⁻¹ (C–C str), 942 cm⁻¹ (imidazole in-plane bending), 1014 cm⁻¹ (C-C-C trigonal bending), 1049 cm⁻¹ (ring deformation in 5-member ring), 1342 cm⁻¹ (C-H deformation in benzene ring), 1613 cm⁻¹ (C=C stretching mode). On the formation of complex, C2 from L2, the disappearance of charges on imidazole ring takes place which leads to diminishing polarity and consequently enhanced polarizability due to which a few peaks become relatively sharp, such as at 281 cm⁻¹ more

 $CH_2$  rocking occurs with C-C deformation at 370 cm⁻¹ and at 1014 cm⁻¹, C–C–C trigonal bending enhances in benzimidazole structure. The peak intensity at 1144 cm⁻¹, 1175 cm⁻¹ and 1354 cm⁻¹ has been increased as well as shifted at higher wavenumbers to 1156  $cm^{-1}$ , 1183  $cm^{-1}$  and 1369  $cm^{-1}$ respectively, predicting strong C-H symmetric, in-plane bending and ring stretching vibrations in imidazole rings respectively. Moreover, the peak shift from 1253  $\text{cm}^{-1}$  to 1269 cm⁻¹ due to an in-plane benzene ring deformation in benzimidazole ring within selenium complex-II (C2). Similarly changes in peak position on the formation of C2 from L2, shift from 1283 cm⁻¹ to 1293 cm⁻¹, have been recognized as CH₂ deformations in the aliphatic chain between two imidazole rings. A new shoulder band appearing at 1342 depicts C-H deformation in the benzene ring.

A strong peak at 1411  $\text{cm}^{-1}$  is assigned to C=Se vibrations in C2 as its counterpart feature is absent in the mean Raman spectrum of L2 which provides us with strong evidence of selenium complex formation while phenyl ring presence in only the second pair of ligand-complex (L2-C2) a peak at 1590 cm⁻¹ has been allotted to C-C-C stretching mode of vibration in phenyl ring of side chains which is absent in other first and third ligand-complex pairs (L1-C1 and L3-C3).

On the other hand, relatively reduced intensities in C2 at some positions such as at 311 cm⁻¹ (ring wag with chain twisting), 434 cm⁻¹ and 474 cm⁻¹ (C–C deformation), 548 cm⁻¹ (wag mode of  $NO_2$ ), 627 cm⁻¹ (in-plane C–C–C deformation), 780 cm⁻¹ (C–H out of plane bending), 887 cm⁻¹ (C–H out-ofplane bending), 942  $\text{cm}^{-1}$  (imidazole in plane bending), 1049 cm⁻¹ (ring deformation in 5-member ring), 1144 cm⁻¹ (C–

#### Paper

H symmetric bending benzimidazole), 1318 cm⁻¹ (C–H deformation of CH₂), 1354 cm⁻¹ (ring stretch imidazole), the shift of 1434 cm⁻¹ to 1448 cm⁻¹ (CH deformation in side chain) and 1613 cm⁻¹ (C=C stretching mode) might be because, in contrast to alkyl side chains (electron-donating effect) in first and third pairs of ligands and complexes (L1–C1 and L3–C3), it has an electron-withdrawing group (nitrobenzene) as a side chain on benzimidazole rings which causes molecule towards somewhat reverse polar effect. Similarly, these Raman spectral features have been found in correspondence with the Raman spectral features calculated from DFT as in ESI as Fig. S2.†

Fig. 5 represents distinct mean Raman spectral features which are characteristic to some specific chemical structural parts of ligand-III (L3) and complex-III (C3). For instance, Raman spectral features at 331  $\text{cm}^{-1}$  (ring wag + chain deformations), 421 cm⁻¹ (ring rotation 5-member ring), 499 cm⁻¹ (C-N out of plane bending), 532 cm⁻¹ (C-N-C deformation aliphatic chain), 572  $\text{cm}^{-1}$  (ring deformation), 609 cm⁻¹ (C-C-C deformation in-plane vibration mode of benzene), 713 cm⁻¹ (C-N out of plane stretching fivemembered ring), 782 cm⁻¹ (C-H out of plane bending), 1016 cm⁻¹ (C-C-C trigonal bending), 1183 cm⁻¹ (in-plane C-H bending), 1225  $\text{cm}^{-1}$  (C-N stretching), 1260  $\text{cm}^{-1}$  (inplane bz ring deformation), 1342 cm⁻¹ (C-H deformation in bz ring), 1354 cm⁻¹ (ring stretch imidazole) and 1611 cm⁻¹ (C=C stretching mode) reflect the structural dissimilarities associated with benzimidazole.

The spectral features at 280  $\text{cm}^{-1}$  and 293  $\text{cm}^{-1}$  (CH₂ rocking),  $331 \text{ cm}^{-1}$  (ring wagging and chain deformation),  $713 \text{ cm}^{-1}$ (C-N out of plane stretching in five-membered ring), 822 and 843 cm⁻¹ (C–C stretching), 1137 cm⁻¹ (C–C stretching, aliphatic vibration), 1183 cm⁻¹ (in-plane C-H bending, benzimidazole), 1294  $\text{cm}^{-1}$  (CH₂ deformation, aliphatic chain) in C3 has been intensified. A similar justification goes for this increase in the intensity of some of the C3 Raman spectral features, as for the L1-C1 and L2-C2 ligand/complex pairs and can be associated with increased polarizability and decreased polarity in the complexes. Additional bands appear as well at 822  $\text{cm}^{-1}$  (C-C stretching), 1342 cm⁻¹ (C-H deformation in benzene ring) and 1412 cm⁻¹ (C=Se vibrations) which are the most distinguished features indicating stable selenium compound formation from its own salt. Changes in peak positions (shifts in peak positions) have also been observed at 1016 to 1023 cm⁻¹ (C-C-C trigonal bending benzimidazole) and 1225 to 1214 cm⁻¹ (C-N stretching benzimidazole) with an increase in intensity from L3 to C3.

Moreover, at some positions, in L3, few features disappeared or decreased in intensity in C3 as well, including 309 cm⁻¹ (deformation vibrations of C–C–C), 532 cm⁻¹ (C–N–C deformation aliphatic chain), 609 cm⁻¹ (C–C–C deformation in-plane vibration mode of benzene), 782 cm⁻¹ (C–H out of plane bending benzimidazole), 890 cm⁻¹ (C–H out-of-plane bending), 1260 cm⁻¹ (in-plane benzene ring deformation), 1313 cm⁻¹ (C– H def of CH₂), 1354 cm⁻¹ (ring stretch imidazole) and shifting from 1433 to 1446 cm⁻¹ (C–H deformation) and 1611 to



Fig. 5 Comparison of mean Raman spectrum of ligand-III (L3) with mean Raman spectrum of complex-III (C3).

1617 cm⁻¹ (C=C stretching mode benzimidazole). In the same way, these Raman spectral features have great correspondence with the Raman spectral features calculated from DFT as shown in ESI in Fig. S3.†

#### 3.3 Principal component analysis (PCA)

Fig. 6–8 are the PCA scatter and loading plots of PCA of Raman spectral data of ligands and their selenium complexes in which both samples have been differentiated in the form of two different clusters indicated by different colors. In each PCA scatter plot, the black color cluster on the positive axis of PC-1 represents the Raman spectral data set of ligand while the green color on the left side is associated with complexes. Mainly PCA is used to decrease the data dimensionality while keeping variability intact in the spectral data.

Fig. 6(a) shows PCA scatter plot of Raman spectral data of ligand-I (black dots) and its respective selenium complex-I (green dots). Maximum variability in data has been explained by PC-1 explaining the dominant source of variability in the data which is 98.42% while PC-2 clarifies the rest of the variability. To explore the ability of PCA to distinguish ligand/salt samples from their respective complexes, PCA is used to identify and confirm the Raman spectral features associated with the differentiation of two different spectral data sets of ligands and their respective complex.

Fig. 7(a) shows PCA scatter plot of Raman spectral data of ligand-II (black dots) and its respective selenium complex-II



Fig. 6 (a) PCA scatter plot of Raman spectral data of ligand-I (black dots) and its respective selenium complex-I (green dots), (b) PCA loadings of Raman spectral data set of ligand-I and complex-I.



Fig. 7 (a) PCA scatter plot of Raman spectral data of ligand-II (black dots) and its respective selenium complex-II (green dots), (b) PCA loadings between Raman spectral data set of ligand-II and complex-II.

(green dots). In this case, the PC-1 predicts 98.88% variability and only 0.14% variability has been explained by PC-2.

Fig. 8(a) shows the PCA Scatter plot of Raman spectral data sets of ligand-III (black dots) and its respective selenium complex-III (green dots). PC-1 explains a maximum of 99.20% variability which indicates clear differentiation between the Raman spectral data sets of ligand-III and its respective complex.

As scatter plot has provided a chromatic display of two sets of data in two different clusters, but it is unable to provide reasons for such variability in data regarding distinct features. To understand this basis of discrimination in the PCA scatter plot, the loadings of PC-1 against wavenumber  $(cm^{-1})$  have been presented in Fig. 6(b), 7(b) and 8(b), for three different ligands and respective complexes, respectively. The reason for taking

PC-1 for loadings is because it has provided maximum variability.

Fig. 6(b) presents the loadings of the PCA of Raman Spectral data sets of ligand-I (L1) compared with the complex-I (C1). The PC loadings are the dimensions of chemical variations, splitting different spectral groups of RS data along their variability. The positive loadings from the top side of the splitting line at zero are associated with the RS data of the ligands which have clusters on the right side in PC-1 of the scatter plot while the negative loadings are associated with the corresponding selenium complexes clustered at the negative side of PC-1. The loading plot expresses the only prominent features of ligands that are distinct from that of complexes. Such spectral loadings re-confirming the structural differences between ligands and their respective complexes as already have been discussed in



Fig. 8 (a) PCA scatter plot of Raman spectral data sets of ligand-III (black dots) and its respective selenium complex-III (green dots), (b) PCA loadings of Raman spectral data set of ligand-III and complex-III.

mean Raman spectra. These differences indicate some kind of structural changes in complexes from their preliminary ligands and hence can be taken as Raman spectral markers of stable complex formation.

The dominant RS features of C1 that appeared in the PCA include 286 cm⁻¹ (CH₂ rocking), 331 cm⁻¹ (imidazole ring wagging accompanied with the attached chains deformation), 713 cm⁻¹ (C–N out of plane stretching vibrations of imidazole rings), 822 cm⁻¹, 1130 cm⁻¹, 1215 cm⁻¹ and 1246 cm⁻¹ (all probably associated with C–C stretching in aliphatic chains on both sides of imidazole rings), 967 cm⁻¹ (CH₃ wagging), 1186 cm⁻¹ (C–H in-plane bending of benzimidazole ring) and 1297 cm⁻¹ (CH₂ deformation in the aliphatic chain). Another and highly differentiating feature at 1416 cm⁻¹ has been

attributed to C=Se vibrations because of its only presence in a complex structure. In contrast, the most prominent positive loadings associated with ligand-I appear at 779 cm⁻¹ (C-H out of plane bending in benzimidazole), 1019 cm⁻¹ (C-C-C trigonal bending inbenzimidazole), 1138 cm⁻¹ (C-C stretching), 1313 cm⁻¹ and 1434 cm⁻¹ (C-H def), 1351 cm⁻¹ (ring stretch in imidazole) and 1610 cm⁻¹ (C=C stretching).

The PCA loadings of Raman spectral data sets of L2–C2 and L3–C3 pairs of ligands and complexes are shown in Fig. 7(b) and 8(b)) which reflect the same reasons of differentiation as has already been elaborated in the above section in the discussion related to Fig. 4 and 5 for ligand-II, &III and their complexes respectively. These unique features in Fig. 7(b) at 311, 345, 434, 474, 548, 627, 680, 780, 887, 942, 1049, 1144, 1318, 1354, 1434

and 1613 cm⁻¹ while in Fig. 8(b) at 309, 532, 609, 782, 890, 1016, 1225, 1260, 1313, 1354, 1433 and 1611 cm⁻¹ act as a fingerprint for a specific structural elucidation of **L2** and **L3**. Similarly visible features at 281, 370, 595, 651, 750, 1014, 1156, 1183, 1269, 1293, 1342, 1369, 1411, 1485, and 1590 cm⁻¹ in Fig. 7(b) and dominant features at 280, 331, 593, 713, 822, 1023, 1137, 1183, 11214, 1294, 1342, 1412 and 1489 cm⁻¹ in Fig. 8(b) belonging to negative loadings enable description of stable metal–ligand bond which means complex formation and these results have also been found compatible with mean Raman spectra results.

## 4. Conclusions

Raman spectroscopy combined with DFT and multivariate data analysis methods is found helpful for the characterization of three different ligands and their respective selenium complexes. The Raman spectral features have been found for each of these ligands and their selenium complexes and it is in ferred that the polarizability of benzimidazolium rings has been increased after complex formation with selenium. This has been shown by the enhanced intensity of the associated Raman peaks, therefore, confirming the formation of newly formed bonds. The unique spectral features have been observed in Raman spectra of complexes I, II and III at 1416, 1411, 1412 cm⁻¹ respectively. As very limited Raman spectroscopic data is available for benzimidazolium salts and their selenium complexes in the literature thus depending upon observations, these bands have been estimated at C=Se. Moreover, complex formation is also indicated by the identification of several new peaks in the spectra of these complexes and these Raman bands were absent in spectra of the ligands. This indicates that the Raman scattering technique with the support of DFT calculations reveals a clear indication of complex formation.

## Conflicts of interest

There are no conflicts of interest to declare.

## Acknowledgements

M. Imran expresses his appreciation to the Deanship of Scientific Research at King Khalid University Saudi Arabia for funding this work through research groups program under grant number R.G.P. 2/522/44.

## References

- 1 J. C. Garrison, C. A. Tessier and W. J. Youngs, *J. Org. Chem.*, 2005, **690**, 6008–6020.
- 2 K. B. Choo, S. M. Lee, W. L. Lee and Y. L. Cheow, *J. Org. Chem.*, 2019, **898**, 120868.
- 3 C. Y.-S. Chung, S.-K. Fung, K.-C. Tong, P.-K. Wan, C.-N. Lok, Y. Huang, T. Chen and C.-M. Che, *Chem. Sci.*, 2017, 8, 1942– 1953.
- 4 R. Fraser, C. G. Van Sittert, P. H. Van Rooyen and M. Landman, *J. Org. Chem.*, 2017, **835**, 60–69.

- 5 A. Kamal, M. Nazari, M. Yaseen, M. A. Iqbal,
   M. B. K. Ahamed, A. S. A. Majid and H. N. Bhatti, *Bioorg. Chem.*, 2019, **90**, 103042.
- 6 T. Klapec, M. L. Mandić, J. Grgić, L. Primorac, A. Perl and V. Krstanović, *Food Chem.*, 2004, **85**, 445–452.
- 7 E. E. Battin, N. R. Perron and J. L. Brumaghim, *Inorg. Chem.*, 2006, **45**, 499–501.
- 8 M. Elsherbini, W. S. Hamama and H. H. Zoorob, *Coord. Chem. Rev.*, 2016, **312**, 149–177.
- 9 M. A. Iqbal, R. A. Haque, S. F. Nasri, A. A. Majid, M. B. K. Ahamed, E. Farsi and T. Fatima, *Chem. Cent. J.*, 2013, 7, 27.
- 10 S. Y. Hussaini, R. A. Haque, P. O. Asekunowo, A. A. Majid, M. T. Agha and M. R. Razali, *J. Org. Chem.*, 2017, 840, 56–62.
- 11 M. N. Ashraf, M. I. Majeed, H. Nawaz, M. A. Iqbal, J. Iqbal, N. Iqbal, A. Hasan, N. Rashid, M. Abubakar and M. Z. Nawaz, Spectrochim. Acta, Part A, 2022, 270, 120823.
- 12 S. Mudi, M. Usman and S. Ibrahim, *Am. J. Chem. Appl.*, 2015, 2, 151–158.
- 13 W. K. Smothers and M. S. Wrighton, *J. Am. Chem. Soc.*, 1983, 105, 1067–1069.
- 14 M. Sayyed, A. H. Abdalsalam, M. M. Taki, M. H. A. Mhareb, B. Alim, A. Baltakesmez and E. Şakar, *Radiat. Phys. Chem.*, 2020, **172**, 108852.
- 15 Analytical applications of Raman spectroscopy, ed. M. J. Pelletier, Oxford: Blackwell science, 1999, vol. 427.
- 16 C. Zong, M. Xu, L.-J. Xu, T. Wei, X. Ma, X.-S. Zheng, R. Hu and B. Ren, *Chem. Rev.*, 2018, **118**, 4946–4980.
- 17 J. R. Ferraro, Introductory Raman Spectroscopy, Elsevier, 2003.
- A. Bankapur, R. S. Krishnamurthy, E. Zachariah, C. Santhosh, B. Chougule, B. Praveen, M. Valiathan and D. Mathur, *PLoS One*, 2012, 7, e35075.
- 19 J. Singh, Res. Chem. Intermed., 2020, 46, 2457-2479.
- 20 C. M. Zoltowski, R. F. Lalisse, C. M. Hadad and Z. D. Schultz, *J. Phys. Chem.*, 2021, **125**, 27596–27606.
- 21 V. Kumar, J. Teotia and A. K. Yadav, *Mater. Today: Proc.*, 2022, **62**, 7137–7141.
- 22 S. Mohan, N. Sundaraganesan and J. Mink, Spectrochim. Acta, Part A, 1991, 47, 1111–1115.
- 23 S. Ali, A. Riaz, H. Nawaz, M. I. Majeed, M. A. Iqbal, H. N. Bhatti, N. Rashid, M. Kashif, M. Tahir and S. Nasir, *Spectrochim. Acta, Part A*, 2020, 232, 118162.
- 24 Y.-J. Mo, Y.-C. Li, J.-G. Zhao, K.-A. Fen, C.-Y. Xiao and M.-X. Wan, *J. Magn. Magn. Mater.*, 2000, **213**, 278–280.
- 25 R. W. Berg, Monatsh. Chem., 2007, 138, 1045-1075.
- 26 N. Sheppard, J. Chem. Phys., 1948, 16, 690-697.
- 27 G. Wang, A. Harrison, X. Li, G. Whittaker, J. Shi, X. Wang,
  H. Yang, P. Cao and Z. Zhang, *J. Raman Spectrosc.*, 2004,
  35, 1016–1022.
- 28 S. Ali, A. Riaz, M. I. Majeed, M. A. Iqbal, H. N. Bhatti, N. Rashid, M. Kashif, M. Tahir, S. Nasir and S. Farooq, *Spectrochim. Acta, Part A*, 2020, 228, 117851.
- 29 S. W. Han and K. Kim, J. Colloid Interface Sci., 2001, 240, 492–497.
- 30 D. Lin-Vien, N. B. Colthup, W. G. Fateley and J. G. Grasselli, *The Handbook of Infrared and Raman Characteristic Frequencies of Organic Molecules*, Elsevier, 1991.

- Open Access Article. Published on 04 December 2023. Downloaded on 7/29/2025 5:27:28 AM. This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.
- 31 V. Renganayaki and S. Srinivasan, *Int. J. PharmTech Res.*, 2011, 3, 1350–1358.
- 32 C. J. Strachan, T. Rades, K. C. Gordon and J. Rantanen, *J. Pharm. Pharmacol.*, 2007, **59**, 179–192.
- 33 X. Zhang, Q. Zhou, Y. Huang, Z. Li and Z. Zhang, *Sensors*, 2011, **11**, 11510–11515.
- 34 C. Muthuselvi, S. Pandiarajan, B. Ravikumar, S. Athimoolam and R. Krishnakumar, *Asian J. Appl. Sci.*, 2018, **11**, 29–37.
- 35 K. Helios, A. Pietraszko, W. Zierkiewicz, H. Wójtowicz and D. Michalska, *Polyhedron*, 2011, **30**, 2466–2472.
- 36 M. Morsy, M. Al-Khaldi and A. Suwaiyan, J. Phys. Chem. A, 2002, **106**, 9196–9203.
- 37 A. Suwaiyan, R. Zwarich and N. Baig, *J. Raman Spectrosc.*, 1990, **21**, 243–249.
- 38 Z. Arp, D. Autrey, J. Laane, S. A. Overman and G. J. Thomas, *Biochem*, 2001, **40**, 2522–2529.
- 39 N. Sundaraganesan, S. Ilakiamani, P. Subramani and B. D. Joshua, *Spectrochim. Acta, Part A*, 2007, **67**, 628–635.

- 40 R. Kengne-Momo, P. Daniel, F. Lagarde, Y. Jeyachandran, J. Pilard, M. Durand-Thouand and G. Thouand, *Int. J. Spectrosc.*, 2012, **2012**, 1–7.
- 41 C. J. Orendorff, M. W. Ducey Jr and J. E. Pemberton, *J. Phys. Chem. A*, 2002, **106**, 6991–6998.
- 42 S. Salama and T. G. Spiro, *J. Am. Chem. Soc.*, 1978, **100**, 1105–1111.
- 43 P. Lasch and D. Naumann, Mol. Cell. Biol., 1998, 44, 189-202.
- 44 R. Wojnarowska, J. Polit, D. Broda, M. Gonchar and E. Sheregii, *Appl. Phys. Lett.*, 2015, **106**, 103701.
- 45 T. Mahmood, H. Nawaz, A. Ditta, M. Majeed, M. Hanif, N. Rashid, H. Bhatti, H. Nargis, M. Saleem and F. Bonnier, *Spectrochim. Acta, Part A*, 2018, 200, 136–142.
- 46 U. Anthoni, L. Henrisksen, P. Nielsen, G. Borch and P. Klaboe, *Spectrochim. Acta, Part A*, 1974, **30**, 1351–1364.
- 47 K. Czamara, K. Majzner, M. Z. Pacia, K. Kochan, A. Kaczor and M. Baranska, *J. Raman Spectrosc.*, 2015, **46**, 4–20.
- 48 N. Stone, C. Kendall, J. Smith, P. Crow and H. Barr, *Faraday Discuss.*, 2004, **126**, 141–157.