

# Chemical Science

Volume 14  
Number 3  
21 January 2023  
Pages 397–724

rsc.li/chemical-science



ISSN 2041-6539

## EDGE ARTICLE

Tatsuya Morofuji, Naokazu Kano *et al.*  
Streptocyanine as an activation mode of amine catalysis  
for the conversion of pyridine rings to benzene rings

Cite this: *Chem. Sci.*, 2023, 14, 485

All publication charges for this article have been paid for by the Royal Society of Chemistry

# Streptocyanine as an activation mode of amine catalysis for the conversion of pyridine rings to benzene rings†

Tatsuya Morofuji,<sup>ID</sup>\* Shota Nagai, Airi Watanabe, Kota Inagawa and Naokazu Kano<sup>ID</sup>\*

Amine catalysts have emerged as an invaluable tool in organic synthesis. Iminium, enamine, and enamine radical cation species are representative activation modes of amine catalysis. However, the development of new amine catalysis activation modes that enable novel synthetic strategies remains highly desirable. Herein, we report streptocyanine as a new amine catalysis activation mode, which enables the skeletal editing of pyridine rings to benzene rings. *N*-Arylation of pyridines bearing an alkenyl substituent at the 3-position generates the corresponding *N*-arylpiperidiniums. The resulting pyridinium reacts with a catalytic amount of piperidine to afford a streptocyanine intermediate. Catalytically generated streptocyanine forms a benzene ring *via* a ring-closing reaction, thereby releasing the amine catalyst. Consequently, the alkene moiety in the starting pyridines is incorporated into the benzene ring of the products. Pyridiniums bearing various alkene moieties were efficiently converted to formyl-substituted benzene derivatives. Mechanistic studies support the postulation that the present catalytic process was intermediated by streptocyanine. In this reaction system, streptocyanine could be regarded as a new activation mode of amine catalysis.

Received 10th November 2022

Accepted 21st December 2022

DOI: 10.1039/d2sc06225a

rsc.li/chemical-science

## Introduction

Amine catalysts are among the most commonly utilized organocatalysts and have become an invaluable organic synthetic tool, as they enable unique and sustainable syntheses of various medicinal compounds and natural products.<sup>1</sup> A representative application of amine catalysts is the activation of carbonyl compounds (Fig. 1a). Amines react with carbonyl compounds to afford iminiums, which facilitate conjugate additions, cycloadditions, and Friedel–Crafts alkylations.<sup>1a,1c,f</sup> Enamines generated by the deprotonation of iminiums are key intermediates in amine-catalyzed aldol reactions,  $\alpha$ -alkylations,  $\alpha$ -aminations, and  $\alpha$ -oxygenations of carbonyl compounds.<sup>1b,1c,f</sup> Enamine radical cations, generated *via* single-electron oxidation, can react with various nucleophiles at the  $\alpha$ -position of the original carbonyl group,<sup>2</sup> thereby expanding the potential of amine catalysis as a new activation mode.<sup>1c,1e</sup> Currently, tertiary amines are used in nucleophilic<sup>3</sup> or hydrogen atom transfer catalysis.<sup>4</sup> The development of new activation modes of amine catalysis remains highly desirable as it would contribute to the advancement of various fields of chemistry, such as drug discovery, agrochemistry, and material science.

Department of Chemistry, Faculty of Science, Gakushuin University, 1-5-1 Mejiro, Toshima-ku, Tokyo 171-8588, Japan. E-mail: tatsuya.morofuji@gakushuin.ac.jp; moro.chemistry@gmail.com; naokazu.kano@gakushuin.ac.jp

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2sc06225a>

In contrast to carbonyl compounds, aromatic rings generally cannot react with amines due to their thermodynamic stability imparted by their aromaticity. An exception is the reaction of *N*-




Fig. 1 Reaction of secondary amines (a) with carbonyl compounds, (b) with *N*-arylpiperidiniums.






Fig. 2 Working hypothesis for streptocyanine catalysis and its application to skeletal editing of pyridine rings to benzene rings. Ar = 2,4-dinitrophenyl.

arylpiperidiniums and secondary amines, which is known to afford the corresponding conjugated methine compounds, streptocyanines (Fig. 1b).<sup>5</sup> Streptocyanines are generally applicable in dye chemistry.<sup>6</sup> In addition, several stoichiometric synthetic methods for generating and converting streptocyanines have been developed.<sup>7</sup> However, to the best of our knowledge, strategies wherein streptocyanines participate in catalytic processes as an active intermediate have not been developed.

Thus, we contemplated the use of streptocyanine as a new amine catalysis activation mode. The utilization of streptocyanines generated from *N*-arylpiperidiniums and amines in a catalytic process would enable unprecedented amine-catalyzed conversions of piperidinium pyridine rings to different aromatic rings. Importantly, the development of such a reaction would provide a new strategy for the skeletal editing of aromatic rings,<sup>8,9</sup> which is one of the hottest topics in current organic chemistry.


Herein, we report the amine-catalyzed skeletal editing of piperidinium pyridine rings *via* streptocyanine intermediates to realize pyridine to benzene conversion. The working hypothesis of the reaction is outlined in Fig. 2. Reactions of pyridines bearing an alkenyl substituent at the 3-position with 2,4-dinitrophenyl tosylate affords the corresponding *N*-arylpiperidiniums. The resulting piperidiniums react with a catalytic amount of piperidine to generate streptocyanine intermediates. The catalytically generated streptocyanine forms a benzene ring *via* ring-closing reaction with concomitant release of the amine catalyst. The process allows for the incorporation of the alkene moiety in the starting pyridine into the benzene ring of the product. The outcome suggests that streptocyanine can be regarded as a new amine catalysis activation mode.

## Results and discussion

### Optimization of reaction conditions

The reaction conditions were optimized for the amine-catalyzed conversion of *N*-arylpiperidinium **1a**, which was readily prepared from the corresponding pyridine (see ESI†), to the


Table 1 Optimization of the reaction conditions<sup>a</sup>



Entry	Catalyst	Temp. (°C)	Time	Yield of <b>2a</b> (%)
1	Piperidine	40	18	58 <sup>b</sup>
2	None	40	18	0 <sup>b</sup>
3	Et <sub>2</sub> NH	40	18	12 <sup>b</sup>
4	Pyrrolidine	40	18	13 <sup>b</sup>
5	Morpholine	40	18	50 <sup>b</sup>
6	Et <sub>3</sub> N	40	18	0 <sup>b</sup>
7	DBU	40	18	0 <sup>b</sup>
8 <sup>c</sup>	Piperidine	40	18	3 <sup>b</sup>
9 <sup>d</sup>	Piperidine	40	18	8 <sup>b</sup>
10	Piperidine	120	42	98 <sup>e</sup>

<sup>a</sup> Pyridinium **1a** (0.1 mmol), amine catalyst (0.02 mmol), K<sub>2</sub>CO<sub>3</sub> (0.2 mmol), H<sub>2</sub>O (10 mmol), and THF were stirred in a pressure tube under argon atmosphere. <sup>b</sup> Yield determined by analysis of <sup>1</sup>H NMR spectroscopy. <sup>c</sup> The reaction was carried out in the absence of K<sub>2</sub>CO<sub>3</sub>. <sup>d</sup> The reaction was carried out in the absence of H<sub>2</sub>O. <sup>e</sup> Isolated yield. Ar = 2,4-dinitrophenyl.

corresponding benzene derivative **2a**. The reaction of **1a**, a catalytic amount of piperidine, potassium carbonate, and water in THF at 40 °C for 18 h delivered **2a** in 58% yield (Table 1, entry 1). The desired products were not formed in the absence



Scheme 1 Scope of benzene ring formation *via* amine-catalyzed activation of pyridine rings. Pyridinium **1** (0.1 mmol), piperidine (0.02 mmol), K<sub>2</sub>CO<sub>3</sub> (0.2 mmol), H<sub>2</sub>O (10 mmol), and THF were stirred in a pressure tube under argon atmosphere. Ar = 2,4-dinitrophenyl. Ad = adamantyl.




of piperidine (entry 2), suggesting the catalytic role of piperidine in the present transformation. The use of several other secondary amines resulted in decreased yields (entries 3–5). Triethylamine and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) did not enable product formation (entries 6 and 7). Potassium carbonate and water were found to be essential for reaction efficiency (entries 8 and 9). A higher reaction temperature and longer reaction time improved the yield of **2a** (entry 10).

### Benzene ring formation *via* amine-catalyzed activation of pyridine rings

The scope of the present benzene ring formation *via* amine-catalyzed activation of pyridine was examined, as shown in Scheme 1. Pyridiniums **1** were readily prepared from the corresponding pyridines (see ESI†). Pyridiniums bearing arylvinyl

groups afforded the corresponding products in good to excellent yields (**2a–2g**). Notably, the synthetically useful bromo-substituted derivative **2a** was tolerated in the present reaction. Benzoylvinyl-substituted pyridinium gave the product **2b** in a good yield. All *para*-, *meta*-, and *ortho*-methyl substituted aryl groups were tolerated in the reaction (**2c–2e**). *para*-Methoxy, an electron-donating group, and *para*-cyano, an electron-withdrawing group, were compatible with the reaction conditions (**2f** and **2g**). Styrylpyridine derivatives were also converted to the corresponding biphenyl derivatives (**2h–2k**). Again, the desired compounds were obtained from substrates substituted with either an electron-withdrawing or electron-donating group. In addition, pyridiniums bearing an alkenyl substituent with a variety of functional groups afforded the corresponding *meta*-substituted benzaldehydes (**2l–2p**).




**Scheme 2** Conversion of pyridiniums to various benzene derivatives using an amine catalyst. Pyridinium **1** (0.1 mmol), piperidine (0.02 mmol),  $K_2CO_3$  (0.2 mmol),  $H_2O$  (10 mmol), and THF were stirred in a pressure tube under argon atmosphere. Ar = 2,4-dinitrophenyl. <sup>a</sup> The reaction was carried out at 40 °C for 16 hours.



The present organocatalytic conversion of pyridiniums was applied to relatively complex substrates, as illustrated in Scheme 2. Treatment of pyridinium **1q**, bearing a cyclohexenone moiety, with a catalytic amount of piperidine afforded 5-formyl substituted tetralone **2q**. Importantly, both the methylene and carbonyl groups that were substituted to the alkenyl carbons in **1q** are located on *ortho*- and *meta*-carbon atoms of the formyl group, respectively, in benzene ring of **2q**. The result confirms the incorporation of the alkene moiety into the benzene ring of the product. In addition, the reported synthesis of **2q** based on electrophilic substitution cannot avoid the formation of undesired 7-formyl substituted tetralone.<sup>10</sup> In contrast, **2q** was obtained as a single regioisomer *via* the present method. Pyridiniums bearing both an aroylvinyl and phenyl group (**1r** and **1s**) were also suitable substrates for the reaction, and the corresponding benzene derivatives (**2r** and **2s**) were efficiently obtained. Notably, the synthesis of benzene derivatives bearing three different substituents at the 1,3,5- or 1,2,4-positions as in **2r** and **2s** would be challenging to achieve using conventional methods. The reaction of pyridinium **1t**, derived from piperonyl acetone, a berry-type flavor, afforded the desired compound **2t** in 78% yield. A derivative of the natural product, dihydro- $\beta$ -ionone, **1u** was transformed into **2u**. The pyridinium synthesized from pregnenolone, an endogenous steroid, was committed to the present catalytic process to obtain the corresponding benzene derivative **2v**.

The synthesis of molecules with two different carbonyl groups often becomes complex and inefficient. By contrast, because the present method enables efficient access to benzene




**Scheme 3** Formal synthesis of an anti-microtubule agent. (a) 4'-Fluoroacetophenone, NaOH, MeOH/H<sub>2</sub>O, r.t., 17 h. (b) 2,4-(NO<sub>2</sub>)<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>OTs, toluene, reflux, 20 h. (c) Piperidine (20 mol%), K<sub>2</sub>CO<sub>3</sub>, H<sub>2</sub>O, THF, 120 °C, 42 h. Ar = 2,4-dinitrophenyl.

derivatives bearing a formyl group, when pyridines bearing an  $\alpha,\beta$ -unsaturated ketone substituent were used as starting materials, benzene derivatives bearing both formyl and acyl groups in *meta*-positions were obtained in a concise manner (**2a–2g**, **2p–2v**). Furthermore, the present method was applied to the formal synthesis of anti-microtubule agent **3** (Scheme 3).<sup>11</sup> Aldol condensation of nicotinaldehyde **4** with 4'-fluoroacetophenone afforded the corresponding pyridine **5**. *N*-arylation of **5** generated pyridinium **1w**, which was subsequently converted to **2w** *via* the developed piperidine-catalyzed reaction. The total yield of the present three-step synthesis of **2w** was 80%. A previously reported synthesis of **2w** entailed four steps with a total yield of 13%.<sup>11</sup> Thus, the synthetic efficiency of **2w** has been dramatically improved by the present method. The conversion of **2w** to **3** has been accomplished in one step.


### Mechanistic studies

To gain mechanistic insights into the present catalytic process, several preliminary experiments were performed. When *N*-arylpyridinium **6** was reacted with 0.5 equiv. of piperidine, the recovery of **6** and the formation of streptocyanine **7** were observed based on <sup>1</sup>H NMR analysis (Scheme 4a). Interestingly, conjugated *N*-arylimine **8**, which has been proposed as a reaction intermediate in the synthesis of streptocyanine,<sup>5a,5e</sup> was not



**Scheme 4** Mechanistic studies. (a) Reaction of *N*-arylpyridinium **6** with 0.5 equiv. of piperidine. (b) HRMS analysis. Ar = 2,4-dinitrophenyl.





Scheme 5 Plausible reaction mechanism for benzene ring formation *via* amine-catalyzed skeletal-editing of the pyridine ring in pyridiniums.

observed. The results support that the active species of the present benzene ring formation is streptocyanine, as described in Fig. 2. To further verify this hypothesis, **1a** was treated with piperidine at 0 °C for 1.5 h and the reaction mixture was analyzed using HRMS, which confirmed the presence of streptocyanine intermediate **9** (Scheme 4b). In addition, an ion peak consistent with protonated cyclohexadiene **10** was found when the reaction mixture was analyzed after stirring at 0 °C for 1 h and then at room temperature for 2 h.

Based on the results, a plausible reaction mechanism for the amine-catalyzed skeletal-editing of pyridinium pyridine rings is described in Scheme 5. Piperidine attacks the 2-position of pyridinium **1** to generate piperidine adduct **A**. 6 $\pi$ -Electrocyclic ring-opening of **A** delivers **B**. A second nucleophilic addition of piperidine to the imine moiety of **B** generates streptocyanine **D** through iminal **C**. Streptocyanine **D** is then converted thermally to the *E/Z* isomer **E** because of the push-pull electronic structure.<sup>5d</sup> Ring closing of **E** *via* a 6 $\pi$ -electrocyclic reaction and hydrolysis of resulting **F** affords cyclohexadiene **G**. It should be noted that streptocyanines are generally stable against hydrolysis under weakly basic conditions because their positive charge is delocalized. Therefore, although the hydrolysis of **D** or **E** could not be excluded, that of readily hydrolysable conjugated imine **F** would be more plausible. Aromatization of **G** results in the production of desired product **2** and regeneration of the

piperidine catalyst. This mechanism could be regarded as an amine-catalyzed variation of ANRORC reactions.<sup>12</sup> Overall, the use of streptocyanine as an amine catalysis activation mode enabled the present process, entailing ring opening of a pyridine ring, thermal *E/Z* isomerization due to the unique electronic structure of streptocyanine, and 6 $\pi$ -electrocyclization to form a benzene ring.

## Conclusions

In summary, we have demonstrated that streptocyanine can be used as a new activation mode of amine catalysis and applied to the conversion of pyridine rings to benzene rings. Pyridinium derivatives bearing various alkene moieties were efficiently converted to formyl benzene derivatives. Moreover, the developed process is applicable to complex molecules derived from a flavor or natural products. The synthetic utility of the method was demonstrated *via* the efficient syntheses of an anti-microtubule agent and benzene derivatives bearing three different substituents. Mechanistic studies supported the hypothesis that streptocyanine is an intermediate in the present catalytic process. It is anticipated that streptocyanine catalysis will have an impact on various chemical areas, just as iminiums, enamines, and radical cations of enamines have done.



## Data availability

All the experimental procedures and characterization data are available in the ESI.†

## Author contributions

T. M. conceptualized the project and designed the experiments. T. M. and N. K. directed the project. S. N., A. W., and K. I. performed the experiments and analyzed the data. T. M. and N. K. co-wrote the manuscript. All authors discussed the experiments and commented on the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- (a) A. Erkkilä, I. Majander and P. M. Pihko, *Chem. Rev.*, 2007, **107**, 5416–5470; (b) S. Mukherjee, J. W. Yang, S. Hoffmann and B. List, *Chem. Rev.*, 2007, **107**, 5471–5569; (c) D. W. C. Macmillan, *Nature*, 2008, **455**, 304–308; (d) C. Grondal, M. Jeanty and D. Enders, *Nat. Chem.*, 2010, **2**, 167–178; (e) M. Mečiarová, P. Tisovský and R. Šebesta, *New J. Chem.*, 2016, **40**, 4855–4864; (f) Y.-Q. Zou, F. M. Hörmann and T. Bach, *Chem. Soc. Rev.*, 2018, **47**, 278–290.
- K. Narasaka, K. Okauchi, K. Tanaka and M. Murakami, *Chem. Lett.*, 1992, **21**, 2099–2102.
- (a) S. France, D. J. Guerin, S. J. Miller and T. Lectka, *Chem. Rev.*, 2003, **103**, 2985–3012; (b) M. Song, J. Zhao and E.-Q. Li, *Chin. Chem. Lett.*, 2022, **33**, 2372–2382.
- (a) J. D. Bell and J. A. Murphy, *Chem. Soc. Rev.*, 2021, **50**, 9540–9685; (b) W. Xiao, X. Wang, R. Liu and J. Wu, *Chin. Chem. Lett.*, 2021, **32**, 1847–1856.
- (a) W.-C. Cheng and M. J. Kurth, *Org. Prep. Proced. Int.*, 2002, **34**, 585–608; (b) T. Zincke, G. Heuser and W. I. Möller, *Justus Liebigs Ann. Chem.*, 1904, **333**, 296–345; (c) S. E. Steinhardt, J. S. Silverston and C. D. Vanderwal, *J. Am. Chem. Soc.*, 2008, **130**, 7560–7561; (d) R. S. Paton, S. E. Steinhardt, C. D. Vanderwal and K. N. Houk, *J. Am. Chem. Soc.*, 2011, **133**, 3895–3905; (e) T. Morofuji, A. Shimizu and J.-I. Yoshida, *J. Am. Chem. Soc.*, 2013, **135**, 5000–5003; (f) B. T. Boyle, J. N. Levy, L. de Lescure, R. S. Paton and A. McNally, *Science*, 2022, **378**, 773–779.
- H. Mustroph, *Phys. Sci. Rev.*, 2021, **6**, 137–147.
- (a) E. h. Ziegler and K. Hafner, *Angew. Chem.*, 1955, **67**, 301; (b) G. Köbrich, *Justus Liebigs Ann. Chem.*, 1961, **648**, 114–123; (c) C. Jutz, R.-M. Wagner, A. Kraatz and H.-G. Löbering, *Justus Liebigs Ann. Chem.*, 1975, **1975**, 874–900; (d) I. Yamaguchi, S. Shingai and M. Sato, *Macromolecules*, 2008, **41**, 6292–6298; (e) A. Colombo, C. Dragonetti, S. Righetto, D. Roberto, A. Valore, T. Benincori, F. Colombo and F. Sannicolò, *J. Mater. Chem.*, 2012, **22**, 19761; (f) T. Morofuji, H. Kinoshita and N. Kano, *Chem. Commun.*, 2019, **55**, 8575–8578; (g) T. Morofuji, K. Inagawa and N. Kano, *Org. Lett.*, 2021, **23**, 6126–6130.
- For reviews of skeletal editing of aromatic rings, see; (a) B. Chattopadhyay and V. Gevorgyan, *Angew. Chem., Int. Ed.*, 2012, **51**, 862–872; (b) A. M. Prokhorov and D. N. Kozhevnikov, *Chem. Heterocycl. Compd.*, 2012, **48**, 1153–1176; (c) P. Anbarasan, D. Yadagiri and S. Rajasekar, *Synthesis*, 2014, **46**, 3004–3023; (d) K. Nogi and H. Yorimitsu, *Chem. Commun.*, 2017, **53**, 4055–4065; (e) Y. Li, H. Wang and X. Li, *Chem. Sci.*, 2020, **11**, 12249–12268; (f) J. Jurczyk, J. Woo, S. F. Kim, B. D. Dherange, R. Sarpong and M. D. Levin, *Nat. Synth.*, 2022, **1**, 352–364.
- For selected examples of skeletal editing of aromatic rings, see; (a) S. Chuprakov, F. W. Hwang and V. G. Gevorgyan, *Angew. Chem., Int. Ed.*, 2007, **46**, 4757–4759; (b) D. Vasu, H. Yorimitsu and A. Osuka, *Angew. Chem., Int. Ed.*, 2015, **54**, 7162–7166; (c) T. Jin, H. Yuan, S. Su, X. Jia, C. Li, J. Li and J. Fang, *Chem. Commun.*, 2018, **54**, 14128–14131; (d) B. D. Dherange, P. Q. Kelly, J. P. Liles, M. S. Sigman and M. D. Levin, *J. Am. Chem. Soc.*, 2021, **143**, 11337–11344; (e) L. Wang, H. Han, L. Gu, W. Zhang, J. Zhao and Q. Wang, *Chem. Sci.*, 2021, **12**, 15389–15398; (f) A. Kaga, H. Iida, S. Tsuchiya, H. Saito, K. Nakano and H. Yorimitsu, *Chem. – Eur. J.*, 2021, **27**, 4567–4572; (g) S. C. Patel and N. Z. Burns, *J. Am. Chem. Soc.*, 2022, **144**, 17797–17802; (h) J. C. Reisenbauer, O. Green, A. Franchino, P. Finkelstein and B. Morandi, *Science*, 2022, **377**, 1104–1109; (i) J. Woo, A. H. Christian, S. A. Burgess, Y. Jiang, U. F. Mansoor and M. D. Levin, *Science*, 2022, **376**, 527–532; (j) E. E. Hyland, P. Q. Kelly, A. M. McKillop, B. D. Dherange and M. D. Levin, *J. Am. Chem. Soc.*, 2022, **144**, 19258–19264.
- Y. Ning, T. Fukuda, H. Ikeda, Y. Otani, M. Kawahata, K. Yamaguchi and T. Ohwada, *Org. Biomol. Chem.*, 2017, **15**, 1381–1392.
- Y. Yamazaki, M. Sumikura, Y. Masuda, Y. Hayashi, H. Yasui, Y. Kiso, T. Chinen, T. Usui, F. Yakushiji, B. Potts, S. Neuteboom, M. Palladino, G. K. Lloyd and Y. Hayashi, *Bioorg. Med. Chem.*, 2012, **20**, 4279–4289.
- (a) T. Fujii and T. Itaya, *Heterocycles*, 1998, **48**, 359–390; (b) E. S. H. E. Ashry, Y. E. Kilany, N. Rashed and H. Assafir, *Adv. Heterocycl. Chem.*, 1999, **75**, 79–165.

