Chemical Science

EDGE ARTICLE

Cite this: Chem. Sci., 2023, 14, 13060

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 24th August 2023 Accepted 25th October 2023

DOI: 10.1039/d3sc04457b

rsc.li/chemical-science

Introduction

Photochemical a-selective radical ring-opening reactions of 1,3-disubstituted acyl bicyclobutanes with alkyl halides: modular access to functionalized cyclobutenes†

Yuanjiu Xiao[,](http://orcid.org/0000-0002-6313-4546) \mathbf{D} \ddagger Tong-Tong Xu, \mathbf{D} \ddagger Jin-Lan Zhou, \mathbf{D} Feng Wu, \mathbf{D} Lei Tang, \mathbf{D} Ruo-Yi Liu, Wen-Biao Wu \mathbb{D}^* and Jian-Jun Feng \mathbb{D}^*

Although ring-opening reactions of bicyclobutanes bearing electron-withdrawing groups, typically with bselectivity, have evolved as a powerful platform for synthesis of cyclobutanes, their application in the synthesis of cyclobutenes remains underdeveloped. Here, a novel visible light induced a-selective radical ring-opening reaction of 1,3-disubstituted acyl bicyclobutanes with alkyl radical precursors for the synthesis of functionalized cyclobutenes is described. In particular, primary, secondary, and tertiary alkyl halides are all suitable substrates for this photocatalytic transformation, providing ready access to cyclobutenes with a single all-carbon quaternary center, or with two contiguous centers under mild reaction conditions. **EDGE ARTICLE**
 (a) Check for undates
 EDGE ARTICLE
 Photochemical α **-selective radical ring-opening

case the clom so 2003.14.1300

CHERE CONTROLLS CONTROLLS CONTROLLS CONTROLLS CONTROLLS**
 CONTROLLS CONTROLLS C

Four-membered rings such as cyclobutanes and cyclobutenes are important frameworks found in an array of natural products and pharmaceuticals with diverse biological and medicinal properties (Scheme 1a).¹ Meanwhile, they also serve as key intermediates in the synthesis of structurally complex targets.² Therefore, the development of new methods for their efficient synthesis has been intensively pursued by the synthetic community. Despite the significant advances made in the area for the synthesis of four-membered rings, modular strategies for the synthesis of these structures remain relatively few in number, particularly in comparison to the available methodologies to synthesize five- and six-membered rings.³ Specifically, synthesis of multisubstituted functionalized cyclobutanes and cyclobutenes featuring quaternary carbon stereocenters remains challenging.⁴

As the smallest of bicyclic hydrocarbons, bicyclo^[1.1.0] butanes (BCBs) are 'spring-loaded' yet bench-stable molecules.⁵ The potential energy stored in the bridging bond (66.3 kcal mol⁻¹ (calculated))⁶ has been harnessed for applications in the

State Key Laboratory of Chemo/Biosensing and Chemometrics, Advanced Catalytic Engineering Research Center of the Ministry of Education, College of Chemistry and Chemical Engineering, Hunan University, Changsha, Hunan 410082, P. R. China. E-mail: 1334154923@qq.com; jianjunfeng@hnu.edu.cn

[†] Electronic supplementary information (ESI) available: Experimental details, characterization, and spectroscopic data. CCDC 2287475. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d3sc04457b>

[‡] These authors contributed equally.

Scheme 1 Ring-opening reactions of BCBs for synthesis of fourmembered rings and their scientific context.

face several challenges such as (a) previous reports on the radical ring-openings of BCBs were limited to monosubstituted BCBs. Again, these reactions occur exclusively at the β -position of the bridging bond of BCBs (Scheme 1b);¹⁴ (b) although α selective ring-opening reactions of monosubstituted bicyclo [1.1.0]butyl boronic ester (BCB-Bpin) with polar heteroatom nucleophiles were reported by Aggarwal and co-workers in 2021, a β -addition pathway was observed when β -methyl-substituted BCB-Bpin was used as the substrate (Scheme 1c).^{13a} To the best of our knowledge, there is no report on the α -selective ringopening reactions of BCBs with radical nucleophiles. (c) So far, most of the transformations have focused on the conjugate addition-type reactivity of BCBs for synthesis of cyclobutanes,¹¹⁻¹⁴ while less work has been reported on the selective construction of cyclobutenes from BCBs via ringopening reactions.¹⁶ As a rare example, the Leitch group reported Lewis acid catalyzed polar electrophilic ring-opening reactions of BCBs with N-alkylimines to generate cyclobutenyl methanamine products with poor to moderate yields (Scheme 1d).^{16a} Therefore, discovery of new catalytic systems for efficient synthesis of valuable cyclobutene products is highly desirable.

As part of our ongoing interest in small ring chemistry¹⁷ and in order to expand the library of known BCB reactivity, we became interested in whether 1,3-disubstituted BCBs could be employed as radical acceptors in radical ring-opening reactions. Along this line, other intriguing questions arise: what type of radical substrate is suitable for the reaction? How to control the regioselectivity? Can this reaction lead to cyclobutenes? Here,

we describe a modular strategy for the synthesis of functionalized cyclobutenes *via* α -selective ring-opening reactions of acyl BCBs with alkyl radicals, which could be generated from the commercially available 1°, 2°, and 3° alkyl bromides by photolysis of C–Br and C–Cl bonds (Scheme 1e).¹⁸

Results and discussion

Initial experiments on a representative α -bromo ketone $2a^{18a}$ and 1.3-disubstituted BCB ester $1a$ under the irradiation of blue LEDs revealed that the commonly used metal-based photocatalysts (PCs) (Ru(bpy)₃Cl₂, Ru(bpy)₃(PF₆)₂, and Ir[dF(CF₃) ppy]₂[dtbbpy]PF₆) and organic-based PCs (Eosin Y, [Acr⁺-Mes] ClO4, and 4-CzlPN) were lowly effective, affording the desired product 3aa with \leq 7% NMR yield (see Table S1, in the ESI†). To our delight, $fac-Ir(ppy)$ ₃ gave a moderate yield of the desired cyclobutene 3aa along with a 5% yield of 4a and cyclobutane 5aa (12% yield) in MeCN (Table 1, entry 1). Importantly, no aimed product 3aa was observed only with either the $fac-Ir(ppy)$ ₃ catalyst or light, which indicated that photoredox catalysis is essential to this ring-opening process (entries 2 and 6). Subsequent investigations demonstrated the importance of solvent (entries 3–5). The use of DMF as the solvent dramatically improved the yield of 3aa from 30% to 58% (entry 5 versus 1). The base also had a substantial effect on the product distribution but no improvement over K_2HPO_4 was seen (entries 7-10).

Table 1 Optimization of reaction conditions⁶

Conditions: 1a (0.2 mmol), 2a (0.3 mmol), $fac-Ir(ppy)_{3}$ (2 mol%) and base (1.0 equiv.) in solvent (2.0 mL) at 25 °C under the irradiation of 12 W blue LEDs for 12 h. b NMR yield with CH₂Br₂ as an internal standard. Isolated yields are indicated in parentheses. \textdegree Without fac-Ir(ppy)₃. ^d Reaction in the absence of light irradiation. ^e H₂O (0.5) equiv.) was added. ^{*f*} Reaction time: 3 h. ^{*g*} Without base. ^{*h*} α -Chloroacetophenone instead of 2a.

Further improvement of the yield was achieved with dimethylacetamide (DMA) instead of DMF as the solvent (entry 11). Notably, the ring-opening reaction can tolerate H_2O (entry 12). Only a 3-h reaction time was needed to reach >99% conversion (entry 13). In the absence of a base, the reaction yield was reduced from 71 to 31% (entry 14). Rather than alkyl bromide 2a, a-chloroacetophenone was also competent to afford the product, albeit with a lower yield (entry 15).

With optimized reaction conditions established, we tested a range of different alkyl bromides (Scheme 2A). The reaction of bromoacetophenones with different substituents on the aryl ring, including electron-donating (4-OMe as in 3ab) and electron-withdrawing groups $(4-CF_3 \text{ as in } 3\text{ae}; 4-CN \text{ as in } 3\text{af})$, at either the para, meta, or ortho position, also proceeded with good efficiency (3aa–3ai). The naphthyl-containing a-bromo ketone 2j furnished 3aj in 91% yield. Additionally, heterocyclic a-bromo ketone 2k underwent the reaction smoothly to deliver

the products in moderate yield (3ak). Aside from aromatic ketones, alkyl ketone 2l containing a cyclopropyl group, can be smoothly transformed into the corresponding product 3al. α -Carbonyl alkyl bromides bearing cyano $(2m)$,¹⁹ amide $(2n)$ and ester (2o) groups are also suitable substrates. Besides primary alkyl bromides, secondary alkyl bromides are still maintaining decent yields (3ap–3aq). Photoredox catalysis has recently demonstrated potential for the construction of congested allcarbon quaternary centers via highly reactive radicals.¹⁸ However, intermolecular construction of two contiguous allcarbon quaternary centers by visible light-induced photoredox catalysis remains underdeveloped.^{18p} To our delight, tertiary-abromoalkyl esters (2r–2t) successfully afforded the corresponding congested cyclobutenes (3ar–3at) in 49–95% yields with exclusive α -selectivity. Importantly, ethyl bromodifluoroacetate also had high reactivity and delivered product 3au bearing a gem-difluoromethylene (CF_2) group, which is popular

Scheme 2 Substrate scope investigation.^{a a} Unless otherwise noted, reactions were performed with 1 (0.2 mmol), 2 (0.30 mmol), fac-Ir(ppy)₃ (2 mol%), K₂HPO₄ (1.0 equiv.), and DMA (2.0 mL), at room temperature under irradiation with 12 W blue LEDs for 3 h (condition A). Isolated yields are indicated. b Condition B: 1 (0.2 mmol)), 2 (0.30 mmol), fac-Ir(ppy)₃ (2 mol%), Na₂CO₃ (1.0 equiv.), Et₃N (0.1 equiv.), and DMF (2.0 mL), at room temperature under irradiation with 12 W blue LEDs for 12 h. c The ring-opening reactions of BCBs (1p-1t) with α -bromo ketone 2a were conducted using condition A. d Combined isolated yield of the diastereomers.</sup>

in drugs and in agrochemicals. Furthermore, tertiary-alkyl bromide 2v with a nitro group smoothly reacted with 1a and gave the product 3av in reasonable yield. Besides α -carbonyl alkyl bromides, less-reactive benzyl bromide derivative 2w was also a suitable substrate in the presence of $fac-Ir(ppy)$ ₃ as the photocatalyst under reoptimized condition B.

The scope and generality of this α -selective ring-opening in terms of 1,3-disubstituted BCBs with representative alkyl bromides (2j, 2s and 2u) is summarized in Scheme 2B. This method is amenable to a series of BCBs bearing different aromatic substituents, including alkyl $(1b, 1f \& 1i)$, CF₃O $(1c)$, halogen (1d, 1e, 1g & 1h) and CF_3 (1l) groups at the para-, metaor ortho-position of aryl rings, and led to the corresponding cyclobutenes in moderate to excellent yields (50–80%). Both naphthyl (1j) and thienyl (1k) substituted BCBs reacted smoothly under condition A. Apart from BCB esters (1b–1m), both BCB ketone 1n and Weinreb amide derived BCB 1o were compatible, giving the corresponding ring-opening products bearing two contiguous quaternary centers and $CF₂$ groups in synthetically useful yield $(1n + 2u \rightarrow 3nu; 1o + 2u \rightarrow 3ou)$. The reaction is not limited to aryl-substituted BCBs. Alkylsubstituted BCB 1p was also found to be compatible, albeit with a low yield (5% NMR yield) under conditions A (for more details, see ESI Scheme S1†). 1,3-Disubstituted sulfonyl BCB 1q did not furnish the desired cyclobutene product. Additionally, we investigated the influence of the substituent at the β -position of BCB in this site-selective reaction. Although the ringopening reaction of monosubstituted BCB ketone 1r (or BCB amide 1s) results in a complex mixture, the reaction of monosubstituted BCB sulfone 1t with 2a produces the cyclobutane product 6 through β -addition (Scheme 2C). Edge Article

in dux and in agreednmical, Furthermore, teriary-ality apony of Jaj afforded de free comparison when are presented with a mode various are ality the mode of the second various are ality the model on a simple

To explore synthetic transformations of these functionalized cyclobutenes, a scale-up synthesis of 3aj (1.0 mmol) was performed without any loss in efficiency and selectivity (Scheme 3). The double bond in 3aj was hydrogenated over Pd/C to produce trisubstituted cyclobutane 7 in 90% yield. The ketone group in 3aj was converted into an alkene in 8. Hydrolysis of the ester group of 3aj afforded the free carboxylic acid 9. Moreover, reduction of the ester group in $3am$ using NaBH₄ provided the primary alcohol 10 in 86% yield. Also, functionalized cyclobutene 11 was synthesized from 10 by hydrolysis of the nitrile under basic conditions. Of note, reduction of the Weinreb amide 3ou with $LiAlH₄$ followed by oxidation allowed for the construction of spirolactone 12 containing a 4-membered ring.

To rationalize the reaction pathway, some experiments were designed and performed. First, the reaction was completely shut down in the presence of TEMPO. In the meantime, the radical trapping product 13 was obtained in 82% yield (Scheme 4A). Second, the reaction of 2-bromoacetonitrile with 1a and 4 methoxystyrene was attempted under condition A, and the cyclobutene 3am together with 14 and 15 was detected by 1 H NMR and HRMS, indicating that the reaction proceeds via a radical pathway with α -selectivity (Scheme 4B). Subsequently, the treatment of 4a with 2a under the standard reaction conditions did not furnish the aimed 3aa (Scheme 4C). This result demonstrated that the reaction was not proceeding via the cyclobutene intermediate 4a. In addition, as shown in Scheme 4D, when 5aa was treated with the photocatalyst under optimized conditions, no desired product was observed.

Moreover, the light–dark experiment was performed using alternative intervals of light and dark (Scheme 5). It has been demonstrated that this photochemical transformation necessitates continuous exposure to visible light, indicating that chain propagation may not be the primary reaction pathway. A Stern–Volmer plot showed strong quenching of $fac-Ir(ppy)_{3}$ ($E^{IV/}$ $I^{\text{III}*}$ = −1.84 V vs. SCE) by alkyl bromide 2a (E_{ox} > 1.6 V vs. SCE, $E_{\text{red}} = -1.2$ V vs. SCE),^{18q} favoring an oxidative quenching cycle (Scheme 6 and Fig. S2–S4, in the ESI†).

Based on these preliminary results, a reasonable mechanistic rationale for our photoredox transformation was formulated, as shown in Scheme 7. Upon photoexcitation, the photocatalyst $fac-Ir(ppy)$ ₃ is converted into the strongly reducing excited state Ir $(m)^*$, which is capable of reducing an alkyl halide 2 to its corresponding alkyl radical **INT-A** and generating $Ir(w)$. Upon α -selective radical addition of the electrophilic species INT-A to BCB 2, the key C–C bond formation step occurred, thus

Scheme 3 Scale-up synthesis and synthetic transformations. Scheme 4 Control experiments.

Scheme 5 Light on/off experiment.

Scheme 6 Stern–Volmer plots of $fac-lr(ppy)_3$ with different quenchers (1a or 2a)

Scheme 7 Proposed mechanism

forming the more stable tertiary radical INT-B. Then, INT-B undergoes a single-electron oxidation with $Ir(w)$ to form the carbocation INT-C with the regeneration of the photocatalyst. Finally, the cyclobutene 3 can be formed through an elimination reaction with the assistance of a base.

Conclusion

In summary, we report the first use of ring-opening reactions of 1,3-disubstituted bicyclobutanes with alkyl radical precursors to construct cyclobutenes via visible light-induced photoredox catalysis. This study provides a concise and modular method for the synthesis of highly valuable alkylated cyclobutenes. A broad range of alkyl halides including 1°, 2°, and 3° alkyl bromides

can be coupled under mild reaction conditions to afford 1,1,3 trisubstituted cyclobutenes with a single all-carbon quaternary center, or with two contiguous (all-carbon quaternary) centers. The synthetic utility of the reaction was showcased by several transformations. Notably, unlike the β -selectivity disclosed in classical polar and radical strain-release reactions of BCBs, an exclusive a-selectivity in current radical ring-openings of acyl-BCBs with carbon-centered radicals is obtained, which is complementary to Aggarwal's a-selective ring-opening reactions of monosubstituted BCB-Bpin with heteroatom-centred polar nucleophiles.^{13a}

Data availability

All detailed procedures, characterization data and NMR spectra are available in the ESI.†

Author contributions

Y. X., T.-T. X., J.-L. Z., F. W., L. T., and R.-Y. L. performed the experiments and conducted the analytical characterization. W.- B. W. and J.-J. F. wrote the manuscript. J.-J. F. conceived the catalytic system.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We are grateful to the Fundamental Research Funds for the Central Universities.

References

- 1 (a) C. Hui, Y. Liu, M. Jiang and P. Wu, Trends Chem., 2022, 4, 677–681; (b) M. R. van der Kolk, M. A. C. H. Janssen, F. P. J. T. Rutjes and D. Blanco-Ania, ChemMedChem, 2022, 17, e202200020; (c) V. M. Dembitsky, J. Nat. Med., 2008, 62, 1–33.
- 2 For reviews, see: (a) J. Chen, Q. Zhu, H. Fang and P. Lu, Chin. J. Chem., 2022, 40, 1346–1358; (b) J. Li, K. Gao, M. Bian and H. Ding, Org. Chem. Front., 2020, 7, 136–154; (c) M. Wang and P. Lu, Org. Chem. Front., 2018, 5, 254–259; (d) T. Seiser, T. Saget, D. N. Tran and N. Cramer, Angew. Chem., Int. Ed., 2011, 50, 7740–7752; (e) E. Lee-Ruff and G. Mladenova, Chem. Rev., 2003, 103, 1449–1483; (f) J. C. Namyslo and D. E. Kaufmann, Chem. Rev., 2003, 103, 1485–1537; (g) R. Guo and M. K. Brown, Acc. Chem. Res., 2023, 56, 2253– 2264.
- 3 (a) D. Didier and F. Reiners, Chem. Rec., 2021, 21, 1144–1160; (b) Y. Xu, M. L. Conner and M. K. Brown, Angew. Chem., Int. Ed., 2015, 54, 11918–11928; (c) N. Gauvry, C. Lescop and F. Huet, Eur. J. Org Chem., 2006, 5207–5218.
- 4 For reviews, see: (a) K.-G. Wen, Y.-Y. Peng and X.-P. Zeng, Org. Chem. Front., 2020, 7, 2576–2597; (b) E. N. Hancock, J. M. Wahl and M. K. Brown, Nat. Prod. Rep., 2019, 36,

1383–1393; For representative examples on synthesis of quaternary carbon containing cyclobutenes, see: (c) H. Xu, W. Zhang, D. Shu, J. B. Werness and W. Tang, Angew. Chem., Int. Ed., 2008, 47, 8933–8936; (d) A. Masarwa, A. Fürstner and I. Marek, Chem. Commun., 2009, 5760– 5762; (e) Y. Odabachian and F. Gagosz, Adv. Synth. Catal., 2009, 351, 379-386; (f) V. López-Carillo and A. M. Echavarren, J. Am. Chem. Soc., 2010, 132, 9292–9294; (g) F. Frébault, M. Luparia, M. T. Oliveira, R. Goddard and N. Maulide, Angew. Chem., Int. Ed., 2010, 49, 5672–5676; (h) C. Schotes and A. Mezzetti, Angew. Chem., Int. Ed., 2011, 50, 3072–3074; (i) S. Niyomchon, D. Audisio, M. Lupariaa and N. Maulide, Org. Lett., 2013, 15, 2318–2321; (j) M. M. Maturi and T. Bach, Angew. Chem., Int. Ed., 2014, 53, 7661–7664; (k) K. Enomoto, H. Oyama and M. Nakada, Chem.–Eur. J., 2015, 21, 2798–2802; (l) T. Kang, S. Ge, L. Lin, Y. Lu, X. Liu and X. Feng, Angew. Chem., Int. Ed., 2016, 55, 5541-5544; (m) C. García-Morales, B. Ranieri, I. Escofet, L. López-Suarez, C. Obradors, A. I. Konovalov and A. M. Echavarren, J. Am. Chem. Soc., 2017, 139, 13628– 13631; (n) F.-G. Zhang and I. Marek, J. Am. Chem. Soc., 2017, 139, 8364–8370; (o) E. N. Hancock, E. L. Kuker, D. Tantillo and M. K. Brown, Angew. Chem., Int. Ed., 2020, 59, 436–441; (p) C. Zhong, Y. Huang, H. Zhang, Q. Zhou, Y. Liu and P. Lu, Angew. Chem., Int. Ed., 2020, 59, 2750– 2754; (q) F. W. Goetzke, A. M. L. Hell, L. van Dijk and S. P. Fletcher, Nat. Chem., 2021, 13, 880–886; (r) M. Yan, Q. Zhou and P. Lu, Angew. Chem., Int. Ed., 2023, 62, e202218008. Edge Article Chemotics Article Chemotics are synthesis of Org. Lett. 2019, 21:32-730-730; [f] Y. Ling, B. Richman are chemotic is licensed under a Creative Commons Attribution-NonCommercial 3.0 American and African and Af

- 5 (a) M. Golfmann and J. C. L. Walker, Commun. Chem., 2023, 6, 9; (b) C. B. Kelly, J. A. Milligan, L. J. Tilley and T. M. Sodano, Chem. Sci., 2022, 13, 11721–11737; (c) A. Fawcett, Pure Appl. Chem., 2020, 92, 751–765; (d) J. Turkowska, J. Durka and D. Gryko, Chem. Commun., 2020, 56, 5718–5734; (e) M. A. A. Walczak, T. Krainz and P. Wipf, Acc. Chem. Res., 2015, 48, 1149–1158.
- 6 K. B. Wiberg, Angew. Chem., Int. Ed., 1986, 25, 312–322.
- 7 (a) A. Cairncross and E. P. Blanchard Jr, J. Am. Chem. Soc., 1966, 88, 496–504; (b) P. Wipf and M. A. A. Walczak, Angew. Chem., Int. Ed., 2006, 45, 4172–4175; (c) R. Kleinmans, T. Pinkert, S. Dutta, T. O. Paulisch, H. Keum, C. G. Daniliuc and F. Glorius, Nature, 2022, 605, 477–482; (d) R. Guo, Y.-C. Chang, L. Herter, C. Salome, S. E. Braley, T. C. Fessard and M. K. Brown, J. Am. Chem. Soc., 2022, 144, 7988–7994; (e) M. Xu, Z. Wang, Z. Sun, Y. Ouyang, Z. Ding, T. Yu, L. Xu and P. Li, Angew. Chem., Int. Ed., 2022, 61, e202214507; (f) Y. Liu, S. Lin, Y. Li, J.-H. Xue, Q. Li and H. Wang, ACS Catal., 2023, 13, 5096–5103; (g) N. Radhoff, C. G. Daniliuc and A. Studer, Angew. Chem., Int. Ed., 2023, e202304771; (h) R. Kleinmans, S. Dutta, K. Ozols, H. Shao, F. Schäfer, R. E. Thielemann, H. T. Chan, C. G. Daniliuc, K. N. Houk and F. Glorius, J. Am. Chem. Soc., 2023, 145, 12324-12332; (i) S. Agasti, F. Beltran, E. Pye, N. Kaltsoyannis, G. Crisenza and D. J. Procter, Nat. Chem., 2023, 15, 535–541; (j) M. Wang, Y. Huang, C. Li and P. Lu, Org. Chem. Front., 2022, 9, 2149– 2153; (k) X. Ma, D. L. Sloman, Y. Han and D. J. Bennett,

Org. Lett., 2019, 21, 7199–7203; (l) Y. Liang, R. Kleinmans, C. G. Daniliuc and F. Glorius, J. Am. Chem. Soc., 2022, 144, 20207–20213; (m) Y. Liang, F. Paulus, C. G. Daniliuc and F. Glorius, Angew. Chem., Int. Ed., 2023, e202305043; (n) Y. Zheng, W. Huang, R. K. Dhungana, A. Granados, S. Keess, M. Makvandi and G. A. Molander, J. Am. Chem. Soc., 2022, 144, 23685-23690; (o) T. V. T. Nguyen, A. Bossonnet and J. Waser, ChemRxiv, 2023, preprint, DOI: [10.26434/chemrxiv-2023-s8j30](https://doi.org/10.26434/chemrxiv-2023-s8j30); (p) T. Yu, J. Yang, Z. Wang, Z. Ding, M. Xu, J. Wen, L. Xu and P. Li, J. Am. Chem. Soc., 2023, 145, 4304–4310; (q) R. Bychek and P. K. Mykhailiuk, Angew. Chem., Int. Ed., 2022, e202205103; (r) H. Wang, H. Shao, A. Das, S. Dutta, H. T. Chan, C. Daniliuc, K. N. Houk and F. Glorius, Science, 2023, 381, 75.

- 8 H. K. Hall Jr and A. B. Padias, J. Polym. Sci., Part A: Polym. Chem., 2003, 41, 625–635.
- 9 (a) K. Tokunaga, M. Sato, K. Kuwata, C. Miura, H. Fuchida, N. Matsunaga, S. Koyanagi, S. Ohdo, N. Shindo and A. Ojida, J. Am. Chem. Soc., 2020, 142, 18522–18531; (b) P. Zhang, R. Zhuang, X. Wang, H. Liu, J. Li, X. Su, X. Chen and X. Zhang, *Bioconjugate Chem.*, 2018, 29, 467-472; (c) B. D. Schwartz, A. P. Smyth, P. E. Nashar, M. G. Gardiner and L. R. Malins, Org. Lett., 2022, 24, 1268–1273.
- 10 A. Kaur, W. Lin, V. Dovhalyuk, L. Driutti, M. L. D. Martino, M. Vujasinovic, J.-M. Löhr, M. E. Sellin and D. Globisch, Chem. Sci., 2023, 14, 5291–5301.
- 11 (a) Y. Gaoni, Tetrahedron Lett., 1982, 23, 5215–5218; (b) R. Panish, S. R. Chintala, D. T. Boruta, Y. Fang, M. T. Taylor and J. M. Fox, J. Am. Chem. Soc., 2013, 135, 9283–9286; (c) R. A. Panish, S. R. Chintala and J. M. Fox, Angew. Chem., Int. Ed., 2016, 55, 4983–4987; (d) M. J. Kerner and P. Wipf, Org. Lett., 2021, 23, 3615–3619; (e) A. Guin, S. Bhattacharjee, M. S. Harariya and A. T. Biju, Chem. Sci., 2023, 14, 6585–6591.
- 12 (a) Y. Gaoni, Tetrahedron Lett., 1988, 29, 1591–1594; (b) S. Hoz, C. Azran and A. Sella, J. Am. Chem. Soc., 1996, 118, 5456–5461; (c) R. Gianatassio, J. M. Lopchuk, J. Wang, C.-M. Pan, L. R. Malins, L. Prieto, T. A. Brandt, M. R. Collins, G. M. Gallego, N. W. Sach, J. E. Spangler, H. Zhu, J. Zhu and P. S. Baran, Science, 2016, 351, 241–246; (d) J. M. Lopchuk, K. Fjelbye, Y. Kawamata, L. R. Malins, C.-M. Pan, R. Gianatassio, J. Wang, L. Prieto, J. Bradow, T. A. Brandt, M. R. Collins, J. Elleraas, J. Ewanicki, W. Farrell, O. O. Fadeyi, G. M. Gallego, J. J. Mousseau, R. Oliver, N. W. Sach, J. K. Smith, J. E. Spangler, H. Zhu, J. Zhu and P. S. Baran, J. Am. Chem. Soc., 2017, 139, 3209– 3226; (e) J. A. Milligan, C. A. Busacca, C. H. Senanayake and P. Wipf, Org. Lett., 2016, 18, 4300–4303; (f) B. D. Schwartz, M. Y. Zhang, R. H. Attard, M. G. Gardiner and L. R. Malins, Chem.–Eur. J., 2020, 26, 2808–2812; (g) M. Takatsuki, H. Aoyama, K. Murai, M. Arisawa and M. Sako, Chem. Commun., 2023, 59, 7467–7470; (h) Z. Zhang and V. Gevorgyan, J. Am. Chem. Soc., 2022, 144, 20875–20883.
- 13 (a) L. Guo, A. Noble and V. K. Aggarwal, Angew. Chem., Int. Ed., 2021, 60, 212-216; (b) S. H. Bennett, A. Fawcett, E. H. Denton, T. Biberger, V. Fasano, N. Winter and

V. K. Aggarwal, J. Am. Chem. Soc., 2020, 142, 16766–16775; (c) B. Wölfl, N. Winter, J. Li, A. Noble and V. K. Aggarwal, Angew. Chem., Int. Ed., 2023, e202217064; (d) H.-C. Shen, M. V. Popescu, Z.-S. Wang, L. de Lescure, A. Noble, R. S. Paton and V. K. Aggarwal, J. Am. Chem. Soc., 2023, 145, 16508–16516; (e) A. Fawcett, T. Biberger and V. K. Aggarwal, Nat. Chem., 2019, 11, 117–122.

- 14 (a) G. Ernouf, E. Chirkin, L. Rhyman, P. Ramasami and J.-C. Cintrat, Angew. Chem., Int. Ed., 2020, 59, 2618–2622; (b) C. J. Pratt, R. A. Aycock, M. D. King and N. T. Jui, Synlett, 2020, 31, 51–54; (c) X. Wu, W. Hao, K.-Y. Ye, B. Jiang, G. Pombar, Z. Song and S. Lin, J. Am. Chem. Soc., 2018, 140, 14836–14843; (d) H. Gao, L. Guo, C. Shi, Y. Zhu, C. Yang and W. Xia, Adv. Synth. Catal., 2022, 364, 2140– 2145; (e) X. Yu, M. Lübbesmeyer and A. Studer, Angew. Chem., Int. Ed., 2021, 60, 675–679; (f) M. Silvi and V. K. Aggarwal, J. Am. Chem. Soc., 2019, 141, 9511–9515.
- 15 M. Ociepa, A. J. Wierzba, J. Turkowska and D. Gryko, J. Am. Chem. Soc., 2020, 142, 5355–5361.
- 16 (a) K. Dhake, K. J. Woelk, J. Becica, A. Un, S. E. Jenny and D. C. Leitch, Angew. Chem., Int. Ed., 2022, 61, e202204719; (b) M. Pomerantz, R. N. Wilke, G. W. Gruber and U. Roy, J. Am. Chem. Soc., 1972, 94, 2752–2758; (c) E. P. Blanchard Jr and A. Cairncross, J. Am. Chem. Soc., 1966, 88, 487–495.
- 17 (a) C.-Z. Zhu, J.-J. Feng and J. Zhang, Angew. Chem., Int. Ed., 2017, 56, 1351–1355; (b) J.-J. Feng and J. Zhang, ACS Catal., 2017, 7, 1533–1542; (c) T.-Y. Lin, H.-H. Wu, J.-J. Feng and J. Zhang, ACS Catal., 2017, 7, 4047–4052; (d) L. Tang, Q.-N. Huang, F. Wu, Y. Xiao, J.-L. Zhou, T.-T. Xu, W.-B. Wu, S. Qu and J.-J. Feng, Chem. Sci., 2023, 14, 9696– 9703; (e) L. Tang, Y. Xiao, F. Wu, J.-L. Zhou, T.-T. Xu and J.-J. Feng, Angew. Chem., Int. Ed., 2023, e202310066.
- 18 For reviews, see: (a) X.-H. Ouyang, R.-J. Song and J.-H. Li, Chem.–Asian J., 2018, 13, 2316–2332; (b) M. M. D. Pramanik, H. Qian, W.-J. Xiao and J.-R. Chen, Org. Chem. Front., 2020, 7, 2531–2537; (c)

A. L. G. Kanegusuku and J. L. Roizen, Angew. Chem., Int. Ed., 2021, 60, 21116–21149; (d) A. Lipp, S. O. Badir and G. A. Molander, Angew. Chem., Int. Ed., 2021, 60, 1714– 1726; (e) S. Poplata, A. Tröster, Y.-Q. Zou and T. Bach, Chem. Rev., 2016, 116, 9748–9815; (f) D. Staveness, I. Bosque and C. R. J. Stephenson, Acc. Chem. Res., 2016, 49, 2295–2306; (g) D. M. Schultz and T. P. Yoon, Science, 2014, 343, 1239176; (h) D. P. Hari and B. König, Angew. Chem., Int. Ed., 2013, 52, 4734–4743; (i) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, Chem. Rev., 2013, 113, 5322–5363; (j) J. Xuan and W.-J. Xiao, Angew. Chem., Int. Ed., 2012, 51, 6828–6838; For selected references, see: ; (k) D. A. Nicewicz and D. W. C. MacMillan, *Science*, 2008, 322, 77–80; (l) J. D. Nguyen, J. W. Tucker, M. D. Konieczynska and C. R. J. Stephenson, J. Am. Chem. Soc., 2011, 133, 4160-4163; (m) C.-J. Wallentin, J. D. Nguyen, P. Finkbeinera and C. R. J. Stephenson, J. Am. Chem. Soc., 2012, 134, 8875–8884; (n) Q. Liu, H. Yi, J. Liu, Y. Yang, X. Zhang, Z. Zeng and A. Lei, Chem.–Eur. J., 2013, 19, 5120–5126; (o) Y. Zhu, L. Zhang and S. Luo, J. Am. Chem. Soc., 2014, 136, 14642–14645; (p) W. Chen, Z. Liu, J. Tian, J. Li, J. Ma, X. Cheng and G. Li, J. Am. Chem. Soc., 2016, 138, 12312–12315; (q) T. Lei, W.-Q. Liu, J. Li, M.-Y. Huang, B. Yang, Q.-Y. Meng, B. Chen, C.-H. Tung and L.-Z. Wu, Org. Lett., 2016, 18, 2479–2482; (r) M. Silvi, C. Sandford and V. K. Aggarwal, J. Am. Chem. Soc., 2017, 139, 5736–5739; (s) D. F. J. Caputo, C. Arroniz, A. B. Dürr, J. J. Mousseau, A. F. Stepan, S. J. Mansfield and E. A. Anderson, Chem. Sci., 2018, 9, 5295–5300; (t) J. Li, M. Kong, B. Qiao, R. Lee, X. Zhao and Z. Jiang, Nat. Commun., 2018, 9, 2445; (u) B. Schweitzer-Chaput, M. A. Horwitz, E. de P. Beato and P. Melchiorre, Nat. Chem., 2019, 11, 129–135; (v) X. Zhang, R. T. Smith, C. Le, S. J. McCarver, B. T. Shireman, N. I. Carruthers and D. W. C. MacMillan, Nature, 2020, 580, 220–226. Chemical Science

Y. K. Aggrewalt, *J.an.* Chem, 36, 3023, 142, 1576-16775; (c) A. L. G. Kanagasaha and J. L. Ruice, Argewalt, and Access Article

A. W. The published on 27 October 2023, 142, 162, 2023, 2023, 2023, 2023,

19 CCDC 2287475 (3am).†