


RSC Chemical Biology

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Y. Hashimoto, S. Shil, M. Tsuruta, K. Kawauchi and D. Miyoshi, *RSC Chem. Biol.*, 2024, DOI: 10.1039/D4CB00287C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Three- and four-stranded nucleic acid structures and their ligands

Yoshiki Hashimoto^a, Sumit Shil^a, Mitsuki Tsuruta^a, Keiko Kawauchi^a, and Daisuke Miyoshi^{*a}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Nucleic acids have the potential to form not only duplexes, but also various non-canonical secondary structures in living cells. Non-canonical structures play regulatory functions mainly in the central dogma. Therefore, nucleic acid targeting molecules are potential novel therapeutic drugs that can target 'undruggable' proteins in various diseases. One of the concerns of small molecules targeting nucleic acids is selectivity, because nucleic acids have only four different building blocks. Three- and four-stranded non-canonical structures, triplexes and quadruplexes, respectively, are promising targets of small molecules because their three-dimensional structures are significantly different from the canonical duplexes, which are the most abundant in cells. Here, we describe some basic properties of the triplexes and quadruplexes and small molecules targeting the triplexes and tetraplexes.

Introduction

Most small-molecule therapeutics in clinical use currently target proteins. Human cells express about 20,000 species of proteins, following the hypothesis of "one gene expresses one protein". Within these proteins, up to 15%, corresponding to 3000 species of proteins, are disease related (Fig. 1)¹⁻³. Within the disease-related proteins, many of the remaining ones are termed 'undruggable', which refers to proteins that are difficult to target pharmacologically. An undruggable protein often lacks a binding site for a small molecule. Intrinsically disordered proteins, lacking fixed or ordered structures, are also often undruggable. The most well-known examples of the former and the latter undruggable proteins are RAS⁴ and MYC⁵, respectively, although these two proteins are important targets for cancer therapeutics. Moreover, recent interactome studies suggest that there are up to 650,000 protein-protein interactions (PPIs) in humans⁶. PPIs are also considered undruggable by small molecules because of the large, flat, and featureless surfaces of PPIs⁷. In fact, only a few percent of these PPIs can be targeted with small drugs^{8,9}. Therefore, alternative strategies for the inactivation of these disease-related undruggable proteins are required.

The 20,000 species of proteins in a human cell are encoded by only 1.5% of the human genome, which contains 3,000,000,000 base pairs in total (Fig. 1)^{10, 11}. Among these proteins, fewer than 700 have been targeted by drugs. This corresponds to only 0.05% of the genetic information encoded in the human genome. However, around 80% of the human genome is transcribed to RNA. As described above, DNA coding amino acid information corresponds to 1.5% of the genome, so the remaining part of the 80% is non-coding RNA (ncRNA)¹², which controls the central dogma at various steps, such as





Fig. 1 Schematic diagram depicting protein expression in the central dogma and the potential nucleic acid-targeted druggable genome.

replication, transcription, processing and localization of mRNA, and translation. When compared with druggable proteins, there is far more opportunity to target RNAs to expand druggability and improve our understanding of biological processes governing the central dogma. If DNA and RNA can be targeted, protein products can be up- or down-regulated at the transcriptional and translational levels.

The canonical structures of DNA and RNA are B-form and A-form duplexes, respectively, with Watson-Crick base pairs. Canonical duplexes are suitable and highly adapted for their functions: storage, inheritance, and the transition of genetic information. However, both DNA and RNA can fold to form various non-canonical and transient structures (Fig. 2a~m), depending on the nucleotide sequence as well as the surrounding molecular environment. Similar to a knot providing a firm place to grip a rope (Fig. 2n), non-canonical structures can be recognition sites and controlling elements for a series of enzymes, including DNA and RNA polymerase, helicase, topoisomerase, nuclease, DNA and RNA modifiers, spliceosome, and ribosome, as well as transcriptional factors¹³⁻¹⁵. In fact, roughly 1,500 proteins are predicted to bind to RNA in a sequence- and structure-selective manner¹⁶. Small molecules targeting DNA and RNA could regulate not only the expression of proteins but also the functions of the DNA- and

^a Frontiers of Innovative Research in Science and Technology, Konan university, 7-1-20 Minatojima-minamimachi, Chuo-ku, Kobe, Hyogo, 650-0047, Japan.





View Article Online
DOI: 10.1039/D4CB00287C

Fig. 2 Schematic illustration of a canonical duplex (a) and non-canonical structures (b-m) of nucleic acids. (b) Mismatch, (c) parallel duplex, (d) hairpin loop, (e) bulge, (f) internal loop, (g) three-way junction, (h) four-way junction, (i) pseudoknot, (j) kissing loop, (k) triplex, (l) G-quadruplex, and (m) i-motif. Black arrows indicate strand orientation. Red and blue bars with circles represent canonical Watson-Crick and non-canonical base pairs, respectively. (n) Schematic illustration of proteins targeting non-canonical structures on the long canonical duplex of nucleic acids.

RNA-binding proteins. Thus, small molecules targeting non-canonical structures have drawn attention as new therapeutic regimens¹⁷⁻¹⁹. Risdiplam, sold under the brand name Evrysdi, represents a milestone given its status as the first small molecule drug that targets RNA²⁰.

One of the concerns of small molecules targeting nucleic acids is selectivity, because nucleic acids have only four different building blocks, whereas proteins are composed of 20 amino acids. In addition, it is generally considered that nucleic acid structures are static and monotonous. In contrast, it has been discussed that there are up to 10,000 unique protein folds in nature²¹⁻²³. From this point of specificity, three- and four-stranded non-canonical structures, triplexes and quadruplexes, respectively, are promising because their three-dimensional structures are significantly different from the canonical duplexes. Noteworthy, recent progresses in bioinformatics, DNA and RNA sequencing, and bioimaging have demonstrated the existence of the multiplexes and their biological roles in living cells of various organisms from virus to human. In this review, we briefly describe the triplex and quadruplex structures of nucleic acids and their potential biological roles. We then describe recent progress in the development of small molecules that target these non-canonical nucleic acid structures.

Triplexes

Structure and biological role of triplexes

The canonical duplex has two grooves formed by the sugar-phosphate backbones of the two strands. The grooves play a vital role in interactions with other molecules, such as protein, small molecules, metal ion, and water. The grooves further play an important role in the sequence-specific recognition of the duplex by single-stranded nucleic acids, either DNA or RNA. These interactions are mediated by the formation of Hoogsteen-type hydrogen bonds between the bases of the third

strand, often called a triplex-forming oligonucleotide (TFO), and the purine bases of the duplex formed by a polypurine-polypyrimidine sequence (Fig. 3a). The first example of a triplex




Fig. 3 (A) Base pairs in parallel and antiparallel triplexes. (B) Schematic illustration of glycosidic conformation for nucleobases in TFO.



was reported by Felsenfeld *et al.* in 1967. They found the formation of an intermolecular RNA triplex by two strands of polyribo(uridylic acid) and one strand of polyribo(adenylic acid)²⁴. After the discovery of the triplex, many triplex structures under various conditions have been identified²⁵. There are two classes of triplex motifs, distinguished by the orientation of the TFO, which binds to the major groove of the target duplex. A polypyrimidine TFO binds to the polypurine strand of the duplex via Hoogsteen hydrogen bonds in a parallel fashion (in the same orientation as the purine strand of the duplex). In contrast, a polypurine TFO binds in an antiparallel orientation to the polypurine strand of the duplex via reverse-Hoogsteen hydrogen bonds^{26, 27}. In an antiparallel triplex, the base pairs are G:G-C and A:A-T (“:” and “-” indicate Hoogsteen and Watson-Crick base pairs, respectively). In a parallel triplex, the base pairs are C⁺:G-C and T:A-T (C⁺ represents a protonated cytosine). Thus, the ideal target of a TFO is a duplex of a homopurine sequence in one strand and a homopyrimidine sequence in the complementary strand. For both parallel and antiparallel motifs, contiguous homopurine-homopyrimidine runs of at least 10 base pairs are required for thermodynamically stable TFO binding²⁸. In addition, all TFOs adopt *anti*-conformations (Fig. 3b)²⁹, although 8-oxoadenin in a TFO can adopt a *syn*-conformation³⁰. The *anti*-conformation in a TFO is the same as that of nucleotides in B- and A-form duplexes.

Large numbers of potential triplex forming sites have been identified in the human genome³¹. Most annotated genes in the human genome contain at least one triplex-forming site, and these sites are enriched in the promoter regions³², indicating a potential role of triplex in controlling gene expression. These TFO-binding sites have attracted much attention because of their therapeutic potential to inhibit the expression of disease-related genes. Evidence for the existence of triplexes in cells comes from triplex-binding antibodies^{33, 34}. However, small molecules probing the triplex are still required to observe the direct dynamic behavior of the triplex over time in living cells.

RNA triplexes

Triplexes have been observed in transcripts such as tRNA^{Phe}, telomerase RNA, self-splicing group II introns, U2-U6 snRNAs, riboswitches, virus RNAs, and RNA stability elements³⁵. These RNA triplexes play roles in telomere synthesis, RNA splicing, binding to ligands and ions in metabolite-sensing riboswitches, and protecting RNA from degradation.

Telomerase RNAs in various eukaryotes have a conserved pseudoknot in which a triplex exists^{36, 37}. The triplex structure is essential for telomerase catalytic activity³⁸, demonstrating that the triplex structure rather than the nucleotide sequence is conserved and required for the ribonucleoprotein function. Triplexes observed in riboswitches are also important as binding sites of a series of ligands. For example, SAM-II riboswitch has triplex-forming sites that serve as structural platforms for ligand binding³⁹. RNA triplexes have also been proposed for various viruses, such as Kaposi's sarcoma-associated herpesvirus (KSHV)⁴⁰. The triplexes protect the viral RNAs and inhibit decay by host nuclear RNA degradation to accumulate in KSHV-




Fig. 4 Secondary structures of MALAT1, PAN, and MEN β RNAs.

infected cells^{41, 42}. Human metastasis-associated lung adenocarcinoma transcript 1 (MALAT1), a long noncoding RNA (lncRNA), folds to form a triplex (Fig. 4). MALAT1 is localized in nuclear speckles and is involved in the regulation of gene expression^{43, 44}. MALAT1 is upregulated in multiple cancer types, depending on the triplex formation, making it a potential drug target^{45, 46}. Unlike many lncRNAs that are rapidly degraded and thus expressed at near-undetectable levels, MALAT1, as well as MEN β and PAN, which are also known to form triplexes (Fig. 4), are stable transcripts with long half-lives⁴⁷. It has been proposed that the triplexes formed by these lncRNAs inhibit nuclear RNA decay⁴⁸. Although the biological roles of RNA stability elements remain unclear, the triplex supports the transport, stability, and translation of an RNA, allowing efficient repression by microRNAs.

Moreover, it was recently reported that lncRNAs bind target DNAs in a sequence-specific manner by forming RNA-DNA triplexes^{49, 50}. This process allows lncRNAs to recruit proteins to specific genomic regions and to regulate gene expression. lncRNAs such as Fendrr⁵¹, MEG3⁵², KHPS1⁵³, PARTICLE^{54, 55}, and HOTAIR⁵⁶ bind to their target DNAs in a sequence-specific manner, utilizing the RNA-DNA triplexes. It has further been proposed by computational analysis that there are many triplex target sites of lncRNAs in the genome⁵⁷. Interestingly, such bioinformatic analysis has revealed a group of genomic regions that may have a very high propensity for triplex formation with a wide range of different RNAs⁵⁸. Excellent reviews have discussed the biological roles of RNA-DNA triplexes^{59, 60}.

Triplex-binding proteins

It has been reported that some proteins such as histones H1 and H2A, and topoisomerase II, bind to AT-rich sequences⁶¹⁻⁶³. A



Table 1. Triplex-binding proteins

Protein	Function / target	Ref
GAGA factor	gene regulation and alteration of chromatin structure / pyrimidine motif	67
Stm1	involved in mitosis / purine motif	68
CDP1	chromosome segregation and histone displacement / purine motif	69
type III intermediate filaments	Gene regulation, DNA repair, and chromatin remodelling / pyrimidine motif	70
loricrin	keratinocyte cornification / purine motif	71
Orc4	origin organization during DNA replication / purine motif	72
hnRNPs	DNA and RNA binding / synthetic oligo DNAs	73
Tn7	transposon insertion / pyrimidine motif	74, 75
XPC-hHR23B, XPA-RPA	DNA damage recognition / TFO-directed psoralen-interstrand crosslink	76
HMG1	chromatin structure, transcriptional regulation / triplex-directed psoralen interstrand crosslinks	77
U2AF65, PSF, p54nrb	splicing factors / purine motif	78
RecQ, BLM, WRN,	3' → 5' helicase / pyrimidine motif with a free 3' tail	79
FANCF	5' → 3' helicase / pyrimidine motif with a free 3' tail	80
DHX9	3' → 5' helicase / purine motif with a free 3' tail	81
SV40 large T-antigen	3' → 5' helicase / pyrimidine motif with a free 3' tail	82
p53	tumor suppressor / intra and intermolecular pyrimidine motif	83

View Article Online

DOI: 10.1039/D4CB00287C

protein that binds preferentially to TAT DNA triplexes was reported for the first time in 1991, although its identification was not possible at that time⁶⁴. Proteins that specifically recognize certain DNA triplex motifs have been reported^{65, 66}. Since the 2000s, many proteins have been identified as DNA triplex-binding proteins⁶⁷⁻⁸³ (Table 1). Although the biological roles of these DNA triplex-binding proteins are highly diverse, it is reasonable that some helicases are triplex-binding proteins. In addition, nuclear proteins are known to bind DNA/RNA triplexes, but their biological functions are unexplored⁸⁴.

Small molecules that target triplexes

The Hoogsteen base pairs between the TFO and the target duplex are generally weaker than the Watson-Crick base pairs in the duplex⁸⁵. The instability of the triplex under physiological conditions limits its biological roles and applications. Of note, a parallel triplex is stable only at acidic pH, because the protonation of cytosines is required for Hoogsteen base pairing, although the stability of antiparallel triplexes does not depend on solution pH⁸⁶. Moreover, multivalent cations such as Mg²⁺ are required to neutralize the electrostatic repulsion between the three nucleic acid strands. The triplex but not the duplex is stabilized under molecular crowding conditions that mimic a cellular environment⁸⁷. To overcome the limited thermal stability, small molecules as well as chemical modifications of TFOs have been applied.

Hélène and co-workers demonstrated the enhancement of triplex thermal stability by attaching small molecules such as acridine, ellipticine, and psoralen to the TFO⁸⁸⁻⁹⁰. Small molecules that bind strongly and selectively to the triplex can also stabilize the triplex without covalent modification. Some triplex-binding small molecules are shown in Fig. 5. Ethidium bromide stabilizes a triplex containing only TAT base pairs⁹¹, whereas it destabilizes a triplex containing both TAT and C⁺GC base pairs because of electrostatic repulsion with C⁺⁹². Some of the earliest discovered small molecules that stabilize the TAT and C⁺GC of triplexes are benzopyrindole derivatives (BePI and

BgPI) and coralyne⁹³⁻⁹⁶. BePI stabilizes TAT-rich triplexes, whereas coralyne does not show sequence selectivity. BePI enhances the sequence-specific inhibition of transcription initiation of a specific gene by RNA polymerase by a TFO. Quinoxaline derivatives BfPQ and BQQ were further developed as analogues of BPI derivatives⁹⁷⁻⁹⁹. These molecules have been shown to significantly stabilize triplexes under physiological conditions, whereas they bind poorly to duplexes. Notably, BQQ promotes a short TFO to bind to its target site in a plasmid and enhances the inhibition of restriction enzyme cleavage of the target site.

Wilson and co-workers proposed that the structural properties of a triplex intercalator include the following: (i) the compounds should be cationic because of the high negative charge density of the triplex. (ii) The compounds should have an aromatic surface to form stacking interactions with the three bases in the triplex. (iii) The aromatic compounds should have torsional flexibility, because the three bases in a triplex have torsional freedom¹⁰⁰. Based on these properties, they synthesized a series of quinoline derivatives and found that a 2-naphthyl derivative could bind to poly(dA)-2poly(dT) to a much greater extent than to the corresponding duplex¹⁰⁰. Other triplex intercalators, such as fisetin¹⁰¹, luteolin¹⁰², palmatine¹⁰³, sanguinarine¹⁰⁴, and ruthenium(II) complexes¹⁰⁵, are also shown in Fig. 5.

Groove binders have also been demonstrated to have affinity with DNA triplexes. It has been shown that netropsin, a drug that interacts with the minor groove of a DNA duplex, binds the minor groove of a triplex and destabilizes it, but stabilizes the duplex^{106, 107}. Umemoto *et al.* showed by using distamycin 2, an analogue of netropsin, that the drug does not destabilize¹⁰⁸. Breslauer and coworkers showed that berenil, DAPI, ethidium, and netropsin induce the formation of a TAT DNA/RNA/DNA triplex¹⁰⁹. In addition, a series of small molecules such as methyl green and mithramycin, which bind and destabilize triplexes, have been reported¹¹⁰⁻¹¹². Aminoglycosides, especially neomycin, stabilize a triplex, while they do not alter the stability






Fig. 5 Chemical structures of triplex-targeting molecules.

of a duplex¹¹³. Triplex stabilization by neomycin is salt- and pH-dependent, and increasing the concentrations of K⁺ and Mg²⁺ and the solution pH decreases the stabilization effects of neomycin^{114, 115}. These aminoglycosides bind with not only DNA triplexes, but also RNA and DNA/RNA triplexes¹¹⁵. A combination of intercalation and groove binding has also been reported. Intercalator–neomycin conjugates have been studied to enhance binding affinity with DNA triplexes. BQQ-neomycin enhances the binding affinity with a TAT DNA triplex up to almost 1000-fold more than that of neomycin through a dual recognition mode: groove binding and intercalation of neomycin and BQQ, respectively¹¹⁶.


As described above, helix-forming MALAT1 lncRNA is a potential anticancer target due to its overexpression in cancer cells, and its knockdown reduces tumor growth. Deletion and mutation studies in the triplex-forming region have shown that alterations in the stability of the triplex lead to significant changes in transcript levels^{117, 118}. Several small molecules, such as diphenylfuran derivatives, have been reported as lncRNA triplex-targeting molecules¹¹⁹. Small molecules targeting the MALAT1 triplex with high selectivity have also been reported, of which a representative example is compound 5 (Fig. 5)¹²⁰. Compound 5 was obtained by small molecule microarray screening of a library comprising approximately 26,000 compounds. Compound 5 regulates MALAT expression. Since NEAT1 has a triplex structure, these results confirm that compound 5 selectively binds to the MALAT1 triplex in living

cells. NMR experiments and MD simulations demonstrated that compound 5 binds within the deep and narrow major groove of the MALAT1 triplex through several van der Waals interactions with the phosphate and nucleobase groups of the Hoogsteen strand and the Watson–Crick strands. An RT-qPCR experiment showed that the MALAT1 expression level in mouse model mammary tumor organoids was suppressed by 51% following treatment with 1 μM of compound 5 whereas NEAT1 expression was unaffected¹²⁰, demonstrating the high sensitivity of compound 5. Molecules such as compound 5, which specifically target the MALAT1 triplex, are a promising new class of anticancer therapeutics and molecular probes for the treatment and investigation, respectively, of cancers driven by MALAT1.

G-quadruplexes

A G-quadruplex (G4) structure of guanylic acid was reported for the first time by Gellert *et al.* in 1962¹²¹. The G4 structure of a guanine-rich telomeric sequence was then revealed by Sen and Gilbert in 1988¹²². These studies made us realize that guanine-rich sequences can form G4s (Fig. 6a). Putative G4-forming sequences are generally considered to have the following consensus: G₂ ≤ L₁ ≤ G₂ ≤ L₁ ≤ G₂ ≤ L₁ ≤ G₂ ≤, in which guanine stretches composing more than two guanines are connected by linker loop sequences (L) including at least one nucleotide¹²³. These putative G4-forming motifs are observed ubiquitously in diverse organisms' genomes. Among the various





View Article Online
DOI: 10.1039/D4CB00287C

Fig. 6 (a) Schematic illustration of the folding of a guanine-rich (G-rich) sequence. The G-rich sequence folds into a G-quadruplex (G4) via a G-quartet. The three-dimensional structure of G4 is adapted from PDB ID: 2HY9. Guanine bases are highlighted in red. (b) Schematic illustration of a telomeric end with a double-stranded G-rich strand (red) and cytosine-rich (C-rich) strand (blue) and with a single-stranded G-rich region. Guanine and cytosine bases are highlighted in red and blue, respectively. (c) Three-dimensional structure of parallel (PDB ID: 2M27, Ref. 135), antiparallel (PDB ID: 6ZX7, Ref. 146), and hybrid (PDB ID: 2HY9, Ref. 137) G4s.

putative G4-forming sequences, the most well-studied G4 is that of telomere DNA. Human telomere DNA comprises repetitive sequences of the hexanucleotide d(TTAGGG)_n repeat unit. In normal human somatic cells, telomere DNA ranges from 5–15 kb in length with an extended single-stranded 3'-overhang of a few hundred bases (Fig. 6b)^{124–126}. In each cell division, the telomere DNA is shortened because of the end replication problem, which leads to apoptosis¹²⁷. In contrast, the telomere DNA length is maintained in cancer cells. In most cancers, telomerase, a ribonucleoprotein, extends the telomere 3' overhang by its reverse transcription activity^{128, 129}. Thus, telomerase inhibitors provide a new approach to cancer chemotherapy¹³⁰. The hybridization between the telomere DNA and its complementary cytosine-rich sequence in the telomerase RNA is essential for the reverse transcription activity¹³¹. Formation and stabilization of the G4 in the telomere overhang by a small molecule can inhibit telomerase activity, and thus cancer cell proliferation^{132, 133}.

A G4 exhibits various topologies, parallel, antiparallel, and hybrid, in an intermolecular and intramolecular manner (Fig. 6c)^{134, 135–137}. G4s show such highly structural polymorphisms depending on the nucleotide sequence as well as molecular environmental factors¹³⁸. Moreover, G4 guanines can adopt either an *anti*- or *syn*-conformation, resulting in different B-form duplexes. The resulting pronounced variations in the glycosidic torsion angles result in four grooves of different sizes, which could be important for the specificity of G4 ligands

binding to target G4s. Parallel G4s comprised exclusively of guanines adopt the *anti*-glycosidic conformation (Fig. 6c)¹³⁹ whereas in an antiparallel G4, guanines adopt both *anti*- and *syn*-conformations and at least one of the four strands must be oriented antiparallel to the other strands (Fig. 6c)¹³⁹. In this chapter, factors affecting the structure and stability of G4 will be briefly explained.

Factors affecting the structure and stability of G4 originating from the nucleotide sequence

Hydrogen bonding: Like the canonical duplex, hydrogen bonding interactions within a G-quartet are critical for the structural stability of G4. The four guanine bases in a coplanar arrangement form eight Hoogsteen-type hydrogen bonds (Fig. 6a). The hydrogen bonds are formed between the N1 on one guanine with the O6 on the neighbor guanine, and the N2 on the first guanine with the N7 on the second guanine, resulting in eight cyclic hydrogen bonds in each G-quartet. By using a series of modified bases, it has been found that the central part of the G-quartet, including the central hydrogen bonds, contributes more than the outer hydrogen bonds to stabilize G4¹⁴⁰.

π - π stacking interactions: Not only hydrogen bonds, but also the π - π stacking interactions between G-quartets, which may be one of the largest planer aromatic surfaces of biomolecules, are central to the structure and stability of G4 (Fig. 7)^{141–144}. A greater number of G-quartets leads to more stable G4s,



although the G4 structures reported so far contain two to four G-quartets. The stacking interactions among intermolecular G-quartets can induce multimerization of G4s. Phan *et al.* reported the dimerization of G4s of (GGGT)₄¹⁴⁵⁻¹⁴⁷. Multimerization of G4s further induces a high-order structure, known as a G-wire^{36, 148-153}. A G-wire-like formation has been proposed as a possible structure of the long telomere DNA, though other high-order structures, such as a bead-on-a-string structure, have also been proposed¹⁵⁴⁻¹⁵⁶.

Loop composition: An intramolecular G4 has three loops connecting the four guanine stretches. An intermolecular G4 of two strands has two loops. Some interactions, such as stacking and hydrogen bonding interactions between bases in the loops and between the bases in the loops and the neighboring G-quartets, are also important for the stability and topology of G4 (Fig. 7). It has been reported that loop length plays a role in the structure and stability of G4^{123, 157-160}. Generally, longer loops result in a more unstable G4. Mergny and co-workers demonstrated that each added base in a loop destabilizes it by 2 °C of melting temperature or by 0.3 kcal/mol of ΔG° in the presence of potassium ions¹⁶¹. Moreover, the same group showed that both the conformation and thermal stability are greatly dependent on loop permutation¹²³. The G4 conformation depending on the loop length has also been observed in other studies. Hazel *et al.* found that a parallel intramolecular G4 is folded when three loops with a single base are present¹⁵⁸. When single thymine loops are present in combination with longer length loops, or when all loops compose more than two bases, both parallel and antiparallel G4s can be formed¹⁵⁸.

Flanking sequences: The flanking sequences next to the first and last guanine stretches also play a role in the structure and stability of G4s. The flanking nucleotides at the 5' and 3' ends can form additional stacking and hydrogen bonding interactions with the neighboring G-quartet or the coalescing loop (Fig. 7). For example, the capping structures formed by the extended flanking nucleotides have been shown to contribute to the stabilization of the c-MYC promoter G4 and the G4 aptamer to the hemagglutinin of influenza A virus^{162, 163}. The flanking sequences further affect the whole G4 structure. The flanking

bases contribute to a particular type of G4 by the specific capping structures formed by the flanking and loop bases¹⁶⁴. In addition, completely truncated sequences without any flanking residues are often prone to form higher-order intermolecular structures¹⁶². These studies lead to the conclusion that each G4 in DNA and RNA has a unique capping structure formed by its specific flanking and loop sequence, and it contributes to the stability and overall structure of a G4¹⁶⁶.

G4-duplex hybrid: Phan and coworkers structurally demonstrated by NMR that thermally stable G4-duplex junction can be formed in various topological combinations (Fig. 8a)¹⁶⁷⁻¹⁶⁹. The G4-duplex hybrid can be identified not only by NMR but also by a fluorescent probe and an intrinsic fluorescence of G4^{170, 171}. Importantly, a combined genomic and structural study of the G4-duplex hybrids identified around 80,000 G4-duplex hybrids-forming sequences in the human genome. About 60% of these sequences were strand-specifically located in genic/promoter regions¹⁷². These G4-duplex hybrids have been reported to play multiple roles in determination of topology, folding and unfolding kinetics, and cation binding of G4¹⁷³⁻¹⁷⁶. Moreover, G4s involving the duplex showed more efficient transcriptional inhibition^{170, 177}. Since the G4-duplex hybrid-forming DNA sequences were identified in the sense and antisense strands¹⁷², corresponding RNA structure can exist in naturally occurring RNAs such as PIM1 mRNA (Fig. 8b)¹⁷⁸ as well as synthetic RNA aptamer such as SC1¹⁷⁹ and Spinach^{180, 181}. The G4-duplex hybrid can be a promising target in diverse biological and pathological processes, because the junction between the G4 and the duplex gives distinct structure from the "canonical" G4, which has shorter loop length. Moreover, the duplex region could be useful for sequence selective binding for a target G4.

Factors affecting the structure and stability of G4 originating from the surrounding environment

Cations: A cation coordinates the O6 carbonyl from the guanines in a G-quartet, resulting in reduced electrostatic repulsion of the central oxygen atoms, enhancing hydrogen bonding interactions, and stabilizing the π - π stacking interactions¹⁸². Therefore, cations are critical for the structure




Fig. 7 Factors affecting the structure and stability of G4. Note that not all factors are shown for clarity. For example, many more hydrogen bonds and stacking interactions are formed in these structures.




Fig. 8 (a) Schematic structure of the G4-duplex hybrids. G-quartet and Watson-Crick base pair are shown in gray and yellow. Green shows non-canonical base pairs. (grey), (b) Three-dimensional G4-duplex hybrid structure in PIM1 gene (PDB: 7CV3). The duplex region is highlighted in yellow.



and stability of G4s. The stabilization effects of monovalent and divalent cations on the thrombin binding aptamer G4 are on the order of $\text{Sr}^{2+} > \text{Ba}^{2+} > \text{K}^+ > \text{Rb}^+ > \text{NH}_4^+ \sim \text{Na}^+ > \text{Cs}^+ (\sim \text{Li}^+)$ (Fig. 9a)¹⁸³⁻¹⁸⁷. In particular, Sr^{2+} and K^+ have a large stabilization effect on G4s¹⁸⁸. These cation effects are common for DNA and RNA G4s with different nucleotide sequences. Thus, the dependency of structural stability on a coexisting cation is often utilized as supporting evidence of G4 formation of a nucleotide sequence of interest. In contrast to the stabilization effects of Sr^{2+} and Ba^{2+} , in the presence of the physiologically relevant monovalent cations K^+ and Na^+ , low concentrations of divalent cations destabilize G4 in the following order: $\text{Zn}^{2+} > \text{Co}^{2+} > \text{Mn}^{2+} > \text{Mg}^{2+} > \text{Ca}^{2+}$ ¹⁸⁹⁻¹⁹¹. Moreover, divalent metal ions at higher concentrations induce a structural transition from an antiparallel G4 to a parallel G4¹⁹¹.

The ion radius of a cation is critical for binding with G4. It is generally considered that this is because there is an optimal size that fits into the cavity of the G-quartet(s). It is also evident that dehydration is required for cation coordination. Thus, the hydration energy is important to determine the structural stability of G4s^{144, 186}. Hud *et al.* studied by NMR the competition between Na^+ and K^+ for coordination with G-quartets using the oligonucleotide $d(\text{G}_3\text{T}_4\text{G}_3)$, which forms an antiparallel G4 in the presence of Na^+ and K^+ (Fig. 9b)¹⁹². They found that the more favored binding of K^+ than Na^+ is driven by the greater dehydration energy of Na^+ than K^+ . Hydration from cations is also important for the structure and stability of G4s under cell-mimicking molecular crowding conditions, as explain in the next section.

Molecular crowding: The molecular environment inside of cells is surprisingly crowded with many biomolecules^{87, 193-197}. The total concentration of biomolecules reaches 400 g/L, and about 40% of the cell volume is occupied by them. These

biomolecules inside cells result in molecular crowding conditions that are totally different from those in a test tube, where most biochemical experiments are carried out. Some key interactions such as hydration and counterion condensation, which are required to form nucleic acid structures, are dependent on the surrounding molecular conditions. Moreover, biochemical reactions are temporally, spatially, and specifically controlled under such complex molecular crowding conditions. Thus, it is necessary to study biomolecules under cell-mimicking conditions to unveil their characteristics inside of cells.

In many cases, molecular crowding conditions induced by the addition of neutral cosolute molecules increase the thermodynamic stability and folding kinetics of G4s¹⁹⁸⁻²⁰⁵. Stabilization effects of molecular crowding have been reported for other non-canonical structures such as triplexes, junctions, dangling ends, and loops^{156, 206, 207}. In contrast to the stabilization effects on non-canonical structures, molecular crowding destabilizes the canonical duplex^{198, 208}. These opposite effects on non-canonical and canonical structures indicate that G4s and other non-canonical structures are thermodynamically more stable under molecularly crowded cellular conditions than has been considered based on test-tube experiments. Thus, G4s as well as other non-canonical structures of nucleic acids are promising candidates as bidding sites for proteins and small molecules. These molecular crowding effects are attributed to the osmotic pressure that stabilizes the less hydrated G4 and other non-canonical structures^{200, 201, 209-212}. In contrast, positive hydration changes through duplex formation led to destabilization under molecular crowding conditions, where the concentration and activity of water molecules decrease^{213, 214}.




Fig. 9 (a) Schematic structure of the thrombin binding aptamer (TBA) (left). Melting temperature (T_m) of the TBA in the presence of various cations (right). Adapted from Ref. 186. Copyright © 2001 American Chemical Society. (b) Schematic structure of $d(\text{G}_3\text{T}_4\text{G}_3)$ (left) and the relationship between the dehydration energy of the cations and the thermodynamic stability of G4 (right). Adapted from Ref. 192. Copyright © 1996 American Chemical Society.




Fig. 10 G4 recognition modes of (a) RHAU, (b) RAP1, and (c) ankyrin derivative. Adapted from Ref. 250. Copyright © 2024 American Chemical Society.



Table 2 Cellular functions of G4-binding proteins.

View Article Online
DOI: 10.1039/D4CB00287C

Protein	Function / Target	Ref
Nucleolin	Regulation of transcript / <i>c-MYC</i>	235
DNA methyltransferase	Methylation of Cytosine at C-5 / <i>c-MYC</i> , <i>CDKN1C</i> and <i>MEST</i>	240
Nucleophosmin	Nucleolar phosphoprotein / <i>c-MYC</i>	242
ADAR	Convert adenosine to inosine / <i>c-MYC</i>	243
PARP1	Gene regulation and DNA repair / <i>BCL-2</i>	256
hnRNP1	Gene regulation, DNA repair, and Chromatin remodelling / <i>KRAS</i>	257
MAZ	DNA repair and Chromatin remodelling / <i>KRAS</i>	285, 259
FANCI	5' → 3' Helicase / Telomeric DNA	255
Pif1	5' → 3' Helicase / <i>c-MYC</i>	241
SMARCA4	Chromatin remodelling / <i>c-KIT</i>	230
UHRF1	Ubiquitin ligase / <i>c-MYC</i> and <i>c-KIT</i>	230
RBM22	pre-mRNA-splicing factor / <i>c-MYC</i> and <i>c-KIT</i>	230
TTF2	Transcription termination factor / <i>c-MYC</i> and <i>c-KIT</i>	230
DDX24	Helicase / <i>c-KIT</i>	230
DDX1	Helicase / <i>c-MYC</i> and <i>c-KIT</i>	230
CIRBP	cold-inducible RNA-binding protein / <i>c-MYC</i>	228
DHX36	Helicase / <i>c-MYC</i>	272
DDX21	Helicase / <i>C9orf72</i>	270
DHX9	3' → 5' Helicase / <i>TP-G4</i>	271
FUS	DNA repair and regulation of splicing / Telomeric DNA	260
TDP-43	Regulation of transcription and splicing / Expanded <i>C9orf72</i> gene	261
α -synuclein	DNA repair and rescuing neurons / Telomeric DNA	262

In addition, structural polymorphisms of G4 are induced and regulated by molecular crowding. For example, molecular crowding induces a structural transition from a duplex to a G4 in a mixture of telomere G-rich and complementary C-rich strands²¹⁵. Molecular crowding further alters the strand orientation of G4s from antiparallel and mixed to parallel^{202, 216-219}, which regulates the activity of enzymes such as telomerase and affects protein and ligand binding to G4s²²⁰⁻²²².

G4 binding proteins: Many G4 binding proteins have been identified by affinity proteomics and computational analysis (Table 2)²²³⁻²³⁰. A database of G4 binding proteins is open to the public (<http://bsbe.iiti.ac.in/bsbe/ipdb/>)^{229, 231}. Here, we briefly introduce proteins that bind to DNA and RNA G4s.

DNA G4 binding proteins: The best studied G4 binding proteins are those that bind to telomere DNA and maintain the telomere length by regulating telomerase activity²³². Shelterin, which is important for homeostasis of telomeric length, also binds to G4 in the *c-MYC* promoter²³³. Because of the importance of *c-MYC* as a transcription factor and thus as a promising target for anticancer drugs²³⁴, proteins targeting *c-MYC* G4s have been focused on. Nucleolin binds to the *c-MYC* promoter G4, promotes the formation of the G4, and inhibits *c-MYC* promoter activity²³⁵. Contrary to these studies, promoter G4s are often found in active genes^{236, 237}. Recently, it was reported that transcription factors can selectively bind G4s, and that G4s and the i-motif, which is another four-stranded non-canonical structure of the complementary cytosine-rich strand as will be discussed later, recruit transcription factors, leading to active transcriptional output²³⁸. Although further studies are required to confirm whether there is a universal up-regulatory or down-regulatory role for G4s in transcription²³⁹, it is now

evident that G4s largely affect transcription. DNA methyltransferase enzymes, which catalyze cytosine methylation at specific locations of the genome, show high affinity with G4s derived from *c-MYC* as well as *CDKN1C* and *MEST*²⁴⁰. Other *c-MYC* G4-binding proteins include Pif1 helicase²⁴¹; nucleophosmin, which is a nucleocytoplasmic shuttling protein²⁴²; ADAR (double-stranded RNA-specific adenosine deaminase), which has a Z-DNA-binding domain^{243, 244}; and DNA polymerase η ²⁴⁵. G4-forming promoter regions of other genes, such as *BCL-2*, *KRAS*, *MYB*, *KIT*, and *VEGF*, are also recognized by proteins including PARP1, hnRNP A1, and MAZ, many of which play a role in transcriptional regulation, chromatin remodeling, and DNA repair²⁴⁶.

Although many G4-binding proteins have been identified, there are still a very limited number of three-dimensional structures of G4-protein complexes. Three proteins, RHAU^{247, 248}, RAP1²⁴⁹, and a designed ankyrin protein utilize an α -helix to recognize a G-quartet (Fig. 10)²⁵⁰. A single α -helix of RHAU and RAP1 stacks on an exposed G-quartet. In contrast, ankyrin derivative uses a bundle of helices and loops to bind an exposed G-quartet. These binding modes lead to the specific recognition of the parallel G4. In these complexes, amino acids such as Arg, Ala, Tyr, Leu, and Met of the proteins form π - π and CH- π interactions with the bases and the sugar ring of the parallel G4. In addition, electrostatic and hydrogen bonding interactions between DNA and polar/basic residues contribute to stabilize the interactions.

Because G4s in the genome inhibit replication and transcription, DNA G4s must be unfolded to proceed with these reactions. Thus, helicases are another category of G4-binding protein²⁵¹⁻²⁵³. A combination of G4 helicases and DNA



replication proteins have therapeutic potential against cancer, because G4 helicases are upregulated in cancer cells²⁵⁴. In fact, cancer cells deficient in a G4 helicase, FANCD1, have higher sensitivity to the G4 ligand telomestatin (Fig. 12)²⁵⁵. A series of 5' to 3' and 3' to 5' helicases proposed to bind G4s is listed in Table 2²⁵⁶⁻²⁶². Recently, Balasubramanian's group reported a co-binding-mediated protein profiling strategy with their own developed G4 ligand, pyridostatin (PDS, Fig. 12), to investigate the interactome of DNA G4s in native chromatin²³⁰. The authors employed this approach in human cells and identified hundreds of putative G4-interacting proteins that comprised diverse functional classes, such as RNA binding, DNA binding, ATP binding, metal ion binding, ribonucleoprotein, hydrolase, repressor, helicase, activator, and chromatin regulator. Among them, SMARCA4, UHRF1, RBM22, TTF2, DDX24, and DDX1 were confirmed to bind to G4. Interestingly, the most frequently observed function of these proteins was RNA binding, even though the target was DNA G4 in these studies. However, this is consistent with the cold-inducible RNA-binding protein (CIRBP) being identified as a new G4 DNA-binding protein both *in vitro* and in cells²²⁸. These results indicate that G4s participate in more diverse biological processes in cells than previously considered via interactions with various proteins.

RNA G4-binding proteins: G4s in mRNAs play important roles in translation²⁶³. For example, a G-rich sequence found in the 5'-untranslated region (UTR) of *NRAS* mRNA folds into a stable parallel intramolecular G4 and inhibits translation²⁶⁴. RNA G4s in the cytoplasm are detected by using a G4 structure-specific antibody^{265, 266}. Moreover, over 13,000 loci where G4s form within the transcriptome in humans have been identified²⁶⁷. RNA G4s are enriched in functionally important regions, including 5'- and 3'-UTRs²⁶⁷⁻²⁶⁹ and introns, although some G4s are observed in exons.

As shown for DNA G4 helicases, several helicases, including DHX36, DDX21, and DHX9, bind to and unwind RNA G4s²⁷⁰⁻²⁷², because RNAs must be at least partly unfolded for their messenger function. Moreover, RNA G4-binding proteins observed in RNA modification reactions, viral pathogenesis, and mitochondria have been reviewed²⁷³. Other classes of RNA G4-binding proteins are ribosomal proteins, the AFF family, and hnRNPs²⁴⁶. Ribosomal RNAs (rRNAs) have many G4-forming sequences with three and two G-quartets^{274, 275}, which exist as rRNA tentacles extending for hundreds of Å from ribosomal surfaces²⁷⁶. Thus, it is reasonable that ribosomal proteins bind to these RNA G4s. The AFF family of genes includes four members: AFF1/AF4, AFF2/FMR2, AFF3/LAF4, and AFF4/AF5q31. hnRNP proteins play critical roles in the packaging, transport, and splicing of mRNAs, and some hnRNP proteins, such as A1 and A2, have been reported to bind to RNA G4s^{277, 278}.

G4 in liquid-liquid phase separation

Interestingly, many of the RNA G4 binding proteins discussed above are involved in liquid-liquid phase separation (LLPS) to form membrane-less organelles (biomolecular condensates and liquid droplets) in cells, which play critical roles in cellular

processes and diseases. LLPS is a hot topic in many research fields from cell biology to soft matter physics^{279, 280}. It is generally considered that LLPS is induced by intrinsically disordered proteins (IDPs) or that include intrinsic disorder regions (IDRs)²⁸¹. Nucleic acids (DNA and RNA) also undergo LLPS with binding to proteins to regulate transcription and translation at various stages coupled with other molecular regulatory mechanisms²⁸². Multimolecular bindings among proteins and nucleic acids via electrostatic, hydrogen bonding, π - π stacking, cation- π , and CH- π interactions are essentials for LLPS²⁸³. Notably, it has been proposed that DNA and RNA G4s are critical for LLPS²⁸⁴⁻²⁸⁷, despite the fact that IDP and IDR are essential for LLPS. Figure 11 shows the LLPS model system with a G4-forming RNA oligonucleotide, derived from repeat expansion sequence in FMR1 mRNA, and Arg- and Gly-rich model peptide, derived from RGG motif of FMRP. It was demonstrated that the RNA-peptide mixture undergoes LLPS, depending on the formation of RNA G4. Moreover, aberrant LLPS may trigger protein aggregation in neurodegenerative diseases such as frontotemporal dementia associated with FUS, amyotrophic lateral sclerosis associated with TDP-43, Parkinson's disease associated with α -synuclein, and Alzheimer's disease associated with tau²⁸⁸. In some of these neurodegenerative diseases, the expansion of G-rich repeat sequences, which may form G4s at the genomic and mRNA levels, participate in the onset mechanism²⁸⁹.




Fig. 11 LLPS model system, consisting of RNA G4 and Arg- and Gly-rich peptide. Adapted from Ref. 289. Copyright © 2022 Royal Society of Chemistry.

G-quadruplex-targeting molecules

The fact that stabilized DNA or RNA G4s inhibit series of enzymes, involving DNA and RNA polymerases, ribosome, and reverse transcriptase, has spurred a huge interest in the structure-based design and synthesis of ligands that bind and stabilize G4s (G4 ligands). As of November 2024, the G4 ligand database (G4LDB) opened²⁹⁰. G4LDB includes over 4800 ligands targeting quadruplexes (G4s and i-motifs of cytosine-rich sequences) entries. Several G4 ligands are in clinical trials, as we discuss later^{291, 292}. The structures of some of these ligands are shown in Fig. 12 and are introduced here. Moreover, most G4 ligands possess cationic functional groups that form electrostatic interactions with anionic phosphates of nucleic acids and that make them water soluble. TMPyP4 is a typical and well-known G4 ligand. TMPyP4 shows high affinity for G4s due to its wide π -plane as well as its positive charges (Fig. 12)²⁹³. TMPyP4 has been shown to inhibit human telomere elongation and gene expression by inhibiting telomerase and RNA



polymerase, respectively²⁹⁴. However, the inhibitory effect on enzymatic activity is not structure selective. In telomerase reverse transcription assays in the presence of DNA duplexes, the inhibition effect on telomerase activity is significantly decreased^{221, 295}. This non-structure-selective binding of G4 ligands may decrease their inhibitory effects on enzymes of interest. At least partly due to their low selectivity, no G4 ligand has advanced beyond phase II trials yet. Quarfloxin (CX-3543, Fig. 12) is the ligand that has reached phase II (NCT00780663)²⁹⁶. CX-3543 binds selectively to G4s formed in rDNA and disrupts rRNA interactions with nucleolin²⁹⁷. Another G4 ligand, QN-302 (Fig. 12), has most recently entered clinical phase I trials (NCT06086522)²⁹¹. We briefly discuss the development of QN-302 later in this manuscript.

Thus, structure-selectivity has also attracted attention in G4 ligand design. Telomestatin, which is isolated from *Streptomyces anulatus* 3533-SV4, selectively binds to and stabilizes telomeric DNA G4s, resulting in highly efficient telomerase inhibition ($IC_{50} = 5$ nM)²⁹⁸. Nagasawa and co-workers reported that L2H2-60TD (Fig. 12), a macrocyclic hexa-oxazole, strongly stabilizes human telomere G4 and shows potent inhibition activity against telomerase²⁹⁹. Recently, a L2H2-60TD derivative has also been used as a G4 imaging tool in living cells³⁰⁰. PDS was designed and synthesized by Balasubramanian in 2008 and has been used in various biological assays^{230, 301-304}. PDS was rationally designed with an expanded planar surface and hydrogen bond sites to efficiently bind to G4 structures. PDS binds and stabilizes telomeric G4 while having no effect on DNA duplexes, indicating high structure-selective binding³⁰¹. Because of this property, a modified PDS analogue has also been used in intracellular G4 imaging applications. For example, Balasubramanian and co-workers reported that SiR-pyPDS, a fluorophore-modified PDS³⁰⁴, exhibits remarkable fluorescence turn-on properties upon binding to G4. PhenDC3, a bisquinolinium family compound, strongly selectively binds to and stabilizes G4s over DNA duplexes^{295, 305, 306}. BRACO-19 is an aromatic molecule containing protonatable sites that was designed based on crystal structure and simulation studies to bind to three G4 grooves. BRACO-19 exhibits 31-fold higher affinity for G4s than duplexes.

Another strategy for obtaining structure-selective G4 ligands is the development of small molecules with anionic charges. Cationic ligands bind nucleic acids with non-structure-selectivity due to the negative charges on their backbones. Therefore, it is useful to develop an anionic ligand that causes electrostatic repulsion with the phosphate groups of nucleic acids. Copper (II) phthalocyanine 3, 4', 4'', 4'''-tetrasulfonic acid (CuAPC, Fig. 12), an anionic ligand, has an IC_{50} of 1.6 μ M against telomerase activity in the presence of excess and competitive DNA duplexes²²¹. Hemin is also a typical negatively charged G4 ligand (Fig. 12)³⁰⁷. Hemin suppresses xlr3b gene expression through its binding to a G4 formed at the CpG island in ATR-X model mice, leading to the recovery of cognitive deficits associated with ATR-X syndrome³⁰⁸. Although many developed G4 ligands aim to inhibit transcription or telomere elongation, ligands involved in other biological processes have been

reported. Tetrandrine binds to G4 selectively and stabilizes it strongly³⁰⁹. Recently, it has been shown that stabilization of RNA G4 formed by an miR-23b/27b/24-1 cluster by using tetrandrine prevents the binding of Drosha to the miR cluster, resulting in the recovery of cardiac function³¹⁰. Thioflavin T (ThT, Fig. 12) is also a selective fluorescent probe for G4, although it




Fig. 12 Chemical structures of G4 ligands: TMPyP4, CX-3543, QN-302, telomestatin, L2H2-60TD, pyridostatin, PhenDC3, telomestatin, BRACO-19, CuAPC, Hemin, Thioflavin T, N-methyl mesoporphyrin IX, ZnAPC, alkylated porphyrin.



has a positive charge³¹¹. N-methyl mesoporphyrin IX (NMM, Fig. 12) is another G4-selective ligand. Bolton *et al.* found that NMM binds to DNA G4 but not to duplexes by fluorescence spectroscopy³¹². NMM enhances fluorescence intensity upon binding to G4s, which is useful for their detection. Later, it was reported that NMM does not bind to other nucleic acid conformations³¹³ and is selective for the parallel topology. The K_d value of NMM for the parallel topology is nearly 10 times less than for the mixed topology and 100 times less than for the antiparallel topology³¹⁴.

Breaking down target RNA G4s is another therapeutic drug strategy. Our group reported that ZnAPC (Fig. 12) selectively binds to RNA G4 formed by the *NRAS* mRNA 5'-UTR. ZnAPC breaks down the target RNA G4 via reactive oxygen species upon photoirradiation, resulting in *NRAS* suppression³¹⁵. Xodo and co-workers also reported that an alkylated porphyrin (Fig. 12) bind to *KRAS* mRNA G4 and cleaves it upon photoirradiation³¹⁶. These results suggest that photodynamic therapy targeting mRNA G4s is promising as a new modality for cancer treatment. A remarkable recent progress in this field is

ribonuclease targeting chimera (RIBOTAC), which can recruit the endogenous ribonuclease RNase L to a RNA target, resulting in the target RNA cleavage in living cells³¹⁷. This strategy has been applied to G4-binding proteins and RNA G4s³¹⁸⁻³²⁰. However, selectivity of G4 ligand is the main issue to apply this technology for a practical and clinical use³²¹.

Systematic comparisons of the three-dimensional structures of G4 ligand/G4 complexes are important for improving the affinity and selectivity of G4 ligands. Yang and colleagues reported the solution structures of c-MYC G4 with phenyl-ethenyl-quinoline (PEQ)³²², a specific c-MYC G4 binder, and of three other G4 ligands³²³. This c-MYC G4 sequence contains G23-to-T and G14-to-T mutations in the 3'-flanking and second loop, respectively. This sequence was used as a model parallel G4 structure. Yang and colleagues found that the flexible flanking regions are recruited in a conserved and sequence-specific way, and that these regions have further potential for selective ligand-G4 hydrogen-bond interactions. For example, a combination of π - π stacking interactions with a G-quartet core and hydrogen bonds with other regions such as the flanking and




Fig. 13 (a) Chemical structures of phenyl-ethenyl-quinoline (PEQ), Quindoline-I (Qi), DC-34, and BMVC. (b) Comparisons of the high-resolution structure of c-MYC G4 with PEQ (PDB ID: 7KCX), Qi (PDB ID: 2L7V), DC-34 (PDB ID: 5W77), and BMVC (PDB ID: 6O2L) binding to the 5'-site and the 3'-site. (c) Three-dimensional structures of the PEQ, Qi, DC-34, and BMVC complexes with c-MYC G4 (PDB ID: 7KBX). Adapted from Ref. 322. Copyright © 2021 Oxford University Press.



loop regions have been observed in other G4 ligands, including PDS and its derivatives³²⁴. Other systematic and detailed examples of G4 ligand/G4 complexes have been reported in the literature^{325–327}, involving contributions of other binding modes, such as cation- π and hydrophobic interactions. A quantitative study of nonselective interactions of G4 ligands with long DNA duplexes using a single molecule stretching technique was recently published³²⁸. Those results showed that five of eight G4 ligands commonly used in diverse studies bind the DNA duplex with a binding affinity similar to that of ethidium bromide, a typical DNA duplex binder. This structure-selectivity of G4 ligands therefore holds promise for practical applications in living cells, since there are many more DNA and RNA bases involved in duplex structures than in G4 structures.

i-Motif

Structure, sequence, and biological role of the i-motif

The i-motif, another tetraplex, is formed by cytosine-rich (C-rich) sequences via cytosine·cytosine⁺ base pairs (Fig. 14a). The first DNA i-motif was reported in 1993 for d(TCCCC) (Fig. 14b, c)³²⁹. Since protonation of cytosine is required, this structure is particularly stable in acidic conditions. Although the pK_a of free cytidine is 4.2, the pK_a is shifted in RNA and DNA^{330, 331}. The value is further shifted to be around 6.5 in the i-motif³³². Moreover, the pK_a value increases as the number of cytosines per tract increases from two to seven³³³. Loop regions connecting cytosine stretches and capping nucleotides next to the stretches also affect the thermodynamics of i-motifs^{334–336}.

The molecular environment also affects the structural stability of i-motifs. In contrast to G4, cation species do not affect i-motifs, but the cation concentration does. It has been reported that a higher cation concentration destabilizes i-motifs^{337, 338}. A lower stability with higher cation concentrations is observed only for the i-motif, but not for other structures of DNA. In contrast to the opposing effects of the cation concentration, molecular crowding stabilizes i-motifs, as with G4s. It has been found that C-rich sequences adopt the i-motif structure at neutral pH under molecular crowding conditions^{339, 340}. Given the stabilization effects of molecular crowding on G4s, an intramolecular structure of an i-motif with a C-rich sequence + a G4 of a complementary G-rich sequence could be more stable than an intermolecular duplex of G- and C-rich sequences, which is generally thermodynamically more stable than the sum of the intramolecular structures (G4 and i-motif) in a physiological buffer³⁴¹. In fact, it has been reported that molecular crowding induces a structural transition of telomeric G-rich and C-rich sequences from an intramolecular G4 and i-motif to an intermolecular duplex²¹⁵. Another factor that affects thermal stability of i-motifs is negative superhelicity, which unwinds the duplex³⁴². Various biological processes such as transcription, replication, recombination, and DNA damage repair produce negative supercoiling³⁴³. Thus, during these processes, the formation of intramolecular structures, including i-motifs, could be accelerated. The C-rich sequence of the human telomere can form an i-motif in the S-phase of

replicating human cells³⁴⁴. In addition, the stabilization of a G4 of a G-rich sequence further aids the complementary C-rich sequence to form an i-motif by reducing structural competition with the intermolecular duplex. Thus, it is possible that the i-motif exists in living cells. In fact, evidence for the existence of i-motifs in living cells has been provided by in-cell NMR and an i-motif antibody^{345, 346}.

As a complementary sequence to putative quadruplex-forming sequences, putative i-motif-forming sequences (PIS) have been identified in various genes. The PIS in the c-MYC gene folds into several different i-motif structures with the stability being remarkable at pH 7³⁴⁷. There are more than 60 structures of i-motifs, with sequences including methylated and other modified cytosines, that are registered in the protein data bank (September 2021). As there are many PIS, there are also proteins that bind to i-motifs and PIS oligonucleotides. Poly-C binding protein is a ubiquitous oligonucleotide-binding protein in eukaryotic cells that regulates gene expression³⁴⁸. This family consists of the hnRNP K (heterogeneous nuclear ribonucleoprotein K) and isoforms of α CP1, including α CP1-4 and α CP-KL. hnRNP K is known as a transcription factor that binds to CT elements in the promoter region of c-MYC³⁴⁹. α CP1 also recognizes the C-rich sequence of human telomere DNA with high affinity³⁵⁰. Lacroix and coworkers studied the interaction of hnRNP K and ASF/SF2 with telomeric C-rich sequences under various pH conditions. Their results suggested that the protein binds to the unfolded sequence³⁵¹. A later study indicated that hnRNA interacts with single-stranded DNAs³⁵². In addition, hnRNP K and hnRNP LL, which are transcriptional factors, have been reported to bind to C-rich sequences of VEGF and bcl-2, respectively^{353, 354}. As the protein unfolds the i-motif structure and forms a stable single-stranded DNA-protein complex, the i-motif may play an important role in controlling gene expression. Moreover, i-motifs have been proposed to have various biological functions through their structure and




Fig. 14 Characteristics of the i-motif structure. (a) Chemical structure of a C:C⁺ base pair. (b) Illustration of the i-motif structure folded by a C-rich sequence, 5'-TCCCC-3'. (c) Crystal structure of the tetrameric i-motif (PDB ID: 2N89).






Fig. 15 Chemical structures of i-motif-targeting molecules phenanthroline, bisacridine, L2H2-4OTD, $[\text{Ru}(\text{bpy})_2(\text{dppz})]^{2+}$, c3, B19, a9, IMC-48, PBP1, mitoxantrone, tilorone, and tobramycin.

protein binding. These biological roles include the inhibition of telomerase activity, a structural element that provides long-range interactions between laterally associated centromeric nucleosomes, transcriptional regulation, and stalling DNA polymerase and thus impeding DNA replication and repair. These biological roles of i-motifs have received thorough recent reviews³⁵⁵⁻³⁵⁷.

i-Motif targeting molecules

The first discovered i-motif ligand was TMPyP4, which was also known as a G4 ligand^{358, 359}. Since it is highly cationic, TMPyP4 binds to duplexes in a non-specific manner, and the binding to G4 was overestimated²⁹⁵. This original study stimulated many researchers to develop i-motif ligands.

As with the G4 ligands discussed above, many of i-motif ligands have a planar structure (Fig. 15). Such ligands include derivatives of phenanthroline³⁶⁰, acridine³⁶¹, macrocyclic poly-oxazole³⁶², and metal complexes^{363, 364}. NMR studies have demonstrated that i-ligands such as L2H2-4OTD and IMC-48 form complexes through stacking interactions with C-C+ base pairs and interactions with loops^{362, 365}. Although the detailed binding modes (in addition to the stacking interactions) were not identified, these findings nonetheless suggest that interactions with the C-C+ base pairs and the loops are critical for stabilizing the i-motif. Phenanthroline derivatives show preferential binding to telomeric i-motifs over duplexes and G4. Acridine derivatives are based on BRACO-19, which is a well-known G4 ligand. This compound has been demonstrated to suppress the proliferation of cancer cells³⁶⁶. One poly-oxazole derivative has been reported to interact with a telomeric i-motif, but not G4³⁶². Interestingly, carboxyl-modified single-walled carbon nanotubes (SWNTs) induce and stabilize human telomeric i-motifs. K562 and HeLa cells were treated with

SWNTs, then the inhibition of telomere elongation was confirmed using a telomeric repeat amplification protocol assay. Colocalization of SWNTs and a telomere restriction fragment, a telomere marker, suggested that SWNTs were located in the telomere region and stabilized telomeric i-motif DNA, causing folding of telomeric G4 and inhibiting telomere elongation^{367, 368}.

It has been shown that small molecules targeting i-motifs regulate the transcription of many genes. B19 and a9 bind to the c-MYC i-motif and downregulate transcription^{369, 370}. Oleanolic acid derivatives interact with the VEGF i-motif and also downregulate transcription^{371, 372}. In contrast, IMC48 and PBP1 bind to the BCL2 i-motif and upregulate transcription^{370, 373}. Although these results suggest that i-motifs are important as *cis*-acting gene regulators, it is still unclear how genes are down- and up-regulated by i-motif formation and ligand binding. The fluorescent intercalator displacement assay is a powerful method to screen ligands that target i-motifs as well as other structures of nucleic acids. Waller and co-workers have discovered i-motif ligands, such as the antitumor agent mitoxantrone and antiviral agents tilorone and tobramycin, by fluorescent intercalator displacement assays utilizing thiazole orange^{374, 375}. However, thiazole orange emits fluorescence not only with i-motifs but also other structures, including G4s and duplexes. Thus, an i-motif fluorescent probe will be useful to develop a more simplified and high-throughput screening system for i-motif-specific ligands.

Targeting three- and four-stranded nucleic acids molecules toward clinical trials

Nucleic acid-targeting small molecule as approved drug.

It is essential to consider clinical applications even at the earliest stages of drug development. For example, risdiplam (Evrysdi®)



is an approved small molecule therapeutic that targets nucleic acids to treat spinal muscular atrophy, and it is the first small molecule to be identified as a splicing modifier³⁷⁶. Risdiplam was identified by phenotypic screening rather than by structure-based screening³⁷⁷. Screening was performed using a human embryonic kidney cell line harboring an SMN2 minigene to increase the inclusion of exon 7 during SMN2 pre-mRNA splicing. Various functional groups were introduced to enhance oral availability³⁷⁸. For example, the insertion of a pyridopyrimidinone moiety reduced the phototoxicity of the compound and a cyclic N-alkyl group prevented nonspecific protein binding. These design steps and modifications allow risdiplam to strongly promote SMN protein expression with low cytotoxicity³⁷⁹⁻³⁸¹. The biological activity of the compounds was confirmed using animal models of SMA. Pharmacokinetic studies showed that the compounds penetrated the brains of adult mice. Finally, the efficiency and safety of risdiplam was confirmed in a clinical development program with SMA patients.

G4 ligands for clinical applications.

Similar problems need to be solved by chemical modifications for small molecules targeting non-canonical structures of nucleic acids. After screening or structure-based rational design the lead compounds must be optimized, as shown by the example of G4 ligands being developed for clinical use. BRACO-19 (Fig. 12), developed by Neidle and coworkers, has been modified, with one derivative exhibiting high selectivity against duplexes and strong inhibition of telomerase activity³⁶⁶. This compound persistently arrested the growth of a breast carcinoma cell line³⁶⁶ and is the first G4 ligand showing an anticancer effect and proven anticancer activity in human tumor xenograft models. Neidle and colleagues also developed and optimized compounds with a naphthalene diimide (NDI) core using structure-based design, medicinal chemistry and pharmacology. QN-302, an NDI derivative with a benzyl-pyrrolidine group, significantly inhibits the growth of pancreatic ductal adenocarcinoma cells in animal models, with GI₅₀ values of 1~2 nM and superior potency compared to derivatives with a methoxy group replacing the benzyl-pyrrolidine³⁸². QN-302 binds tightly to various parallel G4s and is highly specific for G4 compared to duplexes³⁸². QN-302 began clinical trials in 2023^{291, 378}.

The detailed binding mode of QN-302 with G4 has been studied³⁸³⁻³⁸⁵. Molecular modelling studies suggest that QN-302 binds at the duplex-G4 junction through the NDI core via stacking interactions with the G4 terminal G-quartet. The benzyl-pyrrolidine substituent of QN-302 protrudes significantly into the groove. The phenyl ring stacks efficiently onto the adjacent guanine of the lower G-quartet. The benzyl-pyrrolidine group allows enhanced G4 binding affinity and selectivity, enhanced cellular and *in vivo* potency, and greater selectivity in the pattern of downregulated gene expression. Furthermore, the pharmacokinetic properties of QN-302, such as half-life *in vivo* and bioavailability, are not apparently responsible for its superior activity *in vivo*. These results further demonstrate the

importance of the binding affinity and selectivity of G4 ligands for clinical applications.

DOI: 10.1039/D4CB00287C

G4 ligands with higher selectivity.

As the above examples show, the selectivity of the compound for its target is of utmost importance. High selectivity is critical for improving drug efficacy and reducing side effects, as it allows the highly efficient inhibition of G4-targeted enzymes and prevents off-target enzyme inhibition. High selectivity against the duplex is also important to reduce the genotoxicity of G4 ligands and small compounds targeting other three- and four-stranded nucleic acid structures. High selectivity for G4 ligands requires the involvement of directional interactions in addition to stacking interactions. The energy of electrostatic interactions is highest within non-covalent interactions, but it depends only on the distance between the charges, making it difficult to confer selectivity. This can be particularly detrimental to discriminating between G4 and canonical duplex structures. The energy of hydrogen bonding varies with direction, so many G4 ligands have been modified to introduce hydrogen-bonding donors and acceptors to allow hydrogen bonds with the target G4, especially with the loops, grooves, and flanking regions (Fig. 7). The importance of these three regions for specificity is discussed in this subsection.

Loops: There are 21844 possible sequences with three loops in intramolecular G4s with 1~7 nucleotides of which 20492 are found at least once in the human genome¹³⁹. The loop region was recently reported to significantly impact the topology and thermal stability of G4, and the kinetics of G4 formation^{123, 386, 387}. This suggests that the binding of G4 ligands to the loop regions can significantly alter the overall G4 structure and thus modulate protein binding and enzymatic activity. In addition, a G4 structure can adapt other nucleic acid structural motifs, such as a hairpin loop. A duplex can be attached to a G4^{167, 168}. The duplex formed within the loops stabilizes the overall G4 structure compared to an unstructured loop and accelerates G4 folding kinetics^{169, 175-177}.

Grooves: The G4 structure has four grooves: narrow, medium, and wide as defined by the glycosidic bond conformation adopted by any two base-paired 2'-deoxyguanosines of stacked G-quartets³⁸⁸. There are eight combinations of groove types³⁸⁹. The depth and width of these grooves differ significantly from the major and minor grooves of the duplex structure. Moreover, the Hoogsteen base pairs forming the G-quartet use different hydrogen bond acceptors and donors than the Watson-Crick base pairs in the duplex, so the hydrogen bond acceptors and donors exposed in the grooves are also very different in the G4 and in the duplex structures. The hydration pattern in the G4 grooves is also different from in the duplex grooves³⁸⁹. These characteristic grooves are likely important for selective binding to the G4 over the duplex structure.

Flanking: The 5' and 3' flanking regions adjacent to the G4 core region are also targets for hydrogen bonding interactions. The stacking and hydrogen bonding interactions formed between the flanking nucleotide and another flanking nucleotide, the adjacent G-quartet, and the neighboring loop,



all contribute to the overall structure and thermodynamics of G4. The flanking nucleotides reportedly influence the binding affinity of the G4 ligand. As discussed above, the importance of detailed structural information on the loop, groove and flanking regions will increase as the design of G4 ligands progresses.

The three-dimensional structure of the loop regions, and the depth and width of the grooves, depend on the nucleotide sequence. The nucleotide sequences for the 5' and 3'-flanking regions determine the target. These differences are important not only for structural selectivity against the canonical duplex, but also for sequence selectivity towards the many G4s present in cells, although only a few examples of G4 ligands have been reported to show sequence selectivity. Selectivity for DNA G4 and RNA G4 may also be important for precise targeting. More three-dimensional G4 structures are needed to predict the overall topology of G4 from the sequence and also the fine structure of the loops, grooves, and flanking regions.

Small molecules that highly selectively target triplex and i-motifs have been reported, as shown for the triplex of MALAT1 discussed above; however, only a few studies show the effects of these molecules on cellular and organoid functions^{120, 390}. As mentioned above, even small molecules with high selectivity require toxicity testing, improved oral dosing, animal models, and confirmation of compound efficacy and pharmacokinetics. More systematic and quantitative studies of the relationship between structure, selectivity, and function *in vitro* and *in vivo* are required.

Conclusion and perspective

DNA and RNA sequences encode not only genetic information but also higher order information, such as three-dimensional structures and stabilizing factors, and they respond to environmental factors, all of which are critical for gene regulation. Moreover, it has been found that DNA and RNA undergo LLPS in living cells to regulate the central dogma at many steps involving the replication of genomic DNA and viral RNA; the transcription of mRNA, rRNA, and non-coding RNA; the processing and localization of RNAs; and translation from RNA to protein³⁹¹⁻³⁹⁵. One fundamental property of LLPS is responsiveness to cellular stresses, such as high temperatures, nutrient starvation, toxins, and reactive oxygen species. The structure and stability of nucleic acids depend on the molecular environment as we described in this review. Interestingly, recent studies suggest that LLPS of nucleic acids is dependent of their secondary structures^{284, 285}. Thus, DNA and RNA sequences may further encode phase behavior. Despite our knowledge of the biology, chemistry, and physics of non-canonical nucleic acid structures, many aspects are still opening for further investigation. In this regard, researches on the non-canonical structures of DNA and RNA utilizing small ligands *in vitro* and *in vivo* are essential for elucidating their roles in different biological processes. In spite of recent findings of ligands that target non-canonical structures, the number of structure-specific ligands is very limited. Moreover, non-canonical structural ligands that can target a specific sequence are further

needed. To develop structure and sequence-selective ligands, at least the following are required.

(1) Three-dimensional structures of nucleic acid-ligand complexes: Structural information is essential for the rational design of ligands. Even though biochemical and biophysical information for G4 ligand complexes is relatively abundant, there are only a limited number of RNA and DNA G4 ligand complexes with structures that have been solved. Moreover, the structure of an i-motif and ligand complex has not been reported yet.

(2) Dynamic properties of non-canonical structures: Non-canonical structures are not static. Their formation and thermal stability largely depend on molecular environmental factors and protein binding. In the case of DNA, non-canonical structures always compete against the canonical duplex. In the case of RNA, structural switches among the most stable and metastable structures may be important for their biological roles as in the case of riboswitches. Thus, the dynamic and kinetic properties of non-canonical structures should be studied under cell-mimicking conditions and in living cells.

(3) High-throughput screening systems: Since not only high affinity but also high selectivity of ligands are required for cellular applications, screening systems that can evaluate biorthogonality among canonical and non-canonical structures are needed. For example, many i-motif and G4 ligands also bind G4s and i-motifs, respectively. Moreover, it is difficult to eliminate non-specific binding of ligands to the duplex. In addition to structure-specific ligands, sequence-specific ligands are required, because as shown for triplexes and quadruplexes, there are many putative sequences that form certain non-canonical structures.

A wide variety of G4 ligand screening systems, such as fluorescent intercalator displacement (FID), are used to identify ligands³⁹⁶⁻³⁹⁸, but these methods are low-throughput. Obtaining structure- or sequence-selective ligands using such methods requires multiple steps. Our group established new G4 ligand screening systems that are optimized to provide structure- or sequence-selective G4 ligands^{399, 400}, although a molecular probe exhibiting target selectivity is required. Furthermore, cell-based FID assays can provide ligands that bind intracellular G4s efficiently⁴⁰¹. These reports demonstrate that G4 ligands exhibiting structure selectivity can efficiently target G4s within cells. Further structural information on target-selective ligands can enrich target-selective ligand databases such as G4 LDB²⁹⁰ and NALDB⁴⁰². A systematic quantitative structure-activity relationship (QSAR) workflow in 2022 was reported to broadly discover RNA ligands using HIV-1 transactivation response RNA as a model system⁴⁰³. This method can evaluate binding parameters, including affinity and binding kinetics, bypassing the need for structural information. Therefore, the discovery of target-selective ligands not only aids understanding of the scaffolds, but also the rational design of ligands for target-selective binding.

(4) Widely useable databases: Bioinformatic and omics studies have provided a massive amount of data sets to researchers, but it is almost impossible to make effective use of them by examining individual studies. Thus, to unify and utilize



a large amount of data, databases and machine learning systems for wide and open use are desired.

In conclusion, structure- and sequence-selective ligands could be the key to decoding hidden information in nucleic acids and to develop small molecular drugs targeting nucleic acids. Nucleic acids -DNA, RNA, as well as artificial ones- are now coming of age as drug targets and targeting biomolecules.

Data availability

No primary research results, software or code have been included, and no new data were generated or analyzed as part of this review.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by JSPS KAKENHI, grant numbers 24K21801, 21H02062, and 20K21259, a Research Grant of the Asahi Glass Foundation, Japan, the Hirao Taro Foundation and the Tateno Junzo Foundation of Konan Gakuen for Academic Research, Japan, and JST SPRING, grant number JPMJSP2117.

Notes and references

1. A. L. Hopkins and C. R. Groom, *Nat. Rev. Drug Discov.*, 2002, **1**, 727-730.
2. J. P. Overington, B. Al-Lazikani and A. L. Hopkins, *Nat. Rev. Drug Discov.*, 2006, **5**, 993-996.
3. S. J. Dixon and B. R. Stockwell, *Curr. Opin. Chem. Biol.*, 2009, **13**, 549-555.
4. A. D. Cox, S. W. Fesik, A. C. Kimmelman, J. Luo and C. J. Der, *Nat. Rev. Drug Discov.*, 2014, **13**, 828-851.
5. C. V. Dang, *Cell*, 2012, **149**, 22-35.
6. L. Mabonga and A. P. Kappo, *Biophys. Rev.*, 2019, **11**, 559-581.
7. A. Sawyer, *Biotechniques*, 2020, **69**, 239-241.
8. R. Kahan, D. J. Worm, G. V. Castro, S. Ng and A. Barnard, *RSC Chem. Biol.*, 2021, **2**, 387-409.
9. K. S. McPherson and D. M. Korzhnev, *RSC Chem. Biol.*, 2021, **2**, 1167-1195.
10. M. Clamp, B. Fry, M. Kamal, X. Xie, J. Cuff, M. F. Lin, M. Kellis, K. Lindblad-Toh and E. S. Lander, *Proc. Natl. Acad. Sci. U.S.A.*, 2007, **104**, 19428-19433.
11. I. Ezkurdia, D. Juan, J. M. Rodriguez, A. Frankish, M. Diekhans, J. Harrow, J. Vazquez, A. Valencia and M. L. Tress, *Hum. Mol. Genet.*, 2014, **23**, 5866-5878.
12. M. N. Cabili, C. Trapnell, L. Goff, M. Koziol, B. Tazon-Vega, A. Regev and J. L. Rinn, *Genes Dev.*, 2011, **25**, 1915-1927.
13. J. Zell, S. F. Rota, S. Britton and D. Monchaud, *RSC Chem. Biol.*, 2020, **2**, 47-76.
14. G. D. Stormo, *Quant. Biol.*, 2013, **1**, 115-130.
15. W. W. Wasserman and A. Sandelin, *Nat. Rev. Genet.*, 2004, **5**, 276-287.
16. E. L. Van Nostrand, P. Freese, G. A. Pratt, X. Wang, X. Wei, R. Xiao, S. M. Blue, J. Y. Chen, N. A. L. Cody, D. Dominguez, S. Olson, B. Sundararaman, L. Zhan, C. Bazile, L. P. B. Bouvrette, J. Bergalet, M. O. Duff, K. E. Garcia, C. Gelboin-Burkhart, M. Hochman, N. J. Lambert, H. Li, M. P. McGurk, T. B. Nguyen, T. Palden, I. Rabano, S. Sathe, R. Stanton, A. Su, R. Wang, B. A. Yee, B. Zhou, A. L. Louie, S. Aigner, X. D. Fu, E. Lécuycy, C. B. Burge, B. R. Graveley and G. W. Yeo, *Nature*, 2020, **583**, 711-719.
17. H. Tateishi-Karimata and N. Sugimoto, *Chem. Commun.*, 2020, **56**, 2379-2390.
18. H. Tateishi-Karimata and N. Sugimoto, *Nucleic Acids Res.*, 2021, **49**, 7839-7855.
19. J. P. Falese, A. Donlic and A. E. Hargrove, *Chem. Soc. Rev.*, 2021, **50**, 2224-2243.
20. C. Sheridan, *Nat. Biotechnol.*, 2021, **39**, 6-8.
21. C. A. Orengo, D. T. Jones and J. M. Thornton, *Nature*, 1994, **372**, 631-634.
22. S. Govindarajan, R. Recabarren and R. A. Goldstein, *Proteins*, 1999, **35**, 408-414.
23. R. D. Schaeffer and V. Daggett, *Protein Eng. Des. Sel.*, 2011, **24**, 11-19.
24. G. Felsenfeld, D. R. Davies and A. Rich, *J. Am. Chem. Soc.*, 1957, **79**, 2023-2024. DOI: 10.1039/D4CB00267C
25. A. Jain, G. Wang and K. M. Vasquez, *Biochimie*, 2008, **90**, 1117-1130.
26. P. A. Beal and P. B. Dervan, *Science*, 1991, **251**, 1360-1363.
27. A. G. Letai, M. A. Palladino, E. Fromm, V. Rizzo and J. R. Fresco, *Biochemistry*, 1988, **27**, 9108-9112.
28. D. M. Gowers and K. R. Fox, *Nucleic Acids Res.*, 1997, **25**, 3787-3794.
29. D. S. Pilch, C. Levenson and R. H. Shafer, *Biochemistry*, 1991, **30**, 6081-6088.
30. K. R. Fox, *Curr. Med. Chem.*, 2000, **7**, 17-37.
31. F. A. Buske, D. C. Bauer, J. S. Mattick and T. L. Bailey, *Genome Res.*, 2012, **22**, 1372-1381.
32. S. S. Gaddis, Q. Wu, H. D. Thames, J. DiGiovanni, E. F. Walborg, M. C. MacLeod and K. M. Vasquez, *Oligonucleotides*, 2006, **16**, 196-201.
33. Y. M. Agazie, J. S. Lee and G. D. Burkholder, *J. Biol. Chem.*, 1994, **269**, 7019-7023.
34. Y. M. Agazie, G. D. Burkholder and J. S. Lee, *Biochem. J.*, 1996, **316** (Pt 2), 461-466.
35. J. A. Brown, *Wiley Interdiscip. Rev. RNA*, 2020, **11**, e1598.
36. D. Sacks, B. Baxter, B. C. V. Campbell, J. S. Carpenter, C. Cognard, D. Dippel, M. Eesa, U. Fischer, K. Hausegger, J. A. Hirsch, M. Shazam Hussain, O. Jansen, M. V. Jayaraman, A. A. Khalessi, B. W. Kluck, S. Lavine, P. M. Meyers, S. Ramee, D. A. Rüfenacht, C. M. Schirmer and D. Vorwerk, *Int. J. Stroke*, 2018, **13**, 612-632.
37. C. A. Theimer, C. A. Blois and J. Feigon, *Mol. Cell*, 2005, **17**, 671-682.
38. K. Shefer, Y. Brown, V. Gorkovoy, T. Nussbaum, N. B. Ulyanov and Y. Tzfati, *Mol. Cell Biol.*, 2007, **27**, 2130-2143.
39. S. D. Gilbert, R. P. Rambo, D. Van Tyne and R. T. Batey, *Nat. Struct. Mol. Biol.*, 2008, **15**, 177-182.
40. K. T. Tycowski, M. D. Shu, S. Borah, M. Shi and J. A. Steitz, *Cell Rep.*, 2012, **2**, 26-32.
41. C. Cianciolo Cosentino, N. I. Skrypnik, L. L. Brilli, T. Chiba, T. Novitskaya, C. Woods, J. West, V. N. Korotchenko, L. McDermott, B. W. Day, A. J. Davidson, R. C. Harris, M. P. de Caestecker and N. A. Hukriede, *J. Am. Soc. Nephrol.*, 2013, **24**, 943-953.
42. N. K. Conrad, *Virus Res*, 2016, **212**, 53-63.
43. J. N. Hutchinson, A. W. Ensminger, C. M. Clemson, C. R. Lynch, J. B. Lawrence and A. Chess, *BMC Genomics*, 2007, **8**, 39.
44. R. Miyagawa, K. Tano, R. Mizuno, Y. Nakamura, K. Ijiri, R. Rakwal, J. Shibato, Y. Masuo, A. Mayeda, T. Hirose and N. Akimitsu, *Rna*, 2012, **18**, 738-751.
45. T. Gutschner, M. Hämmerle and S. Diederichs, *J. Mol. Med.*, 2013, **91**, 791-801.
46. F. A. Abulwerdi, W. Xu, A. A. Ageeli, M. J. Yonkunas, G. Arun, H. Nam, J. S. Schneekloth, Jr., T. K. Dayie, D. Spector, N. Baird and S. F. J. Le Grice, *ACS Chem. Biol.*, 2019, **14**, 223-235.
47. J. A. Brown, M. L. Valenstein, T. A. Yario, K. T. Tycowski and J. A. Steitz, *Proc. Natl. Acad. Sci. U.S.A.*, 2012, **109**, 19202-19207.
48. J. E. Wilusz, C. K. JnBaptiste, L. Y. Lu, C. D. Kuhn, L. Joshua-Tor and P. A. Sharp, *Genes Dev.*, 2012, **26**, 2392-2407.
49. K. C. Wang and H. Y. Chang, *Mol. Cell*, 2011, **43**, 904-914.
50. T. Li, X. Mo, L. Fu, B. Xiao and J. Guo, *Oncotarget*, 2016, **7**, 8601-8612.
51. P. Grote, L. Wittler, D. Hendrix, F. Koch, S. Währisch, A. Beisaw, K. Macura, G. Bläss, M. Kellis, M. Werber and B. G. Herrmann, *Dev. Cell*, 2013, **24**, 206-214.
52. T. Mondal, S. Subhash, R. Vaid, S. Enroth, S. Uday, B. Reinius, S. Mitra, A. Mohammed, A. R. James, E. Hoberg, A. Moustakas, U. Gyllensten, S. J. Jones, C. M. Gustafsson, A. H. Sims, F. Westerlund, E. Gorab and C. Kanduri, *Nat. Commun.*, 2015, **6**, 7743.
53. A. Postepska-Igielska, A. Giwojna, L. Gasri-Plotnitsky, N. Schmitt, A. Dold, D. Ginsberg and I. Grummt, *Mol. Cell*, 2015, **60**, 626-636.
54. V. B. O'Leary, S. V. Ovsepian, L. G. Carrascosa, F. A. Buske, V. Radulovic, M. Niyazi, S. Moertl, M. Trau, M. J. Atkinson and N. Anastasov, *Cell Rep.*, 2015, **11**, 474-485.
55. V. B. O'Leary, J. Smida, F. A. Buske, L. G. Carrascosa, O. Azimzadeh, D. Maugg, S. Hain, S. Tapio, W. Heidenreich, J. Kerr, M. Trau, S. V. Ovsepian and M. J. Atkinson, *Sci. Rep.*, 2017, **7**, 7163.
56. M. Kalwa, S. Hänzelmann, S. Otto, C. C. Kuo, J. Franzen, S. Jousen, E. Fernandez-Rebollo, B. Rath, C. Koch, A. Hofmann, S. H. Lee, A. E. Teschendorff, B. Denecke, Q. Lin, M. Widschwendter, E. Weinhold, I. G. Costa and W. Wagner, *Nucleic Acids Res.*, 2016, **44**, 10631-10643.
57. C. C. Kuo, S. Hänzelmann, N. Sentürk Cetin, S. Frank, B. Zajzon, J. P. Derks, V. S. Akhade, G. Ahuja, C. Kanduri, I. Grummt, L. Kurian and I. G. Costa, *Nucleic Acids Res.*, 2019, **47**, e32.
58. I. Antonov and Y. A. Medvedeva, *F1000Res*, 2018, **7**, 76.
59. A. Bacolla, G. Wang and K. M. Vasquez, *PLoS Genet.*, 2015, **11**, e1005696.
60. Y. Li, J. Syed and H. Sugiyama, *Cell Chem. Biol.*, 2016, **23**, 1325-1333.
61. E. Käs, E. Izaurralde and U. K. Laemmli, *J. Mol. Biol.*, 1989, **210**, 587-599.
62. E. Winter and A. Varshavsky, *EMBO J.*, 1989, **8**, 1867-1877.
63. Y. Adachi, E. Käs and U. K. Laemmli, *EMBO J.*, 1989, **8**, 3997-4006.



64. R. Kiyama and R. D. Camerini-Otero, *Proc. Natl. Acad. Sci. U.S.A.*, 1991, **88**, 10450-10454.
65. M. Musso, L. D. Nelson and M. W. Van Dyke, *Biochemistry*, 1998, **37**, 3086-3095.
66. A. L. Guiesse, D. Praseuth and C. Hélène, *J. Mol. Biol.*, 1997, **267**, 289-298.
67. E. Jiménez-García, A. Vaquero, M. L. Espinás, R. Soliva, M. Orozco, J. Bernués and F. Azorín, *J. Biol. Chem.*, 1998, **273**, 24640-24648.
68. L. D. Nelson, M. Musso and M. W. Van Dyke, *J. Biol. Chem.*, 2000, **275**, 5573-5581.
69. M. Musso, G. Bianchi-Scarrà and M. W. Van Dyke, *Nucleic Acids Res.*, 2000, **28**, 4090-4096.
70. G. Li, G. V. Tolstonog and P. Traub, *DNA Cell Biol*, 2002, **21**, 163-188.
71. P. Ciotti, M. W. Van Dyke, G. Bianchi-Scarrà and M. Musso, *Eur. J. Biochem.*, 2001, **268**, 225-234.
72. J. Kusic, B. Tomić, A. Divac and S. Kojic, *Mol Biol Rep*, 2010, **37**, 2317-2322.
73. F. Guillonneau, A. L. Guiesse, J. P. Le Caer, J. Rossier and D. Praseuth, *Nucleic Acids Res.*, 2001, **29**, 2427-2436.
74. J. E. Rao and N. L. Craig, *J. Mol. Biol.*, 2001, **307**, 1161-1170.
75. J. E. Rao, P. S. Miller and N. L. Craig, *Proc. Natl. Acad. Sci. U.S.A.*, 2000, **97**, 3936-3941.
76. B. S. Thoma, M. Wakasugi, J. Christensen, M. C. Reddy and K. M. Vasquez, *Nucleic Acids Res.*, 2005, **33**, 2993-3001.
77. S. S. Lange, M. C. Reddy and K. M. Vasquez, *DNA Repair (Amst)*, 2009, **8**, 865-872.
78. L. D. Nelson, C. Bender, H. Mannsperger, D. Buerger, P. Kambakamba, G. Mudduluru, U. Korf, D. Hughes, M. W. Van Dyke and H. Allgayer, *Mol. Cancer*, 2012, **11**, 38.
79. R. M. Brosh, Jr., A. Majumdar, S. Desai, I. D. Hickson, V. A. Bohr and M. M. Seidman, *J. Biol. Chem.*, 2001, **276**, 3024-3030.
80. J. A. Sommers, N. Rawtani, R. Gupta, D. V. Bugreev, A. V. Mazin, S. B. Cantor and R. M. Brosh, Jr., *J. Biol. Chem.*, 2009, **284**, 7505-7517.
81. A. Jain, A. Bacolla, P. Chakraborty, F. Grosse and K. M. Vasquez, *Biochemistry*, 2010, **49**, 6992-6999.
82. V. Kopel, A. Pozner, N. Baran and H. Manor, *Nucleic Acids Res.*, 1996, **24**, 330-335.
83. M. Brázďová, V. Tichý, R. Helma, P. Bažantová, A. Polášková, A. Krejčí, M. Petr, L. Navrátilová, O. Tichá, K. Nejedlý, M. L. Bennink, V. Subramaniam, Z. Bábková, T. Martinek, M. Lexa and M. Adámik, *PLoS One*, 2016, **11**, e0167439.
84. C. Pasquier, S. Agnel and A. Robichon, *G3 (Bethesda)*, 2017, **7**, 2295-2304.
85. G. E. Plum, Y. W. Park, S. F. Singleton, P. B. Dervan and K. J. Breslauer, *Proc. Natl. Acad. Sci. U.S.A.*, 1990, **87**, 9436-9440.
86. J. S. Lee, D. A. Johnson and A. R. Morgan, *Nucleic Acids Res.*, 1979, **6**, 3073-3091.
87. S. Nakano, D. Miyoshi and N. Sugimoto, *Chem. Rev.*, 2014, **114**, 2733-2758.
88. J. S. Sun, J. C. François, T. Montenay-Garestier, T. Saison-Behmoaras, V. Roig, N. T. Thuong and C. Hélène, *Proc. Natl. Acad. Sci. U.S.A.*, 1989, **86**, 9198-9202.
89. L. Perrouault, U. Asseline, C. Rivalle, N. T. Thuong, E. Bisagni, C. Giovannangeli, T. Le Doan and C. Hélène, *Nature*, 1990, **344**, 358-360.
90. M. Takasugi, A. Guendouz, M. Chassignol, J. L. Decout, J. Lhomme, N. T. Thuong and C. Hélène, *Proc. Natl. Acad. Sci. U.S.A.*, 1991, **88**, 5602-5606.
91. P. V. Scaria and R. H. Shafer, *J. Biol. Chem.*, 1991, **266**, 5417-5423.
92. J. L. Mergny, D. Collier, M. Rougée, T. Montenay-Garestier and C. Hélène, *Nucleic Acids Res.*, 1991, **19**, 1521-1526.
93. J. L. Mergny, G. Duval-Valentin, C. H. Nguyen, L. Perrouault, B. Faucon, M. Rougée, T. Montenay-Garestier, E. Bisagni and C. Hélène, *Science*, 1992, **256**, 1681-1684.
94. D. S. Pilch, M. T. Martin, C. H. Nguyen, J. S. Sun, E. Bisagni, T. Garestier and C. Helene, *J. Am. Chem. Soc.*, 1993, **115**, 9941-9951.
95. C. Escude, C. H. Nguyen, J. L. Mergny, J. S. Sun, E. Bisagni, T. Garestier and C. Helene, *J. Am. Chem. Soc.*, 1995, **117**, 10212-10219.
96. T. Biver, A. Boggioni, B. García, J. M. Leal, R. Ruiz, F. Secco and M. Venturini, *Nucleic Acids Res.*, 2010, **38**, 1697-1710.
97. C. Escudé, C. H. Nguyen, S. Kukreti, Y. Janin, J. S. Sun, E. Bisagni, T. Garestier and C. Hélène, *Proc. Natl. Acad. Sci. U.S.A.*, 1998, **95**, 3591-3596.
98. C. H. Nguyen, E. Fan, J. F. Riou, M. C. Bissery, P. Vrignaud, F. Lavelle and E. Bisagni, *Anticancer Drug Des.*, 1995, **10**, 277-297.
99. C. H. Nguyen, C. Marchand, S. Delage, D. Sun, T. Garestier, C. Hélène and E. Bisagni, *J. Am. Chem. Soc.*, 1998, **120**, 2501-2507.
100. W. D. Wilson, F. A. Tanious, S. Mizan, S. Yao, A. S. Kiselyov, G. Zon and L. Strekowski, *Biochemistry*, 1993, **32**, 10614-10621.
101. S. Bhuiya, L. Haque, R. Goswami and S. Das, *J. Phys. Chem. B*, 2017, **121**, 11037-11052.
102. R. Tiwari, L. Haque, S. Bhuiya and S. Das, *Int. J. Biol. Macromol.*, 2017, **103**, 692-700.
103. R. Sinha and G. S. Kumar, *J. Phys. Chem. B*, 2009, **113**, 13410-13420.
104. S. Das, G. S. Kumar, A. Ray and M. Maiti, *J. Biomol. Struct. Dyn.*, 2003, **20**, 703-714.
105. X. J. He and L. F. Tan, *Inorg. Chem.*, 2014, **53**, 11152-11159.
106. Y. W. Park and K. J. Breslauer, *Proc. Natl. Acad. Sci. U.S.A.*, 1992, **89**, 6653-6657.
107. M. Durand, N. T. Thuong and J. C. Maurizot, *J. Biol. Chem.*, 1992, **267**, 24394-24399.
108. K. Umamoto, M. H. Sarma, G. Gupta, J. Luo and R. H. Sarma, *J. Am. Chem. Soc.*, 1990, **112**, 4539-4545.
109. D. S. Pilch and K. J. Breslauer, *Proc. Natl. Acad. Sci. U.S.A.*, 1994, **91**, 9332-9336.
110. R. Wadhwa, S. C. Kaul, Y. Komatsu, Y. Ikawa, A. Sarai and Y. Sugimoto, *FEBS Lett.*, 1993, **315**, 193-196.
111. N. Vigneswaran, C. A. Mayfield, B. Rodu, R. James, H. G. Kim and D. M. Miller, *Biochemistry*, 1996, **35**, 1106-1114.
112. A. K. Jain and S. Bhattacharya, *Bioconjug. Chem.*, 2010, **21**, 1389-1403.
113. D. P. Arya, *Acc. Chem. Res.*, 2011, **44**, 134-146.
114. D. P. Arya and R. L. Coffee, Jr., *Bioorg. Med. Chem. Lett.*, 2000, **10**, 1897-1899.
115. D. P. Arya, R. L. Coffee, Jr., B. Willis and A. I. Abramovitch, *J. Am. Chem. Soc.*, 2001, **123**, 5385-5395.
116. L. Xue, H. Xi, S. Kumar, D. Gray, E. Davis, P. Hamilton, M. Skriba and D. P. Arya, *Biochemistry*, 2010, **49**, 5540-5552.
117. P. Ji, S. Diederichs, W. Wang, S. Böing, R. Metzger, P. M. Schneider, N. Tidow, B. Brandt, H. Buerger, E. Bulik, M. Thomas, W. E. Berdel, H. Serve and C. Müller-Tidow, *Oncogene*, 2003, **22**, 8031-8041.
118. T. Gutschner, M. Hämmerle, M. Eissmann, J. Hsu, Y. Kim, G. Hung, A. Revenko, G. Arun, M. Stentrup, M. Gross, M. Zörnig, A. R. MacLeod, D. L. Spector and S. Diederichs, *Cancer Res.*, 2013, **73**, 1180-1189.
119. A. Donlic, M. Zafferani, G. Padroni, M. Puri and A. E. Hargrove, *Nucleic Acids Res.*, 2020, **48**, 7653-7664.
120. F. A. Abulwerdi, W. Xu, A. A. Ageeli, M. J. Yonkunas, G. Arun, H. Nam, J. S. Schneekloth, Jr., T. K. Dayie, D. Spector, N. Baird and S. F. J. Le Grice, *ACS Chem. Biol.*, 2019, **14**, 223-235.
121. M. Gellert, M. N. Lipsett and D. R. Davies, *Proc. Natl. Acad. Sci. U.S.A.*, 1962, **48**, 2013-2018.
122. D. Sen and W. Gilbert, *Nature*, 1988, **334**, 364-366.
123. M. Cheng, Y. Cheng, J. Hao, G. Jia, J. Zhou, J. L. Mergny and C. Li, *Nucleic Acids Res.*, 2018, **46**, 9264-9275.
124. H. A. Pickett, J. D. Henson, A. Y. Au, A. A. Neumann and R. R. Reddel, *Hum. Mol. Genet.*, 2011, **20**, 4684-4692.
125. V. F. S. Kahl, J. A. M. Allen, C. B. Nelson, A. P. Sobinoff, M. Lee, T. Kilo, R. S. Vasireddy and H. A. Pickett, *Front. Cell Dev. Biol.*, 2020, **8**, 493.
126. W. E. Wright, V. M. Tesmer, K. E. Huffman, S. D. Levene and J. W. Shay, *Genes Dev.*, 1997, **11**, 2801-2809.
127. M. Z. Levy, R. C. Allsopp, A. B. Fletcher, C. W. Greider and C. B. Harley, *J. Mol. Biol.*, 1992, **225**, 951-960.
128. N. W. Kim, M. A. Piatsyzek, K. R. Prowse, C. B. Harley, M. D. West, P. L. Ho, G. M. Coviello, W. E. Wright, S. L. Weinrich and J. W. Shay, *Science*, 1994, **266**, 2011-2015.
129. G. B. Morin, *Cell*, 1989, **59**, 521-529.
130. A. N. Elyadi, D. A. Braasch and D. R. Corey, *Biochemistry*, 2002, **41**, 9973-9981.
131. J. Feng, W. D. Funk, S. S. Wang, S. L. Weinrich, A. A. Avilion, C. P. Chiu, R. R. Adams, E. Chang, R. C. Allsopp, J. Yu, S. Le, D. W. West, C. B. Harley, W. H. Andrews, C. W. Greider and B. Villeponteau, *Science*, 1995, **269**, 1236-1241.
132. A. M. Zahler, J. R. Williamson, T. R. Cech and D. M. Prescott, *Nature*, 1991, **350**, 718-720.
133. S. Neidle, *Fests j.*, 2010, **277**, 1118-1125.
134. H. Yaku, T. Fujimoto, T. Murashima, D. Miyoshi and N. Sugimoto, *Chem. Commun.*, 2012, **48**, 6203-6216.
135. P. Agrawal, E. Hatzakis, K. Guo, M. Carver and D. Yang, *Nucleic Acids Res.*, 2013, **41**, 10584-10592.
136. S. Bielskute, J. Plavec and P. Podbevšek, *Nucleic Acids Res.*, 2021, **49**, 2346-2356.
137. J. Dai, C. Punchedewa, A. Ambrus, D. Chen, R. A. Jones and D. Yang, *Nucleic Acids Res.*, 2007, **35**, 2440-2450.
138. H. Yaku, T. Murashima, D. Miyoshi and N. Sugimoto, *Chem. Commun.*, 2010, **46**, 5740-5742.
139. S. Burge, G. N. Parkinson, P. Hazel, A. K. Todd and S. Neidle, *Nucleic Acids Res.*, 2006, **34**, 5402-5415.
140. J. Gros, F. Rosu, S. Amrane, A. De Cian, V. Gabelica, L. Lacroix and J. L. Mergny, *Nucleic Acids Res.*, 2007, **35**, 3064-3075.
141. C. J. Lech, B. Heddi and A. T. Phan, *Nucleic Acids Res.*, 2013, **41**, 2034-2046.
142. P. Hobza and J. Sponer, *J. Am. Chem. Soc.*, 2002, **124**, 11802-11808.
143. J. Gu and J. Leszczynski, *J. Phys. Chem. A*, 2000, **104**, 6308-6313.
144. C. Roxo, W. Kotkowiak and A. Pasternak, *Int. J. Mol. Sci.*, 2021, **22**.
145. N. Q. Do, K. W. Lim, M. H. Teo, B. Heddi and A. T. Phan, *Nucleic Acids Res.*, 2011, **39**, 9448-9457.
146. N. Q. Do and A. T. Phan, *Chemistry*, 2012, **18**, 14752-14759.
147. M. Adrian, D. J. Ang, C. J. Lech, B. Heddi, A. Nicolas and A. T. Phan, *J. Am. Chem. Soc.*, 2014, **136**, 6297-6305.



148. T. C. Marsh, J. Vesenska and E. Henderson, *Nucleic Acids Res.*, 1995, **23**, 696-700.
149. D. Miyoshi, H. Karimata and N. Sugimoto, *Angew. Chem. Int. Ed. Engl.*, 2005, **44**, 3740-3744.
150. D. Miyoshi, H. Karimata, Z. M. Wang, K. Koumoto and N. Sugimoto, *J. Am. Chem. Soc.*, 2007, **129**, 5919-5925.
151. D. Miyoshi and N. Sugimoto, *Methods Mol. Biol.*, 2011, **749**, 93-104.
152. K. Bose, C. J. Lech, B. Heddi and A. T. Phan, *Nat. Commun.*, 2018, **9**, 1959.
153. A. M. Varzhuk, A. D. Protopopova, V. B. Tsvetkov, N. A. Barinov, V. V. Podgorsky, M. V. Tankevich, M. A. Vlasenok, V. V. Severov, I. P. Smirnov, E. V. Dubrovin, D. V. Klinov and G. E. Pozmogova, *Nucleic Acids Res.*, 2018, **46**, 8978-8992.
154. C. Saintomé, S. Amrane, J. L. Mergny and P. Alberti, *Nucleic Acids Res.*, 2016, **44**, 2926-2935.
155. H. Q. Yu, D. Miyoshi and N. Sugimoto, *J. Am. Chem. Soc.*, 2006, **128**, 15461-15468.
156. H. Yu, X. Gu, S. Nakano, D. Miyoshi and N. Sugimoto, *J. Am. Chem. Soc.*, 2012, **134**, 20060-20069.
157. A. Risitano and K. R. Fox, *Nucleic Acids Res.*, 2004, **32**, 2598-2606.
158. P. Hazel, J. Huppert, S. Balasubramanian and S. Neidle, *J. Am. Chem. Soc.*, 2004, **126**, 16405-16415.
159. N. Kumar and S. Maiti, *Nucleic Acids Res.*, 2008, **36**, 5610-5622.
160. C. Ghimire, S. Park, K. Iida, P. Yangyuoru, H. Otomo, Z. Yu, K. Nagasawa, H. Sugiyama and H. Mao, *J. Am. Chem. Soc.*, 2014, **136**, 15537-15544.
161. A. Guédin, J. Gros, P. Alberti and J. L. Mergny, *Nucleic Acids Res.*, 2010, **38**, 7858-7868.
162. E. Hatzakis, K. Okamoto and D. Yang, *Biochemistry*, 2010, **49**, 9152-9160.
163. A. A. Bizyaeva, D. A. Bunin, V. L. Moiseenko, A. S. Gambaryan, S. Balk, V. N. Tashlitsky, A. M. Arutyunyan, A. M. Kopylov and E. G. Zavyalova, *Int. J. Mol. Sci.*, 2021, **22**.
164. K. W. Lim, S. Amrane, S. Bouaziz, W. Xu, Y. Mu, D. J. Patel, K. N. Luu and A. T. Phan, *J. Am. Chem. Soc.*, 2009, **131**, 4301-4309.
165. Z. Zhang, J. Dai, E. Veliath, R. A. Jones and D. Yang, *Nucleic Acids Res.*, 2010, **38**, 1009-1021.
166. Y. Chen and D. Yang, *Curr Protoc Nucleic Acid Chem*, 2012, **Chapter 17**, Unit17.15.
167. K. W. Lim and A. T. Phan, *Angew. Chem. Int. Ed. Engl.*, 2013, **52**, 8566-8569.
168. K. W. Lim, T. Q. Nguyen and A. T. Phan, *J. Am. Chem. Soc.*, 2014, **136**, 17969-17973.
169. K. W. Lim, Z. J. Khong and A. T. Phan, *Biochemistry*, 2014, **53**, 247-257.
170. S. Ravichandran, M. Razaq, N. Parveen, A. Ghosh and K. K. Kim, *Nucleic Acids Res.*, 2021, **49**, 10689-10706.
171. H. Feng and C. K. Kwok, *RSC Chem. Biol.*, 2022, **3**, 431-435.
172. K. W. Lim, P. Jenjaroenpun, Z. J. Low, Z. J. Khong, Y. S. Ng, V. A. Kuznetsov and A. T. Phan, *Nucleic Acids Res.*, 2015, **43**, 5630-5646.
173. B. Karg, S. Mohr and K. Weisz, *Angew. Chem. Int. Ed. Engl.*, 2019, **58**, 11068-11071.
174. Y. Cheng, Q. Tang, Y. Li, Y. Zhang, C. Zhao, J. Yan and H. You, *J. Biol. Chem.*, 2019, **294**, 5890-5895.
175. T. Q. N. Nguyen, K. W. Lim and A. T. Phan, *J. Phys. Chem. B*, 2020, **124**, 5122-5130.
176. M. Nakata, N. Kosaka, K. Kawauchi and D. Miyoshi, *ACS Omega*, 2024, **9**, 35028-35036.
177. M. Nakata, N. Kosaka, K. Kawauchi and D. Miyoshi, *Biochemistry*, 2024, **64**, 609-619.
178. D. J. Y. Tan, F. R. Winnerdy, K. W. Lim and A. T. Phan, *Nucleic Acids Res.*, 2020, **48**, 11162-11171.
179. A. T. Phan, V. Kuryavyi, J. C. Darnell, A. Serganov, A. Majumdar, S. Ilin, T. Raslin, A. Polonskaia, C. Chen, D. Clain, R. B. Darnell and D. J. Patel, *Nat. Struct. Mol. Biol.*, 2011, **18**, 796-804.
180. K. D. Warner, M. C. Chen, W. Song, R. L. Strack, A. Thorn, S. R. Jaffrey and A. R. Ferré-D'Amaré, *Nat. Struct. Mol. Biol.*, 2014, **21**, 658-663.
181. H. Huang, N. B. Suslov, N. S. Li, S. A. Shelke, M. E. Evans, Y. Koldobskaya, P. A. Rice and J. A. Piccirilli, *Nat. Chem. Biol.*, 2014, **10**, 686-691.
182. D. Bhattacharyya, G. Mirihana Arachchilage and S. Basu, *Front. Chem.*, 2016, **4**, 38.
183. J. S. Lee, *Nucleic Acids Res.*, 1990, **18**, 6057-6060.
184. C. C. Hardin, T. Watson, M. Corregan and C. Bailey, *Biochemistry*, 1992, **31**, 833-841.
185. E. A. Venczel and D. Sen, *Biochemistry*, 1993, **32**, 6220-6228.
186. B. I. Kankia and L. A. Marky, *J. Am. Chem. Soc.*, 2001, **123**, 10799-10804.
187. A. Włodarczyk, P. Grzybowski, A. Patkowski and A. Dobek, *J. Phys. Chem. B*, 2005, **109**, 3594-3605.
188. F. M. Chen, *Biochemistry*, 1992, **31**, 3769-3776.
189. S. W. Blume, V. Guarcello, W. Zacharias and D. M. Miller, *Nucleic Acids Res.*, 1997, **25**, 617-625.
190. C. C. Hardin, A. G. Perry and K. White, *Biopolymers*, 2000, **56**, 147-194.
191. D. Miyoshi, A. Nakao, T. Toda and N. Sugimoto, *FEBS Lett.*, 2001, **496**, 128-133.
192. N. V. Hud, F. W. Smith, F. A. Anet and J. Feigon, *Biochemistry*, 1996, **35**, 15383-15390.
193. M. Beck, V. Lucić, F. Förster, W. Baumeister and O. Medalia, *Nature*, 2007, **449**, 611-615.
194. R. J. Ellis, *Curr. Opin. Struct. Biol.*, 2001, **11**, 114-119.
195. D. S. Goodsell, *Trends Biochem. Sci.*, 1991, **16**, 203-206.
196. S. B. Zimmerman and S. O. Trach, *J. Mol. Biol.*, 1991, **222**, 599-620.
197. K. Luby-Phelps, *Int Rev Cytol*, 2000, **192**, 189-221.
198. D. Miyoshi, H. Karimata and N. Sugimoto, *J. Am. Chem. Soc.*, 2006, **128**, 7957-7963.
199. T. Fujimoto, S. Nakano, N. Sugimoto and D. Miyoshi, *J. Phys. Chem. B*, 2013, **117**, 963-972.
200. D. Miyoshi, K. Nakamura, H. Tateishi-Karimata, T. Ohmichi and N. Sugimoto, *J. Am. Chem. Soc.*, 2009, **131**, 3522-3531.
201. Y. M. Ueda, Y. K. Zouzumi, A. Maruyama, S. I. Nakano, N. Sugimoto and D. Miyoshi, *Sci Technol Adv Mater*, 2016, **17**, 753-759.
202. Y. Xue, Z. Y. Kan, Q. Wang, Y. Yao, J. Liu, Y. H. Hao and Z. Tan, *J. Am. Chem. Soc.*, 2007, **129**, 11185-11191.
203. K. W. Zheng, Z. Chen, Y. H. Hao and Z. Tan, *Nucleic Acids Res.*, 2010, **38**, 327-338.
204. Z. Y. Kan, Y. Yao, P. Wang, X. H. Li, Y. H. Hao and Z. Tan, *Angew. Chem. Int. Ed. Engl.*, 2006, **45**, 1629-1632.
205. A. Marchand, R. Ferreira, H. Tateishi-Karimata, D. Miyoshi, N. Sugimoto and V. Gabelica, *J. Phys. Chem. B*, 2013, **117**, 12391-12401.
206. S. Muhuri, K. Mimura, D. Miyoshi and N. Sugimoto, *J. Am. Chem. Soc.*, 2009, **131**, 9268-9280.
207. H. Tateishi-Karimata, S. Pramanik, S. Nakano, D. Miyoshi and N. Sugimoto, *ChemMedChem*, 2014, **9**, 2150-2155.
208. S. Nakano, H. Karimata, T. Ohmichi, J. Kawakami and N. Sugimoto, *J. Am. Chem. Soc.*, 2004, **126**, 14330-14331.
209. H. Tateishi-Karimata, D. Banerjee, T. Ohyama, S. Matsumoto, D. Miyoshi, S. I. Nakano and N. Sugimoto, *Biochem. Biophys. Res. Commun.*, 2020, DOI: 10.1016/j.bbrc.2020.02.045.
210. D. Miyoshi, T. Fujimoto and N. Sugimoto, *Top. Curr. Chem.*, 2013, **330**, 87-110.
211. M. C. Miller, R. Buscaglia, J. B. Chaires, A. N. Lane and J. O. Trent, *J. Am. Chem. Soc.*, 2010, **132**, 17105-17107.
212. L. Petraccone, B. Pagano and C. Giancola, *Methods*, 2012, **57**, 76-83.
213. D. Miyoshi and N. Sugimoto, *Biochimie*, 2008, **90**, 1040-1051.
214. M. Tsuruta, Y. Sugitani, N. Sugimoto and D. Miyoshi, *Int. J. Mol. Sci.*, 2021, **22**.
215. D. Miyoshi, S. Matsumura, S. Nakano and N. Sugimoto, *J. Am. Chem. Soc.*, 2004, **126**, 165-169.
216. D. Miyoshi, A. Nakao and N. Sugimoto, *Biochemistry*, 2002, **41**, 15017-15024.
217. D. Miyoshi, H. Karimata and N. Sugimoto, *Angew. Chem. Int. Ed. Engl.*, 2005, **44**, 3740-3744.
218. B. Heddi and A. T. Phan, *J. Am. Chem. Soc.*, 2011, **133**, 9824-9833.
219. R. Buscaglia, M. C. Miller, W. L. Dean, R. D. Gray, A. N. Lane, J. O. Trent and J. B. Chaires, *Nucleic Acids Res.*, 2013, **41**, 7934-7946.
220. H. Q. Yu, D. H. Zhang, X. B. Gu, D. Miyoshi and N. Sugimoto, *Angew. Chem. Int. Ed. Engl.*, 2008, **47**, 9034-9038.
221. H. Yaku, T. Murashima, D. Miyoshi and N. Sugimoto, *J. Phys. Chem. B*, 2014, **118**, 2605-2614.
222. H. Yaku, T. Murashima, H. Tateishi-Karimata, S. Nakano, D. Miyoshi and N. Sugimoto, *Methods*, 2013, **64**, 19-27.
223. S. Cogoi, M. Paramasivam, B. Spolaoro and L. E. Xodo, *Nucleic Acids Res.*, 2008, **36**, 3765-3780.
224. B. Pagano, L. Margarucci, P. Zizza, J. Amato, N. Iaccarino, C. Cassiano, E. Salvati, E. Novellino, A. Biroccio, A. Casapullo and A. Randazzo, *Chem. Commun.*, 2015, **51**, 2964-2967.
225. V. González, K. Guo, L. Hurley and D. Sun, *J. Biol. Chem.*, 2009, **284**, 23622-23635.
226. L. T. Gray, A. C. Vallur, J. Eddy and N. Maizels, *Nat. Chem. Biol.*, 2014, **10**, 313-318.
227. V. Brázda, J. Červeň, M. Bartas, N. Mikysková, J. Coufal and P. Pečinka, *Molecules*, 2018, **23**.
228. Z. L. Huang, J. Dai, W. H. Luo, X. G. Wang, J. H. Tan, S. B. Chen and Z. S. Huang, *J. Am. Chem. Soc.*, 2018, **140**, 17945-17955.
229. M. M. Makowski, C. Gräwe, B. M. Foster, N. V. Nguyen, T. Bartke and M. Vermeulen, *Nat. Commun.*, 2018, **9**, 1653.
230. X. Zhang, J. Spiegel, S. Martínez Cuesta, S. Adhikari and S. Balasubramanian, *Nat. Chem.*, 2021, **13**, 626-633.
231. S. K. Mishra, A. Tawani, A. Mishra and A. Kumar, *Sci. Rep.*, 2016, **6**, 38144.
232. T. D. McKnight and D. E. Shippen, *Plant Cell*, 2004, **16**, 794-803.
233. M. A. Islam, S. D. Thomas, V. V. Murty, K. J. Sedoris and D. M. Miller, *J. Biol. Chem.*, 2014, **289**, 8521-8531.
234. S. K. Madden, A. D. de Araujo, M. Gerhardt, D. P. Fairlie and J. M. Mason, *Mol. Cancer*, 2021, **20**, 3.
235. V. González and L. H. Hurley, *Biochemistry*, 2010, **49**, 9706-9714.



236. R. Hänsel-Hertsch, D. Beraldi, S. V. Lensing, G. Marsico, K. Zyner, A. Parry, M. Di Antonio, J. Pike, H. Kimura, M. Narita, D. Tannahill and S. Balasubramanian, *Nat. Genet.*, 2016, **48**, 1267-1272.
237. F. Kouzine, D. Wojtowicz, L. Baranello, A. Yamane, S. Nelson, W. Resch, K. R. Kieffer-Kwon, C. J. Benham, R. Casellas, T. M. Przytycka and D. Levens, *Cell Syst*, 2017, **4**, 344-356.e347.
238. J. Spiegel, S. M. Cuesta, S. Adhikari, R. Hänsel-Hertsch, D. Tannahill and S. Balasubramanian, *Genome Biol.*, 2021, **22**, 117.
239. S. Roychoudhury, S. Pramanik, H. L. Harris, M. Tarpley, A. Sarkar, G. Spagnol, P. L. Sorgen, D. Chowdhury, V. Band, D. Klinkebiel and K. K. Bhakat, *Proc. Natl. Acad. Sci. U.S.A.*, 2020, **117**, 11409-11420.
240. S. L. Cree, R. Fredericks, A. Miller, F. G. Pearce, V. Filichev, C. Fee and M. A. Kennedy, *FEBS Lett.*, 2016, **590**, 2870-2883.
241. A. K. Byrd and K. D. Raney, *J. Biol. Chem.*, 2015, **290**, 6482-6494.
242. A. Gallo, C. Lo Sterzo, M. Mori, A. Di Matteo, I. Bertini, L. Banci, M. Brunori and L. Federici, *J. Biol. Chem.*, 2012, **287**, 26539-26548.
243. H. J. Kang, T. V. Le, K. Kim, J. Hur, K. K. Kim and H. J. Park, *J. Mol. Biol.*, 2014, **426**, 2594-2604.
244. S. C. Ha, J. Choi, H. Y. Hwang, A. Rich, Y. G. Kim and K. K. Kim, *Nucleic Acids Res.*, 2009, **37**, 629-637.
245. S. Eddy, L. Maddukuri, A. Ketkar, M. K. Zafar, E. E. Henninger, Z. F. Pursell and R. L. Eoff, *Biochemistry*, 2015, **54**, 3218-3230.
246. V. Brázda, L. Hároníková, J. C. Liao and M. Fojta, *Int. J. Mol. Sci.*, 2014, **15**, 17493-17517.
247. B. Heddi, V. V. Cheong, H. Martadinata and A. T. Phan, *Proc. Natl. Acad. Sci. U.S.A.*, 2015, **112**, 9608-9613.
248. M. C. Chen, R. Tippiana, N. A. Demeshkina, P. Murat, S. Balasubramanian, S. Myong and A. R. Ferré-D'Amaré, *Nature*, 2018, **558**, 465-469.
249. A. Traczyk, C. W. Liew, D. J. Gill and D. Rhodes, *Nucleic Acids Res.*, 2020, **48**, 4562-4571.
250. K. H. Ngo, C. W. Liew, B. Heddi and A. T. Phan, *J. Am. Chem. Soc.*, 2024, **146**, 13709-13713.
251. M. Sauer and K. Paeschke, *Biochem. Soc. Trans.*, 2017, **45**, 1173-1182.
252. O. Mendoza, A. Bourdoncle, J. B. Boulé, R. M. Brosh, Jr. and J. L. Mergny, *Nucleic Acids Res.*, 2016, **44**, 1989-2006.
253. K. N. Estep, T. J. Butler, J. Ding and R. M. Brosh, *Curr. Med. Chem.*, 2019, **26**, 2881-2897.
254. R. M. Brosh, Jr., *Nat Rev Cancer*, 2013, **13**, 542-558.
255. Y. Wu, K. Shin-ya and R. M. Brosh, Jr., *Mol. Cell. Biol.*, 2008, **28**, 4116-4128.
256. V. A. Soldatenkov, A. A. Vetcher, T. Duka and S. Ladame, *ACS Chem. Biol.*, 2008, **3**, 214-219.
257. M. Paramasivam, A. Membrino, S. Cogoi, H. Fukuda, H. Nakagama and L. E. Xodo, *Nucleic Acids Res.*, 2009, **37**, 2841-2853.
258. S. Cogoi, S. Zorzet, V. Rapozzi, I. Géci, E. B. Pedersen and L. E. Xodo, *Nucleic Acids Res.*, 2013, **41**, 4049-4064.
259. S. Cogoi, M. Paramasivam, A. Membrino, K. K. Yokoyama and L. E. Xodo, *J. Biol. Chem.*, 2010, **285**, 22003-22016.
260. K. Takahama, A. Takada, S. Tada, M. Shimizu, K. Sayama, R. Kurokawa and T. Oyoshi, *Chem. Biol.*, 2013, **20**, 341-350.
261. A. Ishiguro, N. Kimura, Y. Watanabe, S. Watanabe and A. Ishihama, *Genes Cells*, 2016, **21**, 466-481.
262. J. M. Knop, S. K. Mukherjee, R. Oliva, S. Möbitz and R. Winter, *J. Am. Chem. Soc.*, 2020, **142**, 18299-18303.
263. A. Bugaut and S. Balasubramanian, *Nucleic Acids Res.*, 2012, **40**, 4727-4741.
264. S. Kumari, A. Bugaut, J. L. Huppert and S. Balasubramanian, *Nat. Chem. Biol.*, 2007, **3**, 218-221.
265. G. Biffi, M. Di Antonio, D. Tannahill and S. Balasubramanian, *Nat. Chem.*, 2014, **6**, 75-80.
266. E. G. Conlon, L. Lu, A. Sharma, T. Yamazaki, T. Tang, N. A. Shneider and J. L. Manley, *Elife*, 2016, **5**.
267. C. K. Kwok, G. Marsico, A. B. Sahakyan, V. S. Chambers and S. Balasubramanian, *Nat. Methods*, 2016, **13**, 841-844.
268. J. Eddy and N. Maizels, *Nucleic Acids Res.*, 2008, **36**, 1321-1333.
269. J. L. Huppert, A. Bugaut, S. Kumari and S. Balasubramanian, *Nucleic Acids Res.*, 2008, **36**, 6260-6268.
270. E. K. S. McRae, E. P. Booy, A. Moya-Torres, P. Ezzati, J. Stetefeld and S. A. McKenna, *Nucleic Acids Res.*, 2017, **45**, 6656-6668.
271. P. Chakraborty and F. Grosse, *DNA Repair (Amst)*, 2011, **10**, 654-665.
272. B. Giri, P. J. Smaldino, R. G. Thys, S. D. Creacy, E. D. Routh, R. R. Hantgan, S. Lattmann, Y. Nagamine, S. A. Akman and J. P. Vaughn, *Nucleic Acids Res.*, 2011, **39**, 7161-7178.
273. L. Dumas, P. Herviou, E. Dassi, A. Cammas and S. Millevoi, *Trends Biochem. Sci.*, 2021, **46**, 270-283.
274. S. Mestre-Fos, P. I. Penev, S. Suttapitugsakul, M. Hu, C. Ito, A. S. Petrov, R. M. Wartell, R. Wu and L. D. Williams, *J. Mol. Biol.*, 2019, **431**, 1940-1955.
275. S. Mestre-Fos, P. I. Penev, J. C. Richards, W. L. Dean, R. D. Gray, J. B. Chaires and L. D. Williams, *PLoS One*, 2019, **14**, e0226177.
276. J. C. Bowman, A. S. Petrov, M. Frenkel-Pinter, P. I. Penev and L. D. Williams, *Chem. Rev.*, 2020, **120**, 4848-4878.
277. S. Khateb, P. Weisman-Shomer, I. Hershco, L. A. Loeb and M. Fry, *Nucleic Acids Res.*, 2004, **32**, 4145-4154.
278. A. von Hacht, O. Seifert, M. Menger, T. Schütze, A. Arora, Z. Konthür, P. Neubauer, A. Wagner, C. Weise and J. Kurreck, *Nucleic Acids Res.*, 2014, **42**, 6630-6644.
279. S. Boeynaems, S. Alberti, N. L. Fawzi, T. Mittag, M. Polymenidou, F. Rousseau, J. Schymkowitz, J. Shorter, B. Wolozin, L. Van Den Bosch, P. Tompa and M. Fuxreiter, *Trends Cell Biol.*, 2018, **28**, 420-435.
280. S. Alberti and D. Dormann, *Annu. Rev. Genet.*, 2019, **53**, 171-194.
281. Q. Li, X. Peng, Y. Li, W. Tang, J. Zhu, J. Huang, Y. Qi and Z. Zhang, *Nucleic Acids Res.*, 2020, **48**, D320-d327.
282. K. Kohata and D. Miyoshi, *Biophys. Rev.*, 2020, **12**, 669-676.
283. Y. R. Kamimura and M. Kanai, *Bull. Chem. Soc. Jpn.*, 2021, **94**, 1045-1058.
284. Y. Zhang, M. Yang, S. Duncan, X. Yang, M. A. S. Abdelhamid, L. Huang, H. Zhang, P. N. Benfey, Z. A. E. Waller and Y. Ding, *Nucleic Acids Res.*, 2019, **47**, 11746-11754.
285. M. Mimura, S. Tomita, Y. Shinkai, T. Hosokai, H. Kumeta, T. Saio, K. Shiraki and N. Kurita, *J. Am. Chem. Soc.*, 2021, **143**, 9849-9857.
286. M. Tsuruta, T. Torii, K. Kohata, K. Kawauchi, H. Tateishi-Karimata, N. Sugimoto and D. Miyoshi, *Chem. Commun.*, 2022, **58**, 12931-12934.
287. S. Shil, M. Tsuruta, K. Kawauchi and D. Miyoshi, *ChemMedChem*, 2024, DOI: 10.1002/cmcd.202400460, e202400460.
288. A. Zbinden, M. Pérez-Berlanga, P. De Rossi and M. Polymenidou, *Dev. Cell*, 2020, **55**, 45-68.
289. E. Wang, R. Thombre, Y. Shah, R. Latanich and J. Wang, *Nucleic Acids Res.*, 2021, **49**, 4816-4830.
290. Q. F. Yang, X. R. Wang, Y. H. Wang, X. H. Wu, R. Y. Shi, Y. X. Wang, H. N. Zhu, S. Yang, Y. L. Tang and F. Li, *Nucleic Acids Res.*, 2024, DOI: 10.1093/nar/gkae835.
291. S. Neidle, *Molecules*, 2024, **29**.
292. J. Hilton, K. Gelmon, P. L. Bedard, D. Tu, H. Xu, A. V. Tinker, R. Goodwin, S. A. Laurie, D. Jonker, A. R. Hansen, Z. W. Veitch, D. J. Renouf, L. Hagerman, H. Lui, B. Chen, D. Kellar, I. Li, S. E. Lee, T. Kono, B. Y. C. Cheng, D. Yap, D. Lai, S. Beatty, J. Soong, K. I. Pritchard, I. Soria-Bretones, E. Chen, H. Feilottter, M. Rushton, L. Seymour, S. Aparicio and D. W. Cescon, *Nat. Commun.*, 2022, **13**, 3607.
293. F. X. Han, R. T. Wheelhouse and L. H. Hurley, *J. Am. Chem. Soc.*, 1999, **121**, 3561-3570.
294. E. Izbicka, R. T. Wheelhouse, E. Raymond, K. K. Davidson, R. A. Lawrence, D. Sun, B. E. Windle, L. H. Hurley and D. Von Hoff, *Cancer Res.*, 1999, **59**, 639-644.
295. A. De Cian, G. Cristofari, P. Reichenbach, E. De Lemos, D. Monchaud, M. P. Teulade-Fichou, K. Shin-Ya, L. Lacroix, J. Lingner and J. L. Mergny, *Proc. Natl. Acad. Sci. U.S.A.*, 2007, **104**, 17347-17352.
296. Y. X. Yao, B. H. Xu and Y. Zhang, *Biomed. Res. Int.*, 2018, **2018**, 9701957.
297. D. Drygin, A. Siddiqui-Jain, S. O'Brien, M. Schwabe, A. Lin, J. Bliesath, C. B. Ho, C. Proffitt, K. Trent, J. P. Whitten, J. K. Lim, D. Von Hoff, K. Anderes and W. G. Rice, *Cancer Res.*, 2009, **69**, 7653-7661.
298. K. Shin-ya, K. Wierzbka, K. Matsuo, T. Ohtani, Y. Yamada, K. Furihata, Y. Hayakawa and H. Seto, *J. Am. Chem. Soc.*, 2001, **123**, 1262-1263.
299. G. A. Blankson, D. S. Pilch, A. A. Liu, L. F. Liu, J. E. Rice and E. J. LaVoie, *Bioorg. Med. Chem.*, 2013, **21**, 4511-4520.
300. M. Yasuda, Y. Ma, S. Okabe, Y. Wakabayashi, D. Su, Y. T. Chang, H. Seimiya, M. Tera and K. Nagasawa, *Chem. Commun.*, 2020, **56**, 12905-12908.
301. R. Rodriguez, S. Müller, J. A. Yeoman, C. Trentesaux, J. F. Riou and S. Balasubramanian, *J. Am. Chem. Soc.*, 2008, **130**, 15758-15759.
302. D. Koirala, S. Dhakal, B. Ashbridge, Y. Sannohe, R. Rodriguez, H. Sugiyama, S. Balasubramanian and H. Mao, *Nat. Chem.*, 2011, **3**, 782-787.
303. R. Rodriguez, K. M. Miller, J. V. Forment, C. R. Bradshaw, M. Nikan, S. Britton, T. Oelschlaegel, B. Xhemalce, S. Balasubramanian and S. P. Jackson, *Nat. Chem. Biol.*, 2012, **8**, 301-310.
304. M. Di Antonio, A. Ponjavic, A. Radzevičius, R. T. Ranasinghe, M. Catalano, X. Zhang, J. Shen, L. M. Needham, S. F. Lee, D. Klenerman and S. Balasubramanian, *Nat. Chem.*, 2020, **12**, 832-837.
305. D. Monchaud, C. Allain, H. Bertrand, N. Smargiasso, F. Rosu, V. Gabelica, A. De Cian, J. L. Mergny and M. P. Teulade-Fichou, *Biochimie*, 2008, **90**, 1207-1223.
306. A. De Cian, P. Grellier, E. Mouray, D. Depoix, H. Bertrand, D. Monchaud, M. P. Teulade-Fichou, J. L. Mergny and P. Alberti, *ChemBiochem*, 2008, **9**, 2730-2739.
307. X. J. Luo, Q. P. Qin, K. B. Huang, Y. C. Liu and H. Liang, *Asian J. Chem.*, 2013, **25**, 6396-6400.
308. N. Shioda, Y. Yabuki, K. Yamaguchi, M. Onozato, Y. Li, K. Kurosawa, H. Tanabe, N. Okamoto, T. Era, H. Sugiyama, T. Wada and K. Fukunaga, *Nat. Med.*, 2018, **24**, 802-813.
309. Y. Qi, H. Chen, W. Tan, Y. Li, G. Yuan and M. Xu, *Rapid Commun. Mass Spectrom.*, 2015, **29**, 1611-1616.
310. M. Zhu, J. Gao, X. J. Lin, Y. Y. Gong, Y. C. Qi, Y. L. Ma, Y. X. Song, W. Tan, F. Y. Li, M. Ye, J. Gong, Q. H. Cui, Z. H. Huang, Y. Y. Zhang, X. J. Wang, F. Lan, S. Q. Wang, G. Yuan, Y. Feng and M. Xu, *Nucleic Acids Res.*, 2021, **49**, 2522-2536.



311. V. Gabelica, R. Maeda, T. Fujimoto, H. Yaku, T. Murashima, N. Sugimoto and D. Miyoshi, *Biochemistry*, 2013, **52**, 5620-5628.
312. H. Arthanari, S. Basu, T. L. Kawano and P. H. Bolton, *Nucleic Acids Res.*, 1998, **26**, 3724-3728.
313. J. Ren and J. B. Chaires, *Biochemistry*, 1999, **38**, 16067-16075.
314. M. Perenon, H. Bonnet, T. Lavergne, J. Dejeu and E. Defrancq, *Phys. Chem. Chem. Phys.*, 2020, **22**, 4158-4164.
315. K. Kawachi, W. Sugimoto, T. Yasui, K. Murata, K. Itoh, K. Takagi, T. Tsuruoka, K. Akamatsu, H. Tateishi-Karimata, N. Sugimoto and D. Miyoshi, *Nat. Commun.*, 2018, **9**, 2271.
316. A. Ferino, G. Nicoletto, F. D'Este, S. Zorzet, S. Lago, S. N. Richter, A. Tikhomirov, A. Shchekotikhin and L. E. Xodo, *J. Med. Chem.*, 2020, **63**, 1245-1260.
317. M. G. Costales, Y. Matsumoto, S. P. Velagapudi and M. D. Disney, *J. Am. Chem. Soc.*, 2018, **140**, 6741-6744.
318. K. M. Patil, D. Chin, H. L. Seah, Q. Shi, K. W. Lim and A. T. Phan, *Chem. Commun.*, 2021, **57**, 12816-12819.
319. S. Mikutis, M. Rebelo, E. Yankova, M. Gu, C. Tang, A. R. Coelho, M. Yang, M. E. Hazemi, M. Pires de Miranda, M. Eleftheriou, M. Robertson, G. S. Vassiliou, D. J. Adams, J. P. Simas, F. Corzana, J. S. Schneekloth, Jr., K. Tzelepis and G. J. L. Bernardes, *ACS Cent Sci*, 2023, **9**, 892-904.
320. Y. Zhang, L. Wang, F. Wang, X. Chu and J. H. Jiang, *J. Am. Chem. Soc.*, 2024, **146**, 15815-15824.
321. M. Xiao, J. Zhao, Q. Wang, J. Liu and L. Ma, *Biomolecules*, 2022, **12**.
322. J. Dai, M. Carver, L. H. Hurley and D. Yang, *J. Am. Chem. Soc.*, 2011, **133**, 17673-17680.
323. J. Dickerhoff, J. Dai and D. Yang, *Nucleic Acids Res.*, 2021, **49**, 5905-5915.
324. L. Y. Liu, T. Z. Ma, Y. L. Zeng, W. Liu and Z. W. Mao, *J. Am. Chem. Soc.*, 2022, **144**, 11878-11887.
325. F. S. Di Leva, E. Novellino, A. Cavalli, M. Parrinello and V. Limongelli, *Nucleic Acids Res.*, 2014, **42**, 5447-5455.
326. H. J. Sullivan, C. Readmond, C. Radicella, V. Persad, T. J. Fasano and C. Wu, *ACS Omega*, 2018, **3**, 14788-14806.
327. T. Santos, G. F. Salgado, E. J. Cabrita and C. Cruz, *Pharmaceuticals (Basel)*, 2021, **14**.
328. T. Liu, Y. Wu, L. Qin, Q. Luo, W. Li, Y. Cheng, Y. Tu and H. You, *J. Phys. Chem. B*, 2023, **127**, 5859-5868.
329. K. Gehring, J. L. Leroy and M. Guéron, *Nature*, 1993, **363**, 561-565.
330. P. Rajagopal and J. Feigon, *Nature*, 1989, **339**, 637-640.
331. A. Lupták, A. R. Ferré-D'Amare, K. Zhou, K. W. Zilm and J. A. Doudna, *J. Am. Chem. Soc.*, 2001, **123**, 8447-8452.
332. J. L. Leroy, *Nucleic Acids Res.*, 2009, **37**, 4127-4134.
333. E. P. Wright, J. L. Huppert and Z. A. E. Waller, *Nucleic Acids Res.*, 2017, **45**, 2951-2959.
334. T. Fujii and N. Sugimoto, *Phys. Chem. Chem. Phys.*, 2015, **17**, 16719-16722.
335. A. M. Fleming, K. M. Stewart, G. M. Eyring, T. E. Ball and C. J. Burrows, *Org. Biomol. Chem.*, 2018, **16**, 4537-4546.
336. A. L. Lieblein, B. Fürtig and H. Schwalbe, *Chembiochem*, 2013, **14**, 1226-1230.
337. J. L. Mergny, L. Lacroix, X. G. Han, J. L. Leroy and C. Helene, *J. Am. Chem. Soc.*, 1995, **117**, 8887-8898.
338. H. A. Day, P. Pavlou and Z. A. Waller, *Bioorg. Med. Chem.*, 2014, **22**, 4407-4418.
339. A. Rajendran, S. Nakano and N. Sugimoto, *Chem. Commun.*, 2010, **46**, 1299-1301.
340. J. Cui, P. Waltman, V. H. Le and E. A. Lewis, *Molecules*, 2013, **18**, 12751-12767.
341. L. Ying, J. J. Green, H. Li, D. Klenerman and S. Balasubramanian, *Proc. Natl. Acad. Sci. U.S.A.*, 2003, **100**, 14629-14634.
342. D. Sun and L. H. Hurley, *J. Med. Chem.*, 2009, **52**, 2863-2874.
343. N. Gilbert and J. Allan, *Curr. Opin. Genet. Dev.*, 2014, **25**, 15-21.
344. G. Cimino-Reale, E. Pascale, E. Alvino, G. Starace and E. D'Ambrosio, *J. Biol. Chem.*, 2003, **278**, 2136-2140.
345. S. Dzatko, M. Krafčikova, R. Hānsel-Hertsch, T. Fessl, R. Fiala, T. Loja, D. Krafčik, J. L. Mergny, S. Foldynova-Trantirkova and L. Trantirek, *Angew. Chem. Int. Ed. Engl.*, 2018, **57**, 2165-2169.
346. M. Zeraati, D. B. Langley, P. Schofield, A. L. Moye, R. Rouet, W. E. Hughes, T. M. Bryan, M. E. Dinger and D. Christ, *Nat. Chem.*, 2018, **10**, 631-637.
347. T. Simonsson, M. Pribylova and M. Vorlickova, *Biochem. Biophys. Res. Commun.*, 2000, **278**, 158-166.
348. A. V. Makeyev and S. A. Liebhaber, *Rna*, 2002, **8**, 265-278.
349. E. F. Michelotti, G. A. Michelotti, A. I. Aronsohn and D. Levens, *Mol. Cell. Biol.*, 1996, **16**, 2350-2360.
350. Z. Du, J. Yu, Y. Chen, R. Andino and T. L. James, *J. Biol. Chem.*, 2004, **279**, 48126-48134.
351. L. Lacroix, H. Liénard, E. Labourier, M. Djavaheri-Mergny, J. Lacoste, H. Leffers, J. Tazi, C. Hélène and J. L. Mergny, *Nucleic Acids Res.*, 2000, **28**, 1564-1575.
352. A. Bandiera, G. Tell, E. Marsich, A. Scaloni, G. Pocsfalvi, A. Akintunde Akindahunsi, L. Cesaratto and G. Manzini, *Arch. Biochem. Biophys.*, 2003, **409**, 305-314. DOI: 10.1059/D4CB00287C
353. D. J. Uribe, K. Guo, Y. J. Shin and D. Sun, *Biochemistry*, 2011, **50**, 3796-3806.
354. S. Sedghi Masoud, S. M. Hecht and L. H. Hurley, *J. Am. Chem. Soc.*, 2014, **136**, 4172-4185.
355. S. L. Brown and S. Kendrick, *Pharmaceuticals (Basel)*, 2021, **14**.
356. S. Sedghi Masoud and K. Nagasawa, *Chem. Pharm. Bull.*, 2018, **66**, 1091-1103.
357. H. Abou Assi, M. Garavis, C. González and M. J. Damha, *Nucleic Acids Res.*, 2018, **46**, 8038-8056.
358. O. Y. Fedoroff, A. Rangan, V. V. Chemeris and L. H. Hurley, *Biochemistry*, 2000, **39**, 15083-15090.
359. A. Siddiqui-Jain, C. L. Grand, D. J. Bearss and L. H. Hurley, *Proc. Natl. Acad. Sci. U.S.A.*, 2002, **99**, 11593-11598.
360. L. Wang, Y. Wu, T. Chen and C. Wei, *Int. J. Biol. Macromol.*, 2013, **52**, 1-8.
361. P. Alberti, J. Ren, M. P. Teulade-Fichou, L. Guittat, J. F. Riou, J. Chaires, C. Hélène, J. P. Vigneron, J. M. Lehn and J. L. Mergny, *J. Biomol. Struct. Dyn.*, 2001, **19**, 505-513.
362. S. Sedghi Masoud, Y. Yamaoki, Y. Ma, A. Marchand, F. R. Winnerdy, V. Gabelica, A. T. Phan, M. Katahira and K. Nagasawa, *Chembiochem*, 2018, **19**, 2268-2272.
363. H. Xu, H. Zhang and X. Qu, *J. Inorg. Biochem.*, 2006, **100**, 1646-1652.
364. S. Shi, X. Geng, J. Zhao, T. Yao, C. Wang, D. Yang, L. Zheng and L. Ji, *Biochimie*, 2010, **92**, 370-377.
365. S. Kendrick, H. J. Kang, M. P. Alam, M. M. Madathil, P. Agrawal, V. Gokhale, D. Yang, S. M. Hecht and L. H. Hurley, *J. Am. Chem. Soc.*, 2014, **136**, 4161-4171.
366. A. M. Burger, F. Dai, C. M. Schultes, A. P. Reszka, M. J. Moore, J. A. Double and S. Neidle, *Cancer Res.*, 2005, **65**, 1489-1496.
367. X. Li, Y. Peng, J. Ren and X. Qu, *Proc. Natl. Acad. Sci. U.S.A.*, 2006, **103**, 19658-19663.
368. Y. Chen, K. Qu, C. Zhao, L. Wu, J. Ren, J. Wang and X. Qu, *Nat. Commun.*, 2012, **3**, 1074.
369. B. Shu, J. Cao, G. Kuang, J. Qiu, M. Zhang, Y. Zhang, M. Wang, X. Li, S. Kang, T. M. Ou, J. H. Tan, Z. S. Huang and D. Li, *Chem. Commun.*, 2018, **54**, 2036-2039.
370. G. Kuang, M. Zhang, S. Kang, D. Hu, X. Li, Z. Wei, X. Gong, L. K. An, Z. S. Huang, B. Shu and D. Li, *J. Med. Chem.*, 2020, **63**, 9136-9153.
371. L. Meng, Y. Su, F. Yang, S. Xiao, Z. Yin, J. Liu, J. Zhong, D. Zhou and F. Yu, *Bioorg. Med. Chem.*, 2019, **27**, 115147.
372. H. Zeng, S. Kang, Y. Zhang, K. Liu, Q. Yu, D. Li and L. K. An, *Int. J. Mol. Sci.*, 2021, **22**.
373. S. Kendrick, H. J. Kang, M. P. Alam, M. M. Madathil, P. Agrawal, V. Gokhale, D. Yang, S. M. Hecht and L. H. Hurley, *J. Am. Chem. Soc.*, 2016, **138**, 11408.
374. E. P. Wright, H. A. Day, A. M. Ibrahim, J. Kumar, L. J. Boswell, C. Huguin, C. E. Stevenson, K. Pors and Z. A. Waller, *Sci. Rep.*, 2016, **6**, 39456.
375. Q. Sheng, J. C. Neaverson, T. Mahmoud, C. E. M. Stevenson, S. E. Matthews and Z. A. E. Waller, *Org. Biomol. Chem.*, 2017, **15**, 5669-5673.
376. S. Dhillon, *Drugs*, 2020, **80**, 1853-1858.
377. N. A. Naryshkin, M. Weetall, A. Dakka, J. Narasimhan, X. Zhao, Z. Feng, K. K. Ling, G. M. Karp, H. Qi, M. G. Woll, G. Chen, N. Zhang, V. Gabbeta, P. Vazirani, A. Bhattacharyya, B. Furia, N. Risher, J. Sheedy, R. Kong, J. Ma, A. Turpoff, C. S. Lee, X. Zhang, Y. C. Moon, P. Trifillis, E. M. Welch, J. M. Colacino, J. Babiak, N. G. Almstead, S. W. Peltz, L. A. Eng, K. S. Chen, J. L. Mull, M. S. Lynes, L. L. Rubin, P. Fountora, L. Santarelli, D. Haehnke, K. D. McCarthy, R. Schmucki, M. Ebeling, M. Sivaramakrishnan, C. P. Ko, S. V. Paushkin, H. Ratni, I. Gerlach, A. Ghosh and F. Metzger, *Science*, 2014, **345**, 688-693.
378. H. Ratni, G. M. Karp, M. Weetall, N. A. Naryshkin, S. V. Paushkin, K. S. Chen, K. D. McCarthy, H. Qi, A. Turpoff, M. G. Woll, X. Zhang, N. Zhang, T. Yang, A. Dakka, P. Vazirani, X. Zhao, E. Pinard, L. Green, P. David-Pierson, D. Tuerck, A. Poirier, W. Muster, S. Kirchner, L. Mueller, I. Gerlach and F. Metzger, *J. Med. Chem.*, 2016, **59**, 6086-6100.
379. S. Campagne, S. Boigner, S. Rüdiger, A. Moursy, L. Gillioz, A. Knörlein, J. Hall, H. Ratni, A. Cléry and F. H. Allain, *Nat. Chem. Biol.*, 2019, **15**, 1191-1198.
380. H. Ratni, L. Mueller and M. Ebeling, *Prog Med Chem*, 2019, **58**, 119-156.
381. M. Sivaramakrishnan, K. D. McCarthy, S. Campagne, S. Huber, S. Meier, A. Augustin, T. Heckel, H. Meistermann, M. N. Hug, P. Birrer, A. Moursy, B. Khawaja, R. Schmucki, N. Berntsen, N. Giroud, S. Golling, M. Tzouros, B. Banfaj, R. Duran-Pacheco, J. Lamerz, Y. Hsiu Liu, T. Luebbbers, H. Ratni, M. Ebeling, A. Cléry, S. Paushkin, A. R. Krainer, F. H. Allain and F. Metzger, *Nat. Commun.*, 2017, **8**, 1476.
382. A. A. Ahmed, R. Angell, S. Oxenford, J. Worthington, N. Williams, N. Barton, T. G. Fowler, D. E. O'Flynn, M. Sunose, M. McConville, T. Vo, W. D. Wilson, S. A. Karim, J. P. Morton and S. Neidle, *ACS Med Chem Lett*, 2020, **11**, 1634-1644.
383. A. A. Ahmed, S. Chen, M. Roman-Escorza, R. Angell, S. Oxenford, M. McConville, N. Barton, M. Sunose, D. Neidle, S. Haider, T. Arshad and S. Neidle, *Sci. Rep.*, 2024, **14**, 3447.



ARTICLE

Journal Name

384. C. Marchetti, K. G. Zyner, S. A. Ohnmacht, M. Robson, S. M. Haider, J. P. Morton, G. Marsico, T. Vo, S. Laughlin-Toth, A. A. Ahmed, G. Di Vita, I. Pazitna, M. Gunaratnam, R. J. Besser, A. C. G. Andrade, S. Diocou, J. A. Pike, D. Tannahill, R. B. Pedley, T. R. J. Evans, W. D. Wilson, S. Balasubramanian and S. Neidle, *J. Med. Chem.*, 2018, **61**, 2500-2517.
385. A. A. Ahmed, W. Greenhalf, D. H. Palmer, N. Williams, J. Worthington, T. Arshad, S. Haider, E. Alexandrou, D. Guneri, Z. A. E. Waller and S. Neidle, *Molecules*, 2023, **28**.
386. J. Jana, Y. M. Vianney, N. Schröder and K. Weisz, *Nucleic Acids Res.*, 2022, **50**, 7161-7175.
387. A. T. Phan, K. N. Luu and D. J. Patel, *Nucleic Acids Res.*, 2006, **34**, 5715-5719.
388. M. Webba da Silva, M. Trajkovski, Y. Sannohe, N. Ma'ani Hessari, H. Sugiyama and J. Plavec, *Angew. Chem. Int. Ed. Engl.*, 2009, **48**, 9167-9170.
389. K. Li, L. Yatsunyk and S. Neidle, *Nucleic Acids Res.*, 2021, **49**, 519-528.
390. M. Zafferani and A. E. Hargrove, *Cell Chem. Biol.*, 2021, **28**, 594-609.
391. S. F. Banani, H. O. Lee, A. A. Hyman and M. K. Rosen, *Nat. Rev. Mol. Cell Biol.*, 2017, **18**, 285-298.
392. Y. Shin and C. P. Brangwynne, *Science*, 2017, **357**.
393. A. R. Strom, A. V. Emelyanov, M. Mir, D. V. Fyodorov, X. Darzacq and G. H. Karpen, *Nature*, 2017, **547**, 241-245.
394. Y. Lin, D. S. Protter, M. K. Rosen and R. Parker, *Mol. Cell*, 2015, **60**, 208-219.
395. B. R. Sabari, A. Dall'Agnese, A. Boija, I. A. Klein, E. L. Coffey, K. Shrinivas, B. J. Abraham, N. M. Hannett, A. V. Zamudio, J. C. Manteiga, C. H. Li, Y. E. Guo, D. S. Day, J. Schuijers, E. Vasile, S. Malik, D. Hnisz, T. I. Lee, Cisse, II, R. G. Roeder, P. A. Sharp, A. K. Chakraborty and R. A. Young, *Science*, 2018, **361**.
396. D. Monchaud, C. Allain and M. P. Teulade-Fichou, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 4842-4845.
397. A. De Cian, L. Guittat, M. Kaiser, B. Saccà, S. Amrane, A. Bourdoncle, P. Alberti, M. P. Teulade-Fichou, L. Lacroix and J. L. Mergny, *Methods*, 2007, **42**, 183-195.
398. C. Platella, D. Musumeci, A. Arciello, F. Doria, M. Freccero, A. Randazzo, J. Amato, B. Pagano and D. Montesarchio, *Anal. Chim. Acta*, 2018, **1030**, 133-141.
399. Y. Hashimoto, Y. Imagawa, K. Nagano, R. Maeda, N. Nagahama, T. Torii, N. Kinoshita, N. Takamiya, K. Kawauchi, H. Tatesishi-Karimata, N. Sugimoto and D. Miyoshi, *Chem. Commun.*, 2023, **59**, 4891-4894.
400. Y. Hashimoto, H. Kubo, K. Kawauchi and D. Miyoshi, *Chem. Commun.*, 2024, **60**, 13179-13182.
401. S. Zhang, H. Sun, D. Yang, Y. Liu, X. Zhang, H. Chen, Q. Li, A. Guan and Y. Tang, *Anal Chim Acta X*, 2019, **2**, 100017.
402. S. Kumar Mishra and A. Kumar, *Database (Oxford)*, 2016, **2016**.
403. Z. Cai, M. Zafferani, O. M. Akande and A. E. Hargrove, *J. Med. Chem.*, 2022, **65**, 7262-7277.

View Article Online
DOI: 10.1039/D4CB00287C

RSC Chemical Biology Accepted Manuscript



Three- and four-stranded nucleic acid structures and their ligands

Yoshiki Hashimoto, Sumit Shil, Mitsuki Tsuruta, Keiko Kawauchi, and
Daisuke Miyoshi*

Data availability

No primary research results, software or code have been included, and no new data were generated or analyzed as part of this review.

