

Cite this: *Dalton Trans.*, 2024, **53**, 5796

# A frontier-orbital view of the initial steps of lytic polysaccharide monooxygenase reactions†

Erna Katharina Wieduwilt, <sup>a</sup> Leila Lo Leggio <sup>b</sup> and Erik Donovan Hedegård <sup>\*a</sup>

Lytic polysaccharide monooxygenases (LPMOs) are copper enzymes that oxidatively cleave the strong C–H bonds in recalcitrant polysaccharide substrates, thereby playing a crucial role in biomass degradation. Recently, LPMOs have also been shown to be important for several pathogens. It is well established that the Cu(II) resting state of LPMOs is inactive, and the electronic structure of the active site needs to be altered to transform the enzyme into an active form. Whether this transformation occurs due to substrate binding or due to a unique priming reduction has remained speculative. Starting from four different crystal structures of the LPMO LsAA9A with well-defined oxidation states, we use a frontier molecular orbital approach to elucidate the initial steps of the LPMO reaction. We give an explanation for the requirement of the unique priming reduction and analyse electronic structure changes upon substrate binding. We further investigate how the presence of the substrate could facilitate an electron transfer from the copper active site to an H<sub>2</sub>O<sub>2</sub> co-substrate. Our findings could help to control experimental LPMO reactions.

Received 20th December 2023,  
Accepted 23rd February 2024

DOI: 10.1039/d3dt04275h

rsc.li/dalton


## Introduction

Lytic polysaccharide monooxygenases (LPMOs) are copper-dependent enzymes that catalyse the oxidative degradation of several polysaccharide substrates, such as cellulose, chitin, starch, and pectin.<sup>1–9</sup> LPMOs work by activating (otherwise inactive) C–H bonds in the glycosidic bonds that link the monomers in the respective polysaccharides. This activation ultimately leads to oxidation and cleavage of the glycosidic bond.<sup>10–12</sup> Several mechanistic pathways have been suggested for LPMOs, and we have summarized the most recent suggestions in Fig. 1.


Originally, LPMOs were discovered in bacteria and fungi. Today, they have been found across all kingdoms of life (though not yet in mammals) and exhibit a growing number of functions.<sup>10,13</sup> For instance, LPMOs and LPMO-like proteins were recently suggested to also play crucial roles in plant, insect and human pathogenesis such as blood diseases and meningitis.<sup>13,14,15</sup> However, the function that has so far

attracted the most attention is a significant boosting effect on biomass degradation. Nowadays, LPMOs are employed in industrial enzyme cocktails to produce biofuels and other value-added products.<sup>16,17</sup>

To date, eight different LPMO families have been discovered, namely AA9-AA17 (AA12 does not contain LPMOs).<sup>18</sup> All members share a common active site: a copper centre coordinated by two histidine residues in a motif termed the histidine brace (see Fig. 2).<sup>1</sup> Initially, LPMOs were thought to be strictly monooxygenases,<sup>1,19,20</sup> catalysing the reaction



but it remains unclear how protons and electrons can be delivered to the active site after substrate binding (see reactions (3b), (4b) and (7b) in Fig. 1).<sup>12,21</sup> Meanwhile, it has been suggested that LPMOs are in fact peroxygenases,<sup>12,21,22</sup> working according to the reaction




The peroxygenase reaction circumvents the requirement for external protons and electrons<sup>12,21</sup> (see reactions (5b)–(7b) in Fig. 1), and also the requirement by certain monooxygenase reaction mechanisms to reduce the Cu(II) to Cu(I) at every cycle, thus limiting the reducing agent requirement.<sup>22</sup> Furthermore, for the LPMOs where mono- and peroxygenase reactions have been compared, the latter is orders of magnitude faster.<sup>21,22</sup> For a few cases, the catalytic reaction has also been performed anaerobically, showing that these LPMOs indeed exclusively use H<sub>2</sub>O<sub>2</sub> as co-substrate<sup>23,24</sup> (according to

<sup>a</sup>Department of Physics, Chemistry, and Pharmacy, University of Southern Denmark, Campusvej 55, 5230 Odense M, Denmark. E-mail: erdh@sdu.dk

<sup>b</sup>Department of Chemistry, University of Copenhagen, 2100 Copenhagen, Denmark


† Electronic supplementary information (ESI) available: Selected distances and angles for investigated structures. Löwdin spin populations for all open-shell systems. QTAIM atomic charges for copper and summed for all residues. Section S1. Impact of pre-bound oxygen vs. superoxide. Section S2. Impact of chloride. Section S3. Results with TPSS. Section S4. Selected orbital energies and Löwdin reduced orbital populations. See DOI: <https://doi.org/10.1039/d3dt04275h>



**Fig. 1** Simplified reaction mechanism of LPMOs with boxes highlighting the structures that were investigated in this study. Corresponding structures are shown in Fig. 2 and Fig. S1.†

eqn (2) or reactions (5b)–(7b) in Fig. 1). However, whether this is true for all LPMOs is still controversial. Moreover, it is also unknown if some LPMOs can use H<sub>2</sub>O<sub>2</sub> as a “peroxide shunt” pathway as some iron-heme enzymes (see further below).<sup>25</sup>

A complicating factor for experimental investigations concerning the co-substrate is that LPMOs can catalyse the off-path oxidase reaction




in the absence of substrate.<sup>23,26–28</sup> Thus, also in the absence of externally added H<sub>2</sub>O<sub>2</sub>, trace amounts of H<sub>2</sub>O<sub>2</sub> generated by LPMOs may fuel the catalytic reaction. Furthermore, experimental studies are affected by possible non-enzymatic reactions involving reducing agents, free copper including copper leaked from denatured LPMOs, and added or *in situ* produced H<sub>2</sub>O<sub>2</sub>, further complicating the analysis of experimental results.<sup>29–31</sup> These complications have contributed to a situation where the details regarding the LPMOs co-substrate (O<sub>2</sub> or H<sub>2</sub>O<sub>2</sub>) are still controversial.

Regardless of whether LPMOs react according to eqn (1) as monooxygenases or according to eqn (2) as peroxygenases, or variations thereof, it has been firmly demonstrated that an initial reduction of the Cu(II) resting states is required for catalytic turnover (see reactions (1a) and (1b) in Fig. 1).<sup>32</sup> This is also true when LPMOs react as oxidases without substrate and produce H<sub>2</sub>O<sub>2</sub> according to eqn (3). This initial reduction has been denoted a “priming” reduction, for which a variety of reductants can be employed.<sup>2,11,33,34</sup>

It has been shown that both the Cu(I) state and the Cu(II) resting state can bind the substrate (according to reactions (2a) and (2b) in Fig. 1), but some experimental studies show that the reduced LPMO has a 2 to 10-fold higher affinity for cellulose.<sup>23,35</sup> Therefore, it has been proposed that metal reduction precedes substrate binding.

The nature of the priming reduction separates LPMOs from other metalloenzymes that also activate inactive C–H bonds – but whose reaction cycle otherwise shows similarities to the LPMOs’ catalytic cycle:<sup>36</sup> the iron-heme enzyme cytochrome P450 usually forms a highly oxidizing species (Cpd1) from O<sub>2</sub>. The resting state Fe(III)-heme unit is first reduced after substrate binding,<sup>37,38</sup> and O<sub>2</sub> binds to this Fe(II)-heme unit (similar to the O<sub>2</sub> pathway with reactions (3b) and (4b) in Fig. 1). The Fe(III)–O<sub>2</sub><sup>-</sup> species formed from initial O<sub>2</sub> binding is further reduced and protonated into a Fe(III)–OOH intermediate (Cpd0). The HOO<sup>-</sup> unit in Cpd0 is heterolytically cleaved with help from nearby residues that can participate in acid/base chemistry under oxidation of Fe(III) to the oxidative species (Cpd1), responsible for C–H activation. Thus, cytochrome P450 needs a continuous flow of electrons in its catalytic cycle. This is different from LPMOs, which can perform up to 20 catalytic cycles after one priming electron has been added<sup>39</sup> (when they work as peroxygenases). The iron-heme unit is also used by peroxidases, such as cytochrome *c* peroxidase and horseradish peroxidase (HRP). Unlike most cytochrome P450s, the peroxidases form the oxidative species (Cpd1) from H<sub>2</sub>O<sub>2</sub> binding to the ferric Fe(III)-heme resting state. Thus, this is similar to the H<sub>2</sub>O<sub>2</sub> pathway with reactions



**Fig. 2** Left and middle: Substrate-free and substrate-bound structures directly derived from four crystal structures of *LsAA9A*. Right: Structures with pre-bound  $\text{H}_2\text{O}_2$  (see products of reactions (5a) and (5b) in Fig. 1) derived from replacing water or chloride in the two crystal structures of *LsAA9A* with oxidation state I. Blue and green boxes correspond to the ones denoted correspondingly in Fig. 1. Description of the geometry: the active site of LPMOs consists of a copper center coordinated by two histidines (His1 and His78), forming the histidine brace.<sup>1,10,13–17</sup> His1 is the N-terminal residue that binds bidentate through the amine of the N-terminus and the imidazole side chain. His78 binds only through the side chain. In LPMOs of fungal origin, His1 is often methylated (this is also the case for *LsAA9A*). Apart from the histidine brace, there are no residues that are strictly conserved across all LPMOs, however, many LPMOs have a generally conserved tyrosine (Tyr164 in *LsAA9A*) as axial ligand in the oxidation state II. Two water ligands (one in axial and one in equatorial position) complete the octahedral Jahn–Teller distorted active site with axially elongated ligands. In the second coordination sphere, we included Gln162 and His147, which are both important in stabilizing the pre-bound  $\text{H}_2\text{O}_2$  structures.<sup>40–42</sup> If cellobiotetraose is present, it binds from subsite –2 to +2, and forms a hydrogen bond to the so-called “pocket water”<sup>43</sup> that in turn hydrogen bonds to the terminal  $\text{NH}_2$  group of His1. Geometrical changes upon reduction and substrate binding are described in the main text and in detail in ref. 44.

(5a)/(5b) and (6a)/(6b) in Fig. 1, with the important difference that it is the oxidized form of the heme unit that interacts with  $\text{H}_2\text{O}_2$ , and no reduction is required for  $\text{H}_2\text{O}_2$  activation. Instead, the binding of  $\text{H}_2\text{O}_2$  generates the  $\text{Fe(III)-OOH}$  intermediate (after deprotonation of  $\text{H}_2\text{O}_2$ ).<sup>45</sup> The P450 cytochromes can use a similar mechanism known as the “peroxide shunt” pathway, but this generally leads to a one-electron reduced form of Cpd1, denoted Cpd2, and a protein radical.<sup>38</sup>

In general, LPMO reactions with  $\text{O}_2$  and  $\text{H}_2\text{O}_2$  have been intensively studied experimentally and computationally.<sup>10,12,13,43,46</sup> Unlike the iron-heme proteins, computational studies focused mainly on events after binding of the co-substrate,<sup>18,19,47,48</sup> whereas less attention has been paid to the priming reduction,<sup>49,50</sup> substrate binding and the events before the co-substrate interacts with  $\text{Cu(I)}$ . However, a recent combined experimental/theoretical study<sup>51</sup> by Lim *et al.* interrogated the  $\text{Cu(I)}$  species of an AA9 LPMO through core

spectroscopy and density functional theory (DFT) calculations: they found the  $\text{Cu(I)}$  species to have a  $d_{x^2-y^2}$  frontier orbital that was proposed to be optimally oriented to react with an incoming co-substrate ( $\text{H}_2\text{O}_2$  in ref. 51). Lim *et al.* further argue that the  $d_{x^2-y^2}$ -orbital is significantly higher in energy (1.1–1.4 eV) compared to the remaining occupied copper d-orbitals, bringing it closer to the LUMO (the  $\text{H}_2\text{O}_2$   $\sigma^*$ -orbital). This facilitates the one-electron transfer from the copper to  $\text{H}_2\text{O}_2$  in reaction (6b) in Fig. 1.<sup>51</sup> Yet, this investigation did not explicitly include a saccharide substrate. Meanwhile, other studies have suggested that substrate binding has specific impacts on LPMO reactivity: the nature of the substrate has been proposed to take part in determining the mechanism for the generation of the oxidative species<sup>48,52–54</sup> and substrate binding has been proposed to directly influence the orbitals of d-character.<sup>9</sup> Further, several electron paramagnetic resonance (EPR) studies on AA9 and

AA10 LPMOs report the appearance of super-hyperfine couplings and changes in the EPR spectra upon substrate binding.<sup>21,55–57</sup> Different explanations for these observations were given. For example, conformational changes such as a distortion of the copper-coordinating amino acids and a reorganization of coordinating water molecules.<sup>56,57</sup> It has also been proposed that substrate-induced electronic structure changes “pre-dispose” the active site to form a stable Cu(II)-Cell-O<sub>2</sub> intermediate.<sup>56</sup> Some of the discussed electronic structure changes include, for example, a stronger interaction between the copper and the coordinating nitrogen atoms<sup>21,55</sup> and an increase in energy of the  $d_{x^2-y^2}$  frontier orbital.<sup>21,56</sup>

To reveal further details of the first steps of the LPMO reaction and the effects of reduction and substrate binding, we use a frontier orbital approach. Analysing frontier molecular orbitals (MOs) has previously been used to understand the reactivity of both iron-heme enzymes<sup>38,45</sup> and copper enzymes involving a Cu(II)-O<sub>2</sub> moiety.<sup>58–60</sup> The orbitals around the HOMO and LUMO are here taken as acceptors for incoming co-substrates. We analyse changes in the frontier MOs in the presence and absence of substrate and further how reduction and incoming O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> influence the electronic structure. We use the structures from a recent crystallographic study by Tandrup *et al.*<sup>22</sup> to directly link the frontier MOs to accurate experimental structures: In the structures obtained in ref. 22, a controlled photoreduction of the copper in the X-ray beam<sup>61</sup> was exploited to obtain structures that can be assigned Cu(I) and Cu(II) oxidation states,<sup>22,62</sup> both with and without bound cellooligosaccharide substrate (abbreviated as Cell). Thus, with outset in these crystal structures,<sup>22</sup> we investigate how the observed changes in active-site geometry due to reduction and substrate binding influence the electronic structure through analysis of the frontier orbitals. Moreover, we consider also how the frontier orbitals change due to incoming oxidative species in the forms of O<sub>2</sub> or H<sub>2</sub>O<sub>2</sub>.

## Computational details

Our calculations are anchored in the experimental geometry of four different LsAA9A crystal structures measured by Tandrup *et al.* (pdb codes 7PXI, 7PXV, 7PYI and 7PYD, see structures in blue boxes in Fig. 1).<sup>22</sup> In the structures 7PXI and 7PXV no substrate is bound, and they will be labelled as Cu(II) and Cu(I) respectively. Cellotetraose is bound in the structures 7PYD and 7PYI, and they will be labelled Cu(II)-Cell-Cl<sup>-</sup> and Cu(I)-Cell-Cl<sup>-</sup>, respectively. Note that Tandrup *et al.* obtained the Cu(I) crystal structure (pdb code 7PXV) through chemical reduction with ascorbic acid, while the Cu(I)-Cell-Cl<sup>-</sup> structure (pdb code 7PYI) was photoreduced by increasing the radiation dose.<sup>22</sup>

Since all four crystal structures contain only non-hydrogen atoms, hydrogens were added with the program Maestro<sup>63</sup> (which is part of the Schrödinger suite). From the resulting structures, we extracted the copper, the side chains of His1 and His78 (constituting the histidine brace), the side chains of His147, Gln162 and Tyr164, as well as any copper coordinating

water molecules (labelled H<sub>2</sub>O<sub>eq</sub> and H<sub>2</sub>O<sub>ax</sub>). Tyr164 is an axial ligand to the copper in AA9 and other LPMOs, while His147 and Gln162 are important residues in the second coordination sphere (see top left panel in Fig. 2). His1 is methylated<sup>1</sup> and the methylation was always kept. His147 was always protonated in the N<sup>ε2</sup> position.<sup>64</sup> From the crystal structures with the substrate, three units of the cellotetraose, the chloride anion and the so-called pocket water<sup>21</sup> (labelled H<sub>2</sub>O<sub>pkt</sub>) were extracted additionally. The latter binds to the terminal NH<sub>2</sub> group as described in ref. 21. In the Cu(II) and Cu(I) structures without substrate, we further included a water molecule in a similar position to the pocket water (also labelled H<sub>2</sub>O<sub>pkt</sub>). In all extracted structures, His78, His147, Gln162 and Tyr164 were cut between C<sup>α</sup> and C<sup>β</sup>, while the N-terminal coordinating His1 was cut between the backbone carbon atom (C) and C<sup>α</sup>. Except for His1, only the side chain atoms of the residues were kept, and dangling bonds were saturated with hydrogen atoms. For the structures with cellotetraose, the three saccharide units closest to the copper centre were extracted, and the glycosidic bond was cut between the oxygen and the fourth saccharide unit. The resulting systems are shown in Fig. 2 and Fig. S8† (with optimized hydrogen positions, see below).

The hydrogen positions were optimized with the quantum chemistry software ORCA 5.0.1<sup>65</sup> using the DFT functional TPSS<sup>66</sup> and the def2-SV(P) basis set<sup>67</sup> with D3 dispersion correction,<sup>68</sup> Becke–Johnson damping<sup>69</sup> and the resolution of identity (RI) approximation.<sup>70</sup> The auxiliary basis set corresponding to def2-SV(P) was used for the RI approximation (def2/J). The positions of all non-hydrogen atoms were fixed at their crystal structure positions. All energies, orbitals, orbital energies and spin densities were obtained from single-point calculations on these structures with the DFT functionals B3LYP<sup>71–73</sup> and a def2-TZVPP<sup>67</sup> basis set. We use the default version of B3LYP in ORCA, where the calculation of Coulomb (*J*) and exchange (*X*) terms is done with the split RI-*J* and chain-of-spheres for exchange (RIJCOSX) approximation.<sup>74</sup> We additionally calculated the orbital energies with the TPSS<sup>66</sup> functional. Since B3LYP and TPSS generally gave similar trends, we show only the B3LYP results in the main text and refer to section S3 in the ESI† for the TPSS results.

In both substrate-bound structures (Cu(I)-Cell-Cl<sup>-</sup> and Cu(II)-Cell-Cl<sup>-</sup>) a chloride anion is present (it coordinates to copper in the Cu(II)-Cell-Cl<sup>-</sup> structure). The results for the structures with chloride are discussed in section S2 in the ESI.† We substituted chloride with water, keeping the oxygen atom fixed at the position of the chloride atom. The structures with water are labelled Cu(I)-Cell and Cu(II)-Cell (see Fig. 2) and their hydrogen positions were optimized as described above.

We further investigated the effect of H<sub>2</sub>O<sub>2</sub> binding on the electronic structure. H<sub>2</sub>O<sub>2</sub> was added to Cu(I) and Cu(I)-Cell-Cl<sup>-</sup>, replacing either the equatorial<sup>9,18,75</sup> water or chloride (note that with 4.03 Å and 3.84 Å respective distances to copper, neither the water nor the chloride is in coordination distance). In all structures, coordinates of H<sub>2</sub>O<sub>2</sub> were freely optimized together with the hydrogen atoms, employing the

procedure described above. The Cu(I)-Cell...H<sub>2</sub>O<sub>2</sub> structure is shown in Fig. 2. Instead of obtaining a pre-bound Cu(I)...H<sub>2</sub>O<sub>2</sub> (*i.e.* the structure without substrate) with the procedure described above, we could only obtain a structure reminiscent of the “caged” state (indicated as Cu(II)-HO-OH in Fig. 1), where H<sub>2</sub>O<sub>2</sub> coordinates to copper and the O-O distance is elongated to distances of approximately 2 or more ångström.<sup>18–20</sup> The pre-bound form has previously been calculated with both TPSS and B3LYP using a QM/MM embedding scheme to mimic the protein environment<sup>18–20</sup> and it was also obtained by Lim *et al.*<sup>51</sup> who employed a QM-cluster approach. Therefore, we optimized the Cu(I)...H<sub>2</sub>O<sub>2</sub> structure with the same functional (B3LYP) that Lim *et al.*<sup>51</sup> used to obtain the pre-bound state, and obtained a similar orientation for H<sub>2</sub>O<sub>2</sub> (see Fig. 2). Yet, the Cu-H<sub>2</sub>O<sub>2</sub> distance is smaller in our Cu(I)...H<sub>2</sub>O<sub>2</sub> structure (2.54 Å) compared to the structure by Lim *et al.*<sup>51</sup> (2.93 Å), which could result from the fact that we kept the protein non-hydrogen atoms fixed at their crystal structure positions, while Lim *et al.*<sup>51</sup> optimized the complete model.

Prompted by a previous investigation<sup>56</sup> of the influence of the potential coupling of the substrate and the binding of O<sub>2</sub>, we further investigated the binding of O<sub>2</sub> to substrate-free and substrate-bound structures (resulting in the structures in green boxes in the left part of Fig. 1, see Fig. S1†).

We mainly use a frontier MO approach to evaluate electronic structure changes. The characters of the MOs were estimated by Löwdin reduced orbital populations; they provide a percentage for the contribution of the atomic orbitals to each molecular orbital. From a computational perspective, Cu(I) is in a closed-shell singlet state, where all electrons are paired, *i.e.*  $\alpha$ - and  $\beta$ -electrons within a given spatial orbital have the same energy. This is not the case for intermediates with Cu(II) oxidation state (or Cu(I) with O<sub>2</sub>), which are open-shell (doublet or triplet) states. In these states,  $\alpha$ - and  $\beta$ -orbitals may differ in energy, and we refer separately to the  $\alpha$  and  $\beta$  frontier orbitals for these calculations. Structures, individual orbitals and spin densities were plotted using PYMOL<sup>76</sup> and the ORCA\_PLOT tool. Atomic spin populations were calculated using Löwdin partitioning and quantum theory of atoms in molecules (QTAIM) atomic charges were calculated with AIMAll.<sup>77</sup>

## Results & discussion

### Effect of the priming electron on frontier orbitals

We start by investigating the first step of the LPMO mechanism, namely the reduction of the copper centre in the structures without substrate (reaction (1a) in Fig. 1). Upon reduction from Cu(II) to Cu(I), Tandrup *et al.* observe that the copper loses the equatorial and axial water molecules, thereby changing from elongated hexacoordinated to T-shaped geometry. In the T-shaped form, Cu(I) is only coordinated by the nitrogen atoms of the protein ligands. In the here studied structures without substrate (see Cu(II) and Cu(I) structures in Fig. 2 and Table 1), the Cu-O distances increase from 2.2 and 2.7 Å to 4.0 and 3.5 Å for the equatorial and axial water molecules, respectively, upon reduction.<sup>22</sup>

The orbital energy diagrams of Cu(II) and Cu(I) are shown in Fig. 3 for B3LYP (and in Fig. S10† for TPSS). We focus first on the diagrams without substrate, *i.e.* the ones labelled Cu(II) and Cu(I). Concerning the frontier orbitals in the Cu(II) structure, both the  $\alpha$ - and  $\beta$ -HOMOs have the character of tyrosine orbitals (see also Table S7†), and the (occupied) frontier orbitals are all of ligand character. The occupied orbitals of copper d-character are considerably (2.2–8.2 eV) lower in energy than the HOMO. The fact that the orbitals of d-character are removed from the frontier means that they are unlikely to take part in a reaction with an incoming co-substrate. On the other hand, the  $\beta$ -LUMO is mostly (to 62%) a copper d-orbital, namely the  $d_{x^2-y^2}$ -orbital (see Fig. 4), making it the most likely orbital to accept an electron from a reductant.


Moving to the orbital energies of the Cu(I) structure (see Fig. 3), the frontier MOs are dominated by orbitals of copper d-character. HOMO-1, HOMO-7, and HOMO-9 to HOMO-11 are mainly orbitals of copper d-character (40–90%) with some smaller contributions from the surrounding ligands (see Table S8†).

Hence, the priming electron brings the occupied orbitals of copper d-character to the frontier, where they are available for incoming small molecules. This provides a simple frontier-orbital-based explanation for the “priming reduction” requirement for activity.<sup>13,33,34,43</sup>

**Table 1** Selected distances for the structures shown in Fig. 2. Distances with only one atom are between the indicated atom and copper. For the equatorial and axial water ligands, the distance to oxygen is reported. In the case of two oxygen atoms in H<sub>2</sub>O<sub>2</sub>, only the distance to the oxygen closer to copper is reported, while the distance between the two oxygen atoms is given in the last column. A more extensive list of distances and angles for all investigated structures is given in the ESI (Table S1†)

Structure	PDB entry	N <sup>S1</sup> (Å)	N <sup>ter</sup> (Å)	N <sup>e2</sup> (Å)	O <sub>Tyr</sub> (Å)	Equatorial ligand (Å)	Axial ligand (Å)	O-O (Å)
Cu(II)	7pxi	1.87	2.18	1.96	2.74	2.16	2.71	n/a
Cu(I)	7pxv	1.84	2.29	1.99	2.81	4.03	3.49	n/a
Cu(I)...H <sub>2</sub> O <sub>2</sub>	7pxv <sup>a</sup>					2.54		1.50
Cu(II)-Cell	7pyd <sup>b</sup>	2.00	2.26	2.02	2.48	2.25	n/a	n/a
Cu(I)-Cell	7pyi <sup>b</sup>	1.89	2.50	2.00	2.69	3.84	n/a	n/a
Cu(I)-Cell...H <sub>2</sub> O <sub>2</sub>	7pyi <sup>c</sup>					2.39		1.72

<sup>a</sup> The equatorial water (not in coordinating distance) is replaced with H<sub>2</sub>O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> is optimized. <sup>b</sup> Chloride is replaced with H<sub>2</sub>O and oxygen is constrained to the position of chloride. <sup>c</sup> Chloride is replaced with H<sub>2</sub>O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> is optimized.



**Fig. 3** Normalized orbital energies and Löwdin reduced orbital populations per molecular orbital for Cu(II) and Cu(II)-Cell (left panels) as well as Cu(I) and Cu(I)-Cell structures (right panels), obtained from B3LYP/def2-TZVPP calculations (and directly derived from the crystal structures, see blue boxes Fig. 1). The colour indicates the residue with the largest Löwdin reduced orbital population (see legend). Löwdin reduced orbital populations are reported if the population of the copper d-orbitals is larger than 20%. Only these orbitals are shown in the top panels, whereas the bottom panels include all orbitals. The orbital energies are normalized to the HOMO energy.

### Comparison of substrate-free and substrate-bound systems

Having established the effect of the priming electron on the frontier orbitals, we next analyse the influence of substrate binding (corresponding to reactions (2a) and (2b) in Fig. 1), comparing the Cu(II) and Cu(II)-Cell structures as well as the Cu(I) and Cu(I)-Cell structures shown in Fig. 2.

Tandrup *et al.* report two correlated changes to the active site geometries upon substrate binding: first, binding of the substrates induced a loss of planarity in the copper

and the equatorial ligands that are almost coplanar in the Cu(II) structure without substrate. Second, the distance between copper and tyrosine was reduced by *ca.* 0.2 Å upon binding of the substrate to the Cu(II) structure (according to reaction (1b) in Fig. 1, see Table 1 for distances).<sup>22</sup> A third change already described by Frandsen *et al.* is that the axial water molecule is displaced by a hydroxymethyl group of the substrate (see Fig. 2).<sup>21</sup> In the following, we will describe how these structural changes influence the electronic structure.

For the individual oxidation states, the frontier orbital splittings are qualitatively similar before and after the addition of substrate (see Fig. 3 and S10†): the Cu(II)/Cu(II)-Cell systems display occupied frontier MOs mainly of ligand character for Cu(II) and ligand or substrate character for Cu(II)-Cell. The  $\beta$ -LUMOs are in both cases metal-based orbitals of d-character ( $d_{x^2-y^2}$  as depicted in Fig. 4). Thus, even if the substrate binds to the Cu(II) state, *i.e.* before reduction, the reduction of Cu(II)-Cell will likely result in the population of the same  $d_{x^2-y^2}$ -orbital as in Cu(II).

Moving to the reduced Cu(I) and Cu(I)-Cell structures, we see from Fig. 3 (and Fig. S10†) that the orbitals of copper d-character are now in both cases close to the frontier – and that the binding of the substrate does not change this fact. The Cu(I) HOMO–1 is the  $d_{x^2-y^2}$ -orbital shown in Fig. 4. Consistent with Lim *et al.*,<sup>51</sup> we observe that the  $d_{x^2-y^2}$ -orbital in the Cu(I) structure is isolated and 1.3–1.6 eV higher in energy than the other copper d-orbitals (see Table S8†). However, we also observe that after the substrate binds, the copper orbitals of d-character are closer in energy (see Cu(I)-Cell in Fig. 3 and S10†): the energy difference between the  $d_{x^2-y^2}$ -orbital and the next lower copper d-orbital is reduced more than three-fold upon substrate binding, from 1.3 eV in the substrate-free Cu(I) structure to 0.4 eV in the substrate-bound Cu(I)-Cell structure. This was not seen by Lim *et al.* since they did not include a substrate. However, they did investigate the effect of the co-substrate binding (in this case H<sub>2</sub>O<sub>2</sub>) on these energy differences, and we therefore further analyse these differences in more detail in the following section.

Before explicitly including an oxidizing species, we note that the proposal with an isolated  $d_{x^2-y^2}$ -orbital high in energy was also suggested by Courtade *et al.*<sup>56</sup> They argued that substrate binding induced the relative increase of the  $d_{x^2-y^2}$ -orbital. Their study was based on EPR spectra measured for B1AA10A with and without chitin<sup>56</sup> and thus inherently based on the Cu(II) states. For the Cu(II) and Cu(II)-Cell structures, the spin density (Fig. 4) shows that for both substrate-bound and unbound states, the spin density is mostly localized on copper. The spin density closely mimics the  $d_{x^2-y^2}$ -orbital, showing that it is reasonable to interpret this orbital as the single-occupied molecular orbital (SOMO), as done by Courtade *et al.* However, the increase in relative energy of the  $d_{x^2-y^2}$ -orbital cannot be unequivocally confirmed. This may be due to large differences between the LsAA9A LPMO we study here and the AA10 LPMO studied by Courtade *et al.* or it is a consequence of the different methods employed to model the substrate (Courtade *et al.* did not include the substrate explicitly in their calculations but used a four-coordinate species to mimic a substrate-bound intermediate).

We further note that in all four structures (Cu(II), Cu(I), Cu(II)-Cell and Cu(I)-Cell), the HOMO is a tyrosine orbital (see Tables S7–S10†), but its character barely changes upon substrate binding despite the shortening of the Cu–O<sub>Tyr</sub> distance and it is not involved in the reaction with the substrate (or co-substrate as we will see in the next section). Nevertheless, a small part of the spin density is redistributed to Tyr164 (invol-


ving both the oxygen atom and the  $\pi$ -system) upon substrate binding: essentially no spin density is located on tyrosine in Cu(II) (Löwdin spin population of 0.00 for tyrosine, see Table S3†), which increases to 0.02 in Cu(II)-Cell, reflecting the closer distance between copper and tyrosine in the substrate-bound structures.

In summary, the impact of substrate binding on the electronic structure is smaller compared to the impact of the reduction: the reduction brings the occupied copper d-orbitals to the frontier, and this impact of the reduction is not influenced significantly by the presence of the substrate. However, before substrate binding, there is a significant energy difference between the highest occupied copper  $d_{x^2-y^2}$ -orbital and the remaining copper d-orbitals. This difference is reduced by a factor of three when the substrate binds.

### Impact of an oxidizing species

For LPMOs in general (and particularly for LsAA9A), H<sub>2</sub>O<sub>2</sub> has shown to be the (only) co-substrate.<sup>12,23,24</sup> Therefore, we start by discussing substrate-free and substrate-bound structures with H<sub>2</sub>O<sub>2</sub> replacing the equatorial water or chloride in Cu(I) and Cu(I)-Cell-Cl<sup>–</sup>, respectively (note that neither the water nor the chloride are coordinated in these structures). We aim to obtain the same pre-bound state as Lim *et al.*<sup>51</sup> (indicated as Cu(I)⋯H<sub>2</sub>O<sub>2</sub> in Fig. 1), but in one case additionally include a substrate (Cu(I)-Cell⋯H<sub>2</sub>O<sub>2</sub> in Fig. 1). In the pre-bound state, H<sub>2</sub>O<sub>2</sub> is not coordinated to Cu(I), but instead located in the pocket between the substrate and active site, where it is stabilized by interactions with a second-sphere histidine and glutamine (see His147 and Gln162 in Fig. 2).<sup>18–20</sup> We observe that substrate binding brings H<sub>2</sub>O<sub>2</sub> closer to copper (Cu–O distance of 2.54 Å and 2.39 Å in Cu(I)⋯H<sub>2</sub>O<sub>2</sub> and Cu(I)-Cell⋯H<sub>2</sub>O<sub>2</sub>, respectively) and increases the O–O bond in H<sub>2</sub>O<sub>2</sub> from 1.50 Å to 1.72 Å (see Table 1). QM/MM calculations of the pre-bound intermediate in LsAA9A (including the substrate) obtained a Cu–O distance of 2.77–2.98 Å and an O–O distance of 1.44–1.46 Å. In the caged intermediates the electron has been transferred from copper to the  $\sigma^*$ -orbital of H<sub>2</sub>O<sub>2</sub> (reaction (6b) in Fig. 1). These intermediates are in an open-shell singlet state, and they are characterized by long O–O distances of 1.97–2.14 Å and shorter Cu–O distances of approximately 1.9 Å.<sup>18–20</sup> Hence, our Cu(I)-Cell⋯H<sub>2</sub>O<sub>2</sub> intermediate is geometrically somewhat in between a pre-bound and a caged species.


Indeed, upon binding of hydrogen peroxide, a small charge transfer occurs from copper to H<sub>2</sub>O<sub>2</sub> (see Table S4†): the positive charge on copper increases from 0.61 in Cu(I) to 0.68 in Cu(I)⋯H<sub>2</sub>O<sub>2</sub> and from 0.64 in Cu(I)-Cell to 0.77 in Cu(I)-Cell⋯H<sub>2</sub>O<sub>2</sub>. The net charge of the equatorial water (not in coordination distance) is close to neutral (–0.02 in Cu(I) and –0.01 in Cu(I)-Cell), whereas the net charge of H<sub>2</sub>O<sub>2</sub> is more negative with –0.09 in Cu(I)⋯H<sub>2</sub>O<sub>2</sub> and –0.25 in Cu(I)-Cell⋯H<sub>2</sub>O<sub>2</sub>. The larger magnitude of the charge transfer in the presence of substrate is consistent with the copper and H<sub>2</sub>O<sub>2</sub> being closer in the structures with bound substrate, and supports that Cu(I)-Cell⋯H<sub>2</sub>O<sub>2</sub> is somewhere in between a pre-bound and a caged species. Comparison with results for



**Fig. 4** Spin density (isovalue  $0.001 \text{ e bohr}^{-3}$ ) and  $d_{x^2-y^2}$ -orbitals (isovalue  $0.05 \text{ e bohr}^{-3}$ ) for the structures in blue boxes in Fig. 1, obtained from B3LYP/def2-TZVPP calculations. In both cases, green colour corresponds to positive values, and blue to negative ones. The spin density is defined as the difference in the densities contributed by the  $\alpha$  and  $\beta$ -electrons. Shown is the LUMO for Cu(II) and Cu(II)-Cell, as well as the highest occupied orbital of copper d-character for Cu(I) and Cu(I)-Cell, in all cases corresponding to a  $d_{x^2-y^2}$ -orbital.

amyloid  $\beta$ -peptide models coordinated with copper confirms this observation.<sup>78</sup>


Analysing the orbital energy splittings for Cu(I)⋯H<sub>2</sub>O<sub>2</sub> and Cu(I)-Cell⋯H<sub>2</sub>O<sub>2</sub> in Fig. 5 (see Fig. S11† for TPSS results), we observe a similar character and splitting of the occupied frontier orbitals as in Cu(I) and Cu(I)-Cell, with the occupied orbitals of copper d-character close to the frontier. For Cu(I)⋯H<sub>2</sub>O<sub>2</sub>, we obtain an energy difference of 1.2–1.7 eV between the highest occupied  $d_{x^2-y^2}$ -orbital and the remaining MOs of copper d-character (see Table S11†), which is consistent with the difference of 1.1–1.4 eV reported by Lim *et al.*<sup>51</sup> We also find that the  $d_{x^2-y^2}$ -orbital (see Fig. 6) is the HOMO in Cu(I)⋯H<sub>2</sub>O<sub>2</sub>. When the substrate binds, the HOMO is a tyrosine orbital but the  $d_{x^2-y^2}$ -orbital (see Fig. 6) remains the highest-lying orbital with d-character, while the LUMO is always an



**Fig. 5** Normalized orbital energies and Löwdin reduced orbital populations per molecular orbital for substrate-free and substrate-bound structures with pre-bound H<sub>2</sub>O<sub>2</sub>, obtained from B3LYP/def2-TZVPP calculations (and obtained from adding H<sub>2</sub>O<sub>2</sub> to the crystal structures, see right green boxes in Fig. 1). The colour indicates the residue with the largest Löwdin reduced orbital population (see legend). Löwdin reduced orbital population are reported if the population of the copper d-orbitals (here only numbers are given) and H<sub>2</sub>O<sub>2</sub> orbitals is larger than 20%. Only these orbitals are shown in the top panels, whereas the bottom panels include all orbitals. The orbital energies are normalized to the HOMO energy.

orbital of co-substrate (H<sub>2</sub>O<sub>2</sub>) character. Each LUMO is shown in Fig. 6 and has the shape of a  $\sigma^*$ -orbital. This agrees well with the LUMO described by Lim *et al.*<sup>51</sup> However, as in the Cu(I) systems, we observe that substrate binding brings the remaining orbitals of copper d-character closer to the  $d_{x^2-y^2}$ -orbital, decreasing the energy difference between them to 0.8–1.8 eV. Additionally, the substrate lowers the energy difference between the  $d_{x^2-y^2}$ -orbital and the H<sub>2</sub>O<sub>2</sub>  $\sigma^*$ -orbital (from 3.9 eV in Cu(I)⋯H<sub>2</sub>O<sub>2</sub> to 3.1 eV in Cu(I)-Cell⋯H<sub>2</sub>O<sub>2</sub>, see Tables S11 and S12†), making the electron transfer (reaction (6b) in Fig. 1) more favourable with the bound substrate. This is in line with QM/MM and QM-cluster calculations that show that





**Fig. 6**  $d_{x^2-y^2}$ -orbitals and LUMO (isovalue 0.05 e bohr<sup>-3</sup>) for the structures with pre-bound H<sub>2</sub>O<sub>2</sub> obtained from B3LYP/def2-TZVPP calculations (see Cu(I)...H<sub>2</sub>O<sub>2</sub> and Cu(I)-Cell...H<sub>2</sub>O<sub>2</sub> structures in Fig. 1). Green colour corresponds to positive values, and blue to negative ones.

this reaction is overall favourable in *LsAA9A* with and without substrate.<sup>19,20,51</sup>

For comparison with older LPMO papers and to investigate if the conclusions for H<sub>2</sub>O<sub>2</sub> were independent of the co-substrate, we additionally carried out the same experiment with O<sub>2</sub> (see section S1 in the ESI† for further details). However, since several independent studies<sup>23,24</sup> showed that the former is not a co-substrate of *LsAA9A*, we will discuss these results in the ESI.†

## Conclusions

Starting from four different crystal structures of the LPMO *LsAA9A*, we investigated the influence of reduction and substrate binding on the electronic structure of LPMOs.

Our results show that reduction from Cu(II) to Cu(I) has a significant influence on the orbital energies, bringing the occupied orbitals of copper d-character close to the frontier, where they can engage in a reaction. This is independent of the presence or absence of substrate binding and explains why the unique priming reduction is required for LPMO activity.

Further, we observed a similar orbital splitting in Cu(I) as a previous study<sup>51</sup> in pre-bound Cu(I)...H<sub>2</sub>O<sub>2</sub>, where the highest occupied copper d-orbital (the  $d_{x^2-y^2}$ ) is raised by more than 1 eV in energy compared to the other occupied copper d-orbitals. Yet, this energy difference decreases by a factor of three upon substrate binding. Previous studies suggest that a relative rise in energy of the  $d_{x^2-y^2}$ -orbital could be favourable for LPMO reactions, bringing the  $d_{x^2-y^2}$ -orbital closer to the anti-

bonding  $\sigma^*$ - or  $\pi^*$ -orbitals in hydrogen peroxide or dioxygen, respectively.<sup>51,56</sup> Our observation seems to contradict this suggestion. However, the relative energy of the anti-bonding orbitals should also be considered as well as substrate binding. Therefore, we additionally investigated how pre-bound H<sub>2</sub>O<sub>2</sub> (and O<sub>2</sub>) impact the electronic structure of substrate-free and substrate-bound *LsAA9A*. For the substrate-free H<sub>2</sub>O<sub>2</sub> structure, we find that the  $d_{x^2-y^2}$ -orbital is more than 1 eV higher in energy than the lower-lying copper d-orbitals, which is consistent with previous results.<sup>51</sup> This energy difference is decreased by one-third upon substrate binding, which is consistent with our findings without co-substrate. However, substrate binding additionally causes the H<sub>2</sub>O<sub>2</sub>  $\sigma^*$ -orbital to be lower in energy, bringing it closer to the copper  $d_{x^2-y^2}$ -orbital. Based on our calculations, it is this lowering of the H<sub>2</sub>O<sub>2</sub>  $\sigma^*$ -orbital upon substrate binding that facilitates the electron transfer from copper to H<sub>2</sub>O<sub>2</sub>. This does not occur in calculations with O<sub>2</sub> or superoxide (see ESI†).

The above trends are independent of the employed functionals (B3LYP and TPSS). To further generalize our findings, and to compare to EPR data of AA10 LPMOs, substrate-bound structures of this and other LPMO families would be a valuable starting point.

Our findings have significant biological significance, as they explain why the reduction plays an important role in activating *LsAA9A*, independently of the presence or absence of substrate. Rather, the substrate binding has a more subtle impact on the electronic structure, facilitating the reaction between LPMO and H<sub>2</sub>O<sub>2</sub>. From an electronic structure perspective, it could thus be more important to control the supply

of reductant and H<sub>2</sub>O<sub>2</sub> than the substrate supply in experimental settings, since substrate binding alone does not seem to activate LsAA9A.

## Author contributions

E. K. W.: conceptualization, investigation, formal analysis, validation, writing – original draft, writing – review & editing, visualization, project administration. L. L. L.: conceptualization, formal analysis, validation, writing – review & editing, supervision, funding acquisition, project administration. E. D. H.: conceptualization, formal analysis, validation, writing – review & editing, supervision, funding acquisition, project administration.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

EDH thanks The Villum Foundation, Young Investigator Program (grant no. 29412), the Swedish Research Council (grant no. 2019-04205), and Independent Research Fund Denmark (grant no. 2064-00002B) for support. LLL acknowledges funding from the Independent Research Fund Denmark (grant no. 8021-00273B). All of the computing for this project was performed on the GenomeDK cluster. The authors would like to thank GenomeDK and Aarhus University for providing computational resources and support that contributed to these research results.

## References

- R. J. Quinlan, M. D. Sweeney, L. Lo Leggio, H. Otten, J. C. Poulsen, K. S. Johansen, K. B. Krogh, C. I. Jorgensen, M. Tovborg, A. Anthonsen, T. Tryfona, C. P. Walter, P. Dupree, F. Xu, G. J. Davies and P. H. Walton, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 15079–15084.
- G. Vaaje-Kolstad, B. Westereng, S. J. Horn, Z. Liu, H. Zhai, M. Sorlie and V. G. Eijsink, *Science*, 2010, **330**, 219–222.
- P. V. Harris, D. Welner, K. C. McFarland, E. Re, J. C. Navarro Poulsen, K. Brown, R. Salbo, H. Ding, E. Vlasenko, S. Merino, F. Xu, J. Cherry, S. Larsen and L. Lo Leggio, *Biochemistry*, 2010, **49**, 3305–3316.
- V. V. Vu, W. T. Beeson, E. A. Span, E. R. Farquhar and M. A. Marletta, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 13822–13827.
- L. Lo Leggio, T. J. Simmons, J. C. Poulsen, K. E. Frandsen, G. R. Hemsworth, M. A. Stringer, P. von Freiesleben, M. Tovborg, K. S. Johansen, L. De Maria, P. V. Harris, C. L. Soong, P. Dupree, T. Tryfona, N. Lenfant, B. Henrissat, G. J. Davies and P. H. Walton, *Nat. Commun.*, 2015, **6**, 5961.
- M. Couturier, S. Ladeveze, G. Sulzenbacher, L. Ciano, M. Fanuel, C. Moreau, A. Villares, B. Cathala, F. Chaspoul, K. E. Frandsen, A. Labourel, I. Herpoel-Gimbert, S. Grisel, M. Haon, N. Lenfant, H. Rogniaux, D. Ropartz, G. J. Davies, M. N. Rosso, P. H. Walton, B. Henrissat and J. G. Berrin, *Nat. Chem. Biol.*, 2018, **14**, 306–310.
- F. Sabbadin, G. R. Hemsworth, L. Ciano, B. Henrissat, P. Dupree, T. Tryfona, R. D. S. Marques, S. T. Sweeney, K. Besser, L. Elias, G. Pesante, Y. Li, A. A. Dowle, R. Bates, L. D. Gomez, R. Simister, G. J. Davies, P. H. Walton, N. C. Bruce and S. J. McQueen-Mason, *Nat. Commun.*, 2018, **9**, 756.
- C. Filiatrault-Chastel, D. Navarro, M. Haon, S. Grisel, I. Herpoel-Gimbert, D. Chevret, M. Fanuel, B. Henrissat, S. Heiss-Blanquet, A. Margeot and J. G. Berrin, *Biotechnol. Biofuels*, 2019, **12**, 55.
- F. Sabbadin, S. Urresti, B. Henrissat, A. O. Avrova, L. R. J. Welsh, P. J. Lindley, M. Csukai, J. N. Squires, P. H. Walton, G. J. Davies, N. C. Bruce, S. C. Whisson and S. J. McQueen-Mason, *Science*, 2021, **373**, 774–779.
- J. O. Ipsen, M. Hallas-Moller, S. Brander, L. Lo Leggio and K. S. Johansen, *Biochem. Soc. Trans.*, 2021, **49**, 531–540.
- K. K. Meier, S. M. Jones, T. Kaper, H. Hansson, M. J. Koetsier, S. Karkehabadi, E. I. Solomon, M. Sandgren and B. Kelemen, *Chem. Rev.*, 2018, **118**, 2593–2635.
- Z. Forsberg, M. Sorlie, D. Petrovic, G. Courtade, F. L. Aachmann, G. Vaaje-Kolstad, B. Bissaro, A. K. Rohr and V. G. Eijsink, *Curr. Opin. Struct. Biol.*, 2019, **59**, 54–64.
- T. M. Vandhana, J. L. Reyre, D. Sushmaa, J. G. Berrin, B. Bissaro and J. Madhuprakash, *New Phytol.*, 2022, **233**, 2380–2396.
- S. Garcia-Santamarina, C. Probst, R. A. Festa, C. Ding, A. D. Smith, S. E. Conklin, S. Brander, L. N. Kinch, N. V. Grishin, K. J. Franz, P. Riggs-Gelasco, L. Lo Leggio, K. S. Johansen and D. J. Thiele, *Nat. Chem. Biol.*, 2020, **16**, 337–344.
- F. Askarian, S. Uchiyama, H. Masson, H. V. Sorensen, O. Golten, A. C. Bunaes, S. Mekasha, A. K. Rohr, E. Kommedal, J. A. Ludviksen, M. O. Arntzen, B. Schmidt, R. H. Zurich, N. M. van Sorge, V. G. H. Eijsink, U. Krengel, T. E. Mollnes, N. E. Lewis, V. Nizet and G. Vaaje-Kolstad, *Nat. Commun.*, 2021, **12**, 1230.
- K. S. Johansen, *Biochem. Soc. Trans.*, 2016, **44**, 143–149.
- G. Muller, A. Varnai, K. S. Johansen, V. G. Eijsink and S. J. Horn, *Biotechnol. Biofuels*, 2015, **8**, 187.
- E. Drula, M. L. Garron, S. Dogan, V. Lombard, B. Henrissat and N. Terrapon, *Nucleic Acids Res.*, 2022, **50**, D571–D577.
- C. H. Kjaergaard, M. F. Qayyum, S. D. Wong, F. Xu, G. R. Hemsworth, D. J. Walton, N. A. Young, G. J. Davies, P. H. Walton, K. S. Johansen, K. O. Hodgson, B. Hedman and E. I. Solomon, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 8797–8802.
- W. T. Beeson, V. V. Vu, E. A. Span, C. M. Phillips and M. A. Marletta, *Annu. Rev. Biochem.*, 2015, **84**, 923–946.

- 21 B. Bissaro and V. G. H. Eijssink, *Essays Biochem.*, 2023, **67**, 575–584.
- 22 B. Bissaro, A. K. Rohr, G. Muller, P. Chylenski, M. Skaugen, Z. Forsberg, S. J. Horn, G. Vaaje-Kolstad and V. G. H. Eijssink, *Nat. Chem. Biol.*, 2017, **13**, 1123–1128.
- 23 S. Brander, R. Tokin, J. Ø. Ipsen, P. E. Jensen, C. Hernández-Rollán, M. H. H. Nørholm, L. Lo Leggio, P. Dupree and K. S. Johansen, *ACS Catal.*, 2021, **11**, 13848–13859.
- 24 H. Chang, N. Gacias Amengual, A. Botz, L. Schwaiger, D. Kracher, S. Scheiblbrandner, F. Csarman and R. Ludwig, *Nat. Commun.*, 2022, **13**, 6258.
- 25 P. H. Walton and G. J. Davies, *J. Biol. Inorg. Chem.*, 2022, **27**, 705–713.
- 26 T. Isaksen, B. Westereng, F. L. Aachmann, J. W. Agger, D. Kracher, R. Kittl, R. Ludwig, D. Haltrich, V. G. Eijssink and S. J. Horn, *J. Biol. Chem.*, 2014, **289**, 2632–2642.
- 27 R. Kittl, D. Kracher, D. Burgstaller, D. Haltrich and R. Ludwig, *Biotechnol. Biofuels*, 2012, **5**, 79.
- 28 V. G. H. Eijssink, D. Petrovic, Z. Forsberg, S. Mekasha, A. K. Rohr, A. Varnai, B. Bissaro and G. Vaaje-Kolstad, *Biotechnol. Biofuels*, 2019, **12**, 58.
- 29 B. Bissaro, E. Kommedal, A. K. Rohr and V. G. H. Eijssink, *Nat. Commun.*, 2020, **11**, 890.
- 30 A. A. Stepnov, V. G. H. Eijssink and Z. Forsberg, *Sci. Rep.*, 2022, **12**, 6129.
- 31 A. A. Stepnov, Z. Forsberg, M. Sorlie, G. S. Nguyen, A. Wentzel, A. K. Rohr and V. G. H. Eijssink, *Biotechnol. Biofuels*, 2021, **14**, 28.
- 32 O. Golten, I. Ayuso-Fernandez, K. R. Hall, A. A. Stepnov, M. Sorlie, A. K. Rohr and V. G. H. Eijssink, *FEBS Lett.*, 2023, **597**, 1363–1374.
- 33 D. Kracher, S. Scheiblbrandner, A. K. Felice, E. Breslmayr, M. Preims, K. Ludwicka, D. Haltrich, V. G. Eijssink and R. Ludwig, *Science*, 2016, **352**, 1098–1101.
- 34 M. Frommhagen, A. H. Westphal, W. J. H. van Berkel and M. A. Kabel, *Front. Microbiol.*, 2018, **9**, 1080.
- 35 D. Kracher, M. Andlar, P. G. Furtmuller and R. Ludwig, *J. Biol. Chem.*, 2018, **293**, 1676–1687.
- 36 P. H. Walton, G. J. Davies, D. E. Diaz and J. P. Franco-Cairo, *FEBS Lett.*, 2023, **597**, 485–494.
- 37 K. D. Dubey and S. Shaik, *Acc. Chem. Res.*, 2019, **52**, 389–399.
- 38 S. Shaik, S. Cohen, Y. Wang, H. Chen, D. Kumar and W. Thiel, *Chem. Rev.*, 2010, **110**, 949–1017.
- 39 T. M. Hedison, E. Breslmayr, M. Shanmugam, K. Karnpakdee, D. J. Heyes, A. P. Green, R. Ludwig, N. S. Scrutton and D. Kracher, *FEBS J.*, 2021, **288**, 4115–4128.
- 40 M. M. Hagemann and E. D. Hedegard, *Chemistry*, 2023, **29**, e202202379.
- 41 B. Wang, E. M. Johnston, P. Li, S. Shaik, G. J. Davies, P. H. Walton and C. Rovira, *ACS Catal.*, 2018, **8**, 1346–1351.
- 42 E. D. Hedegard and U. Ryde, *Chem. Sci.*, 2018, **9**, 3866–3880.
- 43 K. E. Frandsen, T. J. Simmons, P. Dupree, J. C. Poulsen, G. R. Hemsworth, L. Ciano, E. M. Johnston, M. Tovborg, K. S. Johansen, P. von Freiesleben, L. Marmuse, S. Fort, S. Cottaz, H. Driguez, B. Henrissat, N. Lenfant, F. Tuna, A. Baldansuren, G. J. Davies, L. Lo Leggio and P. H. Walton, *Nat. Chem. Biol.*, 2016, **12**, 298–303.
- 44 T. Tandrup, S. J. Muderspach, S. Banerjee, G. Santoni, J. O. Ipsen, C. Hernandez-Rollan, M. H. H. Norholm, K. S. Johansen, F. Meilleur and L. Lo Leggio, *IUCrJ*, 2022, **9**, 666–681.
- 45 X. Huang and J. T. Groves, *Chem. Rev.*, 2018, **118**, 2491–2553.
- 46 P. Chylenski, B. Bissaro, M. Sorlie, A. K. Rohr, A. Varnai, S. J. Horn and V. G. H. Eijssink, *ACS Catal.*, 2019, **9**, 4970–4991.
- 47 O. Caldararu, E. Oksanen, U. Ryde and E. D. Hedegard, *Chem. Sci.*, 2019, **10**, 576–586.
- 48 B. Wang, Z. Wang, G. J. Davies, P. H. Walton and C. Rovira, *ACS Catal.*, 2020, **10**, 12760–12769.
- 49 C. Laurent, E. Breslmayr, D. Tunega, R. Ludwig and C. Oostenbrink, *Biochemistry*, 2019, **58**, 1226–1235.
- 50 Z. Wang, S. Feng, C. Rovira and B. Wang, *Angew. Chem., Int. Ed.*, 2021, **60**, 2385–2392.
- 51 H. Lim, M. T. Brueggemeyer, W. J. Transue, K. K. Meier, S. M. Jones, T. Kroll, D. Sokaras, B. Kelemen, B. Hedman, K. O. Hodgson and E. I. Solomon, *J. Am. Chem. Soc.*, 2023, **145**, 16015–16025.
- 52 A. Paradisi, E. M. Johnston, M. Tovborg, C. R. Nicoll, L. Ciano, A. Dowle, J. McMaster, Y. Hancock, G. J. Davies and P. H. Walton, *J. Am. Chem. Soc.*, 2019, **141**, 18585–18599.
- 53 T. J. Simmons, K. E. H. Frandsen, L. Ciano, T. Tryfona, N. Lenfant, J. C. Poulsen, L. F. L. Wilson, T. Tandrup, M. Tovborg, K. Schnorr, K. S. Johansen, B. Henrissat, P. H. Walton, L. Lo Leggio and P. Dupree, *Nat. Commun.*, 2017, **8**, 1064.
- 54 L. Bertini, R. Breglia, M. Lambrughini, P. Fantucci, L. De Gioia, M. Borsari, M. Sola, C. A. Bortolotti and M. Bruschi, *Inorg. Chem.*, 2018, **57**, 86–97.
- 55 A. S. Borisova, T. Isaksen, M. Dimarogona, A. A. Kognole, G. Mathiesen, A. Varnai, A. K. Rohr, C. M. Payne, M. Sorlie, M. Sandgren and V. G. Eijssink, *J. Biol. Chem.*, 2015, **290**, 22955–22969.
- 56 G. Courtade, L. Ciano, A. Paradisi, P. J. Lindley, Z. Forsberg, M. Sorlie, R. Wimmer, G. J. Davies, V. G. H. Eijssink, P. H. Walton and F. L. Aachmann, *Proc. Natl. Acad. Sci. U. S. A.*, 2020, **117**, 19178–19189.
- 57 B. Bissaro, I. Isaksen, G. Vaaje-Kolstad, V. G. H. Eijssink and A. K. Rohr, *Biochemistry*, 2018, **57**, 1893–1906.
- 58 P. Chen and E. I. Solomon, *J. Inorg. Biochem.*, 2002, **88**, 368–374.
- 59 R. E. Cowley, J. Cirera, M. F. Qayyum, D. Rokhsana, B. Hedman, K. O. Hodgson, D. M. Dooley and E. I. Solomon, *J. Am. Chem. Soc.*, 2016, **138**, 13219–13229.
- 60 E. I. Solomon, D. E. Heppner, E. M. Johnston, J. W. Ginsbach, J. Cirera, M. Qayyum, M. T. Kieber-

- Emmons, C. H. Kjaergaard, R. G. Hadt and L. Tian, *Chem. Rev.*, 2014, **114**, 3659–3853.
- 61 S. E. Bowman, J. Bridwell-Rabb and C. L. Drennan, *Acc. Chem. Res.*, 2016, **49**, 695–702.
- 62 M. Gudmundsson, S. Kim, M. Wu, T. Ishida, M. H. Momeni, G. Vaaje-Kolstad, D. Lundberg, A. Royant, J. Stahlberg, V. G. Eijssink, G. T. Beckham and M. Sandgren, *J. Biol. Chem.*, 2014, **289**, 18782–18792.
- 63 Schrödinger Release 2021-2, *Maestro*, Schrödinger, LLC, New York, NY, 2021.
- 64 S. Banerjee, S. J. Muderspach, T. Tandrup, K. E. H. Frandsen, R. K. Singh, J. O. Ipsen, C. Hernandez-Rollan, M. H. H. Norholm, M. J. Bjerrum, K. S. Johansen and L. Lo Leggio, *Biomolecules*, 2022, **12**, 194.
- 65 F. Neese, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2022, **12**, e1606.
- 66 J. Tao, J. P. Perdew, V. N. Staroverov and G. E. Scuseria, *Phys. Rev. Lett.*, 2003, **91**, 146401.
- 67 F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305.
- 68 S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- 69 S. Grimme, S. Ehrlich and L. Goerigk, *J. Comput. Chem.*, 2011, **32**, 1456–1465.
- 70 F. Weigend, *Phys. Chem. Chem. Phys.*, 2006, **8**, 1057–1065.
- 71 A. D. Becke, *Phys. Rev. A: Gen. Phys.*, 1988, **38**, 3098–3100.
- 72 C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 1988, **37**, 785–789.
- 73 A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648–5652.
- 74 F. Neese, F. Wennmohs, A. Hansen and U. Becker, *Chem. Phys.*, 2009, **356**, 98–109.
- 75 E. D. Hedegard and U. Ryde, *ACS Omega*, 2017, **2**, 536–545.
- 76 Schrödinger, *The PyMOL Molecular Graphics System, Version 2.6*, 2015.
- 77 T. A. Keith, *TK Gristmill Software*, Overland Park KS, USA, 2019.
- 78 T. Prosdocimi, L. De Gioia, G. Zampella and L. Bertini, *J. Biol. Inorg. Chem.*, 2016, **21**, 197–212.