

# Faraday Discussions

Accepted Manuscript



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

This article can be cited before page numbers have been issued, to do this please use: J. M. M. Sischka, I. Nitschke and A. Voigt, *Faraday Discuss.*, 2024, DOI: 10.1039/D4FD00202D.

Cite this: DOI: 00.0000/xxxxxxxxxx

# The influence of higher order geometric terms on the asymmetry and dynamics of membranes

Jan Magnus Sischka,<sup>a</sup> Ingo Nitschke,<sup>a</sup> and Axel Voigt<sup>\*a,b,c</sup>Received Date  
Accepted Date

DOI: 00.0000/xxxxxxxxxx

We consider membranes as fluid deformable surfaces and allow for higher order geometric terms in the bending energy related to the Gaussian curvature squared and the mean curvature minus the spontaneous curvature to the fourth power. The evolution equations are derived and numerically solved using surface finite elements. The two higher order geometric terms have different effects. While the Gaussian curvature squared term has a tendency to stabilize tubes and enhance the evolution towards equilibrium shapes, thereby facilitating rapid shape changes, the mean curvature minus the spontaneous curvature to the fourth power destabilizes tubes and leads to qualitatively different equilibrium shapes but also enhances the evolution. This is demonstrated in axisymmetric settings and fully three-dimensional simulations. We therefore postulate that not only surface viscosity but also higher order geometric terms in the bending energy contribute to rapid shape changes which are relevant for morphological changes of cells.

## 1 Introduction

Membranes are ubiquitous and essential in biology, they compartmentalize biomaterials, separate the cell from its exterior and organelles from the cytoplasm, dynamically remodel and change conformation. Geometric properties of the membrane have been identified as key players for such processes<sup>1-3</sup>. As the typical thickness of a membrane is orders of magnitude smaller than its lateral extension, treating the membrane as a two-dimensional surface embedded in a three-dimensional space is a reasonable approximation. This separation of length scales allows for a mesoscopic modeling where details related to membrane molecular structure are considered in effective material parameters and geometric quantities and led to the success of the classical Canham-Helfrich model<sup>4,5</sup>, which builds on a bending energy  $\mathcal{F}_{BE}(\mathcal{H}, \mathcal{K}) = \int_{\mathcal{S}} k_{2,0}(\mathcal{H} - \mathcal{H}_0)^2 + k_{2,1} \mathcal{K} d\mathcal{S}$  with mean curvature  $\mathcal{H}$ , Gaussian curvature  $\mathcal{K}$ , bending rigidity parameters  $k_{2,0}$  and  $k_{2,1}$ , a spontaneous curvature  $\mathcal{H}_0$  and additional (local or global) area and volume constraints. For definitions see Section 2.1. Assuming constant values of  $k_{2,1}$ , the second term reduces to a topological measure and thus a constant as long as the topology does not change. We will therefore neglect this term.

Equilibrium shapes, resulting from minimizing the bending en-

ergy  $\mathcal{F}_{BE}$ , have been extensively studied, see<sup>6</sup> for a review. For  $\mathcal{H}_0 = 0$  and wide ranges of the reduced volume, which is the ratio of the volume and the volume of an equivalent sphere with the same area, they are dominated by prolate and oblate shapes<sup>7</sup>. The spontaneous curvature  $\mathcal{H}_0$ , which accounts for the asymmetry of the membrane is able to modify these shapes<sup>7</sup>. A wealth of studies also aims to produce tubes as minimizing shapes. Tubes are ubiquitous in membranes and play crucial roles in trafficking, ion transport, and cellular motility. For idealized situations this is rather simple as  $\mathcal{F}_{BE} = 0$  if  $\mathcal{H} = \mathcal{H}_0$ , which is achieved for an infinite tube of radius  $r$  and  $\mathcal{H}_0 = -1/r$ . However, this solution is not unique, a sphere of radius  $2r$  also leads to  $\mathcal{F}_{BE} = 0$ . More realistic situations with finite volume and area require further considerations, e.g., introducing a spontaneous curvature deviator<sup>8-10</sup>. Also various ideas have been proposed to consider higher order geometric terms in the bending energy  $\mathcal{F}_{BE}$  to enforce the stability of tubes<sup>2,11-14</sup>. E.g., fourth order terms proportional to  $\mathcal{K}^2$  seem plausible as  $\mathcal{K} = 0$  for tubes.

Most of these studies only focus on equilibrium shapes, comparing the bending energy  $\mathcal{F}_{BE}$  of different configurations and addressing their stability. But also the dynamic evolution of the membrane and associated shape changes are of interest. E.g. in the context of the formation of bulges with pinch-offs, which can be associated with endocytosis and exocytosis<sup>15-17</sup> or shape changes associated with cell motility<sup>18-20</sup>. All these processes require additional phenomena, which are not considered in the Canham-Helfrich model. But there is growing interest in the role of membranes in these processes. One striking example, where membranes at least participate are frequently forming and retracting filopodia<sup>21</sup>. This process requires rapid shape changes in

<sup>a</sup> Faculty Mathematics, TU Dresden, 01062 Dresden, Germany.

<sup>b</sup> Center for Systems Biology Dresden (CSBD), Pfotenhauerstr. 108, 01307 Dresden, Germany

<sup>c</sup> Cluster of Excellence, Physics of Life, TU Dresden, Arnoldstr. 18, 01307 Dresden, Germany.

† Supplementary Information available: Corresponding videos to shape evolutions shown in Figs. 5 (a) - (c) and 6 (a) - (c). See DOI: 00.0000/00000000.



cells and according to<sup>22</sup> the membrane and the underlying cortex act as an integrated system to globally coordinate such changes in cell shape. To facilitate these rapid morphological changes, cells maintain an excess of membrane that is organized in membrane reservoirs and is available to the cell in the order of seconds<sup>23</sup>. To understand such processes thus not only requires to unveil the secrets of equilibrium shapes but also to consider the flow of membranes which facilitates the rapid shape changes. To model such processes is a different story and even if various models for the cellular cortex exist<sup>24–27</sup> and also first attempts to couple them with membrane models to form the mentioned integrated system have been proposed<sup>22,28</sup>, we here refrain from such couplings and only aim first for a minimal model of the membrane alone, which facilitates rapid shape changes.

Surface viscosity has been identified as a key player<sup>29</sup> and considering membranes as fluid deformable surfaces<sup>30–33</sup> opened new perspectives on the description of the dynamics of membranes. Fluid deformable surfaces can be viewed as two-dimensional viscous fluids with bending elasticity. Due to this solid-fluid duality, any shape change contributes to tangential flow, and vice versa, any tangential flow on a curved surface induces shape deformations. This tight coupling between shape and flow makes curvature a natural element of the governing equations. As demonstrated by numerical studies of the equations for fluid deformable surfaces, surface hydrodynamics can significantly speed up the evolution<sup>34,35</sup> and can enhance bulging and furrow formation in membranes<sup>17</sup>.

Combined with higher order geometric terms in the bending energy models for fluid deformable surfaces have the potential to enable the rapid shape changes in the release and formation of membrane reservoirs and the formation and retraction of filopodia. We computationally explore the effect of these terms on the equilibrium shapes and the dynamics to reach them and analyze their impact to facilitate rapid shape changes. Any coupling with the cortex<sup>22</sup>, interaction with proteins that induce curvature<sup>36</sup> or to forces exerted on the membrane<sup>37,38</sup> are not considered. We also neglect any interaction with the surrounding bulk phases. This results from the theoretical interest to explore the membrane properties without any additional influence and the limit of a large Saffman–Delbrück number. This number describes the relation between the viscosities of the membrane and the typically less viscous bulk fluid and if large allows the decoupling of surface and bulk flows<sup>39</sup>.

The remaining of the paper is structured as follows. In Section 2 we introduce the full model and briefly mention the considered numerical approach. More details on the derivation of the model and on the numerical approach including convergence studies are provided in Appendices 5.1, 5.2 and 5.3. Computational results are described in Section 3. They contain axisymmetric and full three-dimensional computations addressing the dynamic evolution and the equilibrium shapes. In Section 4 we draw conclusions and mention possible directions to extend the described model to enable simulations of rapid morphological changes of cells.

## 2 Model

### 2.1 Notation

The considered mesoscale model requires basic notation from differential geometry and geometric partial differential equations. Besides classical mathematical text books in these fields we refer to<sup>2</sup> for an introduction in the context of membranes. We follow the same notation as in<sup>17</sup>, which is here repeated for convenience. We consider a time dependent smooth and oriented surface  $\mathcal{S} = \mathcal{S}(t)$  without boundary, embedded in  $\mathbb{R}^3$ . The enclosed volume is denoted by  $\Omega = \Omega(t)$ . We denote by  $\mathbf{v}$  the outward pointing surface normal, the surface projection is  $\mathbf{P} = \mathbf{I} - \mathbf{v} \otimes \mathbf{v}$ , with  $\mathbf{I}$  the identity matrix, the shape operator is  $\mathcal{B} = -\nabla \mathbf{P} \mathbf{v}$ , the mean curvature  $\mathcal{H} = \text{tr} \mathcal{B}$ , and the Gaussian curvature  $\mathcal{K} = \frac{1}{2} (\mathcal{H}^2 - \|\mathcal{B}\|^2)$ . We consider time-dependent Euclidean-based  $n$ -tensor fields in  $T^n \mathbb{R}^3|_{\mathcal{S}}$ . We call  $T^0 \mathbb{R}^3|_{\mathcal{S}} = T^0 \mathcal{S}$  the space of scalar fields,  $T^1 \mathbb{R}^3|_{\mathcal{S}} = T \mathbb{R}^3|_{\mathcal{S}}$  the space of vector fields, and  $T^2 \mathbb{R}^3|_{\mathcal{S}}$  the space of 2-tensor fields. Important subtensor fields are tangential  $n$ -tensor fields in  $T^n \mathcal{S} \leq T^n \mathbb{R}^3|_{\mathcal{S}}$ . Let  $p \in T^0 \mathcal{S}$  be a continuously differentiable scalar field,  $\mathbf{u} \in T \mathbb{R}^3|_{\mathcal{S}}$  a continuously differentiable  $\mathbb{R}^3$ -vector field, and  $\boldsymbol{\sigma} \in T^2 \mathbb{R}^3|_{\mathcal{S}}$  a continuously differentiable  $\mathbb{R}^3 \times \mathbb{R}^3$ -tensor field defined on  $\mathcal{S}$ . We define the different surface gradients by  $\nabla \mathbf{P} p = \mathbf{P} \nabla p^e$ ,  $\nabla \mathbf{P} \mathbf{u} = \mathbf{P} \nabla \mathbf{u}^e \mathbf{P}$  and  $\nabla_{\mathcal{C}} \boldsymbol{\sigma} = \nabla \boldsymbol{\sigma}^e \mathbf{P}$ , where  $p^e$ ,  $\mathbf{u}^e$  and  $\boldsymbol{\sigma}^e$  are arbitrary smooth extensions of  $p$ ,  $\mathbf{u}$  and  $\boldsymbol{\sigma}$  in the normal direction and  $\nabla$  is the gradient of the embedding space  $\mathbb{R}^3$ . The corresponding divergence operators for a vector field  $\mathbf{u}$  and a tensor field  $\boldsymbol{\sigma}$  are  $\text{div} \mathbf{P} \mathbf{u} = \text{tr}(\nabla \mathbf{P} \mathbf{u})$  and  $\text{div}_{\mathcal{C}}(\boldsymbol{\sigma} \mathbf{P}) = \text{tr} \nabla_{\mathcal{C}}(\boldsymbol{\sigma} \mathbf{P})$ , where  $\text{tr}$  is the trace operator. The relations to the covariant derivatives  $\nabla_{\mathcal{S}}$  and the covariant divergence  $\text{div}_{\mathcal{S}}$  on  $\mathcal{S}$ , with  $\Delta_{\mathcal{S}} = \text{div}_{\mathcal{S}} \cdot \nabla_{\mathcal{S}}$  the Laplace-Beltrami operator, read  $\nabla \mathbf{P} p = \nabla_{\mathcal{S}} p$  and  $\text{div} \mathbf{P} \mathbf{u} = \text{div}_{\mathcal{S}}(\mathbf{P} \mathbf{u}) - (\mathbf{u} \cdot \mathbf{v}) \mathcal{H}$ , respectively.

### 2.2 Governing equations

The material velocity  $\mathbf{u} \in T \mathbb{R}^3|_{\mathcal{S}}$  can be decomposed into  $\mathbf{u} = u_N \mathbf{v} + \mathbf{u}_T$ , with  $u_N = \mathbf{u} \cdot \mathbf{v}$  and  $\mathbf{u}_T = \mathbf{P} \mathbf{u}$ , the normal and the tangential part, respectively. The pressure  $p \in T^0 \mathcal{S}$  serves as Lagrange multiplier for the inextensibility constraint. The governing equations for these unknowns read

$$\partial_t \mathbf{u} + \nabla_{\mathbf{w}} \mathbf{u} = -\nabla_{\mathcal{S}} p - p \mathcal{H} \mathbf{v} + \frac{2}{\text{Re}} \text{div}_{\mathcal{C}} \boldsymbol{\sigma} - \gamma \mathbf{u} + \mathbf{b} - \lambda \mathbf{v} \quad (1)$$

$$\text{div} \mathbf{P} \mathbf{u} = 0 \quad (2)$$

$$\int_{\mathcal{S}} \mathbf{u} \cdot \mathbf{v} d\mathcal{S} = 0, \quad (3)$$

where  $[\nabla_{\mathbf{w}} \mathbf{u}]_i = (\nabla_{\mathcal{S}} \mathbf{u}_i, \mathbf{w})$ ,  $i = 1, 2, 3$ , with  $\mathbf{w} = \mathbf{u} - \partial_t \mathbf{X}$  is the relative velocity and  $\mathbf{X}$  a parameterisation of  $\mathcal{S}$ ,  $\boldsymbol{\sigma}(\mathbf{u}) = \frac{1}{2} (\nabla \mathbf{P} \mathbf{u} + (\nabla \mathbf{P} \mathbf{u})^T) \in T^2 \mathbb{R}^3|_{\mathcal{S}}$  is the rate of deformation tensor,  $\text{Re} > 0$  is the Reynolds number,  $\gamma \geq 0$  is a friction coefficient,  $\lambda \in \mathbb{R}$  is a Lagrange multiplier to ensure a constant enclosed volume, and  $\mathbf{b}$



denotes a bending force, defined by

$$\begin{aligned} \mathbf{b} = & -2k_{2,0} \left( \Delta_{\mathcal{S}} \mathcal{H} + (\mathcal{H} - \mathcal{H}_0) \left( \|\mathcal{B}\|^2 - \frac{1}{2} \mathcal{H} (\mathcal{H} - \mathcal{H}_0) \right) \right) \mathbf{v} \\ & - 12k_{4,0} \operatorname{div}_{\mathcal{S}} \left( (\mathcal{H} - \mathcal{H}_0)^2 \nabla_{\mathcal{S}} \mathcal{H} \right) \mathbf{v} \\ & - k_{4,0} \left( (\mathcal{H} - \mathcal{H}_0)^3 ((3\mathcal{H} + \mathcal{H}_0)\mathcal{H} - 8\mathcal{H}) \right) \mathbf{v} \\ & - k_{4,2} \left( 2\operatorname{div}_{\mathcal{S}} ((\mathcal{H}\mathbf{P} - \mathcal{B}) \nabla_{\mathcal{S}} \mathcal{H}) + \mathcal{H} \mathcal{H}^2 \right) \mathbf{v}, \end{aligned} \quad (4)$$

where  $\mathcal{H}_0$  is a spontaneous curvature and  $k_{2,0}, k_{4,0}, k_{4,2} \in \mathbb{R}$  are bending rigidity parameters. The system of equations consider a model for fluid deformable surfaces. Such models consist of incompressible surface Navier-Stokes equations with bending forces and a constraint on the enclosed volume. The highly nonlinear model accounts for the tight interplay between surface evolution, surface curvature and surface hydrodynamics and allows to model membranes with surface viscosity. For derivations of the model (with  $k_{4,0} = k_{4,2} = 0$ ) we refer to<sup>17,31,34,40,41</sup>. They consider different principle and build on a nonlinear Onsager formalism<sup>31</sup>, thin film limits from three-dimensional models<sup>34,40</sup> and a Lagrange-D'Alembert approach<sup>17,41</sup>. For a comparison of derivations for  $\mathbf{b} = 0$  and without the constraint on the enclosed volume we refer to<sup>42-44</sup>.

The mentioned solid-fluid duality of these models, which leads to a tight coupling between shape and flow, and let any shape change contribute to tangential flow, and vice versa, can best be seen by rewriting eqs. (1) - (3) as coupled equations for  $u_N = \mathbf{u} \cdot \mathbf{v}$  and  $\mathbf{u}_T = \mathbf{P}\mathbf{u}$ <sup>34,42,43</sup>. One out of several of the resulting coupling terms results from eq. (2), which leads to  $\operatorname{div} \mathbf{p}\mathbf{u} = u_N \mathcal{H} + \operatorname{div} \mathbf{p}\mathbf{u}_T = 0$  and explicitly demonstrates the relation between the mean curvature and the tangential velocity.

Considering the overdamped limit, formally letting  $\gamma \rightarrow \infty$ , leads to the classical dynamic equations

$$\mathbf{u} = -\nabla_{\mathcal{S}} p - p \mathcal{H} \mathbf{v} + \mathbf{b} - \lambda \mathbf{v} \quad (5)$$

$$\operatorname{div} \mathbf{p}\mathbf{u} = 0 \quad (6)$$

$$\int_{\mathcal{S}} \mathbf{u} \cdot \mathbf{v} d\mathcal{S} = 0, \quad (7)$$

for an inextensible membrane with constant volume. Using (5) in (6) provides the equation for the Lagrange multiplier for the inextensibility constraint  $-\Delta_{\mathcal{S}} p + p \mathcal{H}^2 + \lambda = \mathbf{b} \cdot \mathbf{v} \mathcal{H}$  and corresponds to previous models, if  $\mathbf{b}$  only contains second order geometric terms<sup>45-49</sup>. Further relaxing the constraint on inextensibility leads to the classical Canham-Helfrich models with area and volume constraints<sup>48</sup>.

In contrast with previous approaches for fluid deformable surfaces<sup>17,31,34,35,50</sup> the bending force  $\mathbf{b}$  also contains higher order geometric terms.

## 2.3 Bending forces

The bending force eq. (4) results from the bending energy

$$\mathcal{F}_{BE}(\mathcal{H}, \mathcal{K}) = \int_{\mathcal{S}} f_{BE}(\mathcal{H}, \mathcal{K}) d\mathcal{S} \quad (8)$$

with bending energy density  $f_{BE}$ . Formally we can derive  $f_{BE}$  via Taylor expansion at the spontaneous curvature  $\mathcal{H}_0$  leading to

$$\begin{aligned} f_{BE}(\mathcal{H}, \mathcal{K}) = & \sum_{n=0}^N \sum_{\alpha=0}^{\lfloor \frac{n}{2} \rfloor} k_{n,\alpha} (\mathcal{H} - \mathcal{H}_0)^{n-2\alpha} \mathcal{K}^{\alpha} \\ = & \underbrace{k_{0,0}}_{n=0} + \underbrace{k_{1,0}(\mathcal{H} - \mathcal{H}_0)}_{n=1} + \underbrace{k_{2,0}(\mathcal{H} - \mathcal{H}_0)^2 + k_{2,1}\mathcal{K}}_{n=2} \\ & + \underbrace{k_{3,0}(\mathcal{H} - \mathcal{H}_0)^3 + k_{3,1}(\mathcal{H} - \mathcal{H}_0)\mathcal{K}}_{n=3} \\ & + \underbrace{k_{4,0}(\mathcal{H} - \mathcal{H}_0)^4 + k_{4,1}(\mathcal{H} - \mathcal{H}_0)^2\mathcal{K} + k_{4,2}\mathcal{K}^2}_{n=4} \\ & + \dots \end{aligned}$$

in terms of geometric orders  $n \leq N \in \mathbb{N}$  for different bending rigidity parameters  $k_{n,\alpha} \in \mathbb{R}$ . This expression corresponds to the generalized form of the classical Canham-Helfrich energy ( $n = 2$ ) introduced in<sup>11</sup> and also considered in<sup>2,12,13</sup> if gradient terms are neglected. It can be simplified assuming certain properties of the bending energy: As we are interested in variations of the energy, we can omit the constant contribution  $k_{0,0}$ . We will further not allow for topological changes, thus, we can omit the  $k_{2,1}\mathcal{K}$  term, applying Gauss-Bonnet's theorem. Furthermore, we only allow for terms which guarantee boundedness from below. This excludes all odd geometric orders as well as  $k_{4,1}$ . Considering these points and only contributions up to geometric order  $N = 4$  leads to the following bending energy density

$$f_{BE}(\mathcal{H}, \mathcal{K}) = k_{2,0}(\mathcal{H} - \mathcal{H}_0)^2 + k_{4,0}(\mathcal{H} - \mathcal{H}_0)^4 + k_{4,2}\mathcal{K}^2,$$

which has been considered in<sup>14</sup> to study the stability of tubular shapes considering  $k_{2,0} > 0$ ,  $k_{4,0} = 0$  and  $k_{4,2} \geq 0$  as parameters to stabilize cylindrical shapes ( $k_{4,2} > 0$ ) or to enforce pearling ( $k_{4,2} < 0$ ). The same form is also considered in<sup>12</sup>, arguing that a cylinder with radius  $R$  is stable if  $k_{2,0} < 0$  and  $k_{4,0} = -\frac{R^2}{2} k_{2,0}$  with  $k_{4,2}$  not determined. We here restrict the parameter space to  $k_{2,0} > 0$ ,  $k_{4,0} \geq 0$  and  $k_{4,2} \geq 0$ . We would like to remark that more general parameter combinations are possible, still leading to well-posed bending energies and stable solutions. However,  $\mathcal{K} = 0$  is a property of a tube and thus  $k_{4,2}\mathcal{K}^2$  seems to be the most plausible higher order extension in the bending energy to stabilize tubular structures.  $k_{4,0}(\mathcal{H} - \mathcal{H}_0)^4$  is considered for completeness. In any case the bending force  $\mathbf{b}$  is derived as the negative of the variational derivative of the bending energy (8)

$$\mathbf{b} = -\frac{\delta \mathcal{F}_{BE}}{\delta \mathbf{X}}.$$

A detailed derivation of the bending force is done in Appendix 5.1.



### 2.4 Numerical approach

The numerical approach extends the approach used in<sup>17,35</sup>, which is based on surface finite elements<sup>51,52</sup> and builds on a Taylor-Hood element for the surface Navier-Stokes equations, higher order surface parametrizations, appropriate approximations of the geometric quantities, mesh redistribution, a semi-implicit discretization in time and an iterative approach to deal with the non-local constraint on the enclosed volume. Additional challenges emerge from the higher order geometric terms. In Appendix 5.2 we provide a detailed description and in Appendix 5.3 convergence studies for these terms. The implementation is done in DUNE/AMDiS<sup>53,54</sup>. Throughout the paper we consider  $Re = 1.0$  and  $\gamma = 0$  and only vary  $\mathcal{H}_0$ ,  $k_{2,0}$ ,  $k_{4,0}$  and  $k_{4,2}$ . Numerical parameters, such as mesh size  $h$  and time step  $\tau$  are chosen to resolve the highest curvature values and to meet stability constraints, following<sup>35</sup>.

## 3 Results

### 3.1 Axialymmetric simulation without surface viscosity

We begin by describing the membrane using cylindrical coordinates  $(r, \theta, z)$  and consider a rotational symmetric tube with periodic boundary conditions. With


$$\mathcal{H} = \frac{1}{\sqrt{1+(\partial_z r)^2}} \left( \frac{\partial_{zz} r}{1+(\partial_z r)^2} - \frac{1}{r} \right), \quad \mathcal{K} = -\frac{\partial_{zz} r}{r(1+(\partial_z r)^2)^2} \quad (9)$$

the bending energy density  $f_{BE}(\mathcal{H}, \mathcal{K})$  can be reformulated and considered as a function of  $r(z, t)$ , the radial distance of the axisymmetric membrane from the cylindrical symmetry axis, where  $z$  measures the coordinate along that axis and  $t$  is time. For the evolution we consider the corresponding equations to eq. (5) - (7) but drop the constraints on inextensibility and volume. The resulting equation to solve reads

$$\frac{1}{\sqrt{1+(\partial_z r)^2}} \partial_t r = u_N = -\frac{\delta \mathcal{F}_{BE}}{\delta \mathbf{X}} \cdot \mathbf{v} = b, \quad (10)$$

with  $b = \mathbf{b} \cdot \mathbf{v}$ . While lengthy if fully written down, the resulting model can be solved using standard approaches. We again use finite elements in space, a semi-implicit discretization in time, smoothing of the geometric properties and consider DUNE/AMDiS<sup>53,54</sup> for the realization. Fig. 1 shows the time evolution of a periodically perturbed tube for different values of  $k_{4,2}$  and  $k_{4,0}$  together with the evolution of the bending energy  $\mathcal{F}_{BE}$  over time. For  $k_{4,2} = 0$  and  $k_{4,0} = 0$  this is a well studied problem of the stability of a tube<sup>55-57</sup>. The parameters are chosen to remain within the stability region. The results clearly indicate a stronger damping of the perturbations for increasing values of  $k_{4,2}$  and  $k_{4,0}$ . However, the behaviour slightly differs with respect to the damping of the perturbations and adjustment of  $r(z)$ . This difference is also seen in the decrease of the bending energy.

Results of a linear stability analysis<sup>58</sup> for the case of  $k_{4,2} = 0$  and  $k_{4,0} = 0$  but considering surface viscosity using a Stokes approximation of eq. (1) indicate a similar stability region as in the overdamped limit and we assume this to hold also for  $k_{4,2} > 0$  and  $k_{4,0} > 0$  and the full problem.



5

Fig. 1 (a) Time evolution of axialymmetric simulations for different parameters for  $k_{4,2}$  and  $k_{4,0}$  starting from the same periodic solution  $r_0(z) = \frac{1}{4} \sin(\frac{\pi}{4}z) + 1$  at  $t_0$ . The evolution goes from dark to light converging to a tube with  $r(z) = 0.5$ . The depicted time instances are marked in (b). (b) Corresponding time evolution of the bending energy for different values of  $k_{4,2}$  and  $k_{4,0}$ . Parameters are  $k_{2,0} = 0.05$  and  $\mathcal{H}_0 = -2$ .




Fig. 2 (a) As initial surface we consider a cylinder with hemispherical caps. (b) The geometry is determined by the length of the cylinder  $l$  and the radius  $r$  of the cylinder and hemispherical caps.



### 3.2 Equilibrium shape of tubular cell

The shape of a tubular cell can be approximated by a cylinder with hemispherical caps, as shown in Fig. 2. For this simple shape, we can compute the bending energy as

$$\begin{aligned} \mathcal{F}_{BE} = & k_{2,0} \left( 2\pi r l \left( -\frac{1}{r} - \mathcal{H}_0 \right)^2 + 4\pi r^2 \left( -\frac{2}{r} - \mathcal{H}_0 \right)^2 \right) \\ & + k_{4,0} \left( 2\pi r l \left( -\frac{1}{r} - \mathcal{H}_0 \right)^4 + 4\pi r^2 \left( -\frac{2}{r} - \mathcal{H}_0 \right)^4 \right) \\ & + k_{4,2} \frac{4\pi}{r^2} \end{aligned}$$

where  $r$  is the radius and  $l$  the length of the cylinder. For  $\mathcal{H}_0 = -\frac{1}{r}$ , the energy of the cylindrical part vanishes and the energy simplifies to

$$\mathcal{F}_{BE} = 4\pi \left( k_{2,0} + \frac{k_{4,0}}{r^2} + \frac{k_{4,2}}{r^2} \right).$$

Note that this energy is independent of the length of the cylinder. We consider this shape as the initial configuration and the energy as a quantity for comparison.

Fig. 3 shows the final configurations achieved by solving the full problem, eq. (1) - (3), for three different values  $l$ . In all cases, the shape deviates from the initial configuration. For  $k_{4,2} = 0$  and  $k_{4,0} = 0$  the resulting shapes correspond to the equilibrium prolate shapes in<sup>7</sup>. However, for  $k_{4,2} > 0$  and  $k_{4,0} > 0$ , these shapes deviate. For  $k_{4,2} > 0$  the bending energy is reduced by increasing  $l$  and the Hausdorff distance  $d_H(\mathcal{S}, \mathcal{S}_0)$  to the idealized initial shape of a cylinder with hemispherical caps  $\mathcal{S}_0$  is reduced by increasing  $k_{4,2}$ , see Fig. 4. The Hausdorff distance is a measure of the distance between sets of points, in our case the discretization points of the equilibrium shape and the initial shape. For a mathematical definition and implementational issues see Appendix 5.4. The observed trends are far from being general. This can also not be expected due to the highly nonlinear coupling of the geometric terms and the considered constraints on volume and inextensibility. However, they confirm the intuition of the potential impact of  $k_{4,2}\mathcal{H}^2$  in the bending energy on the emerging equilibrium shapes. The effect of  $k_{4,0} > 0$  differs. The increased tendency to enforce  $\mathcal{H} = \mathcal{H}_0$  leads to undulations with a wavelength related to  $-2/\mathcal{H}_0$ . How well this can be achieved strongly depends on the length  $l$ . Therefore also the Hausdorff distance  $d_H(\mathcal{S}, \mathcal{S}_0)$  to the idealized initial shape of a cylinder with hemispherical caps  $\mathcal{S}_0$  does not decrease but leads to a non-monotonic behaviour if considered as a function of  $l$ . The dependency of  $l$  on the bending energy  $\mathcal{F}_{BE}$  is negligible for the considered parameters. To summarize, within the stability region of the classical Canham-Helfrich model, higher order geometric terms related to  $k_{4,2}\mathcal{H}^2$  seem to stabilize tubular shapes and terms related to  $k_{4,0}(\mathcal{H} - \mathcal{H}_0)^4$  seem to destabilize these shapes.

### 3.3 Dynamic evolution

In order to further explore the influence of the higher order geometric terms related to  $k_{4,2}\mathcal{H}^2$  and  $k_{4,0}(\mathcal{H} - \mathcal{H}_0)^2$  on the dynam-

ics, we consider an initial surface  $\mathcal{S}_0$  as a perturbed unit sphere

$$\mathcal{S}_0 = \{1 + r_0 Y_l^m(\phi, \vartheta) : \phi \in [0, \pi], \vartheta \in [-\pi, \pi]\}, \quad r_0 > 0, \quad (11)$$

$$Y_l^m(\phi, \vartheta) = \sqrt{\frac{2l+1(l-m)!}{4\pi(l+m)!}} P_l^m(\cos \vartheta) e^{im\phi} \quad (12)$$

with spherical harmonics  $Y_l^m$  and Legendre polynomials  $P_l^m$  and the case  $l = 5, m = 3$  and  $r_0 = 0.5$ . The velocity field is initialized with  $\mathbf{u}_0 = 0$ . The initial surface has non-zero mean and Gaussian curvature and is out of equilibrium. It serves as a prototypical examples to study the rapid shape changes featuring moderate geometric properties of a cell with membrane reservoirs. The resulting bending force induces shape deformations in normal direction. However, the curvature terms also induce tangential flows, which also contribute to shape deformations. This coupling between tangential flow and shape deformations is well understood and shown to enhance the evolution towards equilibrium shapes<sup>35</sup>. Here, we explore the evolution for different values of  $k_{4,2}$  and  $k_{4,0}$ , which read  $k_{4,2} = 0, 0.06$  and  $0.12$  and  $k_{4,0} = 0, 0.00625$  and  $0.0125$ , respectively. Figs. 5 and 6 show snapshots of the evolutions. The color coding corresponds to shape deformations (red - movement outwards, blue - movement inwards) and the arrows indicate the tangential velocity, where the length scales with the magnitude. Furthermore, the energy contributions are shown over time. Here, the bending energy is split into the different contributions  $\mathcal{F}_{BE} = \mathcal{F}_{2,0} + \mathcal{F}_{4,0} + \mathcal{F}_{4,2}$  and the total energy  $\mathcal{F} = \mathcal{F}_{BE} + \mathcal{F}_K$  is the sum of the bending energy and the kinetic energy  $\mathcal{F}_K = \int_{\mathcal{S}} \frac{1}{2} \mathbf{u}^2 d\mathcal{S}$ . All evolutions converge to an equilibrium shape, which for the cases  $k_{4,2} = 0$  and  $k_{4,0} = 0$  correspond to the associated Seifert shape<sup>7</sup>, which here corresponds to an oblate. For  $k_{4,2} > 0$  the equilibrium shapes only slightly differ. The difference in orientation might result from the different dynamics. However, due to the decoupling from the surrounding bulk phases and the considered parameter  $\gamma = 0$ , force-free rigid body rotations are also possible<sup>59</sup>. While the equilibrium shapes are similar, significant changes can be observed in the dynamics. The close coupling between the bending energy  $\mathcal{F}_{BE}$  and the kinetic energy  $\mathcal{F}_K$  can be observed and related to significant shape changes. But their appearance differs. The plateau in  $\mathcal{F}_{BE} = \mathcal{F}_{2,0}$  for  $k_{4,2} = 0$  between  $t \approx 2$  and  $t \approx 6$  is reduced for  $k_{4,2} = 0.06$  and  $0.12$  and already ends at  $t \approx 4.5$  and  $t \approx 3$ , respectively. It should be noted that the absolute values of the energies cannot be directly compared as  $k_{4,2}$  varies in the definition. The situation changes for  $k_{4,0} > 0$ . The equilibrium shapes are prolate shapes. Besides this difference in the long time behavior also the short time evolution changes. We observe similar behavior as for  $k_{4,2} > 0$  with an enhanced influence of the kinetic energy and a faster evolution towards intermediate shapes. The final convergence towards the equilibrium shape at late times probably results from similar values for the local minima on the oblate and prolate branch as also known for the classical Canham-Helfrich model<sup>6</sup>. Again, the absolute values of the energies cannot be directly compared as  $k_{4,0}$  varies in the definition.

Qualitatively the higher order geometric terms further enhance the evolution and lead to alternative pathways to dissipate en-






Fig. 3 (a) Equilibrium shapes for  $k_{2,0} = 0.125$ ,  $\mathcal{H}_0 = -2$  and  $k_{4,2} = 0$ . As initial condition we consider a cylinder of radius  $r = 0.5$  with hemispherical caps and vary the length  $l$  of the cylindrical part. From left to right the lengths are  $l = 2, 3$  and  $4$ . The surfaces are colored by mean curvature  $\mathcal{H}$  (upper row) and Gaussian curvature  $\mathcal{K}$  (lower row). (b) Equatorial cuts through the equilibrium shapes for different values of  $k_{4,2}$  and  $l$ , keeping  $k_{2,0} = 0.125$  and  $\mathcal{H}_0 = -2$ . (c) Equatorial cuts through the equilibrium shapes for different values of  $k_{4,0}$  and  $l$ , keeping  $k_{2,0} = 0.125$  and  $\mathcal{H}_0 = -2$ .






Fig. 4 (a), (b) Bending energy of the final shape for fixed values of  $k_{4,2}$  (a) and  $k_{4,0}$  (b) for different length  $l$ . (c), (d) Hausdorff distance between the equilibrium shape and the initial shape for different lengths  $l = 2, 3$  and  $4$  as a function of  $k_{4,2}$  (c) and  $k_{4,0}$  (d).

ergy. This is most pronounced in the inlets highlighting the initial evolution, the drastic decrease of  $\mathcal{F}_{4,2}$  and  $\mathcal{F}_{4,0}$  is associated with large fluctuation of  $\mathcal{F}_K$ . This behavior increases with increasing values of  $k_{4,2}$  and  $k_{4,0}$ . Furthermore, while  $\mathcal{F}_{4,2}$  is roughly one order of magnitude smaller than  $\mathcal{F}_{2,0}$ , the influence on the dynamics is dramatic. This is less pronounced for  $\mathcal{F}_{4,0}$ , which is at the same order as  $\mathcal{F}_{2,0}$ . We expect these fast shape changes to be even enhanced for less regular real geometries of membrane reservoirs, due to the appearance of larger curvature gradients.

## 4 Conclusions

Motivated by rapid shape changes of cells, where an excess of membrane that is organized in membrane reservoirs is made available to the cell in the order of seconds<sup>23</sup>, we formulated a minimal mesoscopic membrane model which helps to facilitate this behavior. This, on the one side includes an extension of the classical Canham-Helfrich model towards higher order geometric terms, and on the other side the explicit treatment of the fluid properties of the membrane by considering membranes as fluid deformable surfaces. The first aspect adds to the classical bending energy  $\mathcal{F}_{2,0} = \int_{\mathcal{S}} k_{2,0} (\mathcal{H} - \mathcal{H}_0)^2 d\mathcal{S}$ , terms proportional to the Gaussian curvature squared,  $\mathcal{F}_{4,2} = \int_{\mathcal{S}} k_{4,2} \mathcal{H}^2 d\mathcal{S}$ , and the mean curvature minus the spontaneous curvature to the fourth power,  $\mathcal{F}_{4,0} = \int_{\mathcal{S}} k_{4,0} (\mathcal{H} - \mathcal{H}_0)^4 d\mathcal{S}$ . The effects of these higher order terms are different. While the one related to the Gaussian curvature squared not only helps to damp perturbations of tubes, it also has a tendency to stabilize them, the term related to the mean curvature minus the spontaneous curvature to the fourth power has a tendency to destabilize tubes. Both effects have been considered in idealized rotational symmetric and full three-dimensional situations by analyzing the evolution and the emerging equilibrium shapes. If combined with the second aspect, which takes the surface viscosity of the membrane into account and combines the bending in normal direction with the proper-

ties of an inextensible surface fluid, as a fluid deformable surface, the dynamic drastically changes. Already for the classical bending energy  $\mathcal{F}_{2,0} = \int_{\mathcal{S}} k_{2,0} (\mathcal{H} - \mathcal{H}_0)^2 d\mathcal{S}$  an enhanced evolution towards the equilibrium shape has been observed if the effects of surface viscosity are taken into account<sup>35</sup>. With the higher order geometric terms this is further enhanced. The considered numerical experiments for the relaxation of a perturbed sphere showed alternative pathways to dissipate energy and strong tangential flows inducing fast shape changes. However, besides enabling rapid shape changes, the evolution and also the emerging equilibrium shapes qualitatively differ for the considered higher order terms. While the model with the Gaussian curvature squared term also converges to similar oblate like shapes as the corresponding Seifert shapes<sup>6</sup> for the classical Canham-Helfrich model, the model with the mean curvature minus the spontaneous curvature to the fourth power term converges to prolate like shapes.

However, the focus of this paper is not to classify the increased phase space of equilibrium shapes resulting from the higher order terms, but to address additional mechanisms which facilitate rapid shape changes. While certainly more research is needed to fully explore the potential of the higher order geometric terms, e.g., with respect to the stability of tubes extending the analysis in<sup>58</sup>, the numerical studies already clearly indicate the potential for rapid shape changes. Even if only passive contributions of a homogeneous membrane are considered, and a full model for morphological changes of a cell requires to take inhomogeneities, active processes of the underlying cortex, adhesion between the membrane and the cortex and probably even more phenomena into account, the study contributes to identifying underlying general mechanical principles which might help to predict and control the dynamics of cells<sup>20</sup>. Surface viscosity and higher order geometric terms in the bending energy provide mechanical cues and probably support active processes to enable rapid shape changes of cells.

A full model able to address the mentioned example of frequently forming and retracting filopodia and the associated fast shape changes of membrane reservoirs will require to couple models for the cellular cortex<sup>24–27</sup> with membrane models of the considered type, which requires additional modeling and numerical efforts. One intermediate step to link this research closer to biology could be efforts to resolve the dynamics of membrane reservoirs and compare them with simulations of the proposed model.

## Conflicts of interest

There are no conflicts to declare.

## Data availability

Data are available from Zenodo at <https://10.5281/zenodo.14503545>.

## Acknowledgements

We acknowledge fruitful discussions with Patrick Zager (UCSF) and Veit Krause (TUD). This work was funded by DFG within FOR3013 "Vector- and tensor-valued surface PDEs". We further acknowledge computing resources at FZ Jülich under Grant No.








Fig. 5 Evolution of a perturbed sphere considering parameters  $k_{2,0} = 0.125$ ,  $\mathcal{H}_0 = 0$  and different values of  $k_{4,2}$ . (a)-(c) Time instances for the evolution for  $k_{4,2} = 0, 0.06$  and  $0.12$ , respectively. The color coding indicates movement in normal direction (red - movement outwards, blue - movement inwards) and the arrows indicate the tangential velocity. The time evolves from left to right, the time instances except for the last one are equal and are depicted in (d). (d) Evolution of the different energy contributions,  $\mathcal{F}_{2,0}$  and  $\mathcal{F}_{4,2}$  are the energies linked to the corresponding bending terms,  $\mathcal{F}_K$  is the kinetic energy. The bending energy is  $\mathcal{F}_{BE} = \mathcal{F}_{2,0} + \mathcal{F}_{4,2}$  and the total energy  $\mathcal{F} = \mathcal{F}_{BE} + \mathcal{F}_K$ . Shown is the evolution for  $k_{4,2} = 0, 0.06$  and  $0.12$  (from left to right). Within †corresponding videos to the evolution in (a) - (c) are provided using a LIC filter for visualization of the tangential flow.

Faraday Discussions Accepted Manuscript






Fig. 6 Evolution of a perturbed sphere considering parameters  $k_{2,0} = 0.125$ ,  $\mathcal{H}_0 = 0$  and different values of  $k_{4,0}$ . (a)-(c) Time instances for the evolution for  $k_{4,0} = 0, 0.00625$  and  $0.0125$ , respectively. The color coding indicates movement in normal direction (red - movement outwards, blue - movement inwards) and the arrows indicate the tangential velocity. The time evolves from left to right, the time instances except for the last one are equal and are depicted in (d). (d) Evolution of the different energy contributions,  $\mathcal{F}_{2,0}$  and  $\mathcal{F}_{4,0}$  are the energies linked to the corresponding bending terms,  $\mathcal{F}_K$  is the kinetic energy. The bending energy is  $\mathcal{F}_{BE} = \mathcal{F}_{2,0} + \mathcal{F}_{4,0}$  and the total energy  $\mathcal{F} = \mathcal{F}_{BE} + \mathcal{F}_K$ . Shown is the evolution for  $k_{4,0} = 0, 0.00625$  and  $0.0125$  (from left to right). Within corresponding videos to the evolution in (a) - (c) are provided using a LIC filter for visualization of the tangential flow.



MORPHO and at ZIH under Grant No. WIR.

## References

- H. T. McMahon and J. L. Gallop, *Nature*, 2005, **438**, 590–596.
- M. Deserno, *Chemistry and Physics of Lipids*, 2015, **185**, 11–45.
- R. Lipowsky and R. Dimova, *Soft Matter*, 2021, **17**, 214–221.
- W. Helfrich, *Zeitschrift für Naturforschung C*, 1973, **28**, 693.
- P. Canham, *J. Theor. Biol.*, 1970, **26**, 61.
- U. Seifert, *Advances in Physics*, 1997, **46**, 13–137.
- U. Seifert, K. Berndl and R. Lipowsky, *Physical Review A*, 1991, **44**, 1182–1202.
- N. Bobrovska, W. Gózdź, V. Kralj-Iglič and A. Iglič, *PloS one*, 2013, **8**, e73941.
- N. Walani, J. Torres and A. Agrawal, *Phys. Rev. E*, 2014, **89**, 062715.
- A. Mahapatra and P. Rangamani, *Soft Matter*, 2023, **19**, 4345–4359.
- M. D. Mitov, *Comptes Rendus de l'Academie Bulgare des Sciences*, 1978, **31**, 513–515.
- J. B. Fournier and P. Galatola, *Europhysics Letters*, 1997, **39**, 225.
- R. Capovilla, J. Guven and J. A. Santiago, *Journal of Physics A: Mathematical and General*, 2003, **36**, 6281.
- T. Shemesh, A. Luini, V. Malhotra, K. N. Burger and M. M. Kozlov, *Biophysical Journal*, 2003, **85**, 3813–3827.
- M. Kaksonen and A. Roux, *Nature Reviews Molecular Cell Biology*, 2018, **19**, 313–326.
- S. C. Al-Izzi, P. Sens and M. S. Turner, *Phys. Rev. Lett.*, 2020, **125**, 018101.
- E. Bachini, V. Krause, I. Nitschke and A. Voigt, *Journal of Fluid Mechanics*, 2023, **977**, A41.
- M. Arroyo, L. Heltai, D. Millán and A. DeSimone, *Proceedings of the National Academy of Sciences*, 2012, **109**, 17874–17879.
- B. Winkler, I. S. Aranson and F. Ziebert, *Communications Physics*, 2019, **2**, 82.
- R. K. Sadhu, A. Iglič and N. S. Gov, *Journal of Cell Science*, 2023, **136**, jcs260744.
- V. T. Ruhoff, N. Leijnse, A. Doostmohammadi and P. M. Bendix, *Trends in Cell Biology*, 2024.
- H. De Belly and O. D. Weiner, *Current Opinion in Cell Biology*, 2024, **89**, 102392.
- H. De Belly, S. Yan, H. B. da Rocha, S. Ichbiah, J. P. Town, P. J. Zager, D. C. Estrada, K. Meyer, H. Turlier, C. Bustamante et al., *Cell*, 2023, **186**, 3049–3061.
- S. R. Naganathan, S. Fürthauer, M. Nishikawa, F. Jülicher and S. W. Grill, *eLife*, 2014, **3**, e04165.
- A.-C. Reymann, F. Staniscia, A. Erzberger, G. Salbreux and S. W. Grill, *eLife*, 2016, **5**, e17807.
- H. B. da Rocha, J. Bleyer and H. Turlier, *Journal of the Mechanics and Physics of Solids*, 2022, **164**, 104876.
- A. Bhatnagar, M. Nestler, P. Gross, M. Kramar, M. Leaver, A. Voigt and S. W. Grill, *Current Biology*, 2023, **33**, 5096–5108.
- T. Itoh and K. Tsujita, *Current Opinion in Cell Biology*, 2023, **81**, 102173.
- H. A. Faizi, R. Dimova and P. M. Vlahovska, *Biophysical Journal*, 2022, **121**, 910–918.
- M. Arroyo and A. DeSimone, *Physical Review E*, 2009, **79**, 031915.
- A. Torres-Sánchez, D. Millán and M. Arroyo, *J. Fluid Mech.*, 2019, **872**, 218–271.
- A. Voigt, *J. Fluid Mech.*, 2019, **878**, 1–4.
- S. C. Al-Izzi and R. G. Morris, *Seminars in Cell & Developmental Biology*, 2021, **120**, 44–52.
- S. Reuther, I. Nitschke and A. Voigt, *J. Fluid Mech.*, 2020, **900**, R8.
- V. Krause and A. Voigt, *J. Comput. Phys.*, 2023, **486**, 112097.
- M. Simunovic, G. A. Voth, A. Callan-Jones and P. Bassereau, *Trends in Cell Biology*, 2015, **25**, 780–792.
- I. Derényi, F. Jülicher and J. Prost, *Phys. Rev. Lett.*, 2002, **88**, 238101.
- A. Roux, G. Cappello, J. Cartaud, J. Prost, B. Goud and P. Bassereau, *Proceedings of the National Academy of Sciences*, 2002, **99**, 5394–5399.
- P. G. Saffman and M. Delbrück, *Proc. Natl. Acad. Sci. USA*, 1975, **72**, 3111–3113.
- I. Nitschke, S. Reuther and A. Voigt, *Physical Review Fluids*, 2019, **4**, 044002.
- I. Nitschke and A. Voigt, *Advances in Differential Equations*, 2025, **30**, 335–420.
- S. Reuther and A. Voigt, *Multiscale Modeling & Simulation*, 2015, **13**, 632–643.
- S. Reuther and A. Voigt, *Multiscale Modeling & Simulation*, 2018, **16**, 1448–1453.
- P. Brandner, A. Reusken and P. Schwering, *Interfaces and Free Boundaries*, 2022, **24**, 533–563.
- T. Biben and C. Misbah, *Physical Review E*, 2003, **67**, 031908.
- Y. Kim and M.-C. Lai, *J. Comput. Phys.*, 2010, **229**, 4840–4853.
- D. Salac and M. J. Miksis, *J. Fluid Mech.*, 2012, **711**, 122–146.
- F. Haußer, W. Marth, S. Li, J. S. Lowengrub, A. Rätz and A. Voigt, *Int. J. Biomath. Biostat.*, 2013, **2**, 19–48.
- S. Aland, S. Egerer, J. Lowengrub and A. Voigt, *J. Comput. Phys.*, 2014, **277**, 32–47.
- V. Krause and A. Voigt, *Journal of the Royal Society Interface*, 2024, **21**, 20240056.
- G. Dziuk and C. M. Elliott, *Acta Numerica*, 2013, **22**, 289–396.
- M. Nestler, I. Nitschke and A. Voigt, *J. Comput. Phys.*, 2019, **389**, 48–61.
- S. Vey and A. Voigt, *Comput. Vis. Sci.*, 2007, **10**, 57–67.
- T. Witkowski, S. Ling, S. Praetorius and A. Voigt, *Adv. Comput. Math.*, 2015, **41**, 1145–1177.
- T. R. Powers, *Rev. Mod. Phys.*, 2010, **82**, 1607–1631.
- R. E. Goldstein, P. Nelson, T. Powers and U. Seifert, *J. Phys. II France*, 1996, **6**, 767–796.
- S. C. Al-Izzi, G. Rowlands, P. Sens and M. S. Turner, *Phys. Rev.*



- Let.*, 2018, **120**, 138102.
- 58 M. Janssen, S. Liese, S. C. Al-Izzi and A. Carlson, *Phys. Rev. E*, 2024, **109**, 044403.
- 59 M. Nestler and A. Voigt, *PAMM*, 2023, **23**, e202300044.
- 60 I. Nitschke, S. Sadik and A. Voigt, *IMA Journal of Applied Mathematics*, 2023, **88**, 917–958.
- 61 J. W. Barrett, H. Garcke and R. Nurnberg, *SIAM Journal on Scientific Computing*, 2008, **31**, 225–253.
- 62 S. Praetorius and F. Stenger, *Archive of Numerical Software*, 2022, **6**, 1–27.
- 63 P. Brandner, T. Jankuhn, S. Praetorius, A. Reusken and A. Voigt, *SIAM Journal on Scientific Computing*, 2022, **44**, A1807–A1832.
- 64 H. Hardering and S. Praetorius, *IMA Journal of Numerical Analysis*, 2023, **43**, 1543–1585.
- 65 F. Commandeur, J. Velut and O. Acosta, *The VTK Journal*, 2011.

## 5 Appendix

### 5.1 Model derivation

We here compute the variational derivatives of the higher order geometric terms in the bending energy. We therefore write the bending energy as

$$\mathcal{F}_{BE} = \int_{\mathcal{S}} k_{2,0} f_{BE}^{2,0} + k_{4,0} f_{BE}^{4,0} + k_{4,2} f_{BE}^{4,2} d\mathcal{S}$$

with bending density components  $f_{BE}^{n,\alpha} = (\mathcal{H} - \mathcal{H}_0)^{n-2\alpha} \mathcal{H}^\alpha$ . The corresponding forces  $\mathbf{b}^{n,\alpha} \in T\mathbb{R}^3|_{\mathcal{S}}$  are given by

$$\langle \mathbf{b}^{n,\alpha}, \mathbf{W} \rangle_{L^2(T\mathbb{R}^3|_{\mathcal{S}})} := - \left\langle \frac{\delta}{\delta \mathbf{X}} \int_{\mathcal{S}} f_{BE}^{n,\alpha} d\mathcal{S}, \mathbf{W} \right\rangle_{L^2(T\mathbb{R}^3|_{\mathcal{S}})}$$

for all  $\mathbf{W} \in T\mathbb{R}^3|_{\mathcal{S}}$ . The result for  $f_{BE}^{2,0}$  is well known

$$\mathbf{b}^{2,0} = -2 \left( \Delta_{\mathcal{S}} \mathcal{H} + (\mathcal{H} - \mathcal{H}_0) \left( \|\mathcal{B}\|^2 - \frac{1}{2} \mathcal{H} (\mathcal{H} - \mathcal{H}_0) \right) \right) \mathbf{v}.$$

For the remaining terms we use the deformation derivative  $\bar{\partial}_{\mathbf{W}}$ <sup>60</sup> and obtain the deformation formula

$$\langle \mathbf{b}^{n,\alpha}, \mathbf{W} \rangle_{L^2(T\mathbb{R}^3|_{\mathcal{S}})} = - \int_{\mathcal{S}} \bar{\partial}_{\mathbf{W}} f_{BE}^{n,\alpha} + f_{BE}^{n,\alpha} \operatorname{div}_C \mathbf{W} d\mathcal{S}. \quad (13)$$

Moreover, it is

$$\mathbf{P}(\bar{\partial}_{\mathbf{W}} \mathcal{B}) \mathbf{P} = \nabla_{\mathcal{S}}(\mathbf{v} \cdot \nabla_C \mathbf{W}) - \mathcal{B} \nabla_C \mathbf{W}, \quad (14)$$

$$\bar{\partial}_{\mathbf{W}} \mathcal{H} = \operatorname{tr}(\mathbf{P}(\bar{\partial}_{\mathbf{W}} \mathcal{B}) \mathbf{P}) = \operatorname{div}_{\mathcal{S}}(\mathbf{v} \cdot \nabla_C \mathbf{W}) - \mathcal{B} : \nabla_C \mathbf{W} \quad (15)$$

valid<sup>60</sup>. As a consequence, the deformation formula (13) and integrations by parts result in

$$\begin{aligned} \langle \mathbf{b}^{4,0}, \mathbf{W} \rangle_{L^2(T\mathbb{R}^3|_{\mathcal{S}})} &= - \int_{\mathcal{S}} 4(\mathcal{H} - \mathcal{H}_0)^3 \bar{\partial}_{\mathbf{W}} \mathcal{H} + (\mathcal{H} - \mathcal{H}_0)^4 \operatorname{div}_C \mathbf{W} d\mathcal{S} \\ &= \int_{\mathcal{S}} 12(\mathcal{H} - \mathcal{H}_0)^2 (\nabla_{\mathcal{S}} \mathcal{H}) \cdot (\mathbf{v} \cdot \nabla_C \mathbf{W}) \\ &\quad + (\mathcal{H} - \mathcal{H}_0)^3 (4\mathcal{B} - (\mathcal{H} - \mathcal{H}_0) \mathbf{P}) : \nabla_C \mathbf{W} d\mathcal{S} \end{aligned}$$

Another application of integrations by parts w. r. t.  $\nabla_C$ , yields

$$\begin{aligned} \mathbf{b}^{4,0} &= - \operatorname{div}_C (12(\mathcal{H} - \mathcal{H}_0)^2 \mathbf{v} \otimes \nabla_{\mathcal{S}} \mathcal{H} \\ &\quad + (\mathcal{H} - \mathcal{H}_0)^3 (4\mathcal{B} - (\mathcal{H} - \mathcal{H}_0) \mathbf{P})), \end{aligned}$$

where

$$\operatorname{div}_C(\boldsymbol{\sigma} \mathbf{P}) = \operatorname{div}_{\mathcal{S}}(\mathbf{P} \boldsymbol{\sigma} \mathbf{P}) - \mathbf{v} \cdot \boldsymbol{\sigma} \mathcal{B} + (\operatorname{div}_{\mathcal{S}}(\mathbf{v} \cdot \boldsymbol{\sigma} \mathbf{P}) + \mathcal{B} : \boldsymbol{\sigma}) \mathbf{v} \quad (16)$$

holds for all  $\boldsymbol{\sigma} \in T^2\mathbb{R}^3|_{\mathcal{S}}$ <sup>41,60</sup>. Therefore, the tangential part of  $\mathbf{b}^{4,0}$  cancels out and we obtain

$$\begin{aligned} \mathbf{b}^{4,0} &= - (12 \operatorname{div}_{\mathcal{S}}((\mathcal{H} - \mathcal{H}_0)^2 \nabla_{\mathcal{S}} \mathcal{H}) \\ &\quad + (\mathcal{H} - \mathcal{H}_0)^3 ((3\mathcal{H} + \mathcal{H}_0) \mathcal{H} - 8\mathcal{H})) \mathbf{v}. \end{aligned}$$

With (14) and (15), the deformation derivative of  $\mathcal{H} = 2(\mathcal{H}^2 - \|\mathcal{B}\|^2)$  reveals

$$\begin{aligned} \bar{\partial}_{\mathbf{W}} \mathcal{H} &= \mathcal{H} \bar{\partial}_{\mathbf{W}} \mathcal{H} - \mathcal{B} : \bar{\partial}_{\mathbf{W}} \mathcal{B} \\ &= (\mathcal{H} \mathbf{P} - \mathcal{B}) : (\nabla_{\mathcal{S}}(\mathbf{v} \cdot \nabla_C \mathbf{W}) - \mathcal{B} \nabla_C \mathbf{W}). \end{aligned}$$

As a consequence, the deformation formula (13), integrations by parts, and  $\mathcal{B}^2 = \mathcal{H} \mathcal{B} - \mathcal{H} \mathbf{P}$ , result in

$$\begin{aligned} \langle \mathbf{b}^{4,2}, \mathbf{W} \rangle_{L^2(T\mathbb{R}^3|_{\mathcal{S}})} &= - \int_{\mathcal{S}} 2\mathcal{H} \bar{\partial}_{\mathbf{W}} \mathcal{H} + \mathcal{H}^2 \operatorname{div}_C \mathbf{W} d\mathcal{S} \\ &= \int_{\mathcal{S}} 2 \operatorname{div}_{\mathcal{S}}(\mathcal{H}(\mathcal{H} \mathbf{P} - \mathcal{B})) \cdot (\mathbf{v} \cdot \nabla_C \mathbf{W}) + \mathcal{H}^2 \mathbf{P} : \nabla_C \mathbf{W} d\mathcal{S}. \end{aligned}$$

Since  $\operatorname{div}_{\mathcal{S}}(\mathcal{H} \mathbf{P} - \mathcal{B}) = 0$  holds, integrations by parts yields

$$\mathbf{b}^{4,2} = - \operatorname{div}_C (2\mathbf{v} \otimes (\mathcal{H} \mathbf{P} - \mathcal{B}) \nabla_{\mathcal{S}} \mathcal{H} + \mathcal{H}^2 \mathbf{P}).$$

Using (16), the tangential part of  $\mathbf{b}^{4,2}$  cancels out and we obtain

$$\mathbf{b}^{4,2} = - (2 \operatorname{div}_{\mathcal{S}}((\mathcal{H} \mathbf{P} - \mathcal{B}) \nabla_{\mathcal{S}} \mathcal{H}) + \mathcal{H} \mathcal{H}^2) \mathbf{v}.$$

Putting everything together yields eq. (4). For the derivation of the other parts of the model we refer to<sup>17,41</sup>.

### 5.2 Numerical method

We consider a surface finite element method (SFEM)<sup>51,52</sup> to solve the highly nonlinear set of geometric and surface partial differen-



tial equations (1)-(3), using the approaches in<sup>17,35</sup>.

We combine the system (1)-(3) with a mesh redistribution approach<sup>61</sup>. These are equations for the parametrization

$$\partial_t \mathbf{X} \cdot \mathbf{v} = \mathbf{u} \cdot \mathbf{v} \quad (17)$$

$$\mathcal{H} \mathbf{v} = \Delta_C \mathbf{X}, \quad (18)$$

which generate a tangential mesh movement to maintain the shape regularity and additionally provide an implicit representation of the mean curvature  $\mathcal{H}$ . We consider a discrete  $k$ -th order approximation  $\mathcal{S}_h^k$  of  $\mathcal{S}$ , with  $h$  the size of the mesh elements, i.e. the longest edge of the mesh. We use the DUNECurvedGrid library<sup>62</sup> and consider each geometrical quantity like the normal vector  $\mathbf{v}_h$ , the shape operator  $\mathcal{B}_h$ , the Gaussian curvature  $\mathcal{K}_h$ , and the inner products  $(\cdot, \cdot)_h$  with respect to  $\mathcal{S}_h^k$ . In the following, we find  $(\mathbf{u}_h^n, p_h^n, \mathcal{H}_h^n, \mathbf{Y}^n) \in [\mathbf{V}_3 \times V_2 \times V_3 \times \mathbf{V}_3](\mathcal{S}_h^{n-1})$  such that:

$$\begin{aligned} \frac{1}{\tau} \left( \mathbf{u}_h^n - \mathbf{u}_h^{n-1}, \mathbf{v}_h \right)_h + \left( \nabla_{\mathbf{w}_h^{n-1}} \mathbf{u}_h^n, \mathbf{v}_h \right)_h &= (p_h^n, \operatorname{div} \mathbf{p} \mathbf{v}_h)_h - \frac{2}{\operatorname{Re}} \left( \boldsymbol{\sigma}(\mathbf{u}_h^n), \nabla \mathbf{p} \mathbf{v}_h \right)_h - \gamma \left( \mathbf{u}_h^n, \mathbf{v}_h \right)_h \\ &+ \left( (2k_{2,0} \nabla_{\mathcal{S}} \mathcal{H}_h^n), \nabla_{\mathcal{S}}(\mathbf{v}_h \cdot \mathbf{v}_h^{n-1}) \right)_h \\ &+ \left( 2k_{2,0}(\mathcal{H}_h^n - \mathcal{H}_0^{n-1}) \mathbf{B}^{n-1} \mathbf{v}_h^{n-1}, \mathbf{v}_h \right)_h + \lambda \left( \mathbf{v}_h^{n-1}, \mathbf{v}_h \right)_h \\ &+ \left( k_{4,0}(\mathcal{H}_h^n - \mathcal{H}_0^{n-1})(\mathcal{H}_h^{n-1} - \mathcal{H}_0^{n-1})^2 \left( 4\mathcal{B}^{n-1} - (\mathcal{H}_h^{n-1} - \mathcal{H}_0^{n-1}) \mathbf{P} \right), \nabla_C \mathbf{v}_h \right)_h \\ &+ \left( 12k_{4,0}(\mathcal{H}_h^{n-1} - \mathcal{H}_0^{n-1})^2 \mathbf{v}_h^{n-1} \otimes \nabla_{\mathcal{S}} \mathcal{H}, \nabla_C \mathbf{v}_h \right)_h \\ &+ \left( 2k_{4,2} \mathcal{H}_h^n \mathbf{v}_h^{n-1} \otimes (\mathbf{P} \nabla_{\mathcal{S}} \mathcal{H}_h^{n-1}), \nabla_C \mathbf{v}_h \right)_h \\ &- \left( 2k_{4,2} \mathbf{v}_h^{n-1} \otimes (\mathcal{B}^{n-1} \nabla_{\mathcal{S}} \mathcal{H}_h^{n-1}), \nabla_C \mathbf{v}_h \right)_h + \left( k_{42}(\mathcal{H}_h^{n-1})^2 \mathbf{P}, \nabla_C \mathbf{v}_h \right)_h \\ (\operatorname{div} \mathbf{p} \mathbf{u}_h^n, q_h)_h &= 0 \\ \frac{1}{\tau} \left( \mathbf{Y}^n \cdot \mathbf{v}_h^{n-1}, h_h \right)_h &= \left( \mathbf{u}_h^n \cdot \mathbf{v}_h^{n-1}, h_h \right)_h \\ \left( \mathcal{H}_h^n \mathbf{v}_h^{n-1}, \mathbf{Z}_h \right)_h + (\nabla_C \mathbf{Y}^n, \nabla_C \mathbf{Z}_h)_h &= - \left( \nabla_C \mathbf{X}^{n-1}, \nabla_C \mathbf{Z}_h \right)_h \end{aligned}$$

for all  $(\mathbf{v}_h, q_h, h_h, \mathbf{Z}_h) \in [\mathbf{V}_3 \times V_2 \times V_3 \times \mathbf{V}_3](\mathcal{S}_h^{n-1})$ , where  $\mathbf{B}^{n-1} = \left( \|\mathcal{B}_h^{n-1}\|^2 - \frac{1}{2} \operatorname{tr} \mathcal{B}_h^{n-1} (\operatorname{tr} \mathcal{B}_h^{n-1} - \mathcal{H}_0(\phi_h^n)) \right)$ .

In the above formulation we used the identity  $(-\nabla_{\mathcal{S}} p_h^n - p_h^n \mathcal{H}_h^n \mathbf{v}_h, \mathbf{v}_h)_h = (p_h^n, \operatorname{div} \mathbf{p} \mathbf{v}_h)_h$ . Note that the Lagrange multiplier  $\lambda$  is unknown, which leads to an underdetermined system. To resolve that problem, we follow the approach introduced in<sup>35</sup>. In order to fulfill the volume constraint,  $\lambda$  has to be chosen such that

$$\Phi(\lambda) := \int_{\mathcal{S}_h^{n-1}} \mathbf{u}_h^n(\lambda) \cdot \mathbf{v}_h d\mathcal{S} = 0. \quad (19)$$

We consider  $\Phi(\lambda) = 0$  as an equation in  $\lambda$  and apply a Newton iteration  $\lambda^{j+1} = \lambda^j - \Phi(\lambda^j)/\Phi'(\lambda^j)$ . After convergence is achieved the new surface  $\mathcal{S}_h^n$  needs to be computed by updating the parametrization  $\mathbf{X}^n = \mathbf{X}^{n-1} + \mathbf{Y}^n$ , lifting the solutions  $\mathbf{u}_h^n$ ,

will drop the index  $k$ . We define the discrete function spaces for scalar functions by  $V_k(\mathcal{S}_h) = \{\psi \in C^0(\mathcal{S}_h) | \psi|_T \in \mathcal{P}_k(T)\}$  and for vector fields by  $\mathbf{V}_k(\mathcal{S}[h]) = [V_k(\mathcal{S}[h])]^3$ . Within these definitions  $T$  is the mesh element and  $\mathcal{P}_k$  are the polynomials of order  $k$ . We consider  $\mathbf{u}_h, \mathbf{X} \in \mathbf{V}_3(\mathcal{S}_h)$ ,  $\mathcal{H}_h \in V_3(\mathcal{S}_h)$ , and  $p_h \in V_2(\mathcal{S}_h)$ , which leads to an isogeometric setting for the velocity and a  $\mathcal{P}_3 - \mathcal{P}_2$  Taylor-Hood element for the surface Navier-Stokes-like equations. We discretize in time using constant time stepping with step size  $\tau$ . In each time step we solve the surface Navier-Stokes-like equations and the mesh redistribution together. We define a discrete surface update variable  $\mathbf{Y}^n = \mathbf{X}^n - \mathbf{X}^{n-1}$ , which is considered as unknown instead of the surface parametrization  $\mathbf{X}^n$ . The system to solve reads:

$p_h^n$  and  $\mathcal{H}_h^n$  to the new surface and computing the remaining geometric quantities  $\mathbf{v}_h^n, \mathcal{B}_h^n, \nabla_{\mathcal{S}} \mathcal{H}_h^n, \mathcal{K}_h^n$  and  $\nabla_{\mathcal{S}} \mathcal{K}_h^n$  for the new surface. While this approach showed the (expected) optimal order of convergence<sup>35</sup>, with respect to computational and numerical analysis results for the underlying surface Stokes equations on stationary surfaces<sup>63,64</sup> for bending terms up to second order, the approach is not sufficient if higher order geometric terms are included.

We therefore introduce a smoothing step of the surface quantities  $\mathcal{H}_h^n, \nabla_{\mathcal{S}} \mathcal{H}_h^n, \mathcal{K}_h^n$  and  $\nabla_{\mathcal{S}} \mathcal{K}_h^n$ . For each surface quantity or its components  $a_h = \mathcal{H}_h^n, [\nabla_{\mathcal{S}} \mathcal{H}_h^n]_i, \mathcal{K}_h^n$  and  $[\nabla_{\mathcal{S}} \mathcal{K}_h^n]_i, i = 1, 2$  we






Fig. 7 (a) Reference surface for the tests. Depicted are the mean curvature  $\mathcal{H}$  and Gaussian curvature  $\mathcal{K}$ . (b) Errors of geometric quantities with and without additional smoothing step for different grid widths  $h$ . The orders of convergence are indicated by the dashed lines and are optimal orders for the numerical implementation.

solve one time step of the diffusion equation

$$a_s - \varepsilon \Delta_{\mathcal{S}} a_s = a_h,$$

where  $\varepsilon > 0$  is a smoothing parameter and  $a_s$  the smoothed surface quantity.

### 5.3 Validation

Instead of a full convergence study of the numerical approach we only test the smoothing of surface quantities. We consider a surface for which the surface quantities can be computed analytically. The surface is parametrized by  $\mathbf{X} : [0, 2\pi) \times [0, \pi) \rightarrow \mathbb{R}^2$ ,

$$\mathbf{X}(\phi, \theta) = \begin{pmatrix} \frac{1}{4} + \frac{3}{4} \cos^2 \theta \sin \phi \sin \theta \\ \cos \phi \left( \frac{1}{4} + \frac{3}{4} \cos^2 \theta \sin \theta \right) \\ \cos \theta \end{pmatrix}.$$

We compute the  $L^2$ -error of the surface quantities  $\mathcal{H}$ ,  $\mathcal{K}$ ,  $\nabla_{\mathcal{S}} \mathcal{H}$  and  $\nabla_{\mathcal{S}} \mathcal{K}$  for different grid width  $h$ . The smoothing parameter  $\varepsilon$  is chosen experimentally such that it shows optimal results. Fig. 7 shows that for this test case the surface quantities converge with the optimal orders and that the additional smoothing step for all quantities improves the approximation. Together with the convergence studies in<sup>17,35</sup> these results provide enough confidence in the numerical approach for the full problem including the higher order geometric terms. They require an appropriate resolution of  $\mathcal{H}$  and  $\nabla_{\mathcal{S}} \mathcal{K}$ , which is achieved in  $O(h^2)$  and  $O(h)$ , respectively. This motivated the considered discrete function spaces.

### 5.4 Hausdorff distance

For two sets  $X, Y \subset \mathbb{R}^3$ , we consider the Hausdorff distance given by

$$d_H(X, Y) = \max \left\{ \sup_{x \in X} d(x, Y), \sup_{y \in Y} d(X, y) \right\},$$

where  $d(x, M) = \sup_{m \in M} \|x - m\|$  and  $\|\cdot\|$  the Euclidean norm. In the implementation we use the *VTK Hausdorff distance point set filter*<sup>65</sup> with the target distance method point to cell.



Data are available from Zenodo at <https://10.5281/zenodo.14503545>.

View Article Online  
DOI: 10.1039/D4FD00202D

Open Access Article. Published on 03 February 2025. Downloaded on 2/21/2025 12:44:58 PM.  
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

