




 Cite this: *RSC Adv.*, 2024, 14, 26233

Fatty aldehyde bisulfite adducts as a purification handle in ionizable lipid synthesis†

 Graham Atwood, Sona Purbiya, Cassandra Reid, Brandon Smith, Kuljit Kaur, Drew Wicks, Peter Gaudet, K. Cory MacLeod  and Jean-Francois Vincent-Rocan *

Rapid access to ALC-0315, a crucial component of the formulated Pfizer Covid vaccine, was obtained by employing solid adduct formation and filtration after an oxidation step in place of the standard chromatographic separation, allowing for a more scalable synthesis. Impurities were removed by formation of this fatty aldehyde bisulfite adduct at the penultimate step and by performing the final reductive amination directly with the fatty aldehyde bisulfite adduct. This eliminates chromatographic separations for all prepared aldehyde containing intermediates. Along with ALC-0315, FTT5 and SM-102 ionizable lipids were prepared utilizing this strategy. This work paves the way for more sustainable access to these critical ionizable lipids that would de-risk the world supply of important vaccines and medicines in the future.

 Received 17th July 2024
 Accepted 14th August 2024

DOI: 10.1039/d4ra05189k

rsc.li/rsc-advances

Introduction

Lipid nanoparticles (LNPs) play a crucial role in the formulation of nucleic acid derived medicines. Ionizable lipids are the major component of these LNPs, and their chemical and physical properties are responsible for the entrapment and delivery of the payload.^{1–5} These lipids are required in significantly larger quantities than the nucleic acids they carry. However, their synthesis includes major drawbacks from a process chemistry perspective such as excessive use of chlorinated solvents and multiple column chromatography operations. ALC-0315 **1** is the ionizable lipid in the Pfizer-BioNTech Covid-19 vaccine, Comirnaty, (0.43 mg per dose) and an estimated 4.6 billion doses were shipped to 181 countries.⁶ Therefore, since 2020 more than 2 MT of ALC-0315 was produced. Herein, we present our effort to make ionizable lipid production more sustainable by developing a route that relies on bisulfite adduct formation and filtration rather than chromatographic purification of intermediates to obtain analytically pure lipid.

There are select few syntheses of ALC-0315 **1** that are reported (Scheme 1).^{7–10} However, they all have significant drawbacks from a process chemistry perspective, using skin sensitizing reagents, chlorinated solvents and having a low control on impurity management requires the use of column chromatography purification at multiple steps. To overcome these challenges and allow for a more sustainable and


scalable synthesis, it was envisioned that the desired lipid could be obtained *via* a fatty aldehyde bisulfite adduct intermediate **4** of the analogous aldehyde **3** which would itself act as a purification handle in the synthesis. Additionally, the bisulfite adduct can be used directly in the next reductive amination step to produce **1**.

To generate the required aldehyde **3**, an acid-catalyzed esterification using the non-symmetrical 6-bromo-1-hexanol could be employed to avoid bis-ester formation. Whereas, prior reports relied on non-selective esterification of symmetrical diols, which typically requires the use of chromatographic separation of the undesired bis-ester byproduct or the use of a large molar excess of diol starting material relative to the carboxylic acid in order to disfavor bis-ester formation. The obtained bromo-ester **2** could then undergo a Ganem type oxidation to furnish the aldehyde **3**.^{11,12} This would allow the aldehyde **3** to be purified *via* bisulfite adduct formation. Alternatively, the direct alkylation of the amine with the bromo-ester **2** to form ALC-0315 **1** is known in the literature but produces the quaternary ammonium salt as a side-product which is not easily removed or controlled (Scheme 1, conditions b).⁷ Finally, a modification of the known reductive amination would provide ALC-0315 **1** by direct reaction of the fatty aldehyde bisulfite adduct **4** in 2-MeTHF as an alternative to the commonly used CH₂Cl₂, increasing the selectivity without needing to add solubilizing groups such as silyl ethers (Scheme 1, conditions e)⁸ which can increase step count as well as unit operations dramatically. These modifications greatly improve the yield of the reductive amination to those reported in the literature.^{7–10} The strategy of purification *via* solid fatty aldehyde bisulfite adduct followed by reductive amination directly with the

BioVectra Inc., Charlottetown, Prince Edward Island C1E 0A1, Canada. E-mail: jvincent-rocan@biovectra.com

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4ra05189k>



Chart 1 Lipids ALC-0315 **1**, FTT5 **5** and SM-102 **6**.

adduct to the free aldehyde and bisulfite anion can cause the degradation of the adduct over time as acetone can also form a bisulfite adduct with the *in situ* generated bisulfite anion. Therefore, alternative anti-solvents were evaluated to avoid loss of the bisulfite adduct and to further improve the purification.

The results of a stability test of the adduct **4** indicate complete degradation of adduct and loss of product in acetone at extended contact times. Changing from acetone to ethyl formate improves the stability of the bisulfite adduct, as determined by ^1H NMR (Fig. S1 \dagger), while still enabling filtration. This is likely due to the similar size and polarity of ethyl formate, while not reacting with the *in situ* generated bisulfite anions as acetone does. Additionally, spiking experiments also indicate that water does not have an impact on bisulfite stability (mechanistic detail Scheme S3 \dagger).

Upon addition of ethyl formate to the solid bisulfite material, a brown solution is obtained along with an easily filterable light-




Chart 2 Structurally similar ALC-0315 impurities.

tan colored solid giving 23.86 g (73% yield) of **4** as a light-brown to tan solid from the starting bromo-ester **2**. This tan solid is noted to be hygroscopic as before, and can be stored either under nitrogen, or in a desiccator. A significant amount of semi-solids are found to have collected on the underside of the filter paper, slowing filtration. This is likely due to the relatively low boiling point of ethyl formate and the low pressure experienced after passing through the filter. A simple change to DMC negates this issue and allows for a more rapidly filtering mixture.

The isolated adduct **4** is readily converted to the free aldehyde *via* washing with 10% sodium carbonate solution and extracting into EtOAc. The free aldehyde product is obtained with good purity (85–88%, HPLC-CAD) and can be used in the final reductive amination step. With the free aldehyde in hand from **4**, the reductive amination to generate ALC-0315 **1** is performed *via* a dosed addition of the free aldehyde with portion-wise addition of the reducing agent, $\text{NaBH}(\text{OAc})_3$. Portion-wise addition of the reducing agent is needed to overcome process limitations.⁸ These conditions give purity of ALC-0315 **1** in the isolated crude oil of 68% as analyzed by HPLC-CAD. Alternatively, the reductive amination can be performed directly from the isolated solid bisulfite adduct **4** (Scheme 1, conditions o) to give an improved crude purity (86% HPLC-CAD), while also allowing 2-MeTHF to replace CH_2Cl_2 as the solvent. Column chromatography gives purified ALC-0315 **1** as a clear slightly yellow oil (53% isolated yield, 94.9% HPLC-CAD). This represents a 37% overall yield from the starting hexyldecanoic acid.

The main impurities in the final isolated ALC-0315 **1** are two structurally related ether impurities, **7** and **8** (Chart 2), which stems from an ether impurity present in the 6-bromo-1-hexanol starting material. This ether impurity in the esterification starting material culminates in two structurally similar compounds to ALC-0315 which are not easily separated *via* column chromatography. Therefore, an alternative synthetic



route was evaluated, avoiding 6-bromo-1-hexanol which contains the impurity.

A straightforward change of the first two steps eliminated the dependence on the problematic starting material and avoids the structurally similar impurities, **7** and **8**, albeit with the introduction of a coupling reagent. An initial EDC coupling of hexyldecanoic acid and 1,6-hexandiol gives the mono-esterified diol **9** in high yield and good purity (94%, 88.0% HPLC-CAD), with the main impurity being the bis-ester byproduct (9.5%), which can be easily removed in a later step during purification of the aldehyde bisulfite-adduct. Alcohol-ester **9** is then oxidized to aldehyde **3** with PIDA in the presence of cat. TEMPO.^{24–26} The crude aldehyde **3** is then subjected to the general bisulfite adduct purification conditions. This gives crystalline off-white solid which filters rapidly to give the purified product as the sodium bisulfite adduct **4** (Fig. 1) in a 90% yield (83% purity by HPLC-CAD) from **9**, and overall yield of 85% from hexyldecanoic acid. This solid isolate eliminates the need for a chromatographic purification prior to its use in the following step, significantly reducing waste that is typically generated in such chromatographic operations (Table S4†). The adduct **4**, as before, is subjected to the improved reductive amination conditions and column purified to give ALC-0315 **1** in a 64% yield for the final reductive amination step and 54% overall yield of ALC-0315 **1** from the starting material, hexyldecanoic acid. This yield is an improvement when compared to other literature preparations of ALC-0315 **1** (20–47%, Scheme 1)^{7–9} as well as patented procedures (50%).¹⁰ The culmination of the process yields ALC-0315 **1** with a purity of 97.3% as analyzed by HPLC-CAD with no structurally similar impurities, **7** or **8**,




Fig. 1 Photo of solid fatty bisulfite adduct **4** produced from **9**.





Fig. 2 Photo of solid fatty bisulfite adduct **13**.

present. This material was subsequently used in a formulation preparation, encapsulating mRNA producing LNPs which match the physicochemical properties of formulation batches that utilize commercially available ALC-0315 (Table S6†).

With the ability to synthesize ALC-0315 **1** in this manner, FTT5 **5**, a proven lipid *in vivo*,¹³ and SM-102 **6** were the next targets for evaluating the route's feasibility across multiple lipids.


Preparation of the bisulfite adduct intermediate **13** of FTT5 **5** follows analogously to **4** (Schemes 2 and S1† for full route) and gives a well behaving flaky tan solid (Fig. 2). This, again, eliminates the need for a chromatographic separation of the aldehyde intermediate. Direct reaction of amine **10** with bisulfite adduct **13** gives FTT5 **5** (26%), closely matching the literature yield (27%).¹³ The inclusion of IPA in the reaction is required to solubilize the amine **10**, otherwise a very slow reaction is observed.

The synthetic route to SM-102 **6** proceeds through bisulfite adduct **19**, after oxidation of bromo-ester **18** (Schemes 3 and S2† for full route). Isolation of the bisulfite adduct **19** gives a well behaving off-white to tan solid (Fig. 3). After direct reaction of bisulfite adduct **19** with amine **15** and column purification, SM-102 **6** is obtained in good yield and purity (67% yield, 96.0% purity by HPLC-CAD). As with ALC-0315 **1** and FTT5 **5**, SM-102 **6** is readily prepared *via* a solid aldehyde bisulfite adduct intermediate, enabling the elimination of chromatographic purification of the aldehyde intermediate, as well as the use of the bisulfite adduct **19** directly in the final step to produce the lipid.



Scheme 2 Synthetic route through solid bisulfite adduct **13** to lipid FTT5 **5**.





Scheme 3 Synthetic route through solid bisulfite adduct 19 to lipid SM-102 6.




Fig. 3 Photo of solid fatty bisulfite adduct 19.

Experimental

Synthetic procedures

General procedure for synthesis of bromo-esters from bromo-alcohols. To a 3-neck RBF with attached Dean–Stark (open at top) and nitrogen inlet is added the acid (1 eq.), followed by toluene (5 V), bromo-alcohol (0.98 eq.), and *p*-TSA (0.1 eq.). This is brought up to a reflux with a very slight nitrogen flow. Upon collection of distillates in the Dean–Stark, the Dean–Stark is filled to just below overflow with toluene and the reaction is allowed to reflux for 16 hours or until the reaction is complete *via* ^1H NMR before it is cooled to room temperature. The reaction mixture is transferred to a separatory funnel and the RBF is washed with 5 V sat. NaHCO_3 . This is transferred into the separatory funnel, shaken, and allowed to separate. The organic layer is separated, dried (Na_2SO_4) and concentrated to give a slightly yellow to light brown oil.

General procedure for synthesis of bromo-esters from bromo-acids. To a 3-neck RBF with attached Dean–Stark (open at top) and nitrogen inlet is added the bromo-acid (0.98 eq.), followed by toluene (5 V), alcohol (1 eq.), and *p*-TSA (0.1 eq.). This is brought up to a reflux with a very slight nitrogen flow. Upon collection of distillates in the Dean–Stark, the Dean–Stark is filled to just below overflow with toluene and the reaction is allowed to reflux for 16 hours or until the reaction is complete *via* ^1H NMR before it is cooled to room temperature. The reaction mixture is transferred to a separatory funnel and the RBF is

washed with 5 V sat. NaHCO_3 . This is transferred into the separatory funnel, shaken, and allowed to separate. The organic layer is separated, dried (Na_2SO_4) and concentrated to give a slightly yellow to light brown oil.

General procedure for synthesis of fatty aldehyde bisulfite adducts. To a RBF with attached condenser is added pyridine *N*-oxide (6.0 eq.), *n*-propyl acetate (9 V), and sodium acetate (2.0 eq.). This is brought to a reflux before the bromo-ester is added, as a solution in *n*-propyl acetate (1 V), to the refluxing solution over 4 hours. After addition is complete the reaction is allowed to reflux for a further 16 hours before cooling to room temperature. To the cooled mixture is added HPW (10 V). This is shaken, separated and organic is washed with sat. NaHCO_3 (10 V) before drying over Na_2SO_4 and filtering. To the *n*-propyl acetate solution at 35 °C is added sodium metabisulfite as a solution in water (0.45 g mL^{-1}) dropwise. This is allowed to stir at 35 °C for 30 minutes before concentrating *in vacuo*. After concentrating an additional 10 V *n*-propyl acetate is charged and concentrated giving a solid material. To the solid is added ethyl formate or DMC (5 V) and the mixture is stirred at room temperature until all the solid material has formed a white to tan solid suspension in the mixture. This mixture is filtered, washed with ethyl formate or DMC (2 V) and the solids are collected and dried in a vacuum oven for 16 hours with a slight nitrogen bleed before weighing the collected bisulfite adduct.

General procedure for synthesis of ionizable lipids *via* reductive amination of fatty aldehyde bisulfite adducts. To a RBF with attached nitrogen inlet is added bisulfite adduct (2.3 eq.), and 2-MeTHF (10 V to the adduct) at RT. This is allowed to stir until a homogeneous mixture is observed. NEt_3 (2.4 eq.) is added to the reaction followed by the reactive amine (1 eq.) and then NaBH(OAc)_3 (4.3 eq.). The reaction is stirred for 16 h at RT before the reaction solution is washed with sat. Na_2CO_3 (10 V to adduct). The organic layer is dried (Na_2SO_4), filtered, and concentrated before being purified *via* column chromatography (SiO_2).

Conclusions

A general route has been developed for multiple commonly used ionizable lipids which have been prepared through solid fatty aldehyde bisulfite adduct intermediates, eliminating the need for column purification of intermediates in these multi-step syntheses. Both ALC-0315 1 and SM-102 6, two ionizable lipids being used in COVID-19 vaccine drug products, as well as FTT5 5, a relatively recent lipid-like compound which has seen



in vivo success, have been prepared by employing fatty aldehyde bisulfite adducts, which allows for easy isolation of the resulting solid intermediates. Additionally, the bisulfite adducts can be used directly in subsequent reductive amination reactions, in which they performed equally or better than the free aldehyde while also eliminating the need for CH_2Cl_2 in the reaction. We believe this to be the first reported isolation of solid fatty aldehyde bisulfite adduct intermediates and their use in the production of ionizable lipids.

The production of ALC-0315 **1** *via* a solid fatty aldehyde bisulfite adduct intermediate enabled an improved purification process to the typical column chromatography, improved yield when compared to other syntheses, and was found to perform equally when formulated with mRNA as other commercial sources of ALC-0315. Isolation of solid fatty aldehyde intermediates *via* bisulfite adduct formation and filtration is viewed as a general approach to improve synthesis of other lipids and the isolation and purification of other fatty alkyl aldehyde intermediates.

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors have no acknowledgments.

Notes and references

- X. Hou, T. Zaks, R. Langer and Y. Dong, *Nat. Rev. Mater.*, 2021, **6**, 1078–1094.
- L. Xu, X. Wang, Y. Liu, G. Yang, R. J. Falconer and C.-X. Zhao, *Adv. NanoBiomed Res.*, 2022, **2**, 2100109.
- R. H. Müller, K. Mäder and S. Gohla, *Eur. J. Pharm. Biopharm.*, 2000, **50**, 161–177.
- R. Paliwal, S. R. Paliwal, R. Kenwat, B. D. Kurmi and M. K. Sahu, *Expert Opin. Ther. Pat.*, 2020, **30**, 179–194.
- M. Jeong, Y. Lee, J. Park, H. Jung and H. Lee, *Adv. Drug Delivery Rev.*, 2023, **200**, 114990.
- L. Schoenmaker, D. Witzgmann, J. A. Kulkarni, R. Verbeke, G. Kersten, W. Jiskoot and D. J. A. Crommelin, *Int. J. Pharm.*, 2021, **601**, 120586.
- I. A. Boldyrev, V. P. Shendrikov, A. G. Vostrova and E. L. Vodovozova, *Russ. J. Bioorg. Chem.*, 2023, **49**, 412–415.
- F. Saadati, S. Cammarone and M. A. Ciufolini, *Chem.–Eur. J.*, 2022, **28**, e202200906.
- J. B. Wolf, J. Weon Lee, M. B. Plutschack, D. Cambié, A. Seidel-Morgenstern and P. H. Seeberger, *React. Chem. Eng.*, 2024, **9**, 959–966.
- N. M. Do, S. A. Eisenbeis and O. A. Salman, Methods for Producing of Lipids, WO2022215002A1, World Intellectual Property Organization, 2022.
- A. G. Godfrey and B. Ganem, *Tetrahedron Lett.*, 1990, **31**, 4825–4826.
- Z. Chen, N. Salehi Marzijarani, S. Quirie, G. F. Pirrone, S. M. Dalby, T. Wang, J. Kim, F. Peng and A. J. Fine, *Org. Process Res. Dev.*, 2022, **26**, 525–532.
- X. Zhang, W. Zhao, G. N. Nguyen, C. Zhang, C. Zeng, J. Yan, S. Du, X. Hou, W. Li, J. Jiang, B. Deng, D. W. McComb, R. Dorkin, A. Shah, L. Barrera, F. Gregoire, M. Singh, D. Chen, D. E. Sabatino and Y. Dong, *Sci. Adv.*, 2020, **6**, eabc2315.
- K. J. McKnelly, W. Sokol and J. S. Nowick, *J. Org. Chem.*, 2020, **85**, 1764–1768.
- E. Fischer and A. Speier, *Ber. Dtsch. Chem. Ges.*, 1895, **28**, 3252–3258.
- Z. Khan, F. Javed, Z. Shamair, A. Hafeez, T. Fazal, A. Aslam, W. B. Zimmerman and F. Rehman, *J. Ind. Eng. Chem.*, 2021, **103**, 80–101.
- J. C. Graham, A. Trejo-Martin, M. L. Chilton, J. Kostal, J. Bercu, G. L. Beutner, U. S. Bruen, D. G. Dolan, S. Gomez, J. Hillegass, J. Nicolette and M. Schmitz, *Chem. Res. Toxicol.*, 2022, **35**, 1011–1022.
- T. M. Bass, D. Zell, S. M. Kelly, T. C. Malig, J. G. Napolitano, L. E. Sirois, C. Han and F. Gosselin, *Org. Process Res. Dev.*, 2024, **28**, 2635–2645.
- C. R. Pandit and N. S. Mani, *Synthesis*, 2009, **2009**, 4032–4036.
- M. Barniol-Xicotá, A. L. Turcu, S. Codony, C. Escolano and S. Vázquez, *Tetrahedron Lett.*, 2014, **55**, 2548–2550.
- X. Li, K. S. Iyer, R. R. Thakore, D. K. Leahy, J. D. Bailey and B. H. Lipshutz, *Org. Lett.*, 2021, **23**, 7205–7208.
- M. G. Kissane, S. A. Frank, G. A. Renner, C. P. Ley, C. A. Alt, P. A. Stroud, R. K. Vaid, S. K. Boini, L. A. McKee, J. T. Vicenzi and G. A. Stephenson, *Tetrahedron Lett.*, 2013, **54**, 6587–6591.
- Y. Itoh, J. Horiuchi and K. Takahashi, *Colloids Surf., A*, 2007, **308**, 118–122.
- L. De Luca, G. Giacomelli and A. Porcheddu, *Org. Lett.*, 2001, **3**, 3041–3043.
- R. Ciriminna and M. Pagliaro, *Org. Process Res. Dev.*, 2010, **14**, 245–251.
- C. Guérin, V. Bellosta, G. Guillamot and J. Cossy, *Org. Lett.*, 2011, **13**, 3534–3537.

