RSC Advances



View Article Online

View Journal | View Issue

PAPER

Check for updates

Cite this: RSC Adv., 2024, 14, 33587

Synthesis of β-enamino malonates through caesium carbonate-promoted reaction of nitro-substituted donor-acceptor cyclopropanes†

A caesium carbonate-promoted reaction of nitro-substituted donor-acceptor cyclopropanes (DACs) with primary aromatic amines in water provides a convenient access to β -enamino malonates under mild

reaction conditions. The transformation takes place through the formation of allene intermediates from

the DACs followed by the conjugate addition of various primary aromatic amines to the intermediates.

The reaction proceeds more efficiently in water as compared with organic solvents and the products

were isolated in good yields by filtering through a silica gel column without any prior extraction procedure.

Sebastin Raj Jeny, Subramani Selvi, Devrugaiya Deerkadharshini and Kannupal Srinivasan *

Received 2nd August 2024 Accepted 16th October 2024

DOI: 10.1039/d4ra05619a

rsc.li/rsc-advances

Introduction

Donor-Acceptor Cyclopropanes (DACs) are one of the types of readily available and easily manipulable building blocks that give access to different classes of organic compounds.1 Simple reagents are often enough to trigger their characteristic reactions such as ring-opening,² annulation³ and rearrangement.⁴ DACs having an extra aroyl or nitro group (apart from the regular donor-acceptor groups) usually display distinct reactivity and afford products that are difficult to obtain by other methods.5 In this context, nitro-substituted DACs are renowned for their unique reactivity and they preferentially undergo ringopening reactions rather than annulation reactions.6 When treated with Lewis acids, nitro-substituted DACs form aroylmethylidene malonates as products or intermediates, which serve as versatile building blocks for a handful of heterocyclic and acyclic compounds.6c,7 On the other hand, when treated with bases, they form arylethylidenes or allenes as intermediates, which too serve as potential precursors for various compounds.8

Recently, we reported that nitro-substituted DACs when reacted with salicylaldehydes in the presence of caesium carbonate and a surfactant in water furnished a series of chromane derivatives through allene intermediates.⁹ During the course of the study, we observed that nitro-substituted DAC **1a** reacts with 2-iodophenol (2) under the same reaction conditions to give the respective Michael adduct **3** (Scheme 1, eqn (1)). The observation prompted us to investigate the reaction of nitrosubstituted DACs **1** with amines **4** for the synthesis of β -enamino malonates **5** (or **6**) (Scheme 1, eqn (2)).

It may be noted that enamines and their derivatives are versatile synthetic intermediates and are usually synthesized by the condensation of amines with carbonyl compounds,¹⁰ hydroamination of alkynes¹¹ or cross-coupling reaction between



Comparison with literature works (eqns. 3 and 4)



Scheme 1 Comparison of the present work with literature works.

School of Chemistry, Bharathidasan University, Tiruchirappalli-620024, Tamil Nadu, India. E-mail: srinivasank@bdu.ac.in; Tel: +91-431-2407053

[†] Electronic supplementary information (ESI) available: Copies of ¹H and ¹³C NMR spectra of all products. See DOI: https://doi.org/10.1039/d4ra05619a

amines and vinyl halides or triflates.¹² Although, a handful of methods are available for the synthesis of enamino esters such as β -enamino malonates,¹³ DACs have been rarely used as starting materials for their synthesis. Sopova and coworkers have used nitro-substituted DACs **1** for the access of enamino malonates **A** using ammonia in methanol (Scheme 1, eqn (3)).¹⁴ We record that our previous work involving synthesis of pyridopyrimidinones 7 from nitro-substituted DACs **1** and 2-aminopyridine (**6**) in water should have also taken place through the formation of the respective enamino malonate intermediates **B** (Scheme 1, eqn (4)).⁸⁶ In this context, the present work

represents a convenient green strategy for the access of β enamino malonates from amines and nitro-substituted DACs using water as solvent. It is interesting to note that water acts as a better solvent than organic solvents for certain reactions of nitro-substituted DACs, likely due to the involvement of polar intermediates.

Results and discussion

We began the study by reacting nitro-substituted DAC **1a** with various primary aromatic amines **4a–f** (Table 1, entries 1–6).



^a Isolated yield. ^b Complicated mixture of products. ^c No reaction.

Paper

Owing to the greenness of the reaction conditions reported in the previous work [caesium carbonate, sodium dodecylsulphate (20 mol%), water, rt],9 we adapted the same reaction conditions for the present transformation. Accordingly, we reacted DAC 1a with aniline 4a under the above reaction conditions for 12 h. Then, the crude reaction mixture (without extraction) was made into slurry by mixing with a little silica gel and charged on a silica gel column. After eluting the column with 10% EtOAc in hexane (filtration), the expected enamino malonate 5a was obtained in 87% yield (Table 1, entry 1). Out of curiosity, we also tested the suitability of organic solvents such as MeOH, EtOH, DCM, 1,2-DCE, THF, acetone, MeCN, DMF, DMSO and toluene for the transformation. Though the reaction takes place in MeOH and EtOH (the yields of 5a were 28% and 23%, respectively, in these solvents), it did not take place in other organic solvents. Next, we investigated the reaction of 1a with anilines having different electron donating, halogen and electron withdrawing substituents. With p-toluidine, the respective product enamino malonate 5b was obtained in 68% yield by adapting the same procedure (entry 2). The presence of strong electron donating or withdrawing groups on aniline was not tolerated in the transformation. Thus, the reaction of 1a with 4methoxyaniline (4c) or *p*-nitroaniline (4d) led to a complicated mixture of products (entries 3 and 4). The reaction proceeded smoothly with o-iodoaniline (4e) and 1-naphthylamine (4f) and the respective enamino malonates 5e and 5f were obtained in 73% and 76% yields, respectively (entries 5 and 6). Next, we investigated the scope of the transformation for nitro-substituted DACs 1b-j having different aryl groups with various primary aromatic amines (Table 1, entries 7-32). The presence of moderately electron donating methyl and isopropyl groups and also halogens on the aryl ring of DACs (1b, 1c and 1e-g) was well tolerated in the transformation and the respective enamino malonates were obtained in 71-94% yields (entries 7-14 and 17-25). At the same time, DACs having a good electron donating group such as methoxy on the aryl ring (1d) as well as an electron-rich aryl ring such as 2-thienyl (1j) were not compatible in the transformation and they formed complex mixtures of products, possibly due to the enhanced reactivity of the aryl rings (entries 15, 16, 31 and 32). Similarly, DAC 1h having a strong electron withdrawing nitro group also did not undergo the expected reaction owing to the destabilizing effect of the aryl ring on the 1,3-zwitterionic intermediate that would be formed from the cyclopropane during the course of the reaction (see the mechanism).^{6c,7a} We also reacted DAC 1i having a 1-naphthyl group with different anilines and the corresponding enamino malonates 5ab-5ad were formed in 78-93% yields (entries 28-30).

Next, we focused our attention on the reaction of DAC **1b** with aromatic secondary amines such as *N*-methylaniline and indoline and aliphatic amines such as benzylamine, diisopropylamine, pyrrolidine and morpholine. Unfortunately, none of them gave the expected enamino malonates. During the investigation of the scope of the transformation with different amines, surprisingly, we observed that when DACs **1b** and **1c** were reacted with benzotriazole (**9**), they afforded the isomeric



Scheme 2 Reaction of DACs 1b and 1c with benzotriazole (9)



Scheme 3 Plausible mechanism of the transformation.

enamines **10a** and **10b** in 79 and 85% yields, respectively (Scheme 2). However, the transformation did not work with benzimidazole.

We propose a mechanism outlined in Scheme 3 for the formation of β -enamino malonates from DACs 1 and aromatic amines. Caesium carbonate removes the acidic proton at C-3 of the DAC, which opens the cyclopropane ring to produce intermediate **P**. The intermediate **P** may loose the nitro group to form the allene intermediate **Q**.^{8c,9} The aza-Michael addition of amine 4 to **Q** followed by migration of double bond to a conjugate position may afford enamino malonate 5. In case of benzotriazole, the final double bond migration is not taking place.

We have also performed a gram scale synthesis of β -enamino malonate **5a** as a model substrate in order to evaluate various green chemistry metrics for the transformation (Table 2). Pleasingly, the transformation has a low *E*-factor (0.3648 kg waste per kg product) and high atom economy (88%), atom efficiency (73%), carbon efficiency (100%) and reaction mass efficiency (73%), indicating the greenness of the methodology.

In order to prove the synthetic utility of the products made in the present study, we subjected enamino malonate 5a to oxidation using SeO₂ in 1,4-dioxane under reflux conditions. Pleasingly, the reaction afforded the aroylmethylidene malonate derivative **11** in 81% yield (Scheme 4).



Scheme 4 Synthetic transformation of 5a into aroylmethylidene malonate 11.

 Table 2
 Evaluation of green chemistry metrics for the gram-scale synthesis of 5a



Entry	Chemicals	Molecular weight (g mol ^{-1})	Weight (mg)
Entry	Chemicais	Molecular weight (g mol)	weight (ing)
Substrate 1a	Diethyl 2-nitro-3-phenylcyclopropane-1,1-dicarboxylate	307.3020 (C ₁₅ H ₁₇ NO ₆)	1228.8
Substrate 4a	Aniline	93.13 (C_6H_7N)	372.52
Product 5a	Diethyl 2-[2-phenyl-1-(phenylamino)ethylidene]malonate	353.4116 (C ₂₁ H ₂₃ NO ₄)	1173.3
Base	Caesium carbonate	$325.82 (Cs_2CO_3)$	1433.61
Surfactant	Sodium dodecyl sulphate	288.38 (C ₁₂ H ₂₅ NaO ₄ S)	230.70
Solvent	Water		
Product yield $= 8$	3%		
E-factor = [(1228.	$(8 + 372.52) - 1173.3] \div 1173.3 = 0.3648$ kg waste per kg product		
Atom economy =	$353.4116 \div (307.3020 + 93.13) \times 100 = 88\%$		
Atom efficiency =	$(83 \times 88) \div 100 = 73\%$		
Carbon efficiency	$= 21 \div 21 = 100\%$		
Reaction mass eff	$iciency = [1173.3 \div (1228.8 + 372.2)] \times 100 = 73\%$		

Conclusions

In summary, we have developed a green procedure for the synthesis of β -enamino malonates from nitro-substituted DACs and primary aromatic amines. Water was found to be an efficient medium for the transformation and the products were isolated in good yields simply by passing the crude reaction mixture through a silica gel column by bypassing the extraction step. The reaction proceeds through caesium carbonate promoted ring-opening of the cyclopropane to form allene intermediates followed by the addition of amines. The operational simplicity, the use of water as medium, mild reaction conditions, simple purification procedure and reasonably wide substrate scope are some of the green aspects of the methodology.

Experimental section

General remarks

Melting points were determined by the open capillary tube method and are uncorrected. The ¹H and ¹³C NMR spectra were recorded on a 400 MHz NMR spectrometer. High-resolution mass spectra (ESI) were recorded on a Q-TOF mass spectrometer. X-ray crystallographic data were collected on a CCD diffractometer using graphite-monochromated Mo K α radiation. Thin layer chromatography (TLC) was performed on precoated alumina sheets and detected under UV light. Silica gel (200–400 mesh) was used for column chromatography. Cyclopropanes **1a–j** were prepared as per our reported procedure^{6c} and all are known compounds.^{6c,8a,15}

General procedure for the synthesis of $\beta\mbox{-enamino}$ malonates 5a–af and 10a–b

To a suspension of nitrocyclopropane dicarboxylates 1 (1.0 mmol) in water (2 mL) were added aromatic primary amine 4

(1.0 mmol), sodium dodecyl sulfate (58 mg, 0.2 mmol), and caesium carbonate (358 mg, 1.1 mmol) and stirred at room temperature for 12 h. Next, the reaction mixture was mixed with silica gel to make slurry, charged on a silica gel column and eluted using ethyl acetate/hexane (1:9 v/v) to obtain the pure product.

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors thank Science and Engineering Research Board (SERB), Council of Scientific and Industrial Research (CSIR) and MoE-RUSA 2.0, India for financial support and DST-FIST for instrumentation facilities at School of Chemistry, Bharathidasan University.

Notes and references

 (a) H. U. Reissig and R. Zimmer, *Chem. Rev.*, 2003, **103**, 1151– 1196; (b) M. Yu and B. L Pagenkopf, *Tetrahedron*, 2005, **61**, 321–347; (c) M. Y. Melnikov, E. M. Budynina, O. A. Ivanova and I. V. Trushkov, *Mendeleev Commun.*, 2011, **21**, 293–301; (d) T. F. Schneider, J. Kaschel and D. B. Werz, *Angew. Chem., Int. Ed.*, 2014, **53**, 5504–5523; (e) H. K. Grover, M. R. Emmett and M. A. Kerr, *Org. Biomol. Chem.*, 2015, **13**, 655–671; (f) P. Singh, R. K. Varshnaya, R. Dey and P. Banerjee, *Adv. Synth. Catal.*, 2020, **362**, 1447–1484; (g) Y. Xia, X. Liu and X. Feng, *Angew. Chem., Int. Ed.*, 2021, **60**, 9192–9204; (*h*) *Donor-Acceptor Cyclopropanes in Organic Synthesis*, ed. P. Banerjee and A. T. Biju, Wiley-VCH, Weinheim, 2024.

- 2 (a) E. M. Budynina, K. L. Ivanov, I. D. Sorokin and M. Y. Melnikov, Synthesis, 2017, 49, 3035–3068; (b)
 V. Pirenne, B. Muriel and J. Waser, Chem. Rev., 2021, 121, 227–263; (c) E. Reyes, U. Uria, L. Prieto, L. Carrillo and J. L. Vicario, Chem. Commun., 2024, 60, 7288–7298.
- 3 (a) V. K. Yadav and V. Sriramurthy, Angew. Chem., Int. Ed., 2004, 43, 2669–2671; (b) A. T. Parsons and J. S. Johnson, J. Am. Chem. Soc., 2009, 131, 3122–3123; (c) G. Sathishkannan and K. Srinivasan, Org. Lett., 2011, 13, 6003–6005; (d) S. Chakrabarty, I. Chatterjee, B. Wibbeling, C. G. Daniliuc and A. Studer, Angew. Chem., Int. Ed., 2014, 53, 5964–5968; (e) M. A. Zotova, R. A. Novikov, E. V. Shulishov and Y. V. Tomilov, J. Org. Chem., 2018, 83, 8193–8207; (f) S. Nicolai and J. Waser, Angew. Chem., Int. Ed., 2022, 61, e202209006; (g) L. Yang, H. Wang, M. Lang and S. Peng, Synthesis, 2023, 56, 389–398.
- 4 (a) J. Kaschel, T. F. Schneider, P. Schirmer, C. Maab, D. Stalke and D. B. Werz, *Eur. J. Org. Chem.*, 2013, 4539–4551; (b) H. Chen, J. Zhang and D. Z. Wang, *Org. Lett.*, 2015, 17, 2098–2101; (c) S. Thangamalar, M. Thangamani and K. Srinivasan, *Org. Biomol. Chem.*, 2022, 20, 3145–3153; (d) S. Jiru, Y. Fu and S. R. Wang, *Org. Lett.*, 2023, 25, 555–559; (e) F. L. Daniel and K. Srinivasan, *J. Org. Chem.*, 2024, 89, 5304–5313.
- 5 T. Selvi and K. Srinivasan, Isr. J. Chem., 2016, 56, 454-462.
- 6 (a) O. Litchits, D. Alberico, I. Zhakharian and A. B. Charette, J. Org. Chem., 2008, 73, 6838–6840; (b) S. S. So, T. J. Auvil,
 V. J. Garza and A. E. Mattson, Org. Lett., 2012, 14, 444–447;
 (c) T. Selvi and K. Srinivasan, J. Org. Chem., 2014, 79, 3653– 3658.
- 7 (a) T. Selvi and K. Srinivasan, Chem. Commun., 2014, 50, 10845–10848; (b) T. Selvi and K. Srinivasan, Adv. Synth. Catal., 2015, 357, 2111–2118; (c) T. Selvi, G. Vanmathi and

K. Srinivasan, *RSC Adv.*, 2015, 5, 49326–49329; (*d*) M. Meenakshi and K. Srinivasan, *Org. Biomol. Chem.*, 2022, **20**, 8741–8746.

- 8 (a) G. Yang, W. Liu, Z. He and Zh. He, Org. Lett., 2016, 18, 4936-4939; (b) S. Selvi and K. Srinivasan, Eur. J. Org. Chem., 2017, 5644-5648; (c) Z.-Y. Zhou, Z.-Y. Xu, Q.-Y. Shen, L.-S. Huang, X. Zhang, A.-B. Xia, D.-Q. Zu and Z.-Y. Xu, Chem. Commun., 2021, 57, 6424-6427.
- 9 S. R. Jeny and K. Srinivasan, J. Org. Chem., 2023, 88, 9395-9400.
- 10 (a) D. Sanchez, D. Bastida, J. Bures, C. Isart, O. Pineda and J. Vilarrasa, *Org. Lett.*, 2012, 14, 536–539; (b) D. Xin and K. Burgess, *Org. Lett.*, 2014, 16, 2108–2110.
- 11 (a) S. Kramer, K. Dooleweerdt, A. T. Lindhardt, M. Rottlander and T. Skrydstrup, *Org. Lett.*, 2009, **11**, 4208–4211; (b)
 J. Bahri, N. Tanbouza, T. Ollevier, M. Taillefer and F. Monnier, *Synthesis*, 2019, **51**, 2086–2090.
- 12 (*a*) J. R. Dehli, J. Legros and C. Bolm, *Chem. Commun.*, 2005, 973–986; (*b*) X. Ji, H. Huang, W. Wu, X. Li and H. Jiang, *J. Org. Chem.*, 2013, **78**, 11155–11162.
- 13 (a) N. D. Koduri, H. Scott, B. Hileman, J. D. Cox, M. Coffin, L. Glicksberg and S. R. Hussaini, Org. Lett., 2012, 14, 440– 443; (b) N. D. Koduri, B. Hileman, J. D. Cox, H. Scott, P. Houng, A. Robbins, K. Bowers, L. Tsebaot, K. Miao, M. Castaneda, M. Coffin, G. Wei, T. D. W. Claridge, K. P. Roberts and S. R. Hussaini, RSC Adv., 2013, 3, 181– 188; (c) N. D. Koduri, Z. Wang, G. Cannell, K. Cooley, T. M. Lemma, K. Miao, M. Nguyen, B. Frohock, M. Castaneda, H. Scott, D. Albinescu and S. R. Hussaini, J. Org. Chem., 2014, 79, 7405–7414; (d) P.-Q. Huang, W. Ou and J.-L. Ye, Org. Chem. Front., 2015, 2, 1094–1106; (e) P.-Q. Huang, W. Ou and J.-L. Ye, Eur. J. Org. Chem., 2017, 582–592.
- 14 A. S. Sopova, O. S. Bakova, K. L. Metelkina and V. V. Perekalin, *Zh. Org. Khim.*, 1975, 11, 68–71; *Chem. Abstr.*, 1975, 83, 9406h.
- 15 S. Thangamalar and K. Srinivasan, *Eur. J. Org. Chem.*, 2023, 26, e202201084.