

Cite this: *Chem. Sci.*, 2024, 15, 16222

All publication charges for this article have been paid for by the Royal Society of Chemistry

# <sup>57</sup>Fe nuclear resonance vibrational spectroscopic studies of tetranuclear iron clusters bearing terminal iron(III)–oxido/hydroxido moieties†

Jin Xiong,<sup>a</sup> Christopher Reed,<sup>b</sup> Barbara Lavina,<sup>cd</sup> Michael Y. Hu,<sup>c</sup> Jiyong Zhao,<sup>c</sup> Esen E. Alp,<sup>c</sup> Theodor Agapie<sup>\*,b</sup> and Yisong Guo<sup>\*,a</sup>

<sup>57</sup>Fe nuclear resonance vibrational spectroscopy (NRVS) has been applied to study a series of tetranuclear iron ([Fe<sub>4</sub>]) clusters based on a multidentate ligand platform (L<sup>3-</sup>) anchored by a 1,3,5-triarylbenzene linker and pyrazolate or (*tert*butylamino)pyrazolate ligand (PzNH<sup>t</sup>Bu<sup>-</sup>). These clusters bear a terminal Fe(III)–O/OH moiety at the apical position and three additional iron centers forming the basal positions. The three basal irons are connected with the apical iron center *via* a μ<sub>4</sub>-oxido ligand. Detailed vibrational analysis *via* density functional theory calculations revealed that strong NRVS spectral features below 400 cm<sup>-1</sup> can be used as an oxidation state marker for the overall [Fe<sub>4</sub>] cluster core. The terminal Fe(III)–O/OH stretching frequencies, which were observed in the range of 500–700 cm<sup>-1</sup>, can be strongly modulated (energy shifts of 20–40 cm<sup>-1</sup> were observed) upon redox events at the three remote basal iron centers of the [Fe<sub>4</sub>] cluster without the change of the terminal Fe(III) oxidation state and its coordination environment. Therefore, the current study provides a quantitative vibrational analysis of how the remote iron centers within the same iron cluster exert exquisite control of the chemical reactivities and thermodynamic properties of the specific iron site that is responsible for small molecule activation.

Received 23rd May 2024  
Accepted 8th September 2024

DOI: 10.1039/d4sc03396e

rsc.li/chemical-science

## Introduction

Metal–oxido/hydroxido (M–O/OH) species are key reactive intermediates involved in challenging oxidation reactions found in numerous biological pathways and organic synthesis.<sup>1–18</sup> Prominent examples include selective carbon–hydrogen (C–H) bond functionalization and water oxidation in biological photosynthesis where metal–oxido/hydroxido species enable the key steps of the reactions, *e.g.* C–H bond cleavage<sup>5,16–19</sup> and O–O bond formation.<sup>8,10,20–23</sup> Given the importance of these M–O/OH species, extensive research efforts have been devoted to reveal the geometric/electronic structural features that define their chemical reactivities. There is no doubt that the metal primary coordination sphere constituted by the direct metal-binding ligand set establishes the geometric and electronic structures of M–O/OH species and further determines their reactivities.<sup>1,16,24–28</sup> It is also increasingly clear in recent years that

the secondary coordination sphere critically tunes the chemical/physical properties of metal centers.<sup>29,30</sup> For example, Lu and coworkers have shown that by changing the composition of amino acids surrounding the copper centers in azurin, the copper redox potentials could be altered to cover the entire physiological range (>1 V).<sup>31</sup> Among the heme-containing proteins, hydrogen bonding interactions surrounding the heme-iron centers have shown to dictate protein function. For instance, in hemoglobins and myoglobins, the proximal protein residues to the heme center stabilize the O<sub>2</sub> binding,<sup>32–37</sup> but in cytochrome P450 dependent enzymes, a different set of proximal protein residues facilitate O–O bond cleavage instead.<sup>7,38–41</sup> In non-heme iron enzymes, H-bonding networks surrounding the iron center have also been shown to lower the energetic barriers for O<sub>2</sub> activation<sup>15</sup> and facilitate organic substrate oxidation.<sup>42,43</sup> For nonheme M–O/OH synthetic complexes, Borovik and coworkers have demonstrated that the basicity of Mn/Fe–O/OH moieties, and subsequently their C–H activation reactivity, can be modulated by tuning the surrounding H-bonding interactions.<sup>16,29,44,45</sup> The influence of the H-bonding interactions to the electronic properties and reactivities of Mn/Fe–O/OH moieties has also been reported by Fout and coworkers,<sup>46–50</sup> Chang and coworkers,<sup>51</sup> and Goldberg and coworkers.<sup>52</sup> In addition, Lewis acidic, redox inactive metal ions, have also been demonstrated to influence the reactivities of (Fe/Mn/Co)–O moieties.<sup>53–58</sup>

Compared to our understanding on the structure and reactivity modulation of M–O/OH species *via* changes from the first

<sup>a</sup>Department of Chemistry, Carnegie Mellon University, Pittsburgh, PA 15213, USA. E-mail: ysguo@andrew.cmu.edu

<sup>b</sup>Division of Chemistry and Chemical Engineering, California Institute of Technology, CA 91125, USA. E-mail: agapie@caltech.edu

<sup>c</sup>Advanced Photon Source, Argonne National Laboratory, Argonne, Illinois 60439, USA

<sup>d</sup>Center for Advanced Radiation Source, University of Chicago, Chicago, Illinois, 60439, USA

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc03396e>



and the second coordination spheres, the understanding of the influence of remote redox active metal centers to M–O/OH moieties is much less developed. Yet, in nature, multi-nuclear metal centers are commonly used to enable essential multi-electron/proton chemistry, such as water oxidation by the oxygen evolving complex (OEC) in photosystem II,<sup>8,59</sup> proton reduction/H<sub>2</sub> oxidation by the H-cluster and the Ni–Fe center in hydrogenases,<sup>60–63</sup> CO<sub>2</sub> reduction by the C-cluster of CODH,<sup>64</sup> O<sub>2</sub> reduction by the copper–iron–heme center in cytochrome c oxidase,<sup>65,66</sup> and N<sub>2</sub> reduction by iron–molybdenum cofactor (FeMco) in nitrogenase.<sup>67–69</sup> Some of these active sites display complex metal clusters with distal metal centers not directly involved in substrate binding. The impact of remote metal centers on substrate activation has been studied by some of us.<sup>70–74</sup> A multidentate ligand platform (L<sup>3–</sup>) anchored by a 1,3,5-triarylbenzene linker and pyrazolate or (*tert*butylamino) pyrazolate ligand (PzNH<sup>t</sup>Bu<sup>–</sup>) were applied to support a [Fe<sub>3</sub>Mn] cluster and a [Fe<sub>4</sub>] cluster bearing a μ<sub>4</sub>-O ligand and a terminal Mn<sup>II/III</sup>–OH<sub>x</sub> or Fe<sup>III</sup>–O/OH moiety at the apical position of the

cluster, respectively (Fig. 1).<sup>73,74</sup> Detailed thermodynamics and reactivity studies demonstrated that by only changing the redox state on the three remote base iron sites of the cluster, the bond dissociation free energy (BDE) of the O–H bond of the apical Mn<sup>II</sup>–OH<sub>2</sub> or Fe<sup>III</sup>–OH moieties can be tuned by a total of ~16 or 12 kcal mol<sup>–1</sup>, respectively.<sup>73,74</sup> It has also been estimated that the BDE of the O–H bond of the terminal Mn<sup>III</sup>–OH moiety could reach >100 kcal mol<sup>–1</sup> by the same remote iron site redox controlling strategy.<sup>73</sup> Clearly, the distal metal centers within the same metal cluster exert exquisite control of the chemical properties of the metal site that is responsible for small molecule binding and activation.

Herein, to further elucidate the impact of the remote iron site redox events towards the geometric and electronic structures of the terminal M–O/OH moiety in these multi-nuclear iron clusters, we have carried out a detailed spectroscopic and computational study on a series of [LFe<sub>3</sub>O(PzNH<sup>t</sup>Bu)<sub>3</sub>Fe(O/OH)]<sup>n+</sup> clusters.<sup>74–77</sup> <sup>57</sup>Fe nuclear resonant vibrational spectroscopy (NRVS) has been used to explore the vibrational features of




Fig. 1 Chemical structures of the [Fe<sub>4</sub>] clusters studied in this work and the selected mononuclear iron(III)–oxido/hydroxide complexes from Borovik and coworkers<sup>75–77</sup> that have been previously studied by <sup>57</sup>Fe NRVS.<sup>77</sup>



$[\text{LFe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{O}(\text{PzNH}^t\text{Bu})_3\text{Fe}^{\text{III}}(\text{O})]^+$  ( $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]-\text{O}$ ) and  $[\text{LFe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{O}(\text{PzNH}^t\text{Bu})_3\text{Fe}^{\text{III}}(\text{OH})]^{2+}$  ( $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]-\text{OH}$ ), and their corresponding one electron reduced clusters,  $[\text{LFe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{O}(\text{PzNH}^t\text{Bu})_3\text{Fe}^{\text{III}}(\text{O})]$  ( $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]-\text{O}$ ) and  $[\text{LFe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{O}(\text{PzNH}^t\text{Bu})_3\text{Fe}^{\text{III}}(\text{O})]^+$  ( $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]-\text{OH}$ ). The subsequent spectral analysis with the assistance of density functional theory (DFT) calculations not only revealed the unique vibrational features of these  $[\text{Fe}_4]$  clusters, but also pointed out the critical role of the  $\text{Fe}-\mu_4\text{O}$  bonding interactions among the four iron centers of the cluster in translating the impact of the redox event on the three remote basal iron sites to the apical terminal  $\text{Fe}^{\text{III}}-\text{O}/\text{OH}$  moiety. Such a redox modulation effect generated by remote iron site redox state change was also compared with the modulation effect of the hydrogen-bonding interactions exerted by the secondary coordination sphere towards  $\text{Fe}-\text{O}/\text{OH}$  moiety in mononuclear iron complexes to provide a broader perspective on the influence of the chemical environment of  $\text{Fe}-\text{O}/\text{OH}$  moieties in modulating their chemical reactivity.

## Results and discussions

We have recorded  $^{57}\text{Fe}$  NRVS spectra on both  $[\text{Fe}_4]$ -oxido ( $[\text{Fe}_4]-\text{O}$ ) and  $[\text{Fe}_4]$ -hydroxido ( $[\text{Fe}_4]-\text{OH}$ ) complexes in two oxidation states ( $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}_2]$  and  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_3]$ ). All the NRVS derived  $^{57}\text{Fe}$

PVDOS spectra are shown in Fig. 2, which exhibit strong spectral features in the region between  $150\text{ cm}^{-1}$  and  $400\text{ cm}^{-1}$  and a series of weak features in the region between  $400\text{ cm}^{-1}$  and  $800\text{ cm}^{-1}$ . Additionally, the strong spectral features show systematic red shifts ( $\Delta\nu \sim 10\text{ cm}^{-1}$ ) in going from the complexes with the oxidized  $[\text{Fe}_4]$  core ( $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]$ ) to the complexes with the one-electron reduced  $[\text{Fe}_4]$  core ( $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]$ ). We start our detailed discussion on the spectroscopic features from the  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]-\text{O}$  complex (Fig. 3). The most intense  $^{57}\text{Fe}$  PVDOS feature is centered at  $215\text{ cm}^{-1}$  with a low energy shoulder at  $190\text{ cm}^{-1}$  and a high energy shoulder at  $250\text{ cm}^{-1}$ . The next strongest feature is observed at  $290\text{ cm}^{-1}$  with a broad high energy shoulder distributed between  $320\text{ cm}^{-1}$  and  $400\text{ cm}^{-1}$ . In the energy range above  $400\text{ cm}^{-1}$ , only relatively weak  $^{57}\text{Fe}$  PVDOS features are observed. To identify the terminal  $\text{Fe}-\text{O}$  stretching vibrations, the measurements for the analogous complex with  $^{18}\text{O}$ -labeling (at both the terminal oxido ligand and the bridging  $\mu_4$ -oxido ligand, see the ESI† for synthesis) have also been performed (Fig. 3A). A clear  $^{18}\text{O}$ -sensitive feature is observed at  $665\text{ cm}^{-1}$  in




Fig. 2  $^{57}\text{Fe}$  PVDOS spectra of the  $[\text{Fe}_4]$ -oxido and  $[\text{Fe}_4]$ -hydroxido complexes. The blue lines represent the spectra from the complexes with a  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]$  core, the red lines represent the spectra from the complexes with a  $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]$  core, the solid lines represent the spectra from the complexes containing a  $\text{Fe}^{\text{III}}-\text{O}$  moiety, and the dash lines represent the spectra from the complexes containing a  $\text{Fe}^{\text{III}}-\text{OH}$  moiety.




Fig. 3  $^{57}\text{Fe}$  PVDOS experimental and DFT calculated spectra of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]-\text{O}$  and the associated vibrational mode assignments. (A)  $^{57}\text{Fe}$  PVDOS spectra (top) and DFT calculated spectra (bottom) of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]-\text{O}$  (black traces) and  $^{18}\text{O}$ -labeled complex,  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2^{18}\text{OFe}^{\text{III}}]-^{18}\text{O}$  (red traces); the simplified structure of this complex is also shown (see Fig. 1 for atom color code definition). See Fig. S2† for additional comparison between the experimental and DFT calculated spectra. (B) The  $\text{Fe}_4$  cluster core geometry with selected atom labeling used in this study (C) the selected normal modes of vibration of a  $\text{OFe}(\text{X}_3)(\text{YZ}_3)$  type molecule.



$[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{-16}\text{O}$ , which is red shifted to  $635\text{ cm}^{-1}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{-18}\text{O}$ . This feature must belong to the stretching vibration of the terminal  $\text{Fe}^{\text{III}}\text{-O}$  moiety ( $\nu(\text{Fe}^{\text{III}}\text{-O})$ ), and the isotope shift ( $\Delta\nu$  ( $^{16}\text{O}/^{18}\text{O}$ ) =  $30\text{ cm}^{-1}$ ) is consistent with the predicted isotope shift of an isolated  $\text{Fe-O}$  stretching vibration based on Hooke's law ( $\Delta\nu = 30\text{ cm}^{-1}$ ). The residual  $665\text{ cm}^{-1}$  peak in the spectrum of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{-18}\text{O}$  is most likely from the  $^{16}\text{O}$  labeled complex due to incomplete  $^{18}\text{O}$  labeling (see more discussion below). Interestingly,  $\nu(\text{Fe}^{\text{III}}\text{-O})$  observed in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$  shows nearly identical energy with the  $\nu(\text{Fe}^{\text{III}}\text{-O})$  observed in a mononuclear ferric-oxido complex,  $[\text{Fe}^{\text{III}}\{\text{H}_3\text{buea}\}(\text{O})]^{2-}$  ( $\nu(\text{Fe}^{\text{III}}\text{-O}) = 660\text{ cm}^{-1}$ ),<sup>77</sup> suggesting that  $\nu(\text{Fe}^{\text{III}}\text{-O})$  is a relatively isolated vibration even in this tetranuclear iron cluster, such as  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$ , and the vibrational frequency is essentially determined by the  $\text{Fe}^{\text{III}}\text{-O}$  bond length ( $d[\text{Fe-O}] = 1.795(8)\text{ \AA}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$  and  $1.813(3)\text{ \AA}$  in  $[\text{Fe}^{\text{III}}\{\text{H}_3\text{buea}\}(\text{O})]^{2-}$  determined by single crystal X-ray diffraction (XRD)<sup>74,75</sup>). We note here that the stabilization of an  $\text{Fe}^{\text{III}}\text{-O}$  moiety is achieved by three hydrogen bonds (H-bonds) provided by the NH groups of the supporting ligand in both complexes. A similar hydrogen bonded  $\text{Fe}^{\text{III}}\text{-O}$  moiety with an  $\text{Fe}^{\text{III}}\text{-O}$  bond length of  $d[\text{Fe-O}] = 1.808(1)\text{ \AA}$  has also been reported by Fout and coworker on an  $\text{Fe}^{\text{III}}\text{-O}$  complex supported by a tripodal ligand, tris(5-cyclohexyl-amineazafulvene-2-methyl)amine ( $\text{N}(\text{afa}^{\text{CY}})_3$ ), with three H-bonds.<sup>46</sup> Roithová and coworkers have reported the  $\nu(\text{Fe}^{\text{III}}\text{-O})$  in gas phase for  $[\text{Fe}^{\text{III}}(\text{O})\text{N}_4\text{Py}]^+$  ( $\text{N}_4\text{Py} = \text{N,N-bis}(2\text{-pyridylmethyl})\text{-N-bis}(2\text{-pyridylmethyl})\text{amine}$ ) and  $[\text{Fe}^{\text{III}}(\text{O})\text{TPA}]^+$  ( $\text{TPA} = \text{tris}(2\text{-pyridylmethyl})\text{amine}$ ) with only a single H-bond provided by a water molecular to the  $\text{Fe}^{\text{III}}\text{-O}$  moieties, which gave  $\nu(\text{Fe}^{\text{III}}\text{-O}) = 787\text{ cm}^{-1}$ , much higher than the  $\nu(\text{Fe}^{\text{III}}\text{-O})$  observed here.<sup>78</sup>

To better understand the  $^{57}\text{Fe}$  PVDOS features exhibited by  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$ , we turned to DFT analysis. For the calculations, a broken symmetry state was used, where the three iron atoms (Fe2,3,4) at the base of the  $[\text{Fe}_4]$  cluster are ferromagnetically coupled and the apical iron atom (Fe1) is antiferromagnetically coupled with all three base iron atoms to form an  $M_S = 9/2$  BS state. The selection of this BS state is supported by the SQUID measurements of this complex, which showed that  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$  exhibits an  $S = 9/2$  ground spin state.<sup>79</sup> By using the functional-basis set combination of B3LYP/TZVP (see the ESI† for the calculation details), the DFT optimized structure using the  $M_S = 9/2$  BS state nicely reproduced the corresponding structure determined by XRD. The comparison of the selected iron-ligand bond lengths and the atom labeling used in this study are shown in Fig. 3B (see Table S1† for additional selected bond lengths based on the XRD results and the DFT results), which demonstrates a good overall agreement between the DFT optimized and the XRD determined structures. Similar with the crystal structure, the DFT optimized structure shows that the three base irons are not structurally equivalent. One of the irons (Fe2) shows longer bond lengths with the coordinating ligands than the other two irons ( $d_{\text{avg}}(\text{Fe2-L}) = 2.136$  (2.186, DFT)  $\text{\AA}$ ,  $d_{\text{avg}}(\text{Fe3-L}) = 2.068$  (2.117, DFT)  $\text{\AA}$ , and  $d_{\text{avg}}(\text{Fe4-L}) = 2.051$  (2.114, DFT)  $\text{\AA}$ ), suggesting that Fe2 is the ferrous center while the other two irons (Fe3 and Fe4) are in the ferric state. Specifically, the  $\text{Fe2-}\mu_4\text{O}$  bond length is  $2.155(7)$

$\text{\AA}$  in crystal structure and  $2.196\text{ \AA}$  in the DFT optimized structure while the  $\text{Fe3-}\mu_4\text{O}$  and  $\text{Fe4-}\mu_4\text{O}$  bond lengths are  $\sim 0.2\text{ \AA}$  shorter. Indeed, the DFT calculated spin population also support this assignment (Table S2†). Fe2 shows a Mulliken spin population of 3.81, which is smaller than those of Fe3 and Fe4 (4.15) and is more consistent with a high-spin ferrous ion. Also, the spin population of the apical iron (Fe1) is 4.01, similar with those of Fe3 and Fe4, thus confirming that Fe1, Fe3, and Fe4 are all in the ferric state. Based on this optimized structure, we carried out frequency calculations and reconstructed  $^{57}\text{Fe}$  PVDOS spectrum of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$ . The calculated spectrum compares well with the experimental data, which allows assignments of the spectral features to specific vibrational modes. The DFT calculated  $\nu(\text{Fe}^{\text{III}}\text{-O})$  is located at  $675\text{ cm}^{-1}$ , which is red shifted to  $646\text{ cm}^{-1}$  with  $^{18}\text{O}$ -labeling on the terminal oxido ligand to give  $\Delta\nu$  ( $^{16}\text{O}/^{18}\text{O}$ ) =  $29\text{ cm}^{-1}$ . Thus, this result supports the assignment of the  $665\text{ cm}^{-1}$   $^{57}\text{Fe}$  PVDOS feature in the experimental data to the same  $\nu(\text{Fe}^{\text{III}}\text{-O})$ , which shows  $\Delta\nu$  ( $^{16}\text{O}/^{18}\text{O}$ ) =  $30\text{ cm}^{-1}$ .

The vibrational mode assignments for the low energy features of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$  can be best understood by comparing with the normal modes of vibration of a simplified  $\text{OFe}(\text{X}_3)(\text{YZ}_3)$  type molecule with a  $C_{3v}$  symmetry,<sup>77</sup> where in our case X represents the equatorial ligand of the apical iron, Y represents the iron ligand trans to the oxido ligand, in this case it is the bridging oxo ligand ( $\mu_4\text{-O}$ ) in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$ , and Z represents the three base iron atoms. The full normal modes of vibration of the  $\text{OFe}(\text{X}_3)(\text{YZ}_3)$  type molecule are listed in Fig. S1,† and selected mode pictures are listed in Fig. 3C. The strongest  $^{57}\text{Fe}$  PVDOS feature in the calculated spectrum of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$  is located at  $\sim 207\text{ cm}^{-1}$  with an unresolved high energy shoulder at  $\sim 218\text{ cm}^{-1}$ , which should correspond to the  $215\text{ cm}^{-1}$  peak observed experimentally. This peak is mainly originated from vibrational modes that highly resembles the  $\nu_4(\text{E})$  and  $\nu_5(\text{E})$  modes (the swing chair motions,  $\nu_4(\text{E})$ , essentially the out of plane bending motions of  $\text{N}_{\text{eq}}\text{-Fe}_1\text{-N}_{\text{eq}}$  and  $\text{Fe}_{\text{base-}\mu_4\text{O}}\text{-Fe}_{\text{base}}$ , and the in-phase rocking motion,  $\nu_5(\text{E})$ , essentially the in-phase combination of the out of plane rocking  $\text{N}_{\text{eq}}\text{-Fe}_1\text{-N}_{\text{eq}}$  with the  $\text{Fe}_{\text{base-}\mu_4\text{O}}\text{-Fe}_{\text{base}}$  bending motions) from the  $\text{OFe}(\text{X}_3)(\text{YZ}_3)$  model. Due to the lower symmetry of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$ , these two E modes split into four major vibrations calculated at  $207\text{ cm}^{-1}$ ,  $212\text{ cm}^{-1}$ ,  $214\text{ cm}^{-1}$ , and  $221\text{ cm}^{-1}$ . The experimentally observed  $190\text{ cm}^{-1}$  peak is calculated at  $\sim 180\text{ cm}^{-1}$ , which is mainly from vibrations resembling the  $\nu_6(\text{E})$  mode (the out-of-phase rocking motion, essentially the out-of-phase combination of  $\nu_5(\text{E})$ ) from the  $\text{OFe}(\text{X}_3)(\text{YZ}_3)$  model calculated at  $\sim 180\text{ cm}^{-1}$  and the symmetric  $\text{N}_{\text{eq}}\text{-Fe}_1\text{-N}_{\text{eq}}$  out-of-plane bending motion calculated at  $185$  ( $\nu_8(\text{A}_1)$  mode). The calculated features at  $300\text{ cm}^{-1}$  with a lower energy shoulder at  $280\text{ cm}^{-1}$  could be assigned to the experimentally observed  $290\text{ cm}^{-1}$  feature. The calculated  $300\text{ cm}^{-1}$  feature mainly originates from  $\text{N}_{\text{eq}}\text{-Fe}_1\text{-N}_{\text{eq}}$  asymmetric stretch motion combined with an expansion motion of the base  $\text{Fe}_{\text{base-}\mu_4\text{O}}$  structural moiety, which is reminiscent to  $\nu_{11}(\text{E})$  mode of the  $\text{OFe}(\text{X}_3)(\text{YZ}_3)$  model. The calculated  $280\text{ cm}^{-1}$  feature is originated from a symmetric  $\text{Fe}_{\text{base-}\mu_4\text{O}}\text{-Fe}_{\text{base}}$  bending motion (the  $\nu_7(\text{A}_1)$  mode). In addition, the calculated  $260\text{ cm}^{-1}$  peak



with a  $244\text{ cm}^{-1}$  shoulder corresponds well with the experimentally observed  $255\text{ cm}^{-1}$  feature. The calculated  $260\text{ cm}^{-1}$  feature can be best described as a symmetric breathing mode involving the  $\text{O1-Fe1-}\mu_4\text{O-Fe}_{\text{base}}$  core moiety of the  $[\text{Fe}_4]$  cluster (the  $\nu_{13}(\text{A}_1)$  mode), while the  $244\text{ cm}^{-1}$  feature mainly represents the asymmetric  $\text{N}_{\text{eq}}\text{-Fe1-N}_{\text{eq}}$  stretching mode (the  $\nu_{11}(\text{E})$  mode). The calculated  $340\text{ cm}^{-1}$  feature could be assigned to the broad  $320\text{ cm}^{-1}$  feature observed experimentally, which is associated with an asymmetric  $\text{Fe}_{\text{base}}\text{-}\mu_4\text{O-Fe}_{\text{base}}$  bending motion involving two  $\text{Fe}(\text{III})$  centers of the base irons (resembling the  $\nu_5(\text{E})$  mode). Finally, the experimental data shows a weak broad feature centered at  $\sim 515\text{ cm}^{-1}$ . The DFT calculated  $^{57}\text{Fe}$  PVDOS spectrum also predicts a similar weak feature centered at  $505\text{ cm}^{-1}$ , which mainly involves the strong movement of  $\mu_4\text{O}$  either along the  $\text{Fe1-O1}$  bond vector (calculated at  $507\text{ cm}^{-1}$ , the  $\nu_{12}(\text{E})$  mode) or perpendicular to it (calculated at  $502\text{ cm}^{-1}$ , the  $\nu_{13}(\text{E})$  mode). The DFT calculated spectrum of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{18}\text{O}$  shows a clear  $^{18}\text{O}$  isotope shift of the  $505\text{ cm}^{-1}$  feature with  $\Delta\nu$  ( $^{16}\text{O}/^{18}\text{O}$ )  $\sim 25\text{ cm}^{-1}$  in addition to the  $^{18}\text{O}$  isotope shift of the terminal  $\text{Fe}^{\text{III}}\text{-O}$  stretching mode mentioned above ( $\nu(\text{Fe}^{\text{III}}\text{-}^{16}\text{O}) = 675\text{ cm}^{-1}$  and  $\nu(\text{Fe}^{\text{III}}\text{-}^{18}\text{O}) = 646\text{ cm}^{-1}$  in the calculated spectra). Such an isotope shift related to  $\mu_4\text{O}$  is not clearly observed for the broad  $515\text{ cm}^{-1}$  feature in the experimental spectrum of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{18}\text{O}$ , which we attribute to incomplete  $^{18}\text{O}$  isotope labeling as evidenced by the mixed  $\nu(\text{Fe}^{\text{III}}\text{-}^{16}\text{O})$  and  $\nu(\text{Fe}^{\text{III}}\text{-}^{18}\text{O})$  observed for the terminal  $\text{Fe}^{\text{III}}\text{-O}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{18}\text{O}$ . Nevertheless, calculations to probe the effect of  $^{16}\text{O}$  vs.  $^{18}\text{O}$  at the  $\mu_4\text{O}$  position show little effect on the terminal  $\text{Fe-O}$  (as well as  $\text{Fe-OH}$ ) vibrational frequencies (see Fig. S2–S5† for all the complexes studied here), which further suggests that the stretching vibration of the terminal  $\text{Fe-O}$  moiety is well isolated from other vibrational modes. The experimentally observed and DFT calculated frequencies of selected normal modes for all four complexes studied here as well as selected bond lengths are listed in Table 1. Overall, the strong  $^{57}\text{Fe}$  PVDOS features in the energy range between  $150\text{ cm}^{-1}$  and  $400\text{ cm}^{-1}$  are mainly the results of vibrations from the  $\text{Fe}_4$  cluster core (the  $\text{O1-Fe1-}\mu_4\text{O-Fe}_{\text{base}}$  moiety), thus reflecting the core structure of this type of complexes.

With a good understanding of the vibrational features from  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O}$ , we turn to the corresponding protonated complex,  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$ . The NRVs derived  $^{57}\text{Fe}$  PVDOS spectrum of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  is almost identical to that of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O}$  (both in peak positions and relative peak intensities) in the energy range below  $400\text{ cm}^{-1}$ , suggesting that the replacement of a terminal oxido to a terminal hydroxido ligand has a negligible impact to the  $\text{Fe}_4$  cluster core structure (Fig. 4). Indeed, the iron–ligand bond lengths of the  $\text{Fe}_4$  cluster core in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  exhibit minimal changes when compared with those in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O}$ . For example, the averaged  $\text{Fe}_1\text{-N}_{\text{eq}}$  bond length is  $2.06\text{ \AA}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  and  $2.09\text{ \AA}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O}$ , and the averaged iron–ligand bond lengths of the base irons ( $d_{\text{avg}}(\text{Fe}_{\text{base}}\text{-L})$ ) are  $2.118\text{ \AA}$ ,  $2.076\text{ \AA}$ , and  $2.058\text{ \AA}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  and  $2.136\text{ \AA}$ ,  $2.068\text{ \AA}$ , and  $2.051\text{ \AA}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O}$ . However, changes in the energy range above  $400\text{ cm}^{-1}$  can be clearly identified. Especially, the

$665\text{ cm}^{-1}$  peak assigned to  $\nu(\text{Fe}^{\text{III}}\text{-O})$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O}$  is not observed in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$ , instead, a broad  $^{16}\text{O}/^{18}\text{O}$  sensitive feature is observed at  $\sim 570\text{ cm}^{-1}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-}^{16}\text{OH}$ , which is shifted to  $\sim 535\text{ cm}^{-1}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-}^{18}\text{OH}$  (Fig. 4). This isotope sensitive feature is assigned to the vibration modes involving the terminal  $\text{Fe}^{\text{III}}\text{-OH}$  moiety. This assignment is consistent with the bond length change in going from the  $\text{Fe1-OH}$  bond ( $d(\text{Fe1-OH}) = 1.879(2)\text{ \AA}$ ) to the  $\text{Fe1-O}$  bond ( $d(\text{Fe1-O}) = 1.796(8)\text{ \AA}$ ). Our DFT calculated  $^{57}\text{Fe}$  PVDOS spectra for  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  support such an assignment. The vibrational mode exhibiting the strongest  $\text{Fe}^{\text{III}}\text{-OH}$  stretching motion ( $\nu(\text{Fe}_1\text{-OH})$ ) is calculated at  $546\text{ cm}^{-1}$ . In comparison,  $\nu(\text{Fe}_1\text{-O})$  is calculated at  $674\text{ cm}^{-1}$  for  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O}$ . This mode ( $\nu(\text{Fe}_1\text{-OH})$ ) is down shifted to  $527\text{ cm}^{-1}$  with  $^{18}\text{O}$  labeling in the DFT calculated results on  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-}^{18}\text{OH}$  (Fig. 3C). It is also worth noting that  $\nu(\text{Fe1-}\mu_4\text{O})$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-}^{16}\text{OH}$ , is calculated at  $520\text{ cm}^{-1}$ , which is blue shifted by  $13\text{ cm}^{-1}$  when compared with the same stretching mode calculated in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O}$  ( $\nu(\text{Fe1-}\mu_4\text{O}) = 507\text{ cm}^{-1}$ ). This reflects the shortening of the  $\text{Fe}_1\text{-}\mu_4\text{O}$  bond by  $\sim 0.1\text{ \AA}$  in going from  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O}$  to  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$ , which is most likely due to the weaker *trans* influence of the OH ligand that in turn strengthens the  $\text{Fe1-}\mu_4\text{O}$  bond. This assignment is further confirmed by the DFT calculated spectrum of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-}^{18}\text{OH}$ , which showed an  $^{18}\text{O}$  isotope shift of  $20\text{ cm}^{-1}$  from  $520\text{ cm}^{-1}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-}^{16}\text{OH}$  to  $500\text{ cm}^{-1}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-}^{18}\text{OH}$ . Compared with the reported  $\nu(\text{Fe}^{\text{III}}\text{-OH})$  frequencies in mononuclear iron complexes, such as in  $[\text{Fe}^{\text{III}}\{\text{H}_3\text{buea}\}(\text{OH})]^-$  ( $\nu(\text{Fe}^{\text{III}}\text{-OH}) = 477\text{ cm}^{-1}$ ) and  $[\text{Fe}^{\text{III}}\{\text{H}_2\text{pout}\}(\text{OH})]^-$  ( $\nu(\text{Fe}^{\text{III}}\text{-OH}) = 556\text{ cm}^{-1}$ ),  $\nu(\text{Fe}^{\text{III}}\text{-OH})$  observed in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  is closer to the frequency of  $\nu(\text{Fe}^{\text{III}}\text{-OH})$  observed in the latter mononuclear iron complex, but is much higher than that in former one. This is consistent with the  $\text{Fe}^{\text{III}}\text{-OH}$  bond lengths found in these complexes determined by XRD.  $d(\text{Fe}^{\text{III}}\text{-OH})$  is  $1.879(2)\text{ \AA}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$ , while it is  $1.893(2)\text{ \AA}$  in  $[\text{Fe}^{\text{III}}\{\text{H}_2\text{pout}\}(\text{OH})]^-$  and  $1.931(2)\text{ \AA}$  in  $[\text{Fe}^{\text{III}}\{\text{H}_3\text{buea}\}(\text{OH})]^-$ . The shorter  $\text{Fe}^{\text{III}}\text{-OH}$  bond in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  or  $[\text{Fe}^{\text{III}}\{\text{H}_2\text{pout}\}(\text{OH})]^-$  is due to the effect of an additional  $\text{Fe-OH}\cdots\text{X}$  H-bond ( $\text{X} = \text{N}$  in  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  and  $\text{X} = \text{O}$  in  $[\text{Fe}^{\text{III}}\{\text{H}_2\text{pout}\}(\text{OH})]^-$ ), which weakens the hydroxide O–H bond strength since the iron-bound hydroxide is the H-bond donor, leading to a strengthening of the  $\text{Fe-O}$  bond. This additional H-bond where the  $\text{Fe}(\text{III})\text{-OH}$  as the H-bond donor is absent in  $[\text{Fe}^{\text{III}}\{\text{H}_3\text{buea}\}(\text{OH})]^-$ .

To further explore the effect of redox event of neighboring iron centers (the three base iron sites,  $\text{Fe}_2$ ,  $\text{Fe}_3$ , and  $\text{Fe}_4$ ) on the vibrational properties of the terminal  $\text{Fe}^{\text{III}}\text{-O/OH}$  moiety and the overall  $[\text{Fe}_4]$  cluster, we recorded NRVs data on  $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]^{16}\text{O}$  and  $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]\text{-OH}$  complexes, the one-electron reduced congeners of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O/OH}$ . All the  $^{57}\text{Fe}$  PVDOS features observed for  $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]^{16}\text{O}$  and  $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]\text{-OH}$  are shifted to lower energies than those of the corresponding one-electron oxidized clusters (Fig. 2, 5 and Table 1),  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]^{16}\text{O}$  and  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  respectively. Specifically, the strong features in the energy range of  $150\text{ cm}^{-1}$  to  $400\text{ cm}^{-1}$  are all red shifted by  $\sim 10\text{--}20\text{ cm}^{-1}$  in



Table 1 Selected vibrational frequencies and bond lengths of the  $[\text{Fe}]_4$  complexes<sup>a,b</sup>

Normal modes	$[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$	$[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$	$[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]\text{-O}$	$[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]\text{-OH}$
v6	190	193	186	180
	<i>180</i>	<i>176</i>	<i>168</i>	<i>174</i>
v8	190	193	186	180
	<i>185</i>	<i>190</i>	<i>177</i>	<i>179</i>
v4	215	218	206	205
	<i>207, 212</i>	<i>206, 208</i>	<i>197, 202</i>	<i>198</i>
v5	215	218	206	205
	<i>214, 221</i>	<i>216, 221</i>	<i>214, 215</i>	<i>208, 216</i>
v3	255	260	246	252
	<i>260</i>	<i>263</i>	<i>256</i>	<i>252</i>
v11	290	295	276	284
	<i>300</i>	<i>290</i>	<i>272</i>	<i>275</i>
v13	515	525	— <sup>c</sup>	— <sup>c</sup>
	<i>507</i>	<i>520</i>	<i>555</i>	<i>580</i>
v14	665	570	640	520
	<i>674</i>	<i>546, 550</i>	<i>670</i>	<i>525</i>
$d(\text{Fe1-O(H)})$	1.796(8)	1.879(2)	1.817(2)	1.907(4)
$d(\text{Fe1-}\mu_4\text{O})$	<i>1.781</i>	<i>1.901</i>	<i>1.785</i>	<i>1.925</i>
	2.049(7)	1.948(2)	1.965(1)	1.889(3)
$d_{\text{avg}}(\text{Fe1-N}_{\text{eq}})$	<i>2.103</i>	<i>1.980</i>	<i>2.058</i>	<i>1.929</i>
	2.090	2.063	2.098	2.084
	<i>2.108</i>	<i>2.097</i>	<i>2.132</i>	<i>2.114</i>

<sup>a</sup> Normal modes are based on a  $\text{OFe}(\text{X}_3)(\text{YZ}_3)$  type molecule (see Fig. 1 and SY). <sup>b</sup> The italic numbers are derived from DFT calculations, the frequencies are in  $\text{cm}^{-1}$ , and the bond lengths are in Å. <sup>c</sup> The feature is too weak to be assigned in the experimental data.




Fig. 4  $^{57}\text{Fe}$  PVDOS spectra of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$  and  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$ . Top: The  $^{57}\text{Fe}$  PVDOS experimental spectra of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$  with the observed frequencies of the strong spectral features highlighted; middle: the  $^{57}\text{Fe}$  PVDOS experimental spectra of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  (black trace) and the  $^{18}\text{O}$ -labeled complex (red trace) with the key observed frequencies indicated; bottom: the DFT calculated spectra of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  (black trace) and the  $^{18}\text{O}$ -labeled complex (red trace) with  $^{18}\text{O}$ -sensitive features indicated. The simplified structures of these two complexes are also shown (see Fig. 1 for the atom color code definition). See Fig. S3† for additional comparisons between the experimental and DFT calculated spectra.

going from the oxidized complexes to the reduced complexes while maintaining the relative intensities of individual peaks (Fig. 2, 5 and Table 1). This strongly suggests that the overall




Fig. 5  $^{57}\text{Fe}$  PVDOS experimental and DFT calculated spectra from different  $[\text{Fe}_4]$  cluster complexes in two different oxidation states. Top: The experimental (top) and DFT calculated (bottom) spectra of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$  (black) and  $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]\text{-O}$  (green). Bottom: The experimental (top) and DFT calculated (bottom) spectra of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$  (black) and  $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]\text{-OH}$  (green). The frequencies of the major spectral features, including  $\nu(\text{Fe}^{\text{III}}\text{-O})$  and  $\nu(\text{Fe}^{\text{III}}\text{-OH})$  in these complexes, are indicated in the figure. The simplified structures of these two complexes are also shown (see Fig. 1 for the atom color code definition). See Fig. S4 and S5† for additional comparisons between the experimental and DFT calculated spectra.



[Fe<sub>4</sub>] cluster core structure is maintained while the iron–ligand bond lengths, particularly related to the base irons (Fe2, Fe3, or Fe4), are elongated due to reduction, thus resulting in the overall red-shift of the vibrational features. This implication is supported by the structures of these complexes determined by XRD. In going from [Fe<sup>II</sup>Fe<sup>III</sup><sub>2</sub>OFe<sup>III</sup>]-O to [Fe<sup>II</sup><sub>2</sub>Fe<sup>III</sup>OFe<sup>III</sup>]-O,  $d_{\text{avg}}(\text{Fe}_{\text{base}}\text{-L})$  (the averaged Fe2/3/4 – ligand bond length) is elongated, particularly for  $d_{\text{avg}}(\text{Fe3-L})$ , which changes from 2.068 Å to 2.112 Å, and for  $d_{\text{avg}}(\text{Fe4-L})$ , which changes from 2.051 Å to 2.093 Å. While the corresponding averaged bond lengths are also elongated in going from [Fe<sup>II</sup>Fe<sup>III</sup><sub>2</sub>OFe<sup>III</sup>]-OH to [Fe<sup>II</sup><sub>2</sub>Fe<sup>III</sup>OFe<sup>III</sup>]-OH, specifically  $d_{\text{avg}}(\text{Fe3-L})$  changes from 2.058 Å to 2.124 Å. However,  $d_{\text{avg}}(\text{Fe1-N}_{\text{eq}})$  (the averaged Fe1–N<sub>eq</sub> bond lengths) only shows minor changes. In [Fe<sup>II</sup>Fe<sup>III</sup><sub>2</sub>OFe<sup>III</sup>]-O and [Fe<sup>II</sup><sub>2</sub>Fe<sup>III</sup>OFe<sup>III</sup>]-O,  $d_{\text{av}}(\text{Fe1-N})$  is 2.090 Å and 2.108 Å respectively, while in [Fe<sup>II</sup>Fe<sup>III</sup><sub>2</sub>OFe<sup>III</sup>]-OH and [Fe<sup>II</sup><sub>2</sub>Fe<sup>III</sup>OFe<sup>III</sup>]-OH,  $d_{\text{avg}}(\text{Fe1-N})$  is 2.098 Å and 2.084 Å respectively. This implies that the reduction does not directly impact the apical iron (Fe1), which is still in the Fe<sup>III</sup> state. All these observations are consistent with the previous report indicating that reduction happens at the base iron sites (Fe2, Fe3, or Fe4).<sup>74</sup> Although such a reduction does not impact the Fe1–N<sub>eq</sub> bond lengths, it strongly impacts the Fe1–μ<sub>4</sub>O bond. The weakening of Fe2/3/4–μ<sub>4</sub>O bonds due to reduction is in turn increasing the bonding interactions between Fe1 and μ<sub>4</sub>O. A 0.08 Å contraction of  $d(\text{Fe1}-\mu_4\text{O})$  (2.049(7) Å to 1.965(1) Å) is observed in the crystal structures from [Fe<sup>II</sup>Fe<sup>III</sup><sub>2</sub>OFe<sup>III</sup>]-O to [Fe<sup>II</sup><sub>2</sub>Fe<sup>III</sup>OFe<sup>III</sup>]-O, while a 0.06 Å contraction of the same bond (1.948(2) Å to 1.889(3) Å) is observed in going from [Fe<sup>II</sup>Fe<sup>III</sup><sub>2</sub>OFe<sup>III</sup>]-OH to [Fe<sup>II</sup><sub>2</sub>Fe<sup>III</sup>OFe<sup>III</sup>]-OH. The strengthening of Fe1–μ<sub>4</sub>O generates a *trans* influence to weaken the Fe1–O/OH bonding interactions, which show a ~0.02–0.03 Å elongation in the redox pair of the oxido complexes and in the redox pair of the hydroxido complexes. Accordingly, the frequencies of  $\nu(\text{Fe}^{\text{III}}\text{-O})$  and  $\nu(\text{Fe}^{\text{III}}\text{-OH})$  show clear red shifts upon reduction.  $\nu(\text{Fe}^{\text{III}}\text{-O})$  is at 665 cm<sup>-1</sup> (DFT: 674 cm<sup>-1</sup>) in [Fe<sup>II</sup>Fe<sup>III</sup><sub>2</sub>OFe<sup>III</sup>]-O, but at ~640 cm<sup>-1</sup> (DFT: 670 cm<sup>-1</sup>) in [Fe<sup>II</sup><sub>2</sub>Fe<sup>III</sup>OFe<sup>III</sup>]-O (Fig. 5 top), which is red shifted to 608 cm<sup>-1</sup> (DFT: 642 cm<sup>-1</sup>) in [Fe<sup>II</sup><sub>2</sub>Fe<sup>III</sup><sup>18</sup>OFe<sup>III</sup>]-<sup>18</sup>O (Fig. S2†).  $\nu(\text{Fe}^{\text{III}}\text{-OH})$  is at 570 cm<sup>-1</sup> (DFT: 546 cm<sup>-1</sup>) in [Fe<sup>II</sup>Fe<sup>III</sup><sub>2</sub>OFe<sup>III</sup>]-OH, but at ~520 cm<sup>-1</sup> (DFT: 525 cm<sup>-1</sup>) in [Fe<sup>II</sup><sub>2</sub>Fe<sup>III</sup>OFe<sup>III</sup>]-OH (Fig. 5 bottom), which is red shifted to ~500 cm<sup>-1</sup> (DFT: 492 cm<sup>-1</sup>) in [Fe<sup>II</sup><sub>2</sub>Fe<sup>III</sup><sup>18</sup>OFe<sup>III</sup>]-<sup>18</sup>OH (Fig. S2†). In summary, the effect of the redox event at distal iron centers is to modulate the bonding interactions between Fe and μ<sub>4</sub>O, thus exerting a *trans* influence to the apical Fe center resulting in changes of the bonding interactions between Fe1 and the terminal oxido or hydroxido ligand. In addition, the <sup>57</sup>Fe PVDOS features between 150 cm<sup>-1</sup> and 400 cm<sup>-1</sup> are good overall oxidation state indicators for the [Fe<sub>4</sub>]-O/OH core (Fig. 2 and 5).

To put the results of the current study into a broader context related to the iron–oxido/hydroxido moieties, we generated the correlation plot between the Fe–O(H) bond length and the NRVs-derived  $\nu(\text{Fe}^{\text{III}}\text{-O(H)})$  frequencies from the four complexes studied here, together with the previously studied mononuclear iron complexes (Fig. 6). Since the DFT method we used in this study reproduced the experimental NRVs data well, we also extended the calculations to predict the vibrational




Fig. 6 The correlation plot between the Fe–O(H) bond length and its stretching frequency. The experimental data points belonging to the [Fe<sub>3</sub>OFe<sup>III</sup>]-O/OH complexes studied in the current work are indicated as red square boxes and the experimental data points belonging to the mononuclear Fe–O/OH complexes reported previously are indicated as black triangles, respectively. The DFT obtained data points of the [Fe<sub>3</sub>OFe<sup>III</sup>]-O/OH complexes experimentally studied in the current work are indicated as the red dots. The DFT calculated data points of three additional [Fe<sub>4</sub>]-O/OH complexes, [Fe<sup>III</sup><sub>3</sub>OFe<sup>III</sup>]-O, [Fe<sup>III</sup><sub>3</sub>OFe<sup>III</sup>]-OH, and [Fe<sup>III</sup><sub>3</sub>OFe<sup>III</sup>]-OH, that are not experimentally studied in this work are shown as red circles. The black curve represents a simulation using Badger's rule with the expression shown in the figure. The constants used are:  $C_{\text{FeO(H)}} = 56.692$ ,  $d_{\text{FeO(H)}} = 1.038$ , which are taken from our previous study.<sup>77</sup> The grey curve presents a simulation by using the results from Tolman and coworkers.<sup>80</sup>

features of the three other [Fe<sub>4</sub>] clusters previously published in this cluster series, namely the all ferric clusters, [Fe<sup>III</sup><sub>3</sub>OFe<sup>III</sup>]-O and [Fe<sup>III</sup><sub>3</sub>OFe<sup>III</sup>]-OH, and the most reduced cluster, [Fe<sup>II</sup><sub>3</sub>OFe<sup>III</sup>]-OH. The calculated  $\nu(\text{Fe}^{\text{III}}\text{-O(H)})$  frequencies of these three complexes are also included in Fig. 6. The present study significantly expands the data available on terminal Fe<sup>III</sup>-O(H) and Fe<sup>IV</sup>=O species. The plot clearly indicates that the Fe–O(H) bond lengths and the corresponding  $\nu(\text{Fe}^{\text{III}}\text{-O(H)})$  frequencies of both mononuclear and tetranuclear iron–oxido/hydroxido complexes follow a similar trend and can be described by Badger's rule (Fig. 6 and S6†).<sup>77,80</sup> This suggests that Fe–O/OH stretching vibrations in these complexes can be well described as the stretching vibration of a diatomic moiety that is relatively isolated from the rest of the molecules. But at the same time the correlation shown in Fig. 6 also suggests that Fe–O/OH bond length and stretching vibration can be strongly influenced by the surrounding environment either by the direct modulation *via* hydrogen bond interactions in the mononuclear Fe–O/OH complexes or by the indirect modulation *via* remote metal redox events in the [Fe<sub>3</sub>OFe<sup>III</sup>]-O/OH complexes, and these two chemical strategies generate a similar effect towards the bond length and the stretching vibration of the Fe–O/OH moiety.

Beyond the structural perturbation of the terminal Fe–O/OH moieties, the basal iron redox event of the [Fe<sub>3</sub>OFe<sup>III</sup>]-O/OH

complexes also impact the chemical reactivity of these moieties. Recent reactivity studies of metal–oxido (M–O) species towards C–H activation in enzymes and bioinspired model complexes strongly indicate that ground state thermodynamics of the M–O species, *e.g.*  $pK_a$ , reduction potential, and the bond dissociation energy (BDE) of the metal bound hydroxide in the corresponding metal–hydroxido (M–OH) species, play key roles in determining M–O reactivity.<sup>81–90</sup> Particularly, a number of model complex studies suggest that the basicity of M–O units ( $pK_a$  of M–OH) is the determining factor in controlling the asynchronous proton-coupled electron transfer (PCET) process during C–H activation, where proton transfer (PT) is dominant within the transition state over electron transfer (ET).<sup>45,91–98</sup> In a previous study by some of us, it has been shown that the C–H activation reactivity demonstrated by the  $[\text{Fe}_3\text{OFe}^{\text{III}}]\text{-O}$  complex following a similar concerted PT-driven process. Furthermore, the redox state change of the basal iron sites that perturbs the bond length and stretching vibrational frequency of the terminal  $\text{Fe}^{\text{III}}\text{-O/OH}$  moiety also modulates the thermodynamic properties, thus the chemical reactivity of this moiety. Specifically, the redox state change of the basal iron sites leads to modulation of the O–H BDE by as much as  $\sim 12$  kcal mol<sup>−1</sup> in going from  $[\text{Fe}^{\text{II}}_3\text{OFe}^{\text{III}}]\text{-OH}$  to  $[\text{Fe}^{\text{III}}_2\text{Fe}^{\text{II}}\text{OFe}^{\text{III}}]\text{-OH}$  (the O–H BDE of  $[\text{Fe}^{\text{II}}_3\text{OFe}^{\text{III}}]\text{-OH}$  was estimated to be 72 kcal mol<sup>−1</sup> while it was estimated to be 84 kcal mol<sup>−1</sup> for  $[\text{Fe}^{\text{III}}_2\text{Fe}^{\text{II}}\text{OFe}^{\text{III}}]\text{-OH}$ ).<sup>74</sup> Thus the direct measurements of Fe–O/OH vibrational frequency by using <sup>57</sup>Fe NRVs provide an useful tool to quantitatively evaluate the structural impact of redox event from the remote iron sites of  $[\text{Fe}_3\text{OFe}^{\text{III}}]\text{-O}$  complexes and help construct the structure–reactivity correlation for these clusters.

## Conclusion

In this study, we have carried out a detailed vibrational analysis on a series of  $[\text{Fe}_4]$  clusters by using <sup>57</sup>Fe NRVs combined with DFT calculations. These clusters bear a terminal  $\text{Fe}^{\text{III}}\text{-O/OH}$  moiety at the apical position and three additional iron centers forming the base of the cluster. The three basal iron centers are connected with the apical iron center *via* a  $\mu_4$ -oxido ligand. Specifically, the complexes studied are  $[\text{LFe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{-O}(\text{PzNH}^t\text{Bu})_3\text{Fe}^{\text{III}}(\text{O})]^+$  ( $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$ ) and  $[\text{LFe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{-O}(\text{PzNH}^t\text{Bu})_3\text{Fe}^{\text{III}}(\text{OH})]^{2+}$  ( $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$ ), and their corresponding one electron reduced clusters,  $[\text{LFe}^{\text{II}}_2\text{-Fe}^{\text{III}}\text{O}(\text{PzNH}^t\text{Bu})_3\text{Fe}^{\text{III}}(\text{O})]$  ( $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]\text{-O}$ ) and  $[\text{LFe}^{\text{II}}_2\text{-Fe}^{\text{III}}\text{O}(\text{PzNH}^t\text{Bu})_3\text{Fe}^{\text{III}}(\text{O})]^+$  ( $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]\text{-OH}$ ). Strong NRVs-derived <sup>57</sup>Fe PVDOS features are observed in the frequency range of 150–400 cm<sup>−1</sup>, which reflects the vibrational features of the overall  $[\text{Fe}_4]$  cluster core and can be used as a good indicator of the overall oxidation state of these clusters. In the frequency range greater than 400 cm<sup>−1</sup>, the stretching vibrations of the terminal  $\text{Fe}^{\text{III}}\text{-O}$  ( $\nu[\text{Fe}^{\text{III}}\text{-O}]$ ) and  $\text{Fe}^{\text{III}}\text{-OH}$  ( $\nu[\text{Fe}^{\text{III}}\text{-OH}]$ ) are also observed. For  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O}$ ,  $\nu[\text{Fe}^{\text{III}}\text{-O}]$  is observed at 660 cm<sup>−1</sup>, while for  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-OH}$ ,  $\nu[\text{Fe}^{\text{III}}\text{-OH}]$  is observed at 570 cm<sup>−1</sup>. These frequency values are highly similar to  $\nu[\text{Fe}^{\text{III}}\text{-O}]$  and  $\nu[\text{Fe}^{\text{III}}\text{-OH}]$  observed in mononuclear  $\text{Fe}^{\text{III}}\text{-O/OH}$  complexes, suggesting that  $\nu[\text{Fe}^{\text{III}}\text{-O}]$  and  $\nu[\text{Fe}^{\text{III}}\text{-OH}]$  are relatively isolated vibrations even in these  $[\text{Fe}_4]$  clusters and

are essentially determined by Fe–O or Fe–OH bond lengths *via* Badger's rule as shown in Fig. 6. However, the redox event in the remote basal iron centers clearly modulates the terminal  $\text{Fe}^{\text{III}}\text{-O/OH}$  vibrations (as well as the bond lengths) as shown by the red shifts of  $\nu[\text{Fe}^{\text{III}}\text{-O}]$  and  $\nu[\text{Fe}^{\text{III}}\text{-OH}]$  (25 cm<sup>−1</sup> and 40 cm<sup>−1</sup> respectively) in  $[\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}]\text{-O/OH}$ , the one-electron reduced complexes of  $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_2\text{OFe}^{\text{III}}]\text{-O/OH}$ . Such a remote structural modulation is also suggested by the shifts observed in the 150–400 cm<sup>−1</sup> region and is due to the weakening of the bonding interactions between  $\mu_4\text{O}$  and the three base iron sites upon reduction, which leads to the strengthening of the apical Fe– $\mu_4\text{O}$  bond and the weakening of the terminal Fe–O/OH bond *via* a *trans* influence. Thus, <sup>57</sup>Fe NRVs data provided comprehensive vibrational characterizations of the structural perturbations of the terminal Fe–O/OH moieties in these  $[\text{Fe}_4]$  clusters due to the redox change of the remote basal iron sites. Overall, this indirect modulation by remote iron redox event in the  $[\text{Fe}_4]$  clusters and the direct modulation *via* hydrogen bond interactions as reported in mononuclear Fe–O/OH complexes generate a similar effect on changing the bond length and the stretching vibration of the terminal Fe–O/OH moieties, which further modify their thermodynamic properties (such as reduction potential,  $pK_a$ , and O–H BDE) and chemical reactivities as previously reported for these complexes.

## Data availability

The data supporting this article have been included as part of the ESI.†

## Author contributions

Conceptualization: T. A., Y. G.; methodology, J. X., C. R., B. L., M. Y. H., J. Z., E. E. A., Y. A., Y. G.; investigation: J. X.; C. R.; T. A., Y. G.; formal analysis: J. X.; C. R., T. A., Y. G.; writing – original draft: J. X., Y. G.; writing – review & editing, J. X., C. R., T. A., Y. G.; funding acquisition: T. A., Y. G.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

The authors acknowledge the National Institute of Health (NIH GM125924 to Y. G., GM127588 to W. c. C.) and the NSF (CHE-1905320 to T. A.) for funding. The NRVs data were recorded at the Advanced Photon Source (APS), Argonne National Laboratory (proposal GUP-60939, GUP-68490, and GUP-73059). This research used resources of the Advanced Photon Source, a U.S. Department of Energy (DOE) Office of Science user facility operated for the DOE Office of Science by Argonne National Laboratory under Contract No. DE-AC02-06CH11357. In addition, this research used Bridges-2 at Pittsburgh Supercomputing Center through allocation [CHE-200003] from the Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS) program, which is supported by





National Science Foundation grants #2138259, #2138286, #2138307, #2137603, and #2138296.

## References

- 1 E. I. Solomon, T. C. Brunold, M. I. Davis, J. N. Kemsley, S.-K. Lee, N. Lehnert, F. Neese, A. J. Skulan, Y.-S. Yang and J. Zhou, Geometric and Electronic Structure/Function Correlations in Non-Heme Iron Enzymes, *Chem. Rev.*, 2000, **100**, 235–350.
- 2 M. Costas, M. P. Mehn, M. P. Jensen and L. Que, Dioxygen Activation at Mononuclear Nonheme Iron Active Sites: Enzymes, Models, and Intermediates, *Chem. Rev.*, 2004, **104**, 939–986.
- 3 M. M. Abu-Omar, A. Loaiza and N. Hontzeas, Reaction Mechanisms of Mononuclear Non-heme Iron Oxygenases, *Chem. Rev.*, 2005, **105**, 2227–2252.
- 4 W. Nam, Dioxygen Activation by Metalloenzymes and Models, *Acc. Chem. Res.*, 2007, **40**, 465.
- 5 L. Que and W. B. Tolman, Biologically inspired oxidation catalysis, *Nature*, 2008, **455**, 333–340.
- 6 E. G. Kovaleva and J. D. Lipscomb, Versatility of biological non-heme Fe(II) centers in oxygen activation reactions, *Nat. Chem. Biol.*, 2008, **4**, 186–193.
- 7 T. L. Poulos, Heme enzyme structure and function, *Chem. Rev.*, 2014, **114**, 3919–3962.
- 8 J. Yano and V. Yachandra, Mn<sub>4</sub>Ca Cluster in Photosynthesis: Where and How Water is Oxidized to Dioxygen, *Chem. Rev.*, 2014, **114**, 4175–4205.
- 9 S. Hematian, I. Garcia-Bosch and K. D. Karlin, Synthetic Heme/Copper Assemblies: Toward an Understanding of Cytochrome c Oxidase Interactions with Dioxygen and Nitrogen Oxides, *Acc. Chem. Res.*, 2015, **48**, 2462–2474.
- 10 J. D. Blakemore, R. H. Crabtree and G. W. Brudvig, Molecular Catalysts for Water Oxidation, *Chem. Rev.*, 2015, **115**, 12974–13005.
- 11 F. Nastro, M. Chino, O. Maglio, A. Bhagi-Damodaran, Y. Lu and A. Lombardi, Design and engineering of artificial oxygen-activating metalloenzymes, *Chem. Soc. Rev.*, 2016, **45**, 5020–5054.
- 12 W. Zhang, W. Lai and R. Cao, Energy-Related Small Molecule Activation Reactions: Oxygen Reduction and Hydrogen and Oxygen Evolution Reactions Catalyzed by Porphyrin- and Corrole-Based Systems, *Chem. Rev.*, 2017, **117**, 3717–3797.
- 13 L. Que, 60 years of dioxygen activation, *JBIC, J. Biol. Inorg. Chem.*, 2017, **22**, 171–173.
- 14 X. Huang and J. T. Groves, Oxygen Activation and Radical Transformations in Heme Proteins and Metalloporphyrins, *Chem. Rev.*, 2018, **118**, 2491–2553.
- 15 A. J. Jasniewski and L. Que, Dioxygen Activation by Nonheme Diiron Enzymes: Diverse Dioxygen Adducts, High-Valent Intermediates, and Related Model Complexes, *Chem. Rev.*, 2018, **118**, 2554–2592.
- 16 J. L. Lee, D. L. Ross, S. K. Barman, J. W. Ziller and A. S. Borovik, C–H Bond Cleavage by Bioinspired Nonheme Metal Complexes, *Inorg. Chem.*, 2021, **60**, 13759–13783.
- 17 M. Guo, T. Corona, K. Ray and W. Nam, Heme and Nonheme High-Valent Iron and Manganese Oxo Cores in Biological and Abiological Oxidation Reactions, *ACS Cent. Sci.*, 2019, **5**, 13–28.
- 18 A. Gunay and K. H. Theopold, C–H Bond Activations by Metal Oxo Compounds, *Chem. Rev.*, 2010, **110**, 1060–1081.
- 19 A. S. Borovik, Role of metal-oxo complexes in the cleavage of C–H bonds, *Chem. Soc. Rev.*, 2011, **40**, 1870–1874.
- 20 J. Limburg, J. S. Vrettos, L. M. Liable-Sands, A. L. Rheingold, R. H. Crabtree and G. W. Brudvig, A Functional Model for O–O Bond Formation by the O<sub>2</sub>-Evolving Complex in Photosystem II, *Science*, 1999, **283**, 1524–1527.
- 21 A. I. Nguyen, M. S. Ziegler, P. Oña-Burgos, M. Sturzbecher-Hohne, W. Kim, D. E. Bellone and T. D. Tilley, Mechanistic Investigations of Water Oxidation by a Molecular Cobalt Oxide Analogue: Evidence for a Highly Oxidized Intermediate and Exclusive Terminal Oxo Participation, *J. Am. Chem. Soc.*, 2015, **137**, 12865–12872.
- 22 B. M. Hunter, N. B. Thompson, A. M. Müller, G. R. Rossman, M. G. Hill, J. R. Winkler and H. B. Gray, Trapping an Iron(VI) Water-Splitting Intermediate in Nonaqueous Media, *Joule*, 2018, **2**, 747–763.
- 23 S. H. Kim, H. Park, M. S. Seo, M. Kubo, T. Ogura, J. Klajn, D. T. Gryko, J. S. Valentine and W. Nam, Reversible O–O Bond Cleavage and Formation between Mn(IV)-Peroxo and Mn(V)-Oxo Corroles, *J. Am. Chem. Soc.*, 2010, **132**, 14030–14032.
- 24 M. Y. M. Pau, J. D. Lipscomb and E. I. Solomon, Substrate activation for O<sub>2</sub> reactions by oxidized metal centers in biology, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 18355–18362.
- 25 R. H. Holm and E. I. Solomon, Introduction: Bioinorganic enzymology II, *Chem. Rev.*, 2014, **114**, 4039–4040.
- 26 Y. Lu, S. M. Berry and T. D. Pfister, Engineering novel metalloproteins: design of metal-binding sites into native protein scaffolds, *Chem. Rev.*, 2001, **101**, 3047–3080.
- 27 J. Liu, S. Chakraborty, P. Hosseinzadeh, Y. Yu, S. Tian, I. Petrik, A. Bhagi and Y. Lu, Metalloproteins containing cytochrome, iron-sulfur, or copper redox centers, *Chem. Rev.*, 2014, **114**, 4366–4369.
- 28 F. Yu, V. M. Cangelosi, M. L. Zastrow, M. Tegoni, J. S. Plegaria, A. G. Tebo, C. S. Mocny, L. Ruckthong, H. Qayyum and V. L. Pecoraro, Protein design: toward functional metalloenzymes, *Chem. Rev.*, 2014, **114**, 3495–3578.
- 29 S. A. Cook and A. S. Borovik, Molecular Designs for Controlling the Local Environments around Metal Ions, *Acc. Chem. Res.*, 2015, **48**, 2407–2414.
- 30 S. A. Cook, E. A. Hill and A. S. Borovik, Lessons from Nature: A Bio-Inspired Approach to Molecular Design, *Biochemistry*, 2015, **54**, 4167–4180.
- 31 P. Hosseinzadeh, N. M. Marshall, K. N. Chacón, Y. Yu, M. J. Nilges, S. Y. New, S. A. Tashkov, N. J. Blackburn and Y. Lu, Design of a single protein that spans the entire 2-V range of physiological redox potentials, *Proc. Natl. Acad. Sci. U. S. A.*, 2016, **113**, 262–267.



- 32 B. A. Springer, S. G. Sligar, J. S. Olson and G. N. Phillips, Mechanisms of Ligand Recognition in Myoglobin, *Chem. Rev.*, 1994, **94**, 699–714.
- 33 J. C. Kendrew, G. Bodo, H. M. Dintzis, R. G. Parrish, H. Wyckoff and D. C. Phillips, A three-dimensional model of the myoglobin molecule obtained by X-ray analysis, *Nature*, 1958, **181**, 662–666.
- 34 J. C. Kendrew, R. E. Dickerson, B. E. Strandberg, R. G. Hart, D. R. Davies, D. C. Phillips and V. C. Shore, Structure of myoglobin: a three-dimensional Fourier synthesis at 2 Å. Resolution, *Nature*, 1960, **185**, 422–427.
- 35 M. F. Perutz, M. G. Rossmann, A. F. Cullis, H. Muirhead, G. Will and A. C. T. North, Structure of haemoglobin: a three-dimensional Fourier synthesis at 5.5-Å. Resolution, obtained by X-ray analysis, *Nature*, 1960, **185**, 416–422.
- 36 B. Shaanan, The iron–oxygen bond in human oxyhaemoglobin, *Nature*, 1982, **296**, 683–684.
- 37 J. Yang, A. P. Kloek, D. E. Goldberg and F. S. Mathews, The structure of *Ascaris* hemoglobin domain I at 2.2 Å resolution: molecular features of oxygen avidity, *Proc. Natl. Acad. Sci. U. S. A.*, 1995, **92**, 4224–4228.
- 38 T. L. Poulos, S. S. T. Freer, R. A. Alden, S. L. Edwards, U. Skogland, B. Eriksson, N. Xuong, T. Yonetani and J. Kraut, The Crystal Structure of Cytochrome c Peroxidase, *J. Biol. Chem.*, 1980, **2**, 575–581.
- 39 T. L. Poulos and J. Kraut, The Stereochemistry of Peroxidase Catalysis, *J. Biol. Chem.*, 1980, **255**, 8199–8205.
- 40 I. Schlichting, J. Bernendzen, K. Chu, A. M. Stock, S. A. Maves, D. E. Benson, R. M. Sweet, D. Ringe, G. A. Petsko and S. G. Sligar, The Catalytic Pathway of Cytochrome P450cam at Atomic Resolution, *Science*, 2000, **287**, 1615–1622.
- 41 S. Nagano and T. L. Poulos, Crystallographic study on the dioxygen complex of wild-type and mutant cytochrome P450cam: implications for the dioxygen activation mechanism, *J. Biol. Chem.*, 2005, **280**, 31659–31663.
- 42 A. J. Fielding, J. D. Lipscomb and L. Que, A Two-Electron-Shell Game: Intermediates of the Extradial-cleaving Catechol Dioxygenases, *JBIC, J. Biol. Inorg. Chem.*, 2014, **19**, 491–504.
- 43 E. G. Kovaleva, M. S. Rogers and J. D. Lipscomb, Structural Basis for Substrate and Oxygen Activation in Homoprotocatechuate 2,3-Dioxygenase: Roles of Conserved Active Site Histidine 200, *Biochemistry*, 2015, **54**, 5329–5339.
- 44 S. K. Barman, M.-Y. Yang, T. H. Parsell, M. T. Green and A. S. Borovik, Semiempirical method for examining asynchronicity in metal–oxido-mediated C–H bond activation, *Proc. Natl. Acad. Sci. U. S. A.*, 2021, **118**, e2108648118.
- 45 S. K. Barman, J. R. Jones, C. Sun, E. A. Hill, J. W. Ziller and A. S. Borovik, Regulating the Basicity of Metal–Oxido Complexes with a Single Hydrogen Bond and Its Effect on C–H Bond Cleavage, *J. Am. Chem. Soc.*, 2019, **141**, 11142–11150.
- 46 E. M. Matson, Y. J. Park and A. R. Fout, Facile Nitrite Reduction in a Non-heme Iron System: Formation of an Iron(III)-Oxo, *J. Am. Chem. Soc.*, 2014, **136**, 17398–17401.
- 47 Y. J. Park, E. M. Matson, M. J. Nilges and A. R. Fout, Exploring Mn–O bonding in the context of an electronically flexible secondary coordination sphere: synthesis of a Mn(III)-oxo, *Chem. Commun.*, 2015, **51**, 5310–5313.
- 48 Z. Gordon, M. J. Drummond, E. M. Matson, J. A. Bogart, E. J. Schelter, R. L. Lord and A. R. Fout, Tuning the Fe(II/III) Redox Potential in Nonheme Fe(II)-Hydroxo Complexes through Primary and Secondary Coordination Sphere Modifications, *Inorg. Chem.*, 2017, **56**, 4852–4863.
- 49 M. J. Drummond, C. L. Ford, D. L. Gray, C. V. Popescu and A. R. Fout, Radical Rebound Hydroxylation Versus H-Atom Transfer in Non-Heme Iron(III)-Hydroxo Complexes: Reactivity and Structural Differentiation, *J. Am. Chem. Soc.*, 2019, **141**, 6639–6650.
- 50 Y. J. Park, M. N. Peñas-Defrutos, M. J. Drummond, Z. Gordon, O. R. Kelly, I. J. Garvey, K. L. Gullett, M. García-Melchor and A. R. Fout, Secondary Coordination Sphere Influences the Formation of Fe(III)-O or Fe(III)-OH in Nitrite Reduction: A Synthetic and Computational Study, *Inorg. Chem.*, 2022, **61**, 8182–8192.
- 51 H. S. Soo, A. C. Komor, A. T. Iavarone and C. J. Chang, A Hydrogen-Bond Facilitated Cycle for Oxygen Reduction by an Acid- and Base-Compatible Iron Platform, *Inorg. Chem.*, 2009, **48**, 10024–10035.
- 52 V. Yadav, J. B. Gordon, M. A. Siegler and D. P. Goldberg, Dioxygen-Derived Nonheme Mononuclear Fe III (OH) Complex and Its Reactivity with Carbon Radicals, *J. Am. Chem. Soc.*, 2019, **141**, 10148–10153.
- 53 T. Devi, Y.-M. Lee, W. Nam and S. Fukuzumi, Metal ion-coupled electron-transfer reactions of metal-oxygen complexes, *Coord. Chem. Rev.*, 2020, **410**, 213219.
- 54 V. F. Oswald, J. L. Lee, S. Biswas, A. C. Weitz, K. Mitra, R. Fan, J. Li, J. Zhao, M. Y. Hu, E. E. Alp, E. L. Bominaar, Y. Guo, M. T. Green, M. P. Hendrich and A. S. Borovik, Effects of Noncovalent Interactions on High-Spin Fe(IV)–Oxido Complexes, *J. Am. Chem. Soc.*, 2020, **142**, 11804–11817.
- 55 S. Banerjee, A. Draksharapu, P. M. Crossland, R. Fan, Y. Guo, M. Swart and L. Que, Sc<sup>3+</sup>-Promoted O–O Bond Cleavage of a (μ-1,2-Peroxo)diiron(III) Species Formed from an Iron(II) Precursor and O<sub>2</sub> to Generate a Complex with an Fe<sup>IV</sup><sub>2</sub>(μ-O)<sub>2</sub> Core, *J. Am. Chem. Soc.*, 2020, **142**, 4285–4297.
- 56 Y. Li, S. Handunneththige, J. Xiong, Y. Guo, M. R. Talipov and D. Wang, Opening the Co<sup>III,IV</sup><sub>2</sub>(μ-O)<sub>2</sub> Diamond Core by Lewis Bases Leads to Enhanced C–H Bond Cleaving Reactivity, *J. Am. Chem. Soc.*, 2020, **142**, 21670–21678.
- 57 D. Lionetti, S. Suseno, E. Y. Tsui, L. Lu, T. A. Stich, K. M. Carsch, R. J. Nielsen, W. A. Goddard, R. D. Britt and T. Agapie, Effects of Lewis Acidic Metal Ions (M) on Oxygen-Atom Transfer Reactivity of Heterometallic Mn<sub>3</sub>MO<sub>4</sub> Cubane and Fe<sub>3</sub>MO(OH) and Mn<sub>3</sub>MO(OH) Clusters, *Inorg. Chem.*, 2019, **58**, 2336–2345.
- 58 D. B. Rice, E. N. Grotemeyer, A. M. Donovan and T. A. Jackson, Effect of Lewis Acids on the Structure and Reactivity of a Mononuclear Hydroxomanganese(III) Complex, *Inorg. Chem.*, 2020, **59**, 2689–2700.



- 59 J. P. McEvoy and G. W. Brudvig, Water-Splitting Chemistry of Photosystem II, *Chem. Rev.*, 2006, **106**, 4455–4483.
- 60 W. Lubitz, H. Ogata, O. Rüdiger and E. Reijerse, Hydrogenases, *Chem. Rev.*, 2014, **114**, 4081–4148.
- 61 A. Volbeda, M.-H. Charon, C. Piras, E. C. Hatchikian, M. Frey and J. C. Fontecilla-Camps, Crystal structure of the nickel-iron hydrogenase from *Desulfovibrio gigas*, *Nature*, 1995, **373**, 580–587.
- 62 Y. Nicolet, C. Piras, P. Legrand, C. E. Hatchikian and J. C. Fontecilla-Camps, *Desulfovibrio desulfuricans* iron hydrogenase: the structure shows unusual coordination to an active site Fe binuclear center, *Structure*, 1999, **7**, 13–23.
- 63 J. W. Peters, W. N. Lanzilotta, B. J. Lemon and L. C. Seefeldt, X-ray Crystal Structure of the Fe-Only Hydrogenase (Cpl) from *Clostridium pasteurianum* to 1.8 Å Resolution, *Science*, 1998, **282**, 1853–1858.
- 64 H. Dobbek, V. Svetlitchnyi, L. Gremer, R. Huber and O. Meyer, Crystal Structure of a Carbon Monoxide Dehydrogenase Reveals a [Ni-4Fe-5S] Cluster, *Science*, 2001, **293**, 1281–1285.
- 65 S. Ferguson-Miller and G. T. Babcock, Heme/Copper Terminal Oxidases, *Chem. Rev.*, 1996, **96**, 2889–2908.
- 66 T. Tsukihara, H. Aoyama, E. Yamashita, T. Tomizaki, H. Yamaguchi, K. Shinzawa-Itōh, R. Nakashima, R. Yaono and S. Yoshikawa, Structures of Metal Sites of Oxidized Bovine Heart Cytochrome c Oxidase at 2.8 Å, *Science*, 1995, **269**, 1069–1074.
- 67 B. M. Hoffman, D. Lukoyanov, Z.-Y. Yang, D. R. Dean and L. C. Seefeldt, Mechanism of Nitrogen Fixation by Nitrogenase: The Next Stage, *Chem. Rev.*, 2014, **114**, 4041–4062.
- 68 T. Spatzal, M. Aksoyoglu, L. Zhang, S. L. a. Andrade, E. Schleicher, S. Weber, D. C. Rees and O. Einsle, Evidence for Interstitial Carbon in Nitrogenase FeMo Cofactor, *Science*, 2011, **334**, 940.
- 69 O. Einsle, F. A. Tezcan, S. L. A. Andrade, B. Schmid, M. Yoshida, J. B. Howard and D. C. Rees, Nitrogenase MoFe-Protein at 1.16 Å Resolution: A Central Ligand in the FeMo-Cofactor, *Science*, 2002, **297**, 1696–1700.
- 70 G. de Ruiter, N. B. Thompson, D. Lionetti and T. Agapie, Nitric Oxide Activation by Distal Redox Modulation in Tetranuclear Iron Nitrosyl Complexes, *J. Am. Chem. Soc.*, 2015, **137**, 14094–14106.
- 71 G. de Ruiter, K. M. Carsch, S. Gul, R. Chatterjee, N. B. Thompson, M. K. Takase, J. Yano and T. Agapie, Accelerated Oxygen Atom Transfer and C–H Bond Oxygenation by Remote Redox Changes in Fe<sub>3</sub>Mn-Iodosobenzene Adducts, *Angew. Chem., Int. Ed.*, 2017, **56**, 4772–4776.
- 72 C. H. Arnett, M. J. Chalkley and T. Agapie, A Thermodynamic Model for Redox-Dependent Binding of Carbon Monoxide at Site-Differentiated, High Spin Iron Clusters, *J. Am. Chem. Soc.*, 2018, **140**, 5569–5578.
- 73 C. J. Reed and T. Agapie, Thermodynamics of Proton and Electron Transfer in Tetranuclear Clusters with Mn–OH<sub>2</sub>/OH Motifs Relevant to H<sub>2</sub>O Activation by the Oxygen Evolving Complex in Photosystem II, *J. Am. Chem. Soc.*, 2018, **140**, 10900–10908.
- 74 C. J. Reed and T. Agapie, A Terminal Fe III-Oxo in a Tetranuclear Cluster: Effects of Distal Metal Centers on Structure and Reactivity, *J. Am. Chem. Soc.*, 2019, **141**, 9479–9484.
- 75 C. E. MacBeth, A. P. Golombek, V. G. Young, C. Yang, K. Kuczera, M. P. Hendrich and A. S. Borovik, O<sub>2</sub> Activation by Nonheme Iron Complexes: A Monomeric Fe(III)-Oxo Complex Derived From O<sub>2</sub>, *Science*, 2000, **289**, 938–941.
- 76 C. E. MacBeth, R. Gupta, K. R. Mitchell-Koch, V. G. Young, G. H. Lushington, W. H. Thompson, M. P. Hendrich and A. S. Borovik, Utilization of Hydrogen Bonds to Stabilize M–O(H) Units: Synthesis and Properties of Monomeric Iron and Manganese Complexes with Terminal Oxo and Hydroxo Ligands, *J. Am. Chem. Soc.*, 2004, **126**, 2556–2567.
- 77 A. C. Weitz, E. A. Hill, V. F. Oswald, E. L. Bominaar, A. S. Borovik, M. P. Hendrich and Y. Guo, Probing Hydrogen Bonding Interactions to Iron-Oxido/Hydroxido Units by <sup>57</sup>Fe Nuclear Resonance Vibrational Spectroscopy, *Angew. Chem., Int. Ed.*, 2018, **57**, 16010–16014.
- 78 E. Andris, R. Navrátil, J. Jašík, M. Puri, M. Costas, L. Que and J. Roithová, Trapping Iron(III)-Oxo Species at the Boundary of the “Oxo Wall”: Insights into the Nature of the Fe(III)-O Bond, *J. Am. Chem. Soc.*, 2018, **140**, 14391–14400.
- 79 C. J. Reed, *Activation of Nitric Oxide and Water by Transition Metal Clusters Relevant to Active Sites in Biology*, California Institute of Technology, 2019.
- 80 A. D. Spaeth, N. L. Gagnon, D. Dhar, G. M. Yee and W. B. Tolman, Determination of the Cu(III)–OH Bond Distance by Resonance Raman Spectroscopy Using a Normalized Version of Badger’s Rule, *J. Am. Chem. Soc.*, 2017, **139**, 4477–4485.
- 81 J. J. Warren, T. A. Tronic and J. M. Mayer, Thermochemistry of Proton-Coupled Electron Transfer Reagents and its Implications, *Chem. Rev.*, 2010, **110**, 6961–7001.
- 82 C. T. Saouma and J. M. Mayer, Do spin state and spin density affect hydrogen atom transfer reactivity?, *Chem. Sci.*, 2014, **5**, 21–31.
- 83 D. Usharani, D. C. Lacy, A. S. Borovik and S. Shaik, Dichotomous Hydrogen Atom Transfer vs Proton-Coupled Electron Transfer during Activation of X–H Bonds (X = C, N, O) by Nonheme Iron–Oxo Complexes of Variable Basicity, *J. Am. Chem. Soc.*, 2013, **135**, 17090–17104.
- 84 M. S. Seo, N. H. Kim, K.-B. Cho, J. E. So, S. K. Park, M. Clémancey, R. Garcia-Serres, J.-M. Latour, S. Shaik and W. Nam, A mononuclear nonheme iron(IV)-oxo complex which is more reactive than cytochrome P450 model compound I, *Chem. Sci.*, 2011, **2**, 1039.
- 85 J. W. Darcy, B. Koronkiewicz, G. A. Parada and J. M. Mayer, A Continuum of Proton-Coupled Electron Transfer Reactivity, *Acc. Chem. Res.*, 2018, **51**, 2391–2399.
- 86 J. J. D. Sacramento and D. P. Goldberg, Factors Affecting Hydrogen Atom Transfer Reactivity of Metal–Oxo Porphyrinoid Complexes, *Acc. Chem. Res.*, 2018, **51**, 2641–2652.



- 87 T. H. Yosca, J. Rittle, C. M. Krest, E. L. Onderko, A. Silakov, J. C. Calixto, R. K. Behan and M. T. Green, Iron(IV) hydroxide pKa and the Role of Thiolate Ligation in C-H Bond Activation by Cytochrome P450, *Science*, 2013, **342**, 825–829.
- 88 C. Kupper, B. Mondal, J. Serrano-Plana, I. Klawitter, F. Neese, M. Costas, S. Ye and F. Meyer, Nonclassical Single-State Reactivity of an Oxo-Iron(IV) Complex Confined to Triplet Pathways, *J. Am. Chem. Soc.*, 2017, **139**, 8939–8949.
- 89 M. A. Ehudin, D. A. Quist and K. D. Karlin, Enhanced Rates of C–H Bond Cleavage by a Hydrogen-Bonded Synthetic Heme High-Valent Iron(IV) Oxo Complex, *J. Am. Chem. Soc.*, 2019, **141**, 12558–12569.
- 90 V. Dantignana, J. Serrano-Plana, A. Draksharapu, C. Magallón, S. Banerjee, R. Fan, I. Gamba, Y. Guo, L. Que, M. Costas and A. Company, Spectroscopic and Reactivity Comparisons between Nonheme Oxoiron(IV) and Oxoiron(V) Species Bearing the Same Ancillary Ligand, *J. Am. Chem. Soc.*, 2019, **141**, 15078–15091.
- 91 R. Gupta and A. S. Borovik, Monomeric Mn III/II and Fe III/II Complexes with Terminal Hydroxo and Oxo Ligands: Probing Reactivity via O–H Bond Dissociation Energies, *J. Am. Chem. Soc.*, 2003, **125**, 13234–13242.
- 92 T. H. Parsell, M.-Y. Yang and A. S. Borovik, C–H Bond Cleavage with Reductants: Re-Investigating the Reactivity of Monomeric Mn III/IV–Oxo Complexes and the Role of Oxo Ligand Basicity, *J. Am. Chem. Soc.*, 2009, **131**, 2762–2763.
- 93 J. W. Darcy, S. S. Kolmar and J. M. Mayer, Transition State Asymmetry in C–H Bond Cleavage by Proton-Coupled Electron Transfer, *J. Am. Chem. Soc.*, 2019, **141**, 10777–10787.
- 94 M. K. Goetz and J. S. Anderson, Experimental Evidence for pKa-Driven Asynchronicity in C–H Activation by a Terminal Co(III)–Oxo Complex, *J. Am. Chem. Soc.*, 2019, **141**, 4051–4062.
- 95 M. K. Goetz and J. S. Anderson, Correction to “Experimental Evidence for pKa-Driven Asynchronicity in C–H Activation by a Terminal Co(III)–Oxo Complex”, *J. Am. Chem. Soc.*, 2020, **142**, 5439–5441.
- 96 M. Mandal, C. E. Elwell, C. J. Bouchey, T. J. Zerk, W. B. Tolman and C. J. Cramer, Mechanisms for Hydrogen-Atom Abstraction by Mononuclear Copper(III) Cores: Hydrogen-Atom Transfer or Concerted Proton-Coupled Electron Transfer?, *J. Am. Chem. Soc.*, 2019, **141**, 17236–17244.
- 97 D. Bím, M. Maldonado-Domínguez, L. Rulišek and M. Srnec, Beyond the classical thermodynamic contributions to hydrogen atom abstraction reactivity, *Proc. Natl. Acad. Sci. U. S. A.*, 2018, **115**, E10287–E10294.
- 98 M. Maldonado-Domínguez and M. Srnec, H-Atom Abstraction Reactivity through the Lens of Asynchronicity and Frustration with Their Counteracting Effects on Barriers, *Inorg. Chem.*, 2022, **61**, 18811–18822.

