

Cite this: *Chem. Sci.*, 2024, 15, 13949

All publication charges for this article have been paid for by the Royal Society of Chemistry

Photosensitizer-free singlet oxygen generation via a charge transfer transition involving molecular O₂ toward highly efficient oxidative coupling of arylamines to azoaromatics†

Shivendra Singh and Tushar Kanti Mukherjee *

Photosensitizer (PS)-mediated generation of singlet oxygen, O₂ (a¹Δ_g) is a well-explored phenomenon in chemistry and biology. However, the requirement of appropriate PSs with optimum excited state properties is a prerequisite for this approach which limits its widespread application. Herein, we report the generation of O₂ (a¹Δ_g) via direct charge-transfer (CT) excitation of the solvent–O₂ (X³Σ_g[−]) collision complex without any PS and utilize it for the catalyst-free oxidative coupling of arylamines to azoaromatics under ambient conditions in aqueous medium. Electron paramagnetic resonance (EPR) spectroscopy revealed the formation of O₂ (a¹Δ_g) upon direct excitation with 370 nm light. The present approach shows broad substrate scope, remarkably fast reaction kinetics (90 and 40 min under an open and O₂ atm, respectively), high selectivity (100%), and excellent yields (up to 100%), and works well for both homo- and hetero-coupling of arylamines. The oxidative coupling of arylamines was found to proceed through the generation of amine radicals via electron transfer (ET) from amines to O₂ (a¹Δ_g). Notably, electron-rich amines show higher yields of azo products compared to electron-deficient amines. Detailed mechanistic investigations using various spectroscopic tools revealed the formation of hydrazobenzene as an intermediate along with superoxide radicals which subsequently transform to hydrogen peroxide. The present study is unique in the way that molecular O₂ simultaneously acts as a light-absorbing chromophore (solvent–O₂ complex) as well as an efficient oxidant (O₂ (a¹Δ_g)) in the same reaction. This is the first report on the efficient, selective, and sustainable synthesis of azo compounds in aqueous medium under an ambient atmosphere without any PCs/PSs and paves the way for further in-depth understanding of the chemical reactivity of O₂ (a¹Δ_g) generated directly via CT excitation of the solvent–O₂ complex toward various photochemical and photobiological transformations.

Received 21st June 2024

Accepted 26th July 2024

DOI: 10.1039/d4sc04115a

rsc.li/chemical-science

Introduction


Molecular oxygen (O₂) plays key roles in many organic transformations due to the generation of various reactive oxygen species (ROS) including singlet O₂ (a¹Δ_g) and superoxide ions (O₂^{•−}).^{1–8} Among these, O₂ (a¹Δ_g) is an efficient and the most utilized ROS and finds applications in other fields such as water disinfection⁹ and photodynamic therapy (PDT).¹⁰ As a consequence of its widespread applications, various sensitive O₂ (a¹Δ_g) detection techniques have been developed over the years.^{11–13} Earlier, Mulliken first predicted three energetically closely lying electronic states of O₂ on the basis of electronic configuration and later assigned them as the triplet ground state (X³Σ_g[−]) and two excited singlet states, a¹Δ_g and b¹Σ_g⁺.¹⁴

Deactivation of O₂ (a¹Δ_g) mainly happens either via radiative or non-radiative pathways and these include the radiative transition from the singlet to triplet ground state via phosphorescence at ~1270 nm and non-radiative quenching by organic molecules, solvents, or photosensitizers.¹⁵

Conventionally, O₂ (a¹Δ_g) is generated using a photosensitizer (PS) upon appropriate light irradiation.^{15–17} The dominant pathway involves initial excitation of the PS to the singlet state (S₁^{*}) followed by intersystem crossing (ISC) to the triplet excited state (T₁^{*}) and finally O₂ (a¹Δ_g) is produced via energy transfer from the T₁^{*} state of the PS to ground state O₂ (X³Σ_g[−]) (Scheme 1A). While energy exchange between PS and O₂ (X³Σ_g[−]) results in the formation of both O₂ (a¹Δ_g) and O₂ (b¹Σ_g⁺), due to rapid spin-allowed deactivation of b¹Σ_g⁺ to a¹Δ_g, only O₂ (a¹Δ_g) persists in the reaction mixture.¹⁵ The requisition of an ideal PS imposes several criteria such as high molar absorption coefficient at the excitation wavelength, triplet state energy greater than 95 kJ mol^{−1} to allow efficient energy transfer to ground state O₂ (X³Σ_g[−]), appreciable triplet state quantum yield, long

Department of Chemistry, Indian Institute of Technology Indore, Indore 453552, Madhya Pradesh, India. E-mail: tusharm@iiti.ac.in

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc04115a>



Scheme 1 Generation of $O_2(a^1\Delta_g)$ via (A) traditional PS-based approach and (B) direct excitation to the CT state.

triplet state lifetime, and high photostability.¹⁷ Previously, a wide range of PSs such as porphyrins and their derivatives,^{18–21} nanoparticles,^{22–24} organic molecules,^{25,26} metal–organic frameworks (MOFs),²⁷ and metal complexes^{28,29} have been utilized to produce $O_2(a^1\Delta_g)$ for various applications. In recent years, heavy-atom-free non-porphyrinoid PSs have emerged as an alternative class of PSs for PDT applications.³⁰ While significant efforts have been made to understand and improve the efficiency of $O_2(a^1\Delta_g)$ generation *via* PS-based systems, the requirement of suitable PSs often raises several challenges. These include unwanted photobleaching of PSs, finite excited state lifetimes of PSs, optimum singlet–triplet energy gap, the presence of other quenchers, competing chemical reactions, and environmental fluctuations. All these limiting factors may influence the efficiency and quantum yield of $O_2(a^1\Delta_g)$ generation.

An alternative route for $O_2(a^1\Delta_g)$ generation was proposed previously *via* direct charge-transfer (CT) excitation involving molecular O_2 and organic solvents without any PS (Scheme 1B).^{31–36} Initially, Tsubomura and Mulliken observed significant changes in the absorption bands of different solvents in the presence of dissolved O_2 upon irradiation.³⁷ The structureless absorption tail observed at longer wavelengths in different oxygen-saturated solvents was attributed to the CT interaction between solvent molecules as electron donors and $O_2(X^3\Sigma_u^-)$ as an electron acceptor. Later, seminal studies of Ogilby and coworkers demonstrated the formation of $O_2(a^1\Delta_g)$ upon irradiation of solvent– O_2 collision complexes.^{31–36} It has been shown that common solvents such as water, methanol, benzene, mesitylene, cyclohexane, acetonitrile, *etc.* under open atmosphere conditions exhibit a structureless long absorption tail due to transition from the solvent– O_2 collision complex to a solvent– O_2 charge transfer (CT) state. Subsequent decay of this CT state results in the production of $O_2(a^1\Delta_g)$. The long wavelength absorption tail appears in a broad wavelength range between 300 and 400 nm, which is solvent-dependent.³⁷ The quantum yield of $O_2(a^1\Delta_g)$ produced upon direct CT band excitation was reported to be ~ 0.2 in organic solvents.³³ In recent work, Bregnhøj and Ogilby utilized a nonlinear two-photon excitation process using visible light (600–730 nm) to

produce an appreciable amount of $O_2(a^1\Delta_g)$.³⁵ Two competing decay channels of the CT state were proposed previously in water.^{34,35} In the first channel, the CT state relaxes to yield solvated $O_2(a^1\Delta_g)$, while the second channel produces $O_2(a^1\Delta_g)$ *via* photosensitization of ground state $O_2(X^3\Sigma_u^-)$ by triplet state water produced upon relaxation. Notably, it has been shown that the nature of the solvent influences the partitioning between these two decay channels.^{34,35} Although these previous studies thoroughly investigated the fundamental photophysical processes involving direct CT excitation and subsequent quantification of $O_2(a^1\Delta_g)$ in various solvents with appreciable yield, the photogenerated $O_2(a^1\Delta_g)$ has not been utilized yet for any chemical transformation. We seek to address whether the $O_2(a^1\Delta_g)$ generated through direct CT excitation in a neat solvent can be utilized for a multistep oxidative coupling reaction. In the present study, we have utilized the $O_2(a^1\Delta_g)$ generated through direct CT excitation in water as an oxidant to drive the oxidative coupling of arylamines to azoaromatics without any PSs/photocatalysts (PCs).

Azo compounds are an important class of functional organic molecules that have enormous potential in molecular materials³⁸ and industrial applications.³⁹ Over the past few decades, several synthetic strategies have been formulated for the synthesis of azoaromatics.⁴⁰ In recent times, efforts were made to convert nitroarenes and aromatic amines to azoaromatics using various metal nanoparticles (NPs) as catalysts;^{41–50} however, most of these methodologies suffer from poor selectivity, slow reaction kinetics, requirement of high temperature, low to moderate yields, and/or inert atmosphere. Similarly, photocatalytic reductive coupling of nitroarenes to azoaromatics with simultaneous production of nitrosobenzene or azoxyaromatics has been demonstrated using various PCs such as supported Au NPs,⁵¹ Cu NPs,⁵² and Au NP functionalized organic cages.⁵³ On the other hand, photocatalytic oxidative coupling (POC) using molecular O_2 as an oxidant has been utilized recently for the synthesis of azo compounds using Ir-based photocatalysts.⁵⁴ Very recently, using a common organic photocatalyst, eosin Y, we formulated a highly efficient and selective POC of arylamines at the charged aqueous interfaces *via* utilizing a non-diffusive ultrafast electron transfer



process at the micellar surface.⁵⁵ Motivated by our previous study, we seek to explore this oxidative coupling reaction under mild reaction conditions without any external PCs/PSs. Notably, most of the earlier studies utilized costly metal-based PCs for the synthesis of azo compounds (Table S1†). To the best of our knowledge, the present study is the first of its kind to demonstrate the utilization of O_2 ($a^1\Delta_g$) generated directly *via* CT excitation toward an industrially important oxidative coupling reaction without any external PCs/PSs.

Results and discussion

PS-free generation and detection of singlet oxygen

First, we recorded the absorption spectra of different solvents such as water, methanol, benzene, mesitylene, and acetonitrile saturated with either N_2 or O_2 to monitor the CT features in the UV-visible region. Notably, a long absorption tail (>350 nm) was clearly evident in all O_2 -saturated solvents compared to those in N_2 -saturated solvents (Fig. 1A–C and S1†). The differences in the spectral signature between N_2 and O_2 saturated solvents suggest the active role of O_2 in the appearance of this long absorption tail. Earlier, Tsubomura and Mulliken assigned this O_2 -dependent absorption feature to a transition from a ground state solvent- O_2 ($X^3\Sigma_u^-$) contact complex to a solvent- O_2 ($X^3\Sigma_u^-$) CT state ($sol^+O_2^{\cdot-}$).³⁷ Moreover, it has also been shown that this long structureless absorption tail can be reversibly removed by purging the solvents with N_2 gas. The molar absorption coefficient (ϵ) of this CT transition in the benzene- O_2 complex is reported to be $\sim 100\text{ M}^{-1}\text{ cm}^{-1}$.³⁷

Previously, it has been shown that these excited CT states subsequently undergo decay or back electron transfer to generate O_2 ($a^1\Delta_g$) *via* mixing of various solvent- O_2 states.^{31–36}

To authenticate the O_2 ($a^1\Delta_g$) formation in our present system, we utilized 9,10-diphenylanthracene (DPA) as a model probe for the detection of singlet oxygen under the irradiation of a 370 nm LED. It has been well-documented in the literature that anthracene and its derivatives are efficient probes for the detection of singlet oxygen.^{56,57} The experiment was carried out in benzene with 200 μM DPA under open air conditions. DPA in benzene exhibits characteristic multiple absorption peaks due to different vibronic transitions (Fig. 1D). Upon irradiation at 370 nm, the absorbance of DPA decreased gradually over time. This decrease in the absorbance indicates the possible transformation of DPA to its endoperoxide form (9,10-diphenyl-9,10-epidioxyanthracene).^{56,57} The formation of 9,10-diphenyl-9,10-epidioxyanthracene was further confirmed using HR-MS measurement (Fig. 1E). A characteristic HR-MS peak for sodium salt of 9,10-diphenyl-9,10-epidioxyanthracene was observed at $385.1188\text{ g mol}^{-1}$, indicating the formation of O_2 ($a^1\Delta_g$) under the irradiation of 370 nm light (Fig. S2†). To further authenticate the formation of O_2 ($a^1\Delta_g$), we performed EPR measurement using 2,2,6,6-tetramethylpiperidine (TEMP) as a singlet oxygen probe in an aqueous medium.² EPR spectra were recorded under dark and 370 nm irradiation. While under dark conditions, no noticeable peaks for O_2 ($a^1\Delta_g$) were observed, three characteristic peaks appeared in the EPR spectrum of the sample kept under 370 nm light irradiation for 30 min, suggesting the formation of TEMP-trapped O_2 ($a^1\Delta_g$) (Fig. 1F).⁵⁸ Taken together, these observations authenticate the formation of O_2 ($a^1\Delta_g$) upon CT band excitation of the solvent- O_2 complex under open air conditions without any external PSs. Next, we seek to address whether this O_2 ($a^1\Delta_g$) can be utilized directly as an efficient oxidant to drive the oxidative coupling of arylamines.




Fig. 1 UV-vis spectra of (A) water, (B) methanol, and (C) benzene under N_2 and O_2 atmosphere. (D) Changes in the absorption spectra of DPA in benzene under the irradiation of a 370 nm LED under an open atmosphere. (E) Schematic showing the conversion of DPA to 9,10-diphenyl-9,10-epidioxyanthracene under the irradiation of a 370 nm LED in benzene under an open atmosphere. (F) Changes in the EPR spectra of TEMP (50 mM) under dark and 370 nm LED irradiation for 30 min in aqueous medium.



Oxidative coupling of arylamines to azoaromatics


We envisaged that the O_2 ($a^1\Delta_g$) generated *via* direct CT band excitation of the solvent- O_2 complex might act as an efficient oxidant to oxidize arylamine to facilitate the oxidative coupling under ambient conditions without any PCs/PSs. Previously, it has been shown that the presence of a base is essential for the conversion of arylamines to azoaromatics as proton abstraction is a critical step in the reaction pathway.^{54,55} Herein, we utilized *p*-toluidine as a model amine substrate and K_3PO_4 as base to check the feasibility of the coupling reaction in aqueous medium under 370 nm light irradiation without any PCs. The optimized reaction conditions for the conversion of *p*-toluidine to 1,2-bis(4-methylphenyl)diazene in the presence of 2 equiv. of K_3PO_4 under 370 nm light irradiation are tabulated in Table 1. Remarkably, a 100% conversion of *p*-toluidine to 1,2-bis(4-methylphenyl)diazene was observed within 90 min of reaction under 370 nm irradiation and open air conditions (Table 1, entry 1). Here it should be noted that a much slower reaction kinetics with reaction time of 24 and 4 h have been reported previously using Ir-based and eosin Y (EY) PCs, respectively.^{54,55} The absence of any desired azo product under dark conditions suggests the essential role of light in the conversion (Table 1, entry 2).

To know the role of atmospheric O_2 , we performed the reaction under a N_2 atmosphere upon 370 nm irradiation. Importantly, no azo product was formed under a N_2 atmosphere, authenticating the direct role of O_2 in the reaction mechanism (Table 1, entry 3). Furthermore, the conversion yield was found to depend strongly on the concentrations of K_3PO_4 used, suggesting that the proton abstraction is also an essential step for the present reaction (Table 1, entries 4 and 5). Interestingly, organic bases such as DABCO and piperidine were also found to be equally effective for the coupling reaction and

show 100% conversion within 90 min of reaction (Table 1, entries 6–8). Replacement of water with methanol as solvent showed 90% conversion efficiency within 90 min of reaction (Table 1, entry 9 and Fig. S3A†). On the other hand, acetonitrile as solvent yielded very slow kinetics and low conversion yield compared to that in water and methanol under standard reaction conditions (Table 1, entry 10 and Fig. S3B†). To check the influence of temperature, we performed the reaction at 60 °C under dark conditions. Notably, no azo product was formed at higher temperature in the absence of 370 nm light (Table 1, entry 11), indicating the negligible role of temperature in the reaction mechanism. Moreover, the conversion yield was found to depend on the wavelength of light used. While 370 nm excitation light resulted in 100% conversion yield, 440 and 525 nm light sources showed 13 and 0% conversions upon 90 min of irradiation (Fig. 2A). It should be noted that *p*-toluidine in water does not absorb 370 nm light (Fig. S4†). In addition, the coupling reaction was found to be more efficient under an O_2 atmosphere and showed 100% conversion within 40 min of irradiation (Fig. 2B). Therefore, we propose that the essential role of 370 nm light irradiation is to provide sufficient energy for the CT band excitation of the water- O_2 complex in the reaction mixture to generate O_2 ($a^1\Delta_g$).

Our present PC-free approach shows a broad substrate scope with excellent conversion yields for a wide range of substituted amines including electron-rich and -deficient amine moieties (Scheme 2). Azo compounds were first isolated and purified using solvent extraction and column chromatography and subsequently characterized using GC-MS and NMR measurements. Arylamines with electron-donating groups (rxns (1)–(6) in Scheme 2A) showed over 90% conversion yield in just 90 min, which is significantly better than the earlier reported studies.^{54,55} Unsubstituted aniline exhibited a conversion yield of 97% under standard reaction conditions (rxn (7) in Scheme 2A). However, *p*-chloroaniline showed a conversion yield of 73% (rxn (8) in Scheme 2A), possibly due to the electron-withdrawing nature of chlorine.^{54,55} Excellent yields were also obtained with arylamines having both electron-donating and -withdrawing substituents (rxns (9)–(11) in Scheme 2A). In addition, 5,6,7,8-tetrahydronaphthalen-1-amine, a fused ring amine substrate also showed an excellent yield of the azo product with a conversion yield of 78% (rxn (12) in Scheme 2A). In contrast, no azo product was obtained for *p*-nitroaniline possibly due to

Table 1 Optimization of reaction conditions



Entry	Deviation from standard conditions	Conversion ^a [%]
1	Standard ^b	100
2	No light	0
3	N_2 atmosphere	0
4	No K_3PO_4	0
5	1 equiv. of K_3PO_4	86
6	2 equiv. of K_2CO_3	100
7	2 equiv. of DABCO	100
8	2 equiv. of piperidine	100
9	Methanol as solvent	90
10	Acetonitrile as solvent	38
11	60 °C	0

^a Determined from GC-MS measurements. ^b Standard conditions: 0.05 mmol *p*-toluidine, 2 equiv. of K_3PO_4 , 5 mL of H_2O , under an ambient air atmosphere with 370 nm (44 W) illumination for 90 min.





Fig. 2 Time-dependent conversions of *p*-toluidine under standard reaction conditions as a function of (A) wavelength of light and (B) atmospheric conditions.






Scheme 2 Substrate scope for (A) homo-coupling and (B) hetero-coupling reactions under standard conditions. Values in parentheses are isolated yields obtained after column chromatography.

the high electron-withdrawing nature of the nitro group (rxn (13) in Scheme 2A), which is similar to that observed previously.⁵⁵

Our present approach can also be extended for the hetero-coupling of different amines to their corresponding azo compounds with the formation of all three possible products as revealed from the GC-MS measurements (Scheme 2B and Fig. S5–S25†). The hetero-coupling of amines was carried out for an extended period (4 h) under the standard conditions mainly to have 100% conversion of reactants to products for all the

cases. No hetero-coupled product was obtained for the hetero-coupling of *p*-nitroaniline with *p*-toluidine possibly due to the highly electron-deficient amine group of *p*-nitroaniline (Scheme 2B and Fig. S26†). This reactivity trend of arylamines correlates well with the previous reports and suggests the possible role of the electron transfer process.^{54,55} Importantly, except for the desired azo product, no unwanted side products such as nitro, nitroso, or azoxy compounds were observed during the course of the reaction (Fig. S27†).



To know whether any oxygenated intermediates such as nitroso or nitro compounds are involved in the reaction mechanism, we performed control experiments with an equal mixture of aniline and nitrosobenzene as starting substrates under standard reaction conditions. The product distributions were monitored using GC-MS measurements. Notably, the reaction with an equal mixture of aniline and nitrosobenzene under standard reaction conditions yielded both azobenzene and azoxybenzene with a 2 : 1 conversion ratio (Scheme 3A). On the other hand, the reaction with nitrosobenzene alone yielded nitrobenzene and azoxybenzene with a 11 : 1 conversion ratio (Scheme 3A). These product distributions are quite different than that observed with aniline alone as the substrate under standard reaction conditions (rxn (7) in Scheme 2A). Therefore, these findings rule out the possible involvement of oxygenated intermediates in the reaction mechanism. Furthermore, the PC- and PS-free approach of our present methodology makes it highly facile to scale it up to the gram scale (Scheme 3B), which is highly promising for the large-scale industrial production of azo compounds.


Mechanistic pathway

We performed several control experiments to understand the mechanism behind this PC- and PS-free synthesis of azo compounds. To rule out the possibility of direct excitation of amine substrates, we recorded the absorption spectra of different arylamines (Fig. S28†). However, none of the amines were found to absorb light ~ 370 nm except 5,6,7,8-tetrahydronaphthalen-1-amine and *p*-nitroaniline, which showed conversion yields of 78 and 0%, respectively.

Our findings indicate that most of these amine substrates do not absorb the excitation light during the reaction. Furthermore, no appreciable changes in the long wavelength region

were observed in the absorption spectrum of *p*-toluidine in the presence of K_3PO_4 , which ruled out the possibility of any light-absorbing complex formation between amine substrates and base (Fig. S29†). Therefore, the only light-absorbing species in the reaction mixture is the water- O_2 complex which absorbs 370 nm light to undergo CT excitation.^{31–37} Our findings revealed that this CT state subsequently produced O_2 ($a^1\Delta_g$) upon relaxation. Previously, it has been reported that amines are good quenchers of O_2 ($a^1\Delta_g$).¹⁵ Generally, amines with low ionization potential readily transfer their electrons to O_2 ($a^1\Delta_g$) and get converted into a radical cation. To check this possibility, we monitored the formation of amine radicals using 2,6-di-*tert*-butyl-4-methylphenol (BHT) as a radical scavenger.⁵⁵ The reaction was set up with 0.05 mmol *p*-toluidine and 2 equiv. K_3PO_4 in 5 mL 1 : 1 MeOH- H_2O mixture due to the low solubility of BHT in aqueous medium. Subsequently, after the addition of 1 equiv. BHT, the reaction mixture was irradiated with 370 nm light for 90 min (Fig. 3A). The product was isolated and characterized using HR-MS and GC-MS measurements. A characteristic HR-MS peak for the *p*-toluidine-BHT adduct was observed at 326.2441 g mol⁻¹ (Fig. S30†).

Moreover, the presence of BHT resulted in only a trace amount of the azo product even after 90 min of irradiation. These observations clearly validate the electron transfer mechanism from amines to O_2 ($a^1\Delta_g$) and authenticates the formation of amine radicals. On the other hand, to confirm the formation of superoxide radicals after the electron transfer to O_2 ($a^1\Delta_g$), we performed EPR measurements using DMPO as a probe under standard reaction conditions (Fig. 3B).⁵⁹ The EPR spectrum of pure DMPO under 370 nm irradiation showed no noticeable EPR peaks. In contrast, a characteristic EPR spectrum of superoxide radicals with six well-resolved peaks was observed upon addition of DMPO into the reaction mixture under 370 nm irradiation (Fig. 3B), which matches well with the earlier literature.⁵⁹ This observation authenticates the



Scheme 3 (A) Control reactions with 0.05 mmol nitrosobenzene in the presence and absence of 0.05 mmol aniline under standard reaction conditions. (B) Scalability experiments for the conversion of *p*-toluidine to its corresponding azo compound.




Fig. 3 (A) Schematic of the radical trap experiment during the oxidative coupling of 0.05 mmol *p*-toluidine. (B) Changes in the EPR spectra of DMPO (50 mM) in the absence and presence of the reaction mixture under the irradiation of a 370 nm LED in the presence of 0.05 mmol *p*-toluidine and 2 equiv. K_3PO_4 . (C) Daylight photographs of peroxide test strips under different conditions (i) control blank strip, (ii) before the reaction, and (iii) after 90 min of reaction with 0.05 mmol *p*-toluidine under standard reaction conditions.



generation of superoxide radicals through single electron transfer from amines to O_2 ($a^1\Delta_g$) during the reaction. Altogether, these results confirm the formation of amine radicals and superoxide ions *via* single electron transfer.


The oxidative coupling of arylamines involves the radical coupling of amines to generate hydrazobenzene as an active intermediate.^{54,55} To check the formation of hydrazobenzene under standard reaction conditions, we performed LC-MS measurement after 10 min of irradiation under a 370 nm LED. Notably, a characteristic LC-MS peak for 1,2-bis(4-methylphenyl)hydrazine was observed at an m/z value of 213.2037 $g\ mol^{-1}$ (Fig. S31†). Finally, to know whether H_2O_2 is generated in the reaction mixture from superoxide radicals,^{54,55} we utilized H_2O_2 sensing strips. These colorless strips were dipped into the reaction mixtures before and after the reaction (Fig. 3C(i)). No color change of the strip was observed upon dipping it inside the reaction mixture before light irradiation (Fig. 3C(ii)). However, a distinct color change of the sensing strip from colorless to blue was noticed after the irradiation of the reaction mixture with 370 nm light (Fig. 3C(iii)). This clearly indicates the formation of H_2O_2 in the reaction mixture.

In accordance with our findings, we propose a plausible reaction mechanism for the oxidative coupling of arylamines to azoaromatics under 370 nm light irradiation (Scheme 4). The solvent (S)- O_2 ($X^3\Sigma_u^-$) ground state complex absorbs 370 nm light to undergo excitation at the CT state ($S^+O_2^{*-}$).^{31–37} This CT state subsequently relaxed to yield O_2 ($a^1\Delta_g$). Owing to its strong oxidizing nature, the photogenerated O_2 ($a^1\Delta_g$) readily accepts a single electron from the amine substrates to form superoxide radicals and amine radicals. These amine radicals form the hydrazobenzene intermediate *via* a coupling reaction after proton abstraction by a base. After another cycle of electron transfer in the presence of O_2 ($a^1\Delta_g$) and subsequent proton abstraction, these hydrazobenzene intermediates ultimately

converted to the desired azo products with the generation of 2 equiv. of H_2O_2 as the byproduct. Therefore, the overall reaction proceeds *via* the production of 4 electrons and 4 protons as reported previously in bulk acetonitrile and micellar medium in the presence of PCs.^{54,55} The estimated quantum yield for the photooxidative conversion of *p*-toluidine to its corresponding azo compound under our standard reaction conditions is estimated to be ~ 0.15 , which substantiates that the present reaction does not follow the radical chain mechanism as proposed previously for photocatalytic Diels–Alder cycloaddition reactions.⁶⁰ To the best of our knowledge, this is the first report where the PS-free generation of O_2 ($a^1\Delta_g$) has been utilized to develop a catalyst-free approach for the selective synthesis of azo compounds in the aqueous medium (Table S1†). The present study paves the way for the development of other sustainable chemical transformations *via* utilizing a direct CT transition involving molecular O_2 in a diverse range of solvents.

Conclusions

In the present study, we have developed an efficient, sustainable, and viable synthetic approach for the oxidative coupling of arylamines to azoaromatics under ambient conditions in aqueous medium without any PCs/PSs. The present approach is based on the efficient generation of O_2 ($a^1\Delta_g$) *via* CT excitation of the water- O_2 complex at 370 nm. The formation of O_2 ($a^1\Delta_g$) has been authenticated using various spectroscopic measurements including the EPR technique. The photogenerated O_2 ($a^1\Delta_g$) has been utilized as a sustainable oxidant for the oxidative coupling of arylamines in the presence of a base under an ambient atmosphere. The reaction proceeds *via* the formation of amine radicals which undergo a coupling reaction to yield hydrazobenzene as an intermediate. These hydrazobenzene intermediates finally transform into azo products after another



Scheme 4 Plausible reaction mechanism for the oxidative coupling of arylamines *via* PS-free singlet oxygen.



cycle of electron transfer and proton abstraction steps with the formation of H₂O₂ as the only byproduct. Our methodology shows broad substrate scope with fast kinetics (90 min) excellent yields (up to 100%), and 100% selectivity under an ambient atmosphere in aqueous medium, and can be easily scaled up to the gram scale without any unwanted toxic byproducts. The present study lays the foundations for further in-depth understanding of the hidden role of O₂ (a¹Δ_g) generated directly *via* CT excitation of the solvent–O₂ complex toward a diverse range of chemical and biochemical transformations.

Data availability

The additional data supporting this article have been included as part of the ESI.†

Author contributions

T. K. M. and S. S. conceptualized the research problem and designed the experiments. S. S. performed the experiments. S. S. and T. K. M. analyzed the data. All authors contributed to the final version of the manuscript (writing – review & editing).

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors acknowledge the Indian Institute of Technology (IIT) Indore for providing financial support and infrastructure. T. K. M. is grateful to the Science and Engineering Research Board (SERB) for a generous research grant (CRG/2023/003004). The authors acknowledge SIC, IIT Indore, and SAIF, IIT Bombay for instrumental facilities. S. S. acknowledges the Prime Minister's Research Fellowship (PMRF), India for research fellowship.

Notes and references

- 1 D. S. Ovoshchnikov, B. G. Donoeva and V. B. Golovko, *ACS Catal.*, 2015, **5**, 34–38.
- 2 S. Suleman, Y. Zhang, Y. Qian, J. Zhang, Z. Lin, Ö. Metin, Z. Meng and H. -L. Jiang, *Angew. Chem., Int. Ed.*, 2024, **63**, e202314988.
- 3 L. Guo, E. Cui, H. Li, C. -H. Tung and Y. Wang, *ACS Sustainable Chem. Eng.*, 2021, **9**, 16670–16677.
- 4 Y. -Z. Chen, Z. U. Wang, H. Wang, J. Lu, S. -H. Yu and H. -L. Jiang, *J. Am. Chem. Soc.*, 2017, **139**, 2035–2044.
- 5 A. Sagadevan, K. C. Hwang and M. -D. Su, *Nat. Commun.*, 2017, **8**, 1812.
- 6 A. A. Ghogare and A. Greer, *Chem. Rev.*, 2016, **116**, 9994–10034.
- 7 D. A. Singleton, C. Hang, M. J. Szymanski, M. P. Meyer, A. G. Leach, K. T. Kuwata, J. S. Chen, A. Greer, C. Foote and K. N. Houk, *J. Am. Chem. Soc.*, 2003, **125**, 1319–1328.
- 8 M. Hayyan, M. A. Hashim and I. M. Al Nashef, *Chem. Rev.*, 2016, **116**, 3029–3085.
- 9 S. Loeb, R. Hofmann and J. -H. Kim, *Environ. Sci. Technol. Lett.*, 2016, **3**, 73–80.
- 10 J. S. Nam, M. -G. Kang, J. Kang, S. -Y. Park, S. J. C. Lee, H. -T. Kim, J. K. Seo, O. -H. Kwon, M. H. Lim, H. -W. Rhee and T.-H. Kwon, *J. Am. Chem. Soc.*, 2016, **138**, 10968–10977.
- 11 M. Niedre, M. S. Patterson and B. C. Wilson, *Photochem. Photobiol.*, 2002, **75**, 382–391.
- 12 C. G. Hübner, A. Renn, I. Renge and U. P. Wild, *J. Chem. Phys.*, 2001, **115**, 9619–9622.
- 13 S. Oelckers, T. Ziegler, I. Michler and B. Röder, *J. Photochem. Photobiol., B*, 1999, **53**, 121–127.
- 14 R. S. Mulliken, *Phys. Rev.*, 1928, **32**, 186–222.
- 15 C. Schweitzer and R. Schmidt, *Chem. Rev.*, 2003, **103**, 1685–1757.
- 16 J. F. Lovell, T. W. B. Liu, J. Chen and G. Zheng, *Chem. Rev.*, 2010, **110**, 2839–2857.
- 17 M. C. DeRosa and R. J. Crutchley, *Coord. Chem. Rev.*, 2002, **233–234**, 351–371.
- 18 P. Maillard, P. Krausz, C. Giannotti and S. Gaspard, *J. Organomet. Chem.*, 1980, **197**, 285–290.
- 19 G. S. Cox, D. G. Whitten and C. Giannotti, *Chem. Phys. Lett.*, 1979, **67**, 511–515.
- 20 C. Tanielian and C. Wolff, *J. Phys. Chem.*, 1995, **99**, 9825–9830.
- 21 R. Schmidt and C. Tanielian, *J. Phys. Chem. A*, 2000, **104**, 3177–3180.
- 22 R. Vankayala, A. Sagadevan, P. Vijayaraghavan, C. -L. Kuo and K. C. Hwang, *Angew. Chem., Int. Ed.*, 2011, **50**, 10640–10644.
- 23 R. Ho-Wu, S. Y. Yau and T. Goodson III, *J. Phys. Chem. B*, 2017, **121**, 10073–10080.
- 24 D. Li, X. Li, T. Zhao, H. Liu, S. Jiang, Q. Zhang, H. Ågren and G. Chen, *ACS Nano*, 2020, **14**, 12596–12604.
- 25 A. A. Gorman and M. A. J. Rodgers, *J. Am. Chem. Soc.*, 1986, **108**, 5074–5078.
- 26 J. Bendig, R. Schmidt and H. -D. Brauer, *Chem. Phys. Lett.*, 1993, **202**, 535–541.
- 27 A. Schlachter, P. Asselin and P. D. Harvey, *ACS Appl. Mater. Interfaces*, 2021, **13**, 26651–26672.
- 28 J. A. Jackson, C. Turro, M. D. Newsham and D. G. Nocera, *J. Phys. Chem.*, 1990, **94**, 4500–4507.
- 29 J. Li and T. Chen, *Coord. Chem. Rev.*, 2020, **418**, 213355.
- 30 V. -N. Nguyen, Y. Yan, J. Zhao and J. Yoon, *Acc. Chem. Res.*, 2021, **54**, 207–220.
- 31 R. D. Scurlock and P. R. Ogilby, *J. Am. Chem. Soc.*, 1988, **110**, 640–641.
- 32 E. A. Gooding, K. R. Serak and P. R. Ogilby, *J. Phys. Chem.*, 1991, **95**, 7868–7871.
- 33 R. D. Scurlock and P. R. Ogilby, *J. Phys. Chem.*, 1989, **93**, 5493–5500.
- 34 P. -G. Jensen, J. Arnbjerg, L. P. Tolbod, R. Toftegaard and P. R. Ogilby, *J. Phys. Chem. A*, 2009, **113**, 9965–9973.
- 35 M. Bregnhøj and P. R. Ogilby, *J. Phys. Chem. A*, 2019, **123**, 7567–7575.



- 36 M. Bregnhøj, M. Westberg, B. F. Minaev and P. R. Ogilby, *Acc. Chem. Res.*, 2017, **50**, 1920–1927.
- 37 H. Tsubomura and R. S. Mulliken, *J. Am. Chem. Soc.*, 1960, **82**, 5966–5974.
- 38 H. Murakami, A. Kawabuchi, K. Kotoo, M. Kunitake and N. Nakashima, *J. Am. Chem. Soc.*, 1997, **119**, 7605–7606.
- 39 A. Bafana, S. S. Devi and T. Chakrabarti, *Environ. Rev.*, 2011, **19**, 350–371.
- 40 E. Merino, *Chem. Soc. Rev.*, 2011, **40**, 3835–3853.
- 41 A. Corma, P. Concepción and P. Serna, *Angew. Chem., Int. Ed.*, 2007, **46**, 7266–7269.
- 42 A. Grierrane, A. Corma and H. García, *Science*, 2008, **322**, 1661–1664.
- 43 C. Zhang and N. Jiao, *Angew. Chem., Int. Ed.*, 2010, **49**, 6174–6177.
- 44 A. Grierrane, A. Corma and H. Garcia, *Nat. Protoc.*, 2010, **5**, 429–438.
- 45 S. Cai, H. Rong, X. Yu, X. Liu, D. Wang, W. He and Y. Li, *ACS Catal.*, 2013, **3**, 478–486.
- 46 B. Dutta, S. Biswas, V. Sharma, N. O. Savage, S. P. Alpay and S. L. Suib, *Angew. Chem., Int. Ed.*, 2016, **55**, 2171–2175.
- 47 M. Wang, J. Ma, M. Yu, Z. Zhang and F. Wang, *Catal. Sci. Technol.*, 2016, **6**, 1940–1945.
- 48 Y. Zou, M. Zhang, F. Cao, J. Li, S. Zhang and Y. Qu, *J. Mater. Chem. A*, 2021, **9**, 19692–19697.
- 49 S. Han, Y. Cheng, S. Liu, C. Tao, A. Wang, W. Wei, H. Yu and Y. Wei, *Angew. Chem., Int. Ed.*, 2021, **60**, 6382–6385.
- 50 J. Qin, Y. Long, F. Sun, P. -P. Zhou, W. D. Wang, N. Luo and J. Ma, *Angew. Chem., Int. Ed.*, 2022, **61**, e202112907.
- 51 H. Zhu, X. Ke, X. Yang, S. Sarina and H. Liu, *Angew. Chem., Int. Ed.*, 2010, **49**, 9657–9661.
- 52 X. Guo, C. Hao, G. Jin, H. -Y. Zhu and X. -Y. Guo, *Angew. Chem., Int. Ed.*, 2014, **53**, 1973–1977.
- 53 B. Mondal and P. S. Mukherjee, *J. Am. Chem. Soc.*, 2018, **140**, 12592–12601.
- 54 J. D. Sitter and A. K. Vannucci, *J. Am. Chem. Soc.*, 2021, **143**, 2938–2943.
- 55 S. Singh, V. Agarwal, T. K. Sarma and T. K. Mukherjee, *Green Chem.*, 2023, **25**, 9109–9114.
- 56 H. H. Wasserman, J. R. Scheffer and J. L. Cooper, *J. Am. Chem. Soc.*, 1972, **94**, 4991–4996.
- 57 I. Jahović, Y. Yang, T. K. Ronson and J. R. Nitschke, *Angew. Chem., Int. Ed.*, 2023, **62**, e202309589.
- 58 Y. Yamakoshi, N. Umezawa, A. Ryu, K. Arakane, N. Miyata, Y. Goda, T. Masumizu and T. Nagano, *J. Am. Chem. Soc.*, 2003, **125**, 12803–12809.
- 59 L. Ren, M. M. Yang, C. H. Tung, L. Z. Wu and H. Cong, *ACS Catal.*, 2017, **7**, 8134–8138.
- 60 M. A. Cismesia and T. P. Yoon, *Chem. Sci.*, 2015, **6**, 5426–5434.

