

Cite this: *Chem. Sci.*, 2024, 15, 17453

All publication charges for this article have been paid for by the Royal Society of Chemistry

Mechanochemical generation of aryl barium nucleophiles from unactivated barium metal†

Koji Kubota,^{ID} *^{ab} Sota Kawamura,^a Julong Jiang,^c Satoshi Maeda^{ID} ^{bc} and Hajime Ito^{ID} *^{ab}

Organobarium reagents are of interest as homologues of the Grignard reagents based on organomagnesium compounds due to their unique reactivity as well as regio- and stereoselectivity. However, reactions involving organobarium reagents are less developed in comparison to reactions involving Grignard reagents due to the lack of a simple and economical synthetic method and their high reactivity. To the best of our knowledge, there is no established method for the direct synthesis of organobarium compounds from commercially available bulk barium metal and organic halides. So far, the generation of organobarium compounds usually requires the use of activated barium (Rieke barium), which significantly reduces the practical utility of organobarium reagents and hinders the development of new organobarium-mediated transformations. Here, we present a mechanochemical strategy based on ball-milling that facilitates the direct generation of various aryl barium nucleophiles from commercially available unactivated barium metal and aryl halides without complicated pre-activation processes. Our simple mechanochemical protocol allows the rapid development of novel carbon–silicon-bond-forming reactions with hydrosilanes mediated by aryl barium nucleophiles; importantly, these reactions are difficult to achieve using other Grignard-type carbon nucleophiles. To the best of our knowledge, this is the first example of a nucleophilic substitution reaction involving an aryl barium species. Furthermore, this mechanochemical strategy established the first example of a nucleophilic addition to a carbonyl compound involving an aryl barium nucleophile. Preliminary theoretical calculations using the artificial force-induced reaction (AFIR) method to reveal the reaction mechanism of the hydrosilane arylation are also described.

Received 10th August 2024
Accepted 21st September 2024

DOI: 10.1039/d4sc05361c

rsc.li/chemical-science

Introduction


Since the pioneering study on the preparation of allyl barium species and their use in the α -selective allylation of carbonyl compounds by Yamamoto and Yanagisawa in 1991, organobarium nucleophiles have attracted substantial attention in synthetic chemistry and natural-product synthesis. Their unique reactivity as well as regio- and stereoselectivity offer several advantages over Grignard reagents prepared from magnesium, which, like barium, is an alkaline-earth-group element.^{1–3} Organobarium nucleophiles are typically prepared *via* the direct insertion of barium metal into a carbon–halogen bond (direct synthesis; Scheme 1A).² However, due to the

relatively high atomization energy of barium, such direct insertions are fundamentally challenging,⁴ and direct reactions with commercial barium metal are usually unsuccessful (Scheme 1A). Instead, the preparation of activated barium metal

A. Conventional method for the generation of organobarium nucleophiles



B. Mechanochemical generation of organobarium nucleophiles



^aDivision of Applied Chemistry and Frontier Chemistry Center, Faculty of Engineering, Hokkaido University, Sapporo, Hokkaido, Japan. E-mail: kbt@eng.hokudai.ac.jp; hajito@eng.hokudai.ac.jp

^bInstitute for Chemical Reaction Design and Discovery (WPI-ICReDD), Hokkaido University, Sapporo, Hokkaido, Japan

^cDepartment of Chemistry, Faculty of Science, Hokkaido University, Sapporo, Hokkaido, 060-0810, Japan

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc05361c>

Scheme 1 Generation of barium-based heavy Grignard reagents *via* the direct insertion of barium metal into carbon–halogen bonds.


(Rieke barium) under harsh conditions using alkali metals is required (Scheme 1A).⁵ Unfortunately, this requirement greatly reduces the practical utility of organobarium nucleophiles. These synthetic difficulties have most likely hampered more extensive investigations into the reactivity of barium-based carbon nucleophiles, and, as a result, their basic reactivity may have remained underappreciated. Although the first synthesis of phenyl barium iodide was reported by Gilman and Schulze almost a century ago,⁶ systematic explorations of the reactivity of aryl barium nucleophiles toward electrophiles have not yet been reported.^{2,7} Therefore, the development of an operationally simple protocol using commercial barium metal would be highly desirable to promote its widespread synthetic applications and enable the rapid development of new reactions that involve organobarium nucleophiles.

Recently, mechanochemical strategies using ball-milling techniques have attracted significant attention as alternatives to conventional solution-based approaches.⁸ The advantages of ball-milling include the reduced usage of harmful organic solvents, faster reaction kinetics, and significantly increased operational simplicity. Recent studies have also revealed that strong mechanical agitation during the ball-milling process can activate zero-valent bulk metals, thus facilitating the direct metalation of organic halides.^{9–15} In the area of the mechanical activation of alkaline earth metals, the groups of Harrowfield,¹¹ Birke,¹² Hanusa,¹³ Bolm,¹⁴ and Kananovich,¹⁵ as well as our own group,^{10e} have independently reported the mechanochemical synthesis of Grignard reagents *via* the direct insertion of magnesium metal into organic halides. More recently, our group has demonstrated that ball-milling allows the generation of heavy Grignard reagents based on calcium from aryl halides and unactivated calcium metal, which is unreactive under solution conditions.^{10e} Importantly, neither pre-activation processes nor the preparation of Rieke metals is required in these cases. These recent achievements inspired us to consider the mechanochemically driven generation of organobarium reagents *via* direct synthesis from unactivated commercial barium metal, which is sparingly reactive in solution (Scheme 1B). Based on the relative atomization energies of alkaline-earth metals, the activation of barium metal (atomization enthalpy $\approx 182 \text{ kJ mol}^{-1}$) is more challenging than that of magnesium ($\sim 146 \text{ kJ mol}^{-1}$) or calcium ($\sim 178 \text{ kJ mol}^{-1}$) metal.⁴ However, we envisaged that the strong mechanical impact during ball-milling could allow sufficient surface activation to overcome the high atomization energy of barium metal and thus enable the direct generation of barium-based heavy Grignard reagents without complicated operating conditions (Scheme 1B).


Results and discussion

As a proof-of-concept study, we focused on the mechanochemically driven generation of aryl barium species from 4-iodo-1,1'-biphenyl (**1a**) (1.0 equiv.) and commercial barium metal (Aldrich, >99.99% purity; 1.0 equiv.), using a Retch MM400 mixer mill (5 mL stainless-steel milling jar with two 10 mm diameter stainless-steel balls; Scheme 2A). The reaction of **1a** was evaluated by monitoring the formation of the

A. Mechanochemical activation of Ba metal



B. Unsuccessful result under solution conditions



Scheme 2 Mechanochemically driven generation of aryl barium nucleophiles. ^a For details of the reaction conditions, see the ESI.†

protonation product (**2a**) by treatment with aqueous HCl (1.0 M) (Scheme 2A). After an optimization study (for details, see the ESI†), we found that the addition of 2.0 equiv. of tetrahydrofuran (THF) was crucial to facilitate the formation of the organobarium nucleophile under mechanochemical conditions; the corresponding protonation product (**2a**) was obtained in 53% yield. Based on a previous study on the isolation of THF-coordinated aryl barium species by Westerhausen *et al.*,^{7a} we anticipated that THF might enhance the thermodynamic stability of the organobarium compound by coordinating to the metal center. However, the fact that full conversion of **1a** was observed despite the moderate yield of **2a** suggests significant decomposition of the organobarium species during the reaction. We found that reactions with aryl bromides and chlorides resulted in poor conversion of the starting materials (for details, see the ESI†). To confirm the effectiveness of mechanical activation *via* ball-milling, we carried out a reaction between barium metal and **1a** in THF at room temperature (Scheme 2B). Under solution conditions, the formation of **2a** was not detected, and a large amount of unreacted **1a** remained (>95% recovery) (Scheme 2B). This result indicates that the strong mechanical agitation provided by ball-milling is crucial for the activation of barium metal,^{9,10} to enable the subsequent generation of the corresponding aryl barium species.

Having demonstrated the ability of this mechanochemical protocol to rapidly generate aryl barium nucleophiles from unactivated barium metal, we next investigated the development of new reactions using these nucleophiles. We hoped that the rapid screening approach enabled by the mechanochemical method would allow the discovery of new reactions with aryl barium nucleophiles.^{10e} First, a series of mechanochemical reactions with various electrophiles was evaluated (Table 1). When benzaldehyde (**3a**) or an aliphatic aldehyde (**3b**) was used as an *in situ* trapping reagent for the formed aryl barium nucleophile, the formation of the desired nucleophilic-addition products was not observed. Instead, a reduction of the carbonyl group or a Tishchenko reaction proceeded, probably triggered by reactions between the carbonyl compounds and barium



Table 1 Reactions with other electrophiles^a

 1a 4.0 equiv	 3
 4	
 3a complex mixture	 3b complex mixture
 3c <1% yield	 3d 24% (NMR)
 3e 84% (isolated)	

^a Conditions: 3 (0.25 mmol), 1a (1.0 mmol), Ba metal (1.0 mmol), and THF (2.0 mmol) in a stainless-steel milling jar (5 mL) with two stainless-steel balls (10 mm).

metal. To prevent these undesired side reactions, we used reduction-resistant trialkoxyboron reagent 3c as the electrophile,¹⁶ albeit that the desired borylation product was not detected. However, we discovered that the use of chlorodimethylphenyl silane (3d) afforded the corresponding silylation product (4a) in 24% yield. The low yield of 4a is most likely due to the competing undesired reduction of a Cl–Si bond by barium metal.¹⁷ Therefore, we then tested dimethylphenylsilane (3e), which is significantly less reactive than the chlorosilane under reductive conditions, and 4a was obtained in 84% yield.¹⁸ Notably, this is the first example of a nucleophilic substitution reaction involving an aryl barium species.^{2,7}

Next, the scope of aryl iodides for the mechanochemically driven generation of aryl barium nucleophiles was investigated (Table 2). Substrates with *ortho*-substituents, such as a methyl (1b), *i*-propyl (1c), or phenyl (1d) group, furnished the desired silylation products (4b–4d) in high yield (82–91%). 1-Iodonaphthalene (1e) was also converted into the corresponding product (4e) in good yield (82%). Substrates with an electron-donating group, such as a methoxy (1f) or amino (1g) group, were tolerated and furnished the desired products (4f and 4g) in moderate yield (61% and 56%, respectively). The reactions of aryl iodides with an electron-withdrawing group, such as trifluoromethyl (1h) or fluorine (1i), afforded the corresponding products (4h and 4i) in low-to-moderate yield (48% and 36%, respectively). Substrates 1h and 1i were fully consumed; the relatively low yield of 4h and 4i is probably due to the lower nucleophilicity of the corresponding aryl barium species. We also carried out the silylation reaction of 4-bromobiphenyl (1j) under mechanochemical conditions, which afforded the desired silylation product (4a) in moderate yield (51%).

Next, we turned our attention to the scope of applicable hydrosilane reagents (Table 3). We found that trialkyl-substituted hydrosilanes (3g–3i) are applicable to this mechanochemical protocol, and the desired products (4k–4m) were obtained in moderate-to-good yield (69%, 43%, and 46% yield, respectively). Trimethylsilyl-substituted hydrosilane (3j) furnished the corresponding product (4n) in 17% yield.

Table 2 Scope of aryl iodides^{a,b}

 1 4.0 equiv	 3e
 4 isolated yield (%)	
 4a 84% yield (51%) ^c	 4b 91% yield
 4c 82% yield	 4d 82% yield
 4e 82% yield	 4f 61% yield
 4g 56% yield	 4h 48% yield
 4i 36% yield	

^a For details of the reaction conditions, see the ESI. ^b Isolated yields are shown. ^c 4-Bromobiphenyl (1j) was used as a substrate and barium metal (8.0 equiv.) was used. The ¹H NMR yield is shown.

To confirm that this reactivity is unique to organobarium species, we conducted the mechanochemical reaction using other alkaline-earth metals (Table 4). The use of magnesium instead of barium showed poor results (4b: <1%; 4j: 18%; 4k: 2%).^{10c,11–15} The reactions in the presence of calcium metal

Table 3 Scope of hydrosilanes^{a,b}

 1 4.0 equiv	 3
 4 isolated yield (%)	
 4j 87% yield	 4k 69% yield
 4l 43% yield	 4m 46% yield
 4n 17% yield	

^a For details of the reaction conditions, see the ESI. ^b Isolated yields are shown.



Table 4 Comparison of the reactivity relative to other alkaline-earth metals in mechanochemically driven hydrosilane arylation.^{a,b}



<p>1b 4.0 equiv</p>	<p>3</p>	<p>4</p>
<p>5 mL milling jar 10 mm ball x 2 (stainless steel) ball-milling (30 Hz) 30 min, in air</p>		
<p>Electronegativity: Mg (1.31), Ca (1.00), Ba (0.89)</p>		
<p>4b</p> <p>Mg: <1% (NMR) Ca: 43% (NMR) Ba: 91% (isolated)</p>	<p>4j</p> <p>Mg: 18% (NMR) Ca: 38% (NMR) Ba: 87% (isolated)</p>	<p>4k</p> <p>Mg: 2% (NMR) Ca: 33% (NMR) Ba: 69% (isolated)</p>

^a Conditions: **3e** (0.25 mmol), **1b** (1.0 mmol), metal (1.0 mmol), and THF (2.0 mmol) in a stainless-steel milling jar (5 mL) with two stainless-steel balls (10 mm). ^b NMR yields are shown.

provided the products in much lower yield (**4b**: 43%; **4j**: 38%; **4k**: 33%) than those in the presence of barium metal (**4b**: 88%; **4j**: 87%; **4k**: 69%).^{10c} Considering the relative electronegativity values of these metals (Table 4), these results suggest that the higher nucleophilicity of the aryl barium nucleophiles compared to those of the other Grignard-type reagents facilitates the substitution of hydrosilanes to form silylation products.

A reaction using Rieke barium under solution-based conditions was also performed for comparison with the mechanochemical method (Scheme 3).⁵ The Rieke method using lithium biphenylide was applied to generate the corresponding aryl barium species from **1b**, and the desired silylation product (**4b**) was obtained in 56% yield (Scheme 3). This result suggests that barium-based heavy Grignard reagents (Ar–BaX) similar to those present in solution are likely generated under mechanochemical conditions and subsequently react with hydrosilanes to form the silylation products.

To obtain mechanistic insight into the mechanochemically driven organobarium-mediated silylation, we investigated the reaction with chiral hydrosilane (*R*)-**3k** (>99% ee) (Scheme 4). (*R*)-**3k** was prepared *via* preparative chiral high-performance liquid chromatography (HPLC) separation.¹⁹ The reaction between **1e** and (*R*)-**3k** in the presence of barium metal under mechanochemical conditions afforded (*S*)-**4o** with 78% ee.

**Scheme 3** Solution-based reaction using Rieke barium.^a For details of the reaction conditions, see the ESI.†**Scheme 4** Mechanochemically driven hydrosilane arylation with chiral hydrosilane **3k**.^a For details of the reaction conditions, see the ESI.†

Although the stereospecificity is not perfect, the reaction mainly proceeded with retention of the configuration of the silicon stereogenic center, excluding a simple S_N2 type mechanism.

To gain deeper insight into the reaction mechanism, we conducted a preliminary computational study using the artificial force-induced reaction (AFIR) method.²⁰ Unlike conventional density-functional-theory (DFT) calculations, in which proposed structures and mechanisms are required, the AFIR method allows the comprehensive and unbiased exploration of chemical structures and reaction pathways.²⁰ Since organobarium compounds often form a dimeric complex,²¹ we first studied the monomeric and dimeric Ar–BaI (Ar = *o*-tolyl) species, as well as their reaction with hydrosilane. For the monomeric Ar–BaI(thf)₂ species, the sampling calculation using the SC-AFIR method revealed 1856 possible structures, while 3729 possible structures were suggested for the dimeric species. Further calculations confirmed that the dimeric species is significantly more stable than the monomeric one (Fig. 1A). The most stable dimeric Ar–BaI geometry was then subjected to reaction with Me₂PhSiH. In the calculated mechanism, when the silane approaches the dimeric Ar–BaI species, it immediately becomes a ligand on the barium species (Fig. 1A). In the optimized structure of intermediate **I**, Me₂PhSiH coordinates to two barium atoms simultaneously, with its phenyl ring acting as a π -ligand to form a half-sandwich complex and its hydride forming a hydrogen-bridged bond with another barium center.²¹ Moreover, the hydride is extracted in a unique way *via* a five-coordinate silicon species (*cf.* the structures of **II** and **TS1**). This is indeed not an S_N2-type reaction, and retention of the configuration of the silicon stereogenic center is assured. A further NBO analysis clearly indicated that the 3d-orbital of the Si atom plays a critical role in forming a penta-coordinated Si complex and therefore allows this unique substitution pathway (for details, see the ESI†). Importantly, these results are consistent with the experimental observations discussed in Scheme 4. In addition, when the silicon–hydrogen bond is completely cleaved *via* **TS1**, the hydride becomes an H-bridge connecting both barium atoms, and the iodide-bridged bonds no longer exist. Interestingly, once the new silicon–carbon bond is formed, the tolyl moiety then becomes another π -ligand to form a half-sandwich complex with a barium center (*cf.* the structure of **III**), suggesting very unique bonding behavior of the barium atom. It is noteworthy that similar single-crystal X-ray structures have already been reported.²²








Fig. 1 Theoretical calculations (B3LYP-D3/6-311G(d,p)-SDD) of the aryl-barium-mediated hydrosilane arylation. Bond lengths are given in Å.

Encouraged by the success of the silylation reactions, we furthermore explored the development of new carbon-carbon bond-forming reactions using the mechanochemically generated aryl barium species. After a quick re-screening of electrophiles, we discovered that the reaction of **1a** with diaryl ketone **5a** proceeded smoothly to give the desired tertiary alcohol **6a** in


83% isolated yield (>99% ^1H NMR yield) (Scheme 5). To the best of our knowledge, this is the first example of a nucleophilic addition to a carbonyl compound involving an aryl barium nucleophile. Notably, other possible byproducts such as a pinacol coupling product **7a** and the simple ketone-reduction product **8a**, which can be generated through a single-electron reduction with barium metal, were not detected in the crude reaction mixture. This Barbier-type arylation of diaryl ketones is a difficult transformation when magnesium metal is used as a reductant due to the formation of these competing byproducts. Indeed, we found that the use of magnesium metal instead of barium metal under mechanochemical conditions resulted in the formation of a mixture of these products (Scheme 5). This result highlights the unique and distinctive reactivity of barium metal under the applied mechanochemical conditions.

Next, the substrate scope regarding the ketone arylation was explored (Table 5). The mechanochemically generated aryl barium nucleophile reacted with various diaryl ketones (**5a–5e**) to deliver the desired alcohols (**6a–6e**) in good to excellent yield (69–94%). Reactions using sterically hindered aryl iodides (**1b** and **1c**) proceeded smoothly to give the corresponding addition products (**6f** and **6g**) in good yield (72% and 66% yield, respectively). It is important to note that the reaction of sterically hindered **1c** in the presence of magnesium metal afforded the desired product **6g** in poor yield (31%) and significant amounts of other byproducts were formed (pinacol coupling byproduct **7g**: 24%; reduction byproduct **8g**: 22%; for details, see the ESI†).


To demonstrate the practical utility of this method, we also investigated a scaled-up reaction (Scheme 6). For that purpose, a ketone-arylation reaction was conducted on the 2.5 mmol scale in a 10 mL stainless-steel jar with two 10 mm stainless-steel balls. The desired product (**6a**) was obtained in 77%



Scheme 5 Barbier-type ketone arylation involving mechanochemically generated aryl barium species. ^a For details of the reaction conditions, see the ESI.†

Table 5 Scope of Barbier-type ketone arylation^a

^a For details of the reaction conditions, see the ESI. ^b 1 (1.5 equiv.) and Ba metal (1.5 equiv.) were used.



Scheme 6 Scaled-up reaction of 5a. ^a For details on the reaction conditions, see the ESI.†

isolated yield, which is comparable to the yield obtained from the reactions on the smaller scale. This result emphasizes the practical utility of this mechanochemical protocol.

Conclusions

In summary, we have demonstrated a mechanochemical strategy using ball-milling for the direct generation of aryl barium nucleophiles *via* the reaction between aryl halides and commercial unactivated barium metal. Importantly, this method does not require complicated pre-activation processes. Using this simple protocol, we discovered a carbon–silicon–bond-forming reaction mediated by an aryl barium species with hydrosilanes; such reactions are rather difficult to achieve using conventional Grignard reagents. To the best of our knowledge, this is the first example of a nucleophilic substitution reaction involving an aryl barium species. In addition, we developed a barium-mediated Barbier-type arylation of diaryl ketones, which is not an easy transformation when magnesium metal is used due to the competing formation of byproducts such as a pinacol coupling product and the simple ketone-reduction product. The details of the reaction mechanism for the mechanochemically driven hydrosilane arylation were discussed based on theoretical calculations using the artificial force-induced reaction (AFIR) method. Further studies to apply this mechanochemical strategy to the generation of allyl-, propargylic-, benzyl-, and simple alkyl barium species are currently in progress in our laboratory.

Data availability

The data supporting this article have been included as part of the ESI.†

Author contributions

K. K. and H. I. conceived and designed the study. K. K., J. J., S. M., and H. I. co-wrote the paper. S. K. performed the chemical experiments and analyzed the data. J. J. and S. M. conducted the theoretical study. All authors discussed the results and the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the Japan Society for the Promotion of Science (JSPS) *via* KAKENHI grants 24H00453, 24H01050, 24H01832, 22H00318, and 22K18333; by the JST *via* CREST grant JPMJCR19R1; by FOREST grant JPMJFR2011; as well as by the Institute for Chemical Reaction Design and Discovery (ICReDD), which was established by the World Premier International Research Initiative (WPI), MEXT, Japan. We thank Mr Keisuke Kondo for his help in cross-checking experiments. We sincerely thank the reviewers for their many constructive comments in revising this paper.



Notes and references

- For pioneering studies on the preparation of allyl barium species and their use in organic synthesis, see: (a) A. Yanagisawa, S. Habaue and H. Yamamoto, *J. Am. Chem. Soc.*, 1991, **113**, 8955; (b) A. Yanagisawa, S. Habaue, K. Yasue and H. Yamamoto, *J. Am. Chem. Soc.*, 1994, **116**, 6130; (c) A. Yanagisawa, K. Ogasawara, K. Yasue and H. Yamamoto, *Chem. Commun.*, 1996, 367; (d) K. Yasue, A. Yanagisawa and H. Yamamoto, *Bull. Chem. Soc. Jpn.*, 1997, **70**, 493; (e) A. Yanagisawa, Y. Yamada and H. Yamamoto, *Synlett*, 1997, **9**, 1090; (f) F. Raeppl and D. Heissler, *Synlett*, 2005, **16**, 2534; (g) A. Yanagisawa, S. Yamafuji and T. Sawae, *Synlett*, 2016, **27**, 2019; (h) A. Yanagisawa, T. Jitsukawa and K. Yoshida, *Synlett*, 2013, **24**, 635.
- For a comprehensive review on organic reactions promoted by barium reagents, see: A. Yanagisawa and K. Yoshida, *Synlett*, 2011, 2929.
- For selected studies on propargylation and benzylation with barium reagents, see: (a) A. Yanagisawa, S. Okitsu and T. Arai, *Synlett*, 2005, 1679; (b) A. Yanagisawa, T. Suzuki, T. Koide, S. Okitsu and T. Arai, *Chem.-Asian J.*, 2008, **3**, 1793; (c) A. Yanagisawa, T. Sawae, S. Yamafuji, T. Heima and K. Yoshida, *Synlett*, 2015, **26**, 1073; (d) A. Yanagisawa, T. Koide and K. Yoshida, *Synlett*, 2010, 1515.
- (a) J. S. Alexander and K. Ruhlandt-Senge, *Eur. J. Inorg. Chem.*, 2002, **2002**, 2761; (b) M. Westerhausen, M. Gärtner, R. Fischer, J. Langer, L. Yu and M. Reiher, *Chem.-Eur. J.*, 2007, **13**, 6292.
- T. C. Wu, H. Xiong and R. D. Rieke, *J. Org. Chem.*, 1990, **55**, 5045.
- H. Gilman and F. Schulze, *Bull. Soc. Chim.*, 1927, **41**, 1333.
- (a) J. Langer, M. Gärtner, R. Fischer, H. Görls and M. Westerhausen, *Inorg. Chem. Commun.*, 2007, **10**, 1001; (b) S.-O. Hauber, F. Lissner, G. B. Deacon and M. Niemeyer, *Angew. Chem., Int. Ed.*, 2005, **44**, 5871.
- For selected reviews on the use of ball-milling for organic synthesis, see: (a) S. L. James, C. J. Adams, C. Bolm, D. Braga, P. Collier, T. Friščić, F. Grepioni, K. D. M. Harris, G. Hyett, W. Jones, A. Krebs, J. Mack, L. Maini, A. G. Orpen, I. P. Parkin, W. C. Shearouse, J. W. Steed and D. C. Waddell, *Chem. Soc. Rev.*, 2012, **41**, 413; (b) G.-W. Wang, *Chem. Soc. Rev.*, 2013, **42**, 7668; (c) J.-L. Do and T. Friščić, *ACS Cent. Sci.*, 2017, **3**, 13; (d) J. G. Hernández and C. Bolm, *J. Org. Chem.*, 2017, **82**, 4007; (e) T.-X. Métro, J. Martinez and F. Lamaty, *ACS Sustainable Chem. Eng.*, 2017, **5**, 9599; (f) T. K. Achar, A. Bose and P. Mal, *Beilstein J. Org. Chem.*, 2017, **13**, 1907; (g) O. Eguagie, J. S. Vyle, P. F. Conlon, M. A. Gilea and Y. Liang, *Beilstein J. Org. Chem.*, 2018, **14**, 955; (h) J. L. Howard, Q. Cao and D. L. Browne, *Chem. Sci.*, 2018, **9**, 3080; (i) J. Andersen and J. Mack, *Green Chem.*, 2018, **20**, 1435; (j) C. Bolm and J. G. Hernández, *Angew. Chem., Int. Ed.*, 2019, **58**, 3285; (k) T. Friščić, C. Mottillo and H. M. Titi, *Angew. Chem., Int. Ed.*, 2020, **59**, 1018; (l) K. Kubota and H. Ito, *Trends Chem.*, 2020, **2**, 1066; (m) A. Porcheddu, E. Colacino, L. De Luca and F. Delogu, *ACS Catal.*, 2020, **10**, 8344; (n) J. A. Leitch and D. L. Browne, *Chem.-Eur. J.*, 2021, **27**, 9721–9726; (o) P. Ying, J. Yu and W. Su, *Adv. Synth. Catal.*, 2021, **363**, 1246; (p) K. J. Ardila-Fierro and J. G. Hernández, *ChemSusChem*, 2021, **14**, 2145; (q) V. Martinez, T. Stolar, B. Karadeniz, I. Brekalo and K. Užarević, *Nat. Rev. Chem.*, 2023, **7**, 51.
- For a tutorial review on the mechanical activation of zero-valent metals, see: A. C. Jones, J. A. Leitch, S. E. Raby-Buck and D. L. Browne, *Nat. Synth.*, 2022, **1**, 763.
- For selected examples of mechanical activation of zero-valent metals, see: (a) Q. Cao, J. L. Howard, E. Wheatley and D. L. Browne, *Angew. Chem., Int. Ed.*, 2018, **57**, 11339; (b) W. I. Nicholson, J. L. Howard, G. Magri, A. C. Seastram, A. Khan, R. R. A. Bolt, L. C. Morrill, E. Richards and D. L. Browne, *Angew. Chem., Int. Ed.*, 2021, **60**, 23128; (c) R. Takahashi, A. Hu, P. Gao, Y. Gao, Y. Pang, T. Seo, J. Jiang, S. Maeda, H. Takaya, K. Kubota and H. Ito, *Nat. Commun.*, 2021, **12**, 6691; (d) V. S. Pfennig, R. C. Villella, J. Nikodemus and C. Bolm, *Angew. Chem., Int. Ed.*, 2022, **61**, e202116514; (e) P. Gao, J. Jiang, S. Maeda, K. Kubota and H. Ito, *Angew. Chem., Int. Ed.*, 2022, **61**, e202207118; (f) R. Takahashi, P. Gao, K. Kubota and H. Ito, *Chem. Sci.*, 2023, **14**, 499; (g) N. Davison, J. A. Quirk, F. Tuna, D. Collison, C. L. McMullin, H. Michaels, G. H. Morritt, P. G. Waddell, J. A. Gould, M. Freitag, J. A. Dawson and E. Lu, *Chem*, 2023, **9**, 576; (h) K. Fujishiro, Y. Morinka, Y. Ono, T. Tanaka, L. T. Scott, H. Ito and K. Itami, *J. Am. Chem. Soc.*, 2023, **145**, 8163.
- J. M. Harrowfield, R. J. Hart and C. R. Whitaker, *Aust. J. Chem.*, 2001, **54**, 423.
- V. Birke, C. Schütt, H. Burmeier and W. K. L. Ruck, *Fresenius Environ. Bull.*, 2011, **20**, 2794.
- I. R. Speight and T. P. Hanusa, *Molecules*, 2020, **25**, 570.
- V. S. Pfennig, R. C. Villella, J. Nikodemus and C. Bolm, *Angew. Chem., Int. Ed.*, 2022, **61**, e202116514.
- J. V. Nallaparaju, T. Nikonovich, T. Jarg, D. Merzhyevski, R. Aav and D. G. Kananovich, *Angew. Chem., Int. Ed.*, 2023, **62**, e202305775.
- M. Fukuzawa, F. Takahashi, K. Nogi, T. Sasamori and H. Yorimitsu, *Org. Lett.*, 2020, **22**, 2303.
- The generation of the corresponding hydrosilane was confirmed by NMR analysis; for details, see the ESI†
- H.-J. Lee, C. Kwak, D.-P. Kim and H. Kim, *Green Chem.*, 2021, **23**, 1193.
- X. Wang, C. Feng, J. Jiang, S. Maeda, K. Kubota and H. Ito, *Nat. Commun.*, 2023, **14**, 5561.
- (a) S. Maeda, K. Ohno and K. Morokuma, *Phys. Chem. Chem. Phys.*, 2013, **15**, 3683; (b) S. Maeda, Y. Harabuchi, M. Takagi, T. Taketsugu and K. Morokuma, *Chem. Rec.*, 2016, **16**, 2232.
- M. Wiesinger, B. Maitland, C. Färber, G. Ballmann, C. Fischer, H. Elsen and S. Harder, *Angew. Chem., Int. Ed.*, 2017, **56**, 16654.
- (a) A. Uresk, N. J. O'Neil, Z. Zhou, Z. Wei and M. A. Petrukhina, *J. Organomet. Chem.*, 2019, **897**, 57; (b) P. M. Chapple and Y. Sarazin, *Eur. J. Inorg. Chem.*, 2020, **2020**, 3321.

