Chemical Science

EDGE ARTICLE

Cite this: Chem. Sci., 2024, 15, 19764

C All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 14th October 2024 Accepted 11th November 2024

DOI: 10.1039/d4sc06989g

rsc.li/chemical-science

Introduction

Atmospheric nitrogen (N_2) fixation is essential for life on Earth because it converts non-reactive N_2 into a reactive form that is suitable for crops and supports their growth.¹ Nitrate $\mathrm{[NO_3^-)}$ is one of the nitrogen fixation products that is widely used to manufacture fertilizers.² Additionally, $NO₃⁻$ is employed as a vital source of nitrogen in the food processing industry and the manufacturing of plastic and medicinal products.³ Traditionally, NO_3^- is prepared from the catalytic oxidation of ammonia (NH₃).⁴ The method suffers from the use of NH₃ which is generally prepared from the Haber–Bosch process. The process involves the use of fossil fuel and the release of large quantities of carbon dioxide $(CO₂)$.⁴ Intense research has been directed toward the goal of producing $NO₃⁻$ in a more ecofriendly way.³

Recently, reactions in water microdroplets and microbubbles have been explored as an alternative approach owing to a significant enhancement in the reaction rates and formation of unexpected products.⁵⁻¹⁷ An exceptional electric field strength $({\sim}10^9 \text{ V m}^{-1})$,¹⁸⁻²¹ the lack of three-dimensional solvation,^{22,23} the orientation of molecules at the interface, $24-26$ evaporation, 27 and pH²⁸ are several factors that drive redox reactions at the airwater interface. Taking advantage of these factors, Bose, Mofidfar, and Zare have shown that $\mathrm{NO_3}^-$ can be successfully prepared by passing N_2 microbubbles through water in the

Understanding the formation of nitrate from nitrogen at the interface of gas–water microbubbles†

Sand[e](http://orcid.org/0000-0001-5266-4253)ep Bose, Yu Xia* and Richard N. Zare D^{*}

Water microbubbles containing Fe²⁺ ions have been found to efficiently transform nitrogen (N₂) to nitrate (NO₃⁻) by initiating Fenton's reaction at the gas–water interface. Herein, we elucidate the mechanism of the formation of nitrate ($NO₃⁻$) from nitrogen (N₂) at the microbubble interface. Several experimental studies were conducted to identify the intermediates formed during the conversion. Our investigation shows the formation of $H_2N_2O_2$, NO, NO₂, and NO₂⁻ intermediates before yielding NO₃⁻ as the final product. Density functional theory (DFT) calculations provide additional support to our observation by providing insights into the energy profiles of the reaction intermediates. We believe that this work not only provides valuable insight into the abiotic nitrogen fixation in microbubbles but also helps in suggesting the modification of parameters to create a more reactive interface that leads to the enhanced production of nitric acid (HNO₃). EDGE ARTICLE
 (A) Check for undeless **Understanding the formation of nitrate from**
 $\frac{1}{2}$
 (A) Check for undeless this space. **Introgen at the interface of gas-water**

Check for understanding the **interface of gas**

presence of Fe²⁺.²⁹ The presence of ferrous ions stimulates OH^{*} production at the microbubble interface by initiating Fenton's reaction as follows: 30,31

$$
Fe^{2+} + H_2O_2 \to Fe^{3+} + OH^{\star} + OH^-
$$
 (1)

The OH^{\cdot} produced from Fenton's reaction activates the N₂ molecule which undergoes a series of transformations to form $HNO₃$ as the final product. Surprisingly, the transformations occur at room temperature and ambient pressure. Additionally, the process does not require any external electric field or radiation, and hence opens up the possibility for an eco-friendly production of HNO₃.

Experimentally, the authors have confirmed that N_2 oxidation takes place at the gas-water interface of a microbubble.²⁹ To ensure the necessity of microbubbles for the transformation, the author simply purged the solution with N_2 to create larger bubbles. No NO_3^- formation was noticed in case of larger bubbles which suggests the requirement of microbubbles for the conversion. Although the final product was analysed and quantified, the exact mechanism for that conversion was not established. The current work involves the understanding of the reaction pathways for the formation of $NO₃⁻$ from $N₂$. We have performed several experiments to recognize the intermediates formed during the process. Based on these identifications, we propose a possible reaction leading to intermediate formation. Finally, we tried to connect all the possible reactions as a sequence of continuous steps that occur to yield the final product. To validate our experimental observations, density functional theory (DFT) calculations were employed. DFT studies unravel the reaction pathways involving the

Department of Chemistry, Stanford University, CA 94305, USA. E-mail: zare@stanford. edu

[†] Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc06989g>

Experimental

The reaction system which was used for $NO₃⁻$ production consists of an air stone submerged in a beaker of aqueous FeSO₄ (50 μ M) solution. The air stone is a cylindrical porous mineral bubbler used to create microbubbles. The beaker is placed inside an ultrasonic bath along with a temperature controller. The air stone is connected to the N_2 gas supply. When N_2 gas (50 psi) is allowed to pass through the air stone, it

Fig. 1 Comparison of the 1 H-NMR spectra of the reaction product after 30 min of microbubbling and the standard $H_2N_2O_2$ solution.

produces microbubbles of N_2 in the solution.²⁹ An air stone is a porous material that is used to diffuse gas bubbles into water. The air stones (Guangzhou Zhushi Aquarium Co., Ltd, ASC-999) are manufactured from quartz sand that has been sintered in a fixed mold. The size of the output bubble depends upon the input gas pressure and the porosity of the air stone. The air stone we employed for our experiment has a porosity of 11%. Through the use of a high-speed camera, we found the average diameter of the microbubbles is \sim 530 µm and the average flux of the output bubble is ∼120 microbubbles per second for a gas pressure of 50 psi. We employed ultrasonication along with bubbling to further split microbubbles to cause them to have smaller sizes. A temperature controller was used to prevent an increase in temperature. However, evaporation still occurs over time, and water was added to the container to maintain the same level of water. An activated carbon filter was inserted in the gas flow to remove any nitrogen oxide impurities present.

Results and discussion

Before identification of intermediates, we detected H_2O_2 formation as it is important for Fenton's reaction to generate OH' for N_2 activation. The quantification of H_2O_2 was done using a potassium titanium oxalate based spectrophotometric method²⁹ which shows a continuous increase in H_2O_2 production with bubbling (Fig. S1†). The N_2 activation leading to $\text{NO}_3^$ as the final product was confirmed by using the method reported by Bose, Mofidfar, and Zare²⁹ which shows a steady increase in $NO₃⁻$ yield with bubbling time (Fig. S2†).

For the identification of intermediates involved in $\mathrm{NO_3}^$ formation, the ¹H-NMR spectrum of the reaction solutions was measured after 30 min of bubbling. The NMR spectrum shows a chemical shift of 2.69 (Fig. 1) which can be attributed to the formation of hyponitrous acid $(H_2N_2O_2)$.³² The generation of $H_2N_2O_2$ in the solution was confirmed by measuring the NMR of a standard $H_2N_2O_2$ solution which shows a peak at the same position. Thus, it can be concluded that N_2 is activated by the in

Fig. 2 (a) Schematic representation of the conversion of non-fluorescent DAF-2 to fluorescent DAF-2T in the presence of NO/NO₂ present in the medium. (b) Change in fluorescence intensity of the DAF-2 containing N₂-bubbling solution with time

situ generated OH' from Fenton's reaction which oxidizes N_2 to yield $H_2N_2O_2$ in a sequence of steps. The reactions can be shown as follows:

$$
N_2 + H' \rightarrow N_2H' + OH' \rightarrow HN_2OH + OH' \rightarrow
$$

$$
N_2OH + H_2O + OH' \rightarrow H_2N_2O_2 + H_2O
$$
 (2)

Other important intermediates identified from the microbubbled solution are nitric oxide (NO') and nitrogen dioxide $(NO₂)$. To obtain direct evidence of NO and NO₂ in the solution, we employed diaminofluorescein (DAF-2) as a fluorescent indicator. DAF-2 itself is a weakly fluorescent molecule which in the presence of NO' and oxygen undergoes a chemical transformation to yield a highly green fluorescent triazole form known as DAF-2T (Fig. 2a).³³ The reaction mechanism is proposed to involve the formation of nitrous anhydride (N_2O_3) which is produced as follows:³⁴

$$
2NO^{+} + O_{2} \rightarrow 2NO_{2}^{'} \tag{3}
$$

$$
NO' + NO'_2 \rightarrow N_2O_3 \tag{4}
$$

The *in situ* produced N_2O_3 reacts with the vicinal diamines of DAF-2 to form the triazole ring compound (DAF-2T).

To check the existence of NO^{\cdot} and NO_2^{\cdot} in our sample, we added 5 mM DAF-2 in DMSO to the microbubbled solution. As the N_2 -bubbling time is increased, we observe an increase in the fluorescence intensity ($\lambda_{\text{ex}} = 495$ nm, $\lambda_{\text{em}} = 515$ nm). The fluorescence intensity of DAF-2 is weak, and it shows substantial enhancement in fluorescence upon N_2 -bubbling for 24 h (Fig. S3†). This suggests that N_2O_3 concentration increases in the solution, and the in situ generated N_2O_3 reacts with the DAF-2 present in the solution to give DAF-2T which enhances the fluorescence (Fig. 2b).³³ Thus, it can be concluded that both NO^{*} and NO $_{\rm 2}^{\rm *}$ are generated in the microbubbles during the reaction. From eqn (2) we showed that the first steps of N_2 activation

136.0715

Fig. 3 Mass spectrum of the reactive oxygen species (ROS) and reactive nitrogen species (RNS) captured by DMPO in the N₂-microbubbled solution.

 m/z

143.0

146.0872 159.0835

170

150

130.0875

130

occurs via $H_2N_2O_2$ formation. The $H_2N_2O_2$ formed possibly reacts with the OH' generated in the system, to produce NO'. The reaction can be written as follow:

$$
H_2N_2O_2 + 2OH' \rightarrow 2NO' + 2H_2O
$$
 (5)

In living cells, NO^{\cdot} reacts with O_2 to form NO \cdot ₂ as shown in eqn (3). However, in our case, the microbubbles were produced using N_2 as feed gas and the system lacks oxygen to produce $NO₂$. Thus, a possible way by which $NO₂$ could be produced from NO' is shown by the following reaction:

$$
NO' + OH' \rightarrow HNO_2 \tag{6}
$$

$$
HNO2 + OH' \rightarrow NO2' + H2O \t(7)
$$

Fig. 4 (a) Mass spectrum of the N_2 -bubbled sample after 8 h. The mass spectrum shows a tiny peak of $NO₂⁻$ along with $NO₃⁻$. (b) A comparison of the mass spectrum of the control NO_2^- (20 μ M) before bubbling and after bubbling for 2 h and 4 h.

114,0912

 $2.0x10^4$

1.5x10

 $1.0x10^4$

 $5.0x10^3$

 0.0

 $11C$

Absolute intensity

Additionally, we employed the spin trap, 5,5-dimethyl-1-pyrroline-N-oxide (DMPO), to capture the reactive oxygen species (ROS) and reactive nitrogen species (RNS) as shown previously by Song et al.³⁵ During the microbubbling experiment, DMPO (2 mM) was added to the solution and bubbling was continued for a sufficiently longer time to capture any radical species generated in the solution. Finally, the solution was analyzed using mass spectrometry (Orbitrap MS). DMPO captures ROS such as OH' and OOH' which were characterized by the formation of DMPO-OH $(m/z$ 130.0875) and DMPO-OOH $(m/z$ 146.0872). The OH captured by DMPO was generated from Fenton's reaction in the presence of Fe^{2+} in the solution whereas the OOH $^{\circ}$ detected possibly results from the catalytic regeneration of Fe^{2+} from Fe^{3+} at the gas–water interface as follows:

$$
Fe^{3+} + H_2O_2 \to Fe^{2+} + HO_2^+ + H^+ \tag{8}
$$

In addition, we observed features at m/z 114.0912 and 136.0715 corresponding to DMPO-H and DMPO-Na, respectively. Apart from ROS, we also observed RNS such as NO' and $NO₂$ which were detected as DMPO-NO $(m/z 143.0847)$ and DMPO-NO₂ $(m/z 159.0835)$ in the mass spectrum (Fig. 3). As both the NO' and NO₂ are radical species, they form a covalent bond at the C(sp^2)-2 position of the DMPO resulting in the formation of DMPO-NO and DMPO-NO₂. From both the fluorescence measurement and mass spectral interpretation we concluded that NO' and $NO₂$ intermediates are produced during the bubbling.

When the mass spectrum of the N2-microbubbled sample was recorded aer 4 h, it showed a feature at m/z 46 due to NO2 − (negative ion mode) along with the product peak of NO3 − at m/z 63. This suggests that NO2 − is formed during the reaction and possibly works as an intermediate to obtain NO3 [−] as the nal product (Fig. 4a). To conrm NO3 [−] conversion occurs via NO2 −, we externally prepare a NO2 − solution of known concentration (20 mM) and the solution was microbubbled. When the mass spectra were recorded (LTQ MS), we observed an increase in the NO3 − intensity with increasing bubbling time (Fig. 4b). This suggests that the NO2 [−] generated in the microbubble is involved in NO3 − formation which possibly occurs via oxidation in the presence of OHc. Because N2 as a feed gas can cause NO3 − formation, we employed argon gas for this particular experiment to create microbubbles. Also, NO2 − in aqueous solution when kept for long and exposed to light undergoes slow oxidation to NO3 −. To check NO2 − oxidation in aqueous solution we kept the solution without bubbling for 2 h and 4 h and measured the mass spectrum. We did not observe any oxidation of the NO2 − solution when kept without bubbling (Fig. S4†). This suggests that the NO2 [−] oxidation to NO3 − takes place only in microbubbles. As eqn (7) shows that NO2 − reacts with OHc to form NO2, so we believe the conversion of NO2 − to NO3 [−] occurs via NO2 generation in the microbubbles as follows: Open Access Article. Published on 11 November 2024. Downloaded on 4/29/2025 1:25:42 PM. This article is licensed under a [Creative Commons Attribution-NonCommercial 3.0 Unported Licence.](http://creativecommons.org/licenses/by-nc/3.0/) **[View Article Online](https://doi.org/10.1039/d4sc06989g)**

$$
HNO2 + OH' \rightarrow NO2' + H2O
$$
 (9)

$$
NO2 + OH' \rightarrow HNO3
$$
 (10)

$$
N_2^{\star} \rightarrow H_2N_2O_2 \rightarrow 2NO + 2H_2O \rightarrow 2HNO_2 + 2H_2O \rightarrow 2NO_2 + 4H_2O \rightarrow 2HNO_3 + 4H_2O \tag{11}
$$

 $\overline{2}$ $\cdot N_{2}H$ $-N₂OH$ Ω \overline{M} -2 $H₂O$ Free Energy (eV) -4 •OH -6 •OH ·HON₂O HON₂OH -8 \cdot OH $H₂$ C **NO** -10 $H₂O$ \cdot OH -12 \cdot OH **HONO** NO: -14 HNO₃ -16

Fig. 5 DFT calculation showing the energy profile of the intermediates involved in the N₂ oxidation to NO_3^- in the microbubbles. The experimentally identified intermediates are shown in green color dashed rectangles.

Chemical Science **Edge Article**

From all the above experiments we identified that $H_2N_2O_2$, NO', NO₂, and NO₂[–] are the intermediates formed when N₂ is converted to $NO₃⁻$ in the microbubbles. Based on our experimental evidence and combining eqn (1) – (9) we suggest the oxidation of N_2 to NO_3^- by OH $\dot{\ }$ activation takes place through a series of sequential transformations as follows:

To validate the proposed reaction pathway for the conversion of N₂ to NO₃ $^-$ *via* the gas–water interface, we conducted further investigations into the reaction mechanism using DFT calculations. The designed interfacial reaction system typically includes active radicals, such as hydrogen and hydroxyl radicals. As we previously reported, hydrogen radicals at the gas– water interface can activate nitrogen molecules, allowing them to react with other radicals.³⁶ In subsequent steps, the nitrogen– nitrogen triple bond $(N=N)$ breaks progressively, eventually leading to the formation of nitrate through oxidation (Fig. 5). Compared to the high activation energy of 0.95 eV for O_2 oxidation, the further oxidation of NO' by OH' to form $HNO₃$ proceeds without an energy barrier. Chemical Science

From all the show experimentation direction and the N¹²₂² **E. S.** 20001 **E.** (2000) **Commonstrate Commonstrate Commonstrate Commonstrate Commonstrate Commonstrate Science Article is licensed unde**

We do not claim this is the only pathway through which $NO₃⁻$ is formed, and it is possible that the reaction might involve some unknown intermediates and reaction pathways that we failed to identify. But the experimental and theoretical evidence we have found points to the reaction pathways we have presented as being operative.

Conclusions

In summary, we have experimentally detected different intermediates namely $H_2N_2O_2$, NO', NO', and NO₂⁻ produced in microbubbles during the conversion of N_2 to NO_3^- . ¹H-NMR spectroscopy was utilized to identify $H_2N_2O_2$. DAF-2 was employed to confirm the existence of NO^{\cdot} and NO₂ in the microbubbled solution spectrophotometrically. In addition, DMPO-based spin trap experiments captured the RNS (NO' and NO $_2^{\cdot}$) in the mass spectrum. The presence of NO $_2^{-}$ along with $NO₃⁻$ was observed in the mass spectrum. DFT calculations support our experimental evidence by providing insights into the energy profile of the reaction intermediates.

Data availability

The data supporting this article have been included as part of the ESI.†

Author contributions

S. B. performed most of the experiments, analysis, and drafting of the first version of the manuscript. Y. X. helped in the DFT calculation. R. N. Z., the principal investigator, proposed the project, supervised all co-authors, and completely revised the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

S. B. would like to thank the Nehru-Fulbright program for the fellowship. We thank the Air Force Office of Scientific Research through the Multidisciplinary University Research Initiative (MURI) program (AFOSR FA9550-21-1-0170) for supporting this project.

Notes and references

- 1 N. Cherkasov, A. O. Ibhadon and P. Fitzpatrick, Chem. Eng. Process. Process Intensif., 2015, 90, 24–33.
- 2 2015–2032 Nitric Acid Market Analysis: Industry Market Size, Plant Capacity, Production, Operating Efficiency, Demand & Supply, Type, End-User Industries, Sales Channel, Regional Demand, Foreign Trade, Company Share, Manufacturing Process, Policy and Regulatory Landscape, 2024, [https://](https://www.chemanalyst.com/industry-report/nitric-acid-market-615) [www.chemanalyst.com/industry-report/nitric-acid-market-](https://www.chemanalyst.com/industry-report/nitric-acid-market-615)[615](https://www.chemanalyst.com/industry-report/nitric-acid-market-615).
- 3 Y. Tang, J. Dai, P. Zhang, G. Niu, Y. Duan and Y.-H. Tian, ACS Sustainable Chem. Eng., 2024, 12, 11319–11326.
- 4 US EPA, AP 42 Chapter 8.8 Nitric acid Production, [https://](https://gaftp.epa.gov/ap42/ch08/s08/final/c08s08_feb1998.pdf) gaft[p.epa.gov/ap42/ch08/s08/](https://gaftp.epa.gov/ap42/ch08/s08/final/c08s08_feb1998.pdf)final/c08s08_feb1998.pdf.
- 5 X. Song, Y. Meng and R. N. Zare, J. Am. Chem. Soc., 2022, 144, 16744–16748.
- 6 S. Jin, H. Chen, X. Yuan, D. Xing, R. Wang, L. Zhao, D. Zhang, C. Gong, C. Zhu, X. Gao, Y. Chen and X. Zhang, JACS Au, 2023, 3, 1563–1571.
- 7 Y. B. Vogel, C. W. Evans, M. Belotti, L. Xu, I. C. Russell, L.-J. Yu, A. K. K. Fung, N. S. Hill, N. Darwish, V. R. Gonçales, M. L. Coote, K. Swaminathan Iyer and S. Ciampi, Nat. Commun., 2020, 11, 6323.
- 8 J. K. Lee, D. Samanta, H. G. Nam and R. N. Zare, Nat. Commun., 2018, 9, 1562, DOI: [10.1038/s41467-018-04023-z](https://doi.org/10.1038/s41467-018-04023-z).
- 9 A. Nandy, A. Kumar, S. Mondal, D. Koner and S. Banerjee, J. Am. Chem. Soc., 2023, 145, 15674–15679.
- 10 D. Satyabola, T. Ahuja, S. Bose, B. Mondal, P. Srikrishnarka, M. P. Kannan, B. K. Spoorthi and T. Pradeep, J. Phys. Chem. C, 2021, 125, 10998–11006.
- 11 A. Ray Chowdhuri, B. K. Spoorthi, B. Mondal, P. Bose, S. Bose and T. Pradeep, Chem. Sci., 2021, 12, 6370–6377.
- 12 S. Bose, A. Chatterjee, S. K. Jenifer, B. Mondal, P. Srikrishnarka, D. Ghosh, A. R. Chowdhuri, M. P. Kannan, S. V Elchuri and T. Pradeep, ACS Sustainable Chem. Eng., 2021, 9, 4554–4563.
- 13 J. Ghosh and R. G. Cooks, TrAC, Trends Anal. Chem., 2023, 161, 117010.
- 14 S. Banerjee, E. Gnanamani, X. Yan and R. N. Zare, Analyst, 2017, 142, 1399–1402.
- 15 X. Yan, R. M. Bain and R. G. Cooks, Angew. Chem., Int. Ed., 2016, 55, 12960–12972.
- 16 D. Zhang, J. Wang, H. Chen, C. Gong, D. Xing, Z. Liu, I. Gladich, J. S. Francisco and X. Zhang, J. Am. Chem. Soc., 2023, 145, 6462–6470.
- 17 B. K. Spoorthi, K. Debnath, P. Basuri, A. Nagar, U. V Waghmare and T. Pradeep, Science, 2024, 384, 1012–1017.
- 18 D. Zhang, X. Yuan, C. Gong and X. Zhang, J. Am. Chem. Soc., 2022, 144, 16184–16190.
- 19 Z. Song, C. Liang, K. Gong, S. Zhao, X. Yuan, X. Zhang and J. Xie, J. Am. Chem. Soc., 2023, 145, 26003–26008.
- 20 H. Xiong, J. K. Lee, R. N. Zare and W. Min, J. Phys. Chem. Lett., 2020, 11, 7423–7428.
- 21 C. Zhu, L. N. Pham, X. Yuan, H. Ouyang, M. L. Coote and X. Zhang, J. Am. Chem. Soc., 2023, 145, 21207–21212.
- 22 L. Qiu, Z. Wei, H. Nie and R. G. Cooks, Chempluschem, 2021, 86, 1362–1365.
- 23 Z. Wei, Y. Li, R. G. Cooks and X. Yan, Annu. Rev. Phys. Chem., 2020, 71, 31–51.
- 24 K. Inoue, M. Ahmed, S. Nihonyanagi and T. Tahara, Nat. Commun., 2020, 11, 5344.
- 25 G. R. Medders and F. Paesani, J. Am. Chem. Soc., 2016, 138, 3912–3919.
- 26 Z. Zhou, X. Yan, Y.-H. Lai and R. N. Zare, J. Phys. Chem. Lett., 2018, 9, 2928–2932.
- 27 G. Rovelli, M. I. Jacobs, M. D. Willis, R. J. Rapf, A. M. Prophet and K. R. Wilson, Chem. Sci., 2020, 11, 13026–13043.
- 28 H. Wei, E. P. Vejerano, W. Leng, Q. Huang, M. R. Willner, L. C. Marr and P. J. Vikesland, Proc. Natl. Acad. Sci. U. S. A., 2018, 115, 7272–7277. Edge Article

10 D. Zhang, X. Yuan, C. Grog and X. Zhang, J. Am. Chem, Soc., 28 H. Wei, F. P. Vejerino, W. Lewin, Grow Michael Science

10 D. Zhang, A. Wei, H. W. Stein, N. Yuan, X. Non, X. Zhang, N. Wei, This article is l
	- 29 S. Bose, M. Mofidfar and R. N. Zare, J. Am. Chem. Soc., 2024, 146, 27964–27971, DOI: [10.1021/jacs.4c11899](https://doi.org/10.1021/jacs.4c11899).
	- 30 M. Umar, H. A. Aziz and M. S. Yusoff, Waste Manag., 2010, 30, 2113–2121.
	- 31 S. O. Ganiyu, M. Zhou and C. A. Martínez-Huitle, Appl. Catal., B, 2018, 235, 103–129.
	- 32 S. Chen, S. Liang, R. Huang, M. Zhang, Y. Song, Y. Zhang, S. Tao, L. Yu and D. Deng, Nat. Synth., 2024, 3, 76–84.
	- 33 H. Kojima, N. Nakatsubo, K. Kikuchi, S. Kawahara, Y. Kirino, H. Nagoshi, Y. Hirata and T. Nagano, Anal. Chem., 1998, 70, 2446–2453.
	- 34 L. J. Ignarro, J. M. Fukuto, J. M. Griscavage, N. E. Rogers and R. E. Byrns, Proc. Natl. Acad. Sci. U. S. A., 1993, 90, 8103–8107.
	- 35 X. Song, C. Basheer, Y. Xia and R. N. Zare, Environ. Sci. Technol., 2024, 58, 16196–16203.
	- 36 J. Li, Y. Xia, X. Song, B. Chen and R. N. Zare, Proc. Natl. Acad. Sci. U. S. A., 2024, 121, e2318408121.