

# Sensors & Diagnostics

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Z. Zhang, Md. A. Ahamed and D. Yang, *Sens. Diagn.*, 2024, DOI: 10.1039/D4SD00373J.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

# Biological Properties and DNA Nanomaterials Biosensors of Exosomal miRNAs in Disease Diagnosis

Zhikun Zhang<sup>1</sup>, Md. Ahasan Ahamed<sup>4</sup> and Dayong Yang<sup>2,3,\*</sup>

<sup>1</sup> School of Chemical and Pharmaceutical Engineering, Hebei University of Science and Technology, Shijiazhuang, 050018, PR China

<sup>2</sup> Department of Chemistry, State Key Laboratory of Molecular Engineering of Polymers, Shanghai Key Laboratory of Molecular Catalysis and Innovative Materials, College of Chemistry and Materials, Fudan University, Shanghai, 200438, PR China

<sup>3</sup> Bioinformatics Center of AMMS, 100850, Beijing, PR China

<sup>4</sup> Department of Electrical Engineering, Pennsylvania State University, University Park, 16802, USA

\* Corresponding Author, Email: dayongyang@fudan.edu.cn.



## Abstract

MicroRNAs (miRNAs) regulate gene expression and are important biomarkers in molecular diagnostics, prognosis, and personalized medicine. The miRNAs that are found within exosomes, also known as exo-miRs, have been shown to demonstrate increased levels of both abundance and stability. Thus, exo-miRs show potential as a reliable biomarker for further investigation. Due to the programmable nanostructures, biocompatibility, and excellent molecular recognition ability, biosensing platforms based on DNA nanomaterial are considered promising for detecting exo-miRs in clinical analysis, including cancer, neurodegenerative disorders, and infectious diseases. Although considerable advancements have been achieved in exo-miR-based testing, there are ongoing challenges in accurately detecting and analyzing multiple targets concurrently at low concentrations in complex biological samples. The primary focus of our research is to thoroughly analyze the biogenesis of exo-miRs, carefully assess their levels of expression in various clinical diseases, and comprehensively investigate their correlations with a wide range of diseases, including cancer, infection, and neurodegenerative disorders. We also examined recent progress in DNA nanomaterials-based detection methods for exo-miRs. This study explores the challenges and intricacies faced during the creation and execution of exo-miR tests within a clinical setting to diagnose diseases. The successful development and implementation of DNA nanomaterials for exo-miRs detection can significantly revolutionize the early detection, monitoring, and management of various medical conditions, leading to enhanced healthcare outcomes.

Keywords: disease diagnostics, exosome, miRNAs, DNA nanomaterials, sensor



## 1 Introduction


MicroRNAs (miRNAs or miRs) are a group of endogenous, small molecules (typically 22 nucleotides long), non-coding, and structurally functional RNAs.<sup>1,2</sup> miRs have become increasingly recognized as essential biomarkers in the fields of molecular diagnostics, disease progression monitoring, and individualized medical treatment because of their notable impact on the regulation of gene expression.<sup>3</sup> Dysregulated miRNAs are strongly associated with the development of various,<sup>4,5</sup> such as tumor,<sup>5,6</sup> cardiovascular diseases,<sup>7,8</sup> central nervous system diseases,<sup>9</sup> kidney injury,<sup>10</sup> and other conditions. Different miRNAs may be present in bodily fluids, including freely circulating miRNAs, miRNAs bound to ribonucleoproteins, and miRNAs enclosed within various exosomes. Exosomes are extracellular vesicles released by all types of cells, serving a crucial function in transporting biomolecules, including proteins, metabolites, and assorted nucleic acids in biological processes.<sup>3,11,12</sup> miRs contained within exosomes (so-called exo-miRs) exhibit higher levels of abundance and stability under the protection of lipid bilayer-membrane when contrasted with freely circulating miRs in the bloodstream.<sup>13</sup> They exhibit greater value in liquid biopsy due to enhanced preservation when exposed to enzymes commonly found in bodily fluids. Therefore, exo-miRs have been the subject of thorough investigation in the past few years as they emerge as a promising biomarker for liquid biopsy, attracting considerable attention in disease diagnostics.

The conventional methods for analyzing exo-miRs are reverse transcription-polymerase chain reaction (RT-PCR),<sup>14,15</sup> digital PCR,<sup>16</sup> and next-generation sequencing technology (NGS).<sup>17</sup> Despite their excellent analytical performance, these methods still exhibit limitations for clinical applications. Reverse transcription-polymerase chain reaction (RT-PCR) as the gold standard technique can be utilized to detect exo-miRs; however, it exhibits a comparatively restricted need for costly equipment, involving multiple manual steps that prolong the assay time.<sup>18</sup> Digital PCR can detect very low levels of miRNAs. However, it requires a significant financial investment and involves a complex operational process.<sup>16</sup> Next-generation sequencing typically involves extensive pretreatment and multiple procedural steps.<sup>19</sup> Therefore, the direct and accurate detection of exo-miRs derived from tumors presents a significant challenge, requiring the precise identification and the quantitative assessment of exo-miRs amidst complex background interference, diverse subtypes of extracellular vesicles (EVs), and varying expression levels of



various exo-miRs. Therefore, developing a sensitive, accurate, and simplified bioassay and platform to quantify diseases-derived exo-miRs in complicated biofluid samples is highly desirable to promote the development of clinically viable miRNA biomarkers of disease diagnostics.

Currently, there has been a significant focus on the advancement of multiplexed, sensitive, and straightforward techniques for exo-miRs detection, including surface-enhanced Raman scattering (SERS),<sup>20</sup> surface plasmon resonance (SPR),<sup>21</sup> microfluidics,<sup>22</sup> fluorescence,<sup>23,24</sup> and electrochemistry.<sup>25</sup> DNA nanotechnology generates a wide array of nanostructures due to its precise design, unique ability for programming, and natural compatibility with biological systems.<sup>26–28</sup> DNA nanomaterial-based biosensors have garnered significant interest as potential diagnostic tools within this field. The attributes of DNA nanostructures, such as their diminutive scale, precise molecular recognition abilities, and inherent self-assembly stability, render them a feasible biosensing platform fabrication for facilitating the integration of nanotechnology and diagnostics. This allows DNA nanoassemblies to fulfill a crucial function in advancing the development of highly specific and sensitive biosensors with improved performance for exo-miRs detection. This review offers an in-depth analysis of the biogenesis of exo-miRs, emphasizing the linkages between miRs and various diseases. We examined the latest developments in DNA nanomaterials, focusing on their design and mechanisms in exo-miRs detection (**Figure 1**). This review article also explores the challenges faced and offers suggestions for future progress. The goal is to bridge the gap between research on exo-miRs and their application in disease diagnostics to facilitate the development of novel, clinically useful diagnostic tools.



**Figure 1.** Various DNA nanomaterials for detecting exo-miRs associated with diseases by fluorescent, electrochemical, SERS, and Uv-vis sensing platforms.

## 2 Biosynthesis of exo-miRs

Exo-miRs biosynthesis starts with the traditional miRNA biogenesis pathway inside the cell.<sup>29</sup> Their size ranges from 40 to 120 nm, which provides valuable information for cancer diagnosis, brain diseases, infectious diseases, and therapeutic treatment. Their origins can be traced back to the intraluminal budding of multivesicular bodies (MVBs), a fusion of MVBs with cell membranes, and endolysosomal pathways. Generally, it contains a variety of membrane, cytosolic, metabolites, DNA, proteins, circular RNA (circRNA), messenger RNAs (mRNAs), and microRNAs (miRNAs), as shown in **Figure 2A**. Among these, miRNAs are non-coding RNAs that regulate genetic expression, which is transcribed from DNA and converted to primary miRNAs, processed into precursor forms, and mature miRNAs.<sup>30</sup> Generally, miRNAs bind with the 3' untranslated region (UTR) region of a gene target, which causes miRNA degradation or




translation inhibition. They can also interact with 5' UTR and gene promoters.<sup>29</sup> Normally, miRNA activity depends on the subcellular target site, quantity, and interaction strength.<sup>31</sup> One way of intraluminal vesicles (ILVs) by miRNAs is secretion in extracellular fluids via vesicles like exosomes or bound to proteins like Argonaute 2 (AGO2).<sup>32–34</sup> In other studies, exo-miRs can be found in biological fluids such as saliva and serum, which is now widely used for nucleic acid extraction.<sup>35–38</sup> Primarily, miRNAs are transcribed in the nucleus as primary miRNAs (pri-miRNAs) by RNA polymerase II, followed by cleavage activity by Drosha-DGCR8 and create precursor miRNAs (pre-miRNAs), which Exportin-5 exports to the cytoplasm.<sup>31,35</sup> Then, the Dicer enzyme comes into play to process pre-miRNAs to mature miRNA duplexes in the cytoplasm.<sup>39</sup> One strand of the duplex can regulate a gene's expression by inhibiting or degrading the translation of target mRNAs after being loaded into the RNA-induced silencing complex (RISC).<sup>40</sup> On the other hand, mature miRNAs are selectively encapsulated into exosomes, which facilitate intercellular communication.<sup>41</sup> Following general miRNA synthesis, the packed exo-miRs is a selective process.<sup>35</sup> miRNA sorting and packaging is very complex process where Heterogeneous nuclear ribonucleoproteins (hnRNPs), ESCRT (Endosomal Sorting Complex Required for Transport) and AGO2 proteins are responsible for sorting, identifying and managing specific miRNAs for exosome packaging.<sup>42</sup> Finally, these exosomes and Microvesicles are released from the MVBs into extracellular space, and Lysosomes are formed inside of the cell through the process of Golgi apparatus, as shown in **Figure 2B**.<sup>43</sup>

Exosomes travel toward the recipient cells by bodily fluids such as blood and serum to interact with the cell and outside exo-miRs. Recipient cells can take exosomes and reflect on their unique miRNAs, which control gene expression and facilitate intercellular communication.<sup>44</sup> Then, it fuses inside the cell membrane or receptor-ligand interaction called endocytosis.<sup>45</sup> After endocytosis, the exosomal cargo miRNA is trapped. Once the exo-miRs is inside an early endosome containing ILVs, the ILVs are small vesicles inside MVBs and contain miRNA, DNA, RNA, proteins, and lipids. They are created through the inward budding of the MVBs' membrane, and the membrane fusion process facilitates the release of miRNA to the cytosol.<sup>35,46</sup> Moving forward, RISC interacts with miRNA, and ESCRT facilitates the sorting process.<sup>47</sup> On the other hand, the ILVs are released into the extracellular environment as exosomes when the MVBs and plasma membrane fuse. To create diagnostic instruments and treatment plans, it is essential to find



out the miRNA type that is primarily responsible for early disease detection and the amount of miRNA present at that stage, where exo-miRs act as significant biomarkers.<sup>39</sup>



**Figure 2: Exosome structure and biosynthesis.** A) Exosome size and functional components such as miRNA, mRNA, circRNA, proteins, and lipids.<sup>48</sup> B) miRNA is transcribed in the nucleus, processed by Drosha and Dicer, and loaded into RISC. Exosomes are released from MVBs and carry miRNA to recipient cells.<sup>40</sup>

### 3 Correlation between exo-miRs and various diseases

Exo-miRs have emerged as potential biomarkers for disease detection and monitoring and can be found in various non-invasive biological fluids such as blood, saliva, and urine. Previously, we provided deep insight into miRNA biogenesis, where they serve as a primary cargo content of exosomes in early diagnosis of cancer, neurological diseases such as Parkinson's disease (PD) and Alzheimer's disease (AD), cardiovascular diseases such as heart diseases, pulmonary arterial hypertension, and aortic aneurysms, infectious diseases, and respiratory conditions. Modulating immune responses and inflammation of infectious and respiratory diseases, potential drug resistance for cancer and neurological disease, and miRNA detection are valuable tools for early





diagnosis and setting up specific treatment plans. Their role in disease pathogenesis makes them attractive targets for developing diagnostic and therapeutic interventions.

### 3.1 Cancer

Exo-miRs are crucial in regulating cancer drug resistance through gene expression modulation, drug efflux, metabolic reprogramming, DNA damage repair, apoptosis, and epithelial-to-mesenchymal transition (EMT).<sup>40</sup> MicroRNAs are integral to a intricate network that includes their target genes (RhoA, FIH and ACOX1, etc.) and signalling cascades (the EGF-AKT signalling axis, the ERK-MMP9 cascade and the Hippo signalling pathway, etc.), which plays a crucial role in the progression of cancer.<sup>49</sup> These miRNAs are transferred between cancer cells and the tumor microenvironment, influencing tumor growth, metastasis, and therapeutic resistance. For example, Guo et al. explained in their study that exo-miRs like miR-1246, secreted by ovarian cancer (OC) cells, regulate drug efflux transporters like ATP-binding cassette subfamily B member 1 (ABCB1), contributing to chemotherapy resistance by decreasing drug efficacy.<sup>40,50</sup> Similarly, miR-365 induces resistance to gemcitabine by increasing cytidine deaminase activity in pancreatic cancer (PC) cells.<sup>51</sup> Exo-miRs like miR-21 and miR-155 also play the most significant roles in cancer progression by modulating apoptosis, promoting tumor survival, and drug resistance when carried by exosomes.<sup>52</sup> miR-21, often overexpressed in breast cancers (BC), colorectal cancers, and OC, is transferred by cancer-associated fibroblasts (CAFs) to nearby cancer cells. It reduces the expression of pro-apoptotic factors like Phosphatase and Tensin Homolog (PTEN) and allows cells to survive chemotherapy.<sup>40</sup> The transfer of miR-21 via exosomes has explicitly been noted to reduce the expression of pro-apoptotic factors such as PTEN, leading to enhanced cell survival and drug resistance.<sup>53</sup> On the other hand, miR-155 reprograms tumor-associated macrophages (TAMs) into a pro-tumorigenic phenotype, aiding in tumor proliferation and immune evasion.<sup>54</sup> These macrophages, in turn, secrete factors that promote tumor cell proliferation and immune evasion. In addition, exosomal miR-155 contributes to chemotherapy resistance by downregulating genes involved in DNA repair and apoptosis, allowing cancer cells to thrive even in chemotherapeutic agents, as shown in **Figure 3A**. This dynamic interaction between exo-miRs and immune cells in the tumor microenvironment highlights the sophisticated mechanisms by which tumors evade therapeutic interventions.<sup>40</sup>



Exo-miRs also significantly impact cancer metastasis. In BC, for instance, Zhang et al. showed that miR-9 induces fibroblast cells to acquire cancer-associated fibroblast-like properties, supporting tumor growth and metastasis. At the same time, miR-200b promotes proliferation and invasion in colorectal cancer by downregulating tumor suppressors.<sup>55</sup> Modani et al. note that miR-10b achieves this by targeting genes that regulate cell adhesion and migration, thus facilitating the spread of cancer to distant organs.<sup>39</sup> miR-10b, secreted by highly metastatic BC cells, enhances recipient cells' migratory and invasive capabilities, facilitating cancer spread to distant organs.<sup>40</sup> These exo-miRs drive metastasis and create a supportive tumor microenvironment by influencing gene expression and facilitating communication between cancer and stromal cells. In addition to miRNAs, these exosomes transfer proteins, lipids, and other molecules that modify the behavior of both cancerous and non-cancerous cells in the microenvironment. Additionally, exo-miRs are crucial for maintaining cancer stem cell (CSC) populations, which are inherently resistant to conventional therapies.<sup>56</sup> For example, Santos et al. miR-155, secreted by CSCs, triggers the EMT process, making recipient cells more resistant to chemotherapy.<sup>57</sup> Meanwhile, miR-210, transferred by exosomes in gemcitabine-resistant PC stem cells, inhibits apoptosis and promotes survival.<sup>39</sup>

Furthermore, circulating exo-miRs offer a minimally invasive method for early cancer detection and monitoring treatment response. For example, miR-1290 and miR-375 elevated in prostate cancer patients potentially serve as reliable biomarkers for early detection.<sup>58</sup> Another example is that miR-122 reduces glucose uptake in liver cells, providing metabolic advantages to cancer cells, while miR-126-5p alters lipid metabolism, essential for cancer cell proliferation. This metabolic reprogramming can exacerbate conditions such as cachexia, diabetes, and obesity, highlighting the intricate relationship between cancer, exosomes, and systemic metabolic dysfunction.<sup>59</sup> Exo-miRs, such as miR-19, linked to brain cancer metastasis, and miR-25-3p promoting liver metastasis in colorectal cancer via the CXCL12/CXCR4 pathway, are key diagnostic markers.<sup>60</sup> Since exo-miRs are available in saliva, Zhang et al. highlight the critical role of salivary miRNAs in the early diagnosis and progression of various cancers, including oral squamous cell carcinoma (OSCC), esophageal cancer (EC), gastric cancer (GC), pancreatic cancer (PC), and breast cancer (BC).<sup>61</sup> The miRNAs, such as miR-200 and miR-21, are significantly altered in OSCC, making them promising biomarkers for early detection.<sup>62,63</sup> Similarly, elevated



levels of miR-10, miR-144, and miR-451 are associated with EC,<sup>64–66</sup> while miR-140-5p and miR-301a are linked to GC. For PC, miR-1246, miR-4644, miR-103, and miR-107 in salivary exosomes show promise for differentiating patients from normal controls.<sup>61,67</sup> In BC, miR-21 is a significant marker, showing reduced levels post-treatment.<sup>68</sup> The potential of salivary miRNAs as non-invasive biomarkers for the early detection and monitoring of various cancers. Understanding the roles of exo-miRs in cancer drug resistance, metastasis, and tumor-stroma interactions can help develop new therapeutic strategies to improve cancer diagnosis and treatment outcomes.

### 3.2 Concussion and Parkinson's disease

Exo-miRs are also implicated in the early stages of Concussion, neurodegenerative diseases, Parkinson's disease (PD), and Alzheimer's disease (AD). Specific miRNAs, like miR-124 and miR-132, are found to be dysregulated in exosomes derived from neural cells, contributing to disease pathogenesis.<sup>69</sup> Concussions, especially prevalent in children and adolescents, represent about two-thirds of mild Traumatic Brain Injuries (TBI) cases and involve the regulation of gene expression through miRNA. While most children recover within two weeks, around one-third may experience prolonged concussion symptoms (PCS). miR-27 has been identified as a potential biomarker for concussions due to its accuracy in distinguishing concussed individuals. Additionally, miRNAs such as miR-320, miR-133, and let-7a have been linked to prolonged symptoms, while specific miRNAs (miR-320, miR-629, let-7b-5p) correlate with symptoms like memory issues, headaches, and fatigue.<sup>61</sup> In the central nervous system (CNS), exosomes facilitate communication between neurons and glial cells, contributing to the progression of various diseases. They carry miRNAs, such as miR-150, that regulate neuroinflammatory responses and contribute to disease progression.<sup>70</sup> Exosomal release from various CNS cells, including neurons, astrocytes, and microglial cells, influences synaptic communication and neuro-glial interactions.<sup>71</sup> miR-150, miR-223, and miR-23a have been identified in the serum and plasma of Multiple Sclerosis (MS) patients. These biomarkers have shown high diagnostic accuracy, with miR-150 demonstrating 85% sensitivity and 90% specificity, making them valuable tools for MS diagnosis.<sup>72</sup> Other examples are miR-126, miR-223, miR-9, and miR-124, which have shown potential in diagnosing ischemic stroke with higher sensitivity and specificity compared to free circulating miRNAs. For instance, miR-223 demonstrated an AUC of 0.859 with 84.0% sensitivity



and 78.8% specificity for stroke diagnosis.<sup>73</sup> miR-21, miR-222, and miR-124-3p are significantly elevated in the serum of GBM patients and have shown high diagnostic accuracy in distinguishing GBM from healthy controls. ROC curve analysis for miR-21 revealed an AUC of 0.927, demonstrating its strong potential as a biomarker for GBM.<sup>74</sup> Additionally, miR-301a levels were found to correlate with tumor recurrence and surgical resection outcomes, highlighting their role in monitoring disease progression.<sup>72</sup>

Parkinson's disease (PD) is a progressive neurodegenerative disorder that primarily affects individuals over the age of 65 and is the second most common neurodegenerative disorder after Alzheimer's disease (AD).<sup>61</sup> Its diagnosis is typically based on clinical symptoms such as tremors, rigidity, and non-motor issues like depression and dementia. Vital pathological features of PD include the loss of dopaminergic neurons and the accumulation of  $\alpha$ -synuclein.<sup>71</sup> Recent studies have highlighted alterations in miRNA levels in saliva, cerebrospinal fluid (CSF), and serum of PD patients, which affect genes linked to the disease.<sup>75</sup> Research has indicated that miRNAs can regulate genes linked to Parkinson's disease, such as the SNCA gene, PRKN gene, and PARK7 gene, etc.<sup>76,77</sup> Notably, miR-153, miR-409-3p, miR-375, miR-1468-5p, miR-19b, miR-195, and miR-24, found in CSF and plasma have emerged as potential diagnostic biomarkers.<sup>78</sup> Among these, miR-153 and miR-223 have been associated with  $\alpha$ -synuclein accumulation, showing moderate diagnostic accuracy.<sup>79</sup> Other miRNAs, including miR-29, miR-28, miR-19b-3p, and elevated levels of miR-409-3p and let-7g-3p, exhibit high sensitivity and specificity as diagnostic biomarkers.<sup>72</sup> Additionally, miRNAs such as miR-7, miR-34b/c, miR-124, miR-221/222, miR-137, and miR-433 play roles in PD pathology, contributing to neuroinflammation, mitochondrial dysfunction, and dopamine metabolism, as depicted in **Figure 3B**. Exo-miRs like miR-193b, miR-135a, and miR-384 have been identified as potential biomarkers for early AD detection. miR-384 has been particularly effective in distinguishing AD from other neurological diseases like PD with dementia and vascular dementia.<sup>72</sup> Furthermore, studies report that while miR-1 and miR-19b-3p are reduced, miR-153 and miR-409-3p are elevated in PD patients' CSF exosomes. Exo-miRs may also help detect mild cognitive impairment (MCI), an early stage of AD, underscoring their potential as noninvasive tools for early diagnosis and treatment strategies in both PD and AD.<sup>78,80-</sup>

85



### 3.3 Infectious diseases

Exo-miRs serve as a potential biomarker for diagnosing various infectious diseases, significantly influencing cell signaling and directing the host immune response. changes in their expression profiles often appear early, even before the pathogen is detectable. these small, non-coding RNAs encapsulated within exosomes are protected from degradation and can be transported to recipient cells, which modulate gene expression and immune responses. Below, we discuss the role of exo-miRs in various infectious diseases, focusing on their functions and mechanisms of action in SARS-CoV-2, HIV, HCV, Enterovirus, Epstein-Barr virus (EBV), HBV, and Ebola virus.<sup>86</sup>

Several exo-miRs play a role in inhibiting viral replication and modulating immune responses in SARS-CoV-2 infection. For example, Drury et al. noted their significance in diseases caused by various pathogens, including *Helicobacter pylori* and SARS-CoV-2.<sup>87</sup> Zhou and Xu found that downregulation of miR-204 in pulp tissue, serum, and saliva indicated *H. pylori* infection, highlighting the role of miRNA in regulating inflammatory pathways and their potential for early gastric cancer detection.<sup>88</sup> Hicks et al. observed altered salivary miRNA levels, such as decreases in miR-4495, miR-296, miR-548, and miR-1273, in children with severe SARS-CoV-2 infection.<sup>89</sup> Saulle et al. identified significant downregulation of let-7a, let-7b, and let-7c, and upregulation of miR-23a, miR-23b, and miR-29c, alongside three immunomodulatory miRNA in SARS-CoV-2.<sup>90</sup> miR-223-3p, miR-24-3p, miR-145-5p, and miR-75p have been shown to inhibit SARS-CoV-2 replication by targeting viral proteins and mediating membrane fusion and viral entry into host cells. Additionally, miR-148a and miR-590 modulate the immune response during SARS-CoV-2 infection by targeting IRF9. miR-148a inhibits USP33, reducing IRF9 protein expression and impacting the immune response to viral infection. Similarly, miR-590 directly targets IRF9, regulating pro-inflammatory pathways. Both miRNAs are key in lowering viral load and controlling immune reactions.<sup>91-93</sup> **Figure 3D** depicts the mechanism of exosomal miRNA involvement in SARS-CoV-2 infection. Infected cells release exosomes containing viral RNA, miRNAs, proteins, and other molecules, facilitating immune system activation. These exosomes contribute to inflammation and can lead to severe immune responses, including sepsis and multi-organ failure, while also serving as potential biomarkers for disease progression. However, challenges remain, such as the lack of consensus on specific miRNAs to be used as biomarkers,



the influence of factors like age, gender, and lifestyle on miRNA expression, and variations in laboratory practices for detecting these miRNAs. The specificity issue can be resolved by integrating a clustered, regularly interspaced short palindromic repeats (CRISPR) assay<sup>18,94–96</sup>. Despite these challenges, exosomal miRNA offers excellent potential for improving early diagnosis and therapeutic interventions in infectious diseases.<sup>86,87</sup> Influenza infection is characterized by the modulation of miRNAs like miR-155 and miR-323, which influence the host's antiviral response and inhibit viral replication.<sup>97</sup>

For HIV and HIV-1, vmiR-TAR reduces Cdk9 and Bcl-2, delaying apoptosis and boosting viral production. vmiR88 and vmiR99 activate TLR8 in macrophages, increasing pro-inflammatory cytokines. Exosomal miR-155-5p from HIV-infected T cells activates NF- $\kappa$ B in cervical cancer cells, promoting invasion.<sup>98,99</sup> In HIV infection, miRNAs like miR-223, miR-21, and miR-29b are involved in modulating viral replication and immune responses, with specific miRNAs downregulated during infection, affecting disease progression and immune function. Hepatitis viruses, including HBV and HCV, deregulate miRNAs like miR-122 and miR-29b, linked to liver fibrosis and viral replication.<sup>97</sup> In Hepatitis B Virus (HBV) infections, exosomal miR-122 promotes viral replication, while miR-574-5p inhibits HBV polymerase, reducing replication. These miRNAs serve as potential targets for antiviral therapies and disease monitoring.<sup>100,101</sup> miR-122 promotes anti-Hepatitis C Virus (HCV) antibody production by stimulating B cell proliferation. Exosomal let-7b and miR-206 activate macrophages via TLR7, enhancing BAFF release. These miRNAs help control viral replication and support the immune response during HCV infection.<sup>102,103</sup> During Enterovirus 71 (EV71) infection, miR-146a suppresses type I interferon responses, boosting viral replication. Similarly, miR-30a from EV71-infected cells targets MyD88 in macrophages, inhibiting immune responses and enhancing replication.<sup>104,105</sup> In Epstein-Barr Virus (EBV) infection, miR-BHRF1-1 plays a role in immune evasion. At the same time, miR-BART3 is known for modulating gene expression linked to apoptosis and inflammation.<sup>106</sup> miR-BART3 targets IPO7 to increase IL-6 expression, while miR-BHRF1 down-regulates p53, enhancing cell survival and viral replication.<sup>107,108</sup> In Ebola virus (EBOV) infections, exosomal miR-21 and miR-222-3p enhance cell survival and viral replication by promoting anti-apoptotic and pro-inflammatory responses, making them potential therapeutic targets for controlling EBOV infections.<sup>109</sup> miR-VP-3p and miR-155 are linked to immune



response modulation and interferon signaling, and their presence in serum can be used for early diagnosis of infection.<sup>97</sup> Chikungunya virus infection leads to differential expression of miRNAs in mosquito saliva, such as miR-184 and miR-125, which regulate inflammation and immune responses, influencing the transmission and severity of the disease in humans.<sup>97</sup> Bayer et al. found that infection with varicella zoster virus could be attenuated in cells that had been pre-exposed to exosome-packaged miRNA clusters from chromosome 19; this mechanism offers broader antiviral resistance, especially against rubella and togavirus and HIV-1.<sup>110</sup> Kaposi sarcoma-associated herpesvirus (KSHV), human papillomavirus (HPV), and BK polyomavirus (BKV) use exo-miRs to manipulate host cells. KSHV promotes the Warburg effect via miR-K12-11,<sup>111</sup> shifting metabolism to aerobic glycolysis and enhancing angiogenesis and cell migration. HPV infection leads to the upregulation of miR-20a-5p, miR-423-3p, and let-7d-5p, which promote cancer progression by influencing cell survival and proliferation. BKV-associated bkv-miR-B1-3p and bkv-miR-B1-5p are detected in urine and blood as biomarkers for diagnosing BK virus nephropathy and monitoring viral replication in kidney transplant patients.<sup>106</sup> In tuberculosis, miRNAs such as miR-144 and miR-361-5p are differentially expressed, mediating immune response modulation and disease progression.<sup>112</sup> Filariasis involves miRNAs like miR-223 and miR-71, which regulate immune responses and are critical in the development and persistence of the infection. These miRNAs offer insights into the pathophysiological conditions associated with filariasis and could be used as diagnostic markers to evaluate disease severity and treatment efficacy.<sup>97</sup>

### 3.4 Other diseases

In cardiovascular diseases, exo-miRs like miR-1 and miR-133a are associated with early cardiac dysfunction, regulate gene expression between cells and myocardial injury, and facilitate communication between cells involved in heart function.<sup>113</sup> Exo-miRs have been linked to the progression of atherosclerosis, myocardial infarction, and heart failure, influencing inflammation, cell death, and tissue repair.<sup>114</sup> **Figure 3c** illustrates cell communication involving exo-miRs in the context of heart disease and their interaction with damaged heart muscle. The figure shows how different cell types, including cardiomyocytes, mesenchymal stem cells (MSC), adipocytes, macrophages, and cardiac stromal cells, release or respond to exo-miRs that regulate critical




processes like cell proliferation (e.g., miR-29a, miR-34a), fibrosis (e.g., miR-29a), and inflammation (e.g., miR-155). These miRNAs communicate between cells and influence damaged heart tissue's repair and response mechanisms, contributing to pathological processes like fibrosis, hypertrophy, and inflammatory responses in heart disease. Atherosclerosis is a leading cause of cardiovascular disease, characterized by the thickening and hardening of arterial walls due to the accumulation of lipids and fibrous tissues. Exo-miRs, such as miR-143 and miR-145, regulate vascular smooth muscle cells (VSMCs) and endothelial cells, mediating intercellular communication to prevent or promote atherosclerotic plaque formation.<sup>115</sup> These miRNAs help reduce atherosclerosis by modulating inflammatory responses and promoting the repair of damaged endothelial cells.<sup>116,117</sup> Acute Coronary Syndrome (ACS), which includes myocardial infarction and unstable angina, is marked by the rupture or erosion of atherosclerotic plaques. ACS patients, miRNAs such as miR-133a/b, the miR-30 family, miR-19, miR-20, miR-499, miR-1, miR-146a-5p, miR-208a, miR-126, and miR-21 are elevated, correlating with ACS and coronary artery disease (CAD).<sup>118,119</sup> Ischemia-reperfusion injury (MIRI) occurs when blood flow is restored to the previously ischemic myocardial tissue, exacerbating oxidative stress and inflammation damage. miR-126 and miR-423-3p have reduced MIRI by targeting specific signaling pathways to inhibit cardiomyocyte apoptosis and promote cardiac cell survival. These exo-miRs offer therapeutic potential for reducing myocardial damage after ischemic events.<sup>120</sup> In heart failure (HF), exo-miRs contribute to myocardial remodeling, leading to heart muscle thickening and dysfunction. Elevated levels of miRNAs such as miR-425 and miR-744 are linked to cardiac fibroblast activation and collagen synthesis, both critical in myocardial fibrosis and the progression of HF.<sup>121</sup>

Pulmonary Arterial Hypertension (PAH) is characterized by increased pulmonary vascular resistance due to remodeling of the pulmonary arteries. miR-143 and miR-145 establish the communication between pulmonary artery smooth muscle cells and endothelial cells, promoting vascular remodeling.<sup>122</sup> Aortic Aneurysm, an abnormal enlargement of the aorta, is caused by miR-106a and miR-24, which have been implicated in the pathogenesis of aortic aneurysm by promoting the apoptosis of vascular smooth muscle cells and inflammation.<sup>123</sup> Vascular calcification (VC) involves abnormal calcium deposition in blood vessels linked to atherosclerosis and increased cardiovascular risk. miR-324-3p regulates calcification pathways, offering potential





therapeutic targets to reverse calcification and stabilize plaques.<sup>124</sup> In Rheumatic Valvular Disease (RVD), miR-155-5p modulates inflammation and fibrosis in heart valves via the S1PR1/STAT3 pathway, providing potential treatment options for valve dysfunction.<sup>114</sup>



**Figure 3: Exo-miRs role for diseases.** A) **Cancer:** miR-21 and miR-210 (upregulated) promote tumor growth and angiogenesis, while miR-27a and miR-125b enhance treatment resistance. Downregulated miR-451 impairs DNA repair.<sup>52</sup> B) **Brain Disease:** miR-153 and miR-409-3p (upregulated) are associated with neuroinflammation and  $\alpha$ -synuclein accumulation, while miR-124 and miR-133b (downregulated) impair neuronal repair. In AD, miR-384 and miR-193b are upregulated, contributing to disease progression, while miR-29a/b and miR-137 (downregulated) affect synaptic function.<sup>82</sup> C) **Cardiovascular Disease:** miR-29a and miR-34a (upregulated) drive cell proliferation and fibrosis, while miR-155 promotes inflammation. miR-21 contributes to cardiac fibrosis and hypertrophy.<sup>120,125,126</sup> D) **Infectious Disease (SARS-CoV-2):** Exosomes from infected cells carry miRNAs like miR-21, triggering immune responses that lead to severe inflammation and potentially multi-organ failure.<sup>127</sup>



## 4 DNA nanomaterials-based sensing platform for exo-miRs detection

Exo-miRs present in saliva, blood, plasma, and urine have been identified at detectable levels, offering potential benefits for disease diagnosis using liquid biopsy.<sup>4,128–130</sup> The efficient and prompt monitoring of exo-miRs allows for the convenient diagnosis of diseases in patients. Nevertheless, there are various challenges in detecting exo-miRs in situ for molecule biosensors,<sup>131</sup> stemming from their complex compositions, small size, membrane obstruction, and low concentration.<sup>131,132</sup> The designed molecular biosensors must be capable of entering a confined space to identify the target and maintain a consistent and stable signal output within a nanoscale environment in exosomes. Biosensors incorporating nucleic acid nanomaterials possess distinctive qualities that allow them to effectively address the limitations associated with detecting exo-miRs, owing to their small size, unsurpassed programmability, natural biocompatibility, and superior molecular recognition capacity.<sup>133,134</sup> In current state-of-art, various well-established DNA nanomaterials are utilized as biosensors for detecting exo-miRs, such as tetrahedral DNA nanostructures (TDNs), branched DNA, spherical DNA, and hairpin DNA.<sup>135,136</sup> This selection explores the recent advancements in nucleic acid synthesis and their application in detecting exo-miRs, providing valuable insights for precise biological analysis and clinical diagnostics.

### 4.1 Tetrahedral DNA nanomaterials

Tetrahedral DNA nanostructures (TDNs) were initially developed by Turberfield in 2004,<sup>137,138</sup> exhibiting structural stability, easy internalization, biocompatibility, and versatile functionality.<sup>137,139–143</sup> Briefly, TDNs could be prepared by four specifically designed single DNA strands using the complementary base pairing rules. In exosomes, TDN-scaffold biosensors are utilized to prevent nucleases from degrading them. Maintaining a consistent spatial distance can help minimize the entanglement and local overcrowding among neighboring probes.<sup>144</sup> As a result of these benefits, biosensors based on TDN can effectively penetrate exosomes spatially oriented, which enhances the ability to detect miRs by accurately adjusting the lateral distance between the biosensors and miRs.<sup>145,146</sup> Furthermore, they have exhibited potential as biosensors for accurately identifying extracellular microRNAs.

To attain simple, quick detection abilities and highly sensitive detection, methods utilizing TDNs-scaffold for electrochemical and fluorescent detection of exo-miRs were developed.<sup>147,148</sup> Yang et al. utilized TDNs nanomaterials to form the sandwich structure with exo-miRs and the




capture DNA on an electronic chemsensor for ultrasensitive detection exo-miRs (**Figure 4A**).<sup>149</sup> In the detecting process, TDNs nanomaterials containing the spatial structure was capability of linking with multiple electroactive compounds in stoichiometry on electrochemical sensors, which increased the sensitivity. The recommended sensing system illustrated remarkable anti-interference characteristics, specificity, and sensitivity, capable of determining low level of 34 aM in serum derived from patients with breast cancer without the need for enzymes. Motivated by this mechanism, Chao et al. utilized the flowered nickel-iron layered double hydroxide@AuNPs and TDNs to design a reusable electrochemiluminescence biosensor for miR-27a.<sup>150</sup> In the presence of miRNA, TDN-Ru(bpy)<sub>3</sub><sup>2+</sup> functions as an electrochemiluminescence sensor, established a steady sandwich structure with miR-27a and hairpin DNA through complementary base pairing, enabling the detection of miRNA detection. This platform exhibited heightened sensitivity, exceptional selectivity, and consistent reproducibility. The limit of detection (LoD) decreased to 11.7 aM with a broad linear dynamic range. Furthermore, Miao et al. designed TDNs nanomaterials with a walker strand on top and multiple track strands around it by assembling the triplex-forming oligonucleotide for miR-141 in cell and serum samples.<sup>151</sup> The platform enabled the regeneration of the detecting interface by manipulating pH conditions. Using the tetrahedral DNA-supported walking nanomachine allowed for high sensitive miR-141 assay by electrochemical method. The LoD were found to be as low as 4.9 aM, indicating a relatively low value. Simultaneously, The designed platform was effectively utilized for miRNA detection in cell and serum samples, allowing for differentiation between pathological data and those of healthy individuals. Based on the aforementioned reports, the TDNs-based electronic chemosensors exhibited high detection sensitivity, selectivity, and specificity, along with ease of operation. However, combining these methods requires a two-step process that includes the extraction of exosomes and the detection of miRNA, ultimately making it overly complex and impractical for practical applications. Within exosomes, TDNs were able to efficiently attach to specific microRNAs as a result of their spatial configuration, enabling the identification of target molecules. Furthermore, TDNs can readily be linked with electroactive compounds to generate electrochemical signals, which effectively amplified the detection signal. Currently, TDNs-based various electrochemical detection systems for exo-miRs have been developed based on the mechanism described above, with the successful detection of aM-level exo-miRs (**Table 1**).



To simplify and expedite the detection process, TDNs-based optical platforms were developed for exo-miR detection in situ for diagnostic purposes. Fluorescence resonance energy transfer (FRET) is one approach that enables sensitive, real-time monitoring of molecular interactions, enhancing the specificity of diagnostic assays.<sup>152</sup> Wang et al. created a TDNs-based FRET sensing platform to sensitively detect tumor-related miR-146b by DNA-assisted cyclic amplification (**Figure 4B**).<sup>153</sup> When there was no hsa-miR-146b as target, the 5' terminal and the 3' terminal of fluorescent DNA (HP) were functionalized with FAM and TAMRA did not change the hairpin structure due to the hybridization with the extended DNA strand of the TDNs. In that condition, the detection system exhibited a diminished FRET signal. The target hsa-miR-146b existed and induced the hybridization with the extended DNA strand of the DNA tetrahedron, inducing release of HP and produced the significant FRET. Therefore, the level of hsa-miR-146b can be determined through the alteration in the fluorescent value. Furthermore, another DNA was designed to replace hsa-miR-146b for cyclic signal amplification, increasing the sensitivity of assay in various cells. The LoD for hsa-miR-146b-5p was found to be 6 pM under optimal experimental conditions. Meanwhile, Hu et al. created a fluorescent TDNs probe using catalytic hairpin assembly (CHA) to enable the sensitive and specific detection of miR-21 and miR-155 in living cells through fluorescence signaling, as depicted in **Figure 4C**.<sup>154</sup> The DNA tetrahedron underwent functionalization with elements such as H1, H2, and Protector, each incorporating two pairs of fluorophores and quenching molecules. Exo-miR-21 triggered the chain displacement effect and emitted Cy3 fluorescence. The amplification of the signal in the CHA between H1 and H2 on FDTp was observed in the presence of miR-155, resulting in increased sensitivity of FAM fluorescence to miR-155. The miRNAs miR-21 and miR-155 were effectively visualized within viable cells, with a LoD of 5 pM. Meanwhile, Electrochemical and fluorescent sensors can detect exo-miRs in TDNs-based sensors. Electrochemical methods have excellent sensitivity. When comparing electrochemical methods to fluorescence, it is important to note that fluorescence possesses the unique ability to identify exo-miRs precisely in situ, making it a straightforward and practical approach for application. The optical approaches of TDNs usually requires the utilization of fluorescent molecules, demonstrating sensitivity typically within the picomolar range. The detection sensitivity is lower compared to electrochemical methods. One of the primary benefits is that the fluorescence detection method enables the in vivo detection of exosomal microRNAs



through a simple procedure.



**Figure 4.** (A) Illustrative diagram of TDNs synthesis for nanolabel supercharge-based electrochemical platform for exo-miR analysis,<sup>149</sup> (B) the diagrammatic representation of the designed DTNP-based FRET detecting sensor for miRNA analysis in living cells,<sup>142</sup> (C) synthesis of FDTp sensors for two types of miR-21 and miR-155 imaging in living cells.<sup>154</sup>

#### 4.2 Branched DNA nanomaterials

Branched DNA with multi-arm junctions was first created by Seeman, including 3-, 4-, 5-, 6-, 8-, and 12-arm junctions.<sup>155–157</sup> These nanomaterials have shown promise in serving as a flexible element in producing multifunctional nanomaterial biosensors by strategically adjusting their locations.<sup>26,28,158</sup> Luo's group utilized branched DNA labeled with fluorescent dyes (green and red) to create a system capable of detecting multiple targets.<sup>159,160</sup> Um et al. designed the fluorescence-coded DNA nanostructure probe system to detect the intracellular miR-21 and miR-22 in cell




(**Figure 5A**).<sup>161</sup> The system is capable of detecting multiple miRNAs across various breast cancer cell lines. Flow cytometry results illustrated the feasibility of directly comparing miRNA expression levels among different breast cancer cell lines with distinct characteristics. Additionally, Yang et al. detailed various branched DNA nanostructures used as scaffolds for creating super-branched silver nanoclusters (super-AgNCs), which exhibited adjustable spatial structures and fluorescent characteristics for biosensing applications.<sup>162,163</sup> Branched DNA-AgNCs displayed remarkable biocompatibility with living organisms and minimal harmful effects, which allows for their versatile use in a wide range of biomedical purposes. Meanwhile, Yang et al. designed thermostable DNA nanostructures to construct nucleic acids detection platform,<sup>164–166</sup> which was used to ultra-sensitively detect miR-223 on the Surface-enhanced Raman spectroscopy (SERS) platform (**Figure 5B**)<sup>166</sup>. This work successfully detected multiple liver cancer biomarkers, involving miR-223 and protein. The intricately structured branched-DNA utilized various complementary single strand DNA as captures for detecting miR-223, along with a double-stranded steady branched centre to enhance the distance of captures and expose a greater number of DNA captures to minimizing lateral interactions for improved sensitivity.<sup>167</sup> The incorporation of branched DNA results in 2 orders of magnitude increase in sensitivity for miRNA detection compared to using single-stranded DNA. LoD for miR-223 and AFP were 10 aM and  $10^{-12}$  M (S/N=3), respectively. Lin et al. created a localized branched catalytic hairpin assembly (bCHA) strategy for miRNAs in living cells by spatial-confinement approach of branched DNA structure without enzymes (**Figure 5C**).<sup>168</sup> This platform utilized three separate nanoprobe, each containing a distinct bCHA hairpin structure. The specific miRNA triggers bCHA among the LRN probes, resulting in the fluorescent signals amplification. Improvements in detection sensitivity and speed were achieved by anchoring each bCHA hairpin to individual nanospheres, while maintaining accuracy through the reduction of non-specific interactions. The LoD of miR-21 in vitro can reach as low as 3.66 pM in live cells and clinical tissues.<sup>169</sup> One target-induced connection of probes orderly to each other enhances the contrast in a cell by avoiding signal dispersion in amplification. This strategy was first programmed to implement miR-21 detection MCF-7 cells by target-triggered hairpin-free chain-branching assembly (HFCBA). Following the introduction of probes, the signal in cells exhibited a gradual increase in intensity in tandem with the growth of DNA dendrimers. This led to a significant enhancement in contrast within HFCBA through the



accumulation of in situ fluorescence, while also preventing signal dispersion during amplification. In 2020, Wang et al. developed an all-in-one biosensor to detect multiplexed exo-miRs in serum in situ for breast cancer diagnosis.<sup>170</sup> They utilized a branched DNA to construct a simple all-in-one biosensor for simultaneous determination of several miRNAs in situ by strand displacement reaction. Branched DNA included three oligonucleotides, which were extended at their 5' ends by incorporating three single-stranded recognition sequences with quenchers, involving BHQ1, BHQ2, and BHQ2, respectively. Following this, three reporter sequences were attached to the corresponding recognition sequences using different fluorophores (FAM, Cy3, and Cy5) to create a multicolor DNA biosensor capable of emitting self-quenched fluorescence. The biosensor is capable of entering exosomes, hybridizing with complementary miRNA targets to create longer duplexes, and releasing reporter sequences to activate significant fluorescent signals for the simultaneous assay of several miRNAs in exosomes. MiR-21, miR-27a, and miR-375 were selected as representative targets due to their elevated expression levels in breast cancer cells (MCF-7). Fluorescent characteristics of the detecting platform in the presence of MCF-7 exosomes illustrated the positive correlations to their level, and the LoD were found to be  $0.116 \mu\text{g mL}^{-1}$ ,  $0.125 \mu\text{g mL}^{-1}$ , and  $0.287 \mu\text{g mL}^{-1}$  for exosomes through detection of miR-21, miR-27a and miR-375, respectively. Conversely, no apparent correlations were found between fluorescence intensities and controlled concentrations of MCF-10A exosomes. Chen developed an aratiometric electrochemical DNA biosensor for the detection of exo-miR-21 (**Figure 5D**).<sup>171</sup> They created a locked nucleic acid (LNA)-modified branched DNA nanostructure to construct a ratiometric electrochemical DNA platform for exo miR-21 detection. miR-21 triggered the LNA-assisted strand displacement reaction on the the branched DNA, results in a structure alternation and the signal ratio change. This indicates the varying distances between the electrode surface and the two electroactive molecules attached to the branched DNA. The biosensor demonstrated high accuracy and sensitivity using a dual signal ratiometric method, achieving a limit of detection as low as 2.3 fM. Furthermore, due to the linear correlation between the logarithm of the signal ratio and the logarithm of the miR-21 concentration, the biosensor demonstrated sufficient stability in detecting miR-21 in MCF-7 cell-derived exosomes with effective selectivity. Branched DNA presents multiple modification sites that uphold a specific spatial arrangement. Various signaling molecules can be linked with different sites to enable multiplexing of exo-miRs. Currently, the integration of



branched DNA with SERS, electrochemical, and fluorescence sensors has led to the development of various detection platforms for Exo-miRs in situ, achieving a high sensitivity at the attomole (aM) level.



**Figure 5.** (A) Branched DNA nanomaterials with fluorescent dyes for intracellular miR-22 and miR-21 detection,<sup>161</sup> (B) branched DNA-based AgNF for detecting AFP and miRNA on SERS,<sup>166</sup> (C) synthesis of branched DNA nanomaterials for miR-21 detection in living cell,<sup>168</sup> (D) schematic diagram of a 1:L platform by target triggered synthesis of DNA dendrimers for contrast-increased imaging of miRNA in living cells.<sup>169</sup>

### 4.3 Cubic DNA nanocage sensors


The cubic DNA nanocage exhibits improved structural stability against enzymatic degradation, remarkable cellular permeability, high biocompatibility, and size-specific discrimination capability through the spatial confinement phenomenon.<sup>28</sup> Based on cubic DNA nanocages, the sensing platform primarily uses a fluorescence beacon to create an optical detection





assay directly on the samples. For example, Ye et al. created a cubic DNA nanocage probe designed to infiltrate EVs originating from MDA-MB-231 cells to monitor miRNA-10b in situ using a combination of fluorescence and quenching.<sup>172</sup> Within the exosome, miRNA-10b selectively binds with the quencher, restoring fluorescence for the cubic DNA nanocage. A strong linear relationship was found between fluorescent value and the concentration of extracellular vesicles (EVs) within the range of  $1.11 \times 10^8$  to  $5.70 \times 10^{10}$  particles  $\text{mL}^{-1}$  in 120 min. Meanwhile, Mi et al. developed a cubic DNA nanocage three-dimensional molecular beacon to accurately detect multiplexed exo-miRs in confined spaces (**Figure 6A**).<sup>140</sup> They devised a precise and adaptable cubic DNA nanostructure-based three-dimensional molecular beacon (ncMB), effectively overcoming the aforementioned challenge and achieving the detection of exo-miRNA. In this detection platform, steric hindrance and electrostatic repulsion caused by the distinctive three-dimensional arrangement of ncMB created a barrier between fluorescent probes, preventing unintended fluorescence quenching in single exo-miRs detection and unexpected FRET in dual exo-miRs detection. In comparison to conventional molecular beacons used for detecting exosomal miRNA in both single and dual modes, the larger DNA nanocage of ncMB was found to shield the fluorophore from interactions with the quencher and acceptor fluorophore by employing steric hindrance and electrostatic repulsion. Additionally, Li et al. utilized a spatial confinement strategy to design an accelerated DNA nanoprobe for fast, in situ monitoring of exo-miRs (**Figure 6B**).<sup>173</sup> This nanoprobe is capable of detecting exo-miRs and differentiating tumor-derived exosomes from those originating from normal cells to a high degree of accuracy, thereby establishing a foundation for bioimaging exo-miRs and diagnosing diseases. Sun et al. developed DNA cage-based exo-miRs detection for diagnostics of various stage in cancer progression using the characteristics of the size-selective ability and polyethylene glycol (PEG)-enhanced thermophoretic accumulation.<sup>174</sup> This method was highly sensitive, selective, and in situ detection of exo-miRs with a LoD of 2.05 fM in serum samples. Exo-miR-21 or miR155 demonstrated a discrimination accuracy of 90% in clinical tests when distinguishing between BC patients and healthy people, surpassing the performance of conventional molecular probes capable of detecting miRs. The cubic DNA nanocage combined with a fluorescence beacon can effectively transfer into the exosome and detect the miRs in site. The main focus of the detection process of the cubic DNA nanostructure primarily centers on the use of the optical detection system with a cage structure.

This structure serves to efficiently bind fluorescent molecules, safeguard them within the confines of the cage, and effectively shield them from interference by the fluorescence of neighboring molecules within the surrounding exosome. This analysis strategy increases the accuracy and precision of the detection system, allowing for the in situ detection of exo-miRs with fM levels of LoD.



**Figure 6.** (A) molecular beacon-based cubic DNA nanocage for exosomal miRNA detection in confined spaces,<sup>140</sup> (B) spatial confinement-based cubic DNA nanocage exosomal miRNA detection in situ,<sup>173</sup> (C) functional components and equivalent SSDC for the cyclic amplification detection of miRNAs,<sup>175</sup> (D) thermophoretic sensor based on NanoFlare (TSN) for in situ determination of exo-miRs,<sup>176</sup> (E) in situ simultaneous determination of several miRNAs in exosomes using MBs for diagnosis. F: fluorescent dye. Q: quencher,<sup>177</sup> (F) Simultaneous exosomal miR-21 and miR-155 detection by nucleic acid functionalized disposable paper-based sensors.<sup>178</sup>

#### 4.4 Spherical DNA nanomaterials

Spherical nucleic acids (SNA) consist of a core nanoparticle surrounded by a densely packed layer of oligonucleotides. In 1996, Mirkin et al. first developed SNA, which has since



demonstrated notable effects in diverse disciplines.<sup>179,180</sup> First, regarding the sandwich structure for miRNA detection, Liu et al. developed “sandwich” structure of the SNA and peptide nucleic acid (PNA) to construct an electrochemical exo-miRs detection platform.<sup>181</sup> PNA with a neutrally charged peptide skeleton serves as a capture probe, leading to a reduced charge and increased the binding efficiency with SNA-miRNA on the electrode surface. RuHex, being an electroactive compound, has the capability to interact with PNA-miRNA-SNA “sandwich” structures that have been assembled on the electrode surface via electrostatic adsorption. Chronocoulometric was utilized to measure the charges of RuHex, indicating the level of exo-miR. The LoD can reach as low as 49 aM with a detection range from 100 aM to 1 nM. The method demonstrates good sensitivity, but its complexity can be attributed to the initial lysis step. For achieving the exo-miRs detection in site, in 2023, yang et al. created a precise and reliable one-step method named dual-surface-protein-guided orthogonal recognition of tumor by profiling of miRs, which effectively increased the detecting accuracy of prostate cancer.<sup>182</sup> This platform employed two allosteric aptamers against exosomal marker CD63 and tumor marker EpCAM to build barcode for tumor-specific exosome subtypes. Meanwhile, the tumor-derived exosomes with barcode triggered targeted membrane fusion with liposome probes to introduce miRNA detection reagents, allowing for in situ sensitive detection of tumor-derived exo-miRs. With a signature of six miRs, this sensors differentiated PCa and benign prostatic hyperplasia with an accuracy of 100%. Additionally, the diagnostic accuracy were found to be 90.6% in classifying metastatic and nonmetastatic PCa. Wu et al. developed spherical nucleic acid-mediated spatial matching-guided nonenzymatic DNA circuits for miRNA detection to predict and prevent Malignant tumor (**Figure 6C**).<sup>175</sup> In this detection system, matching-guided nonenzymatic DNA circuit (SSDC) were created using AuNPs-based spherical nucleic acid to mediate spatial for effective detection of miRNAs. Because of the significantly improved ability to resist nuclease degradation, the occurrence of false positive signals is prevented even when present in a complex biological environment. The miRNA target has the capability to align the hairpin DNA probe into a linear structure, enabling the probe to enter the inner section of a core/shell DNA-functionalized signal nanoamplifier and trigger a strand displacement reaction that leads to the production of an enhanced fluorescence signal. The LoD was measured to be as low as 17 pM, and the imaging of microRNA was found to be well aligned with the widely accepted gold standard polymerase chain reaction (PCR) technique. The capability



to image intracellular miRNAs is greatly improved compared to traditional fluorescence in situ hybridization methods, allowing in vivo SSDC-based imaging to be proficient in accurately predicting the development of tumors. The use of intratumoral chemotherapy, directed by SSDC-based imaging, effectively managed the growth and advancement of tumors prior to the presentation of noticeable clinical symptoms. Sun et al. created a thermophoretic sensor known as NanoFlares (TSN) to detect exosomal exo-miRNAs in situ without the need for RNA extraction or target amplification (**Figure 6D**).<sup>176</sup> The accumulation of NanoFlare treated exosomes through thermophoresis results in an increased fluorescence signal when exo-miRs bind to NanoFlares, enabling the direct and quantitative measurement of exo-miRs at concentrations as low as 0.36 fM in 0.5  $\mu$ L serum samples within a 15-minute timeframe. One of the most reliable indicators, exo-miR-375, demonstrated an accuracy rate of 85% in identifying estrogen receptor-positive breast cancer in its early stages (stage I-II). This study offers a practical tool for enhancing cancer diagnosis. Duan et al. synthesized Au-cored spherical nucleic acid biosensor to detect multiplex exosomal RNAs in situ.<sup>183</sup> The probes' structures have been specifically tailored for the large targets of the PIWI-interacting RNA complex to prevent any steric hindrance in the analytical process, ensuring reliable sensitivity. Moreover, the strategic adjustment of the input during the synthesis of the anchor DNA-report DNA reactants with substantial concentration variations promotes the consistency of the product probes in both structure and detection sensitivity. The probes demonstrate the ability to detect piRNAs in exosomes in real human plasma samples with high sensitivity and specificity, allowing for differentiation between breast cancer patients and normal controls. This indicates that they may have the potential to serve as a straightforward and precise liquid biopsy tool for cancer diagnostics. Although the methods based on SNAs have high sensitivity, they exhibit significant noise in detecting exo-miRs. Spherical DNA interacts with various metal nanoparticles to synthesize sizable ball structures with nucleic acids surface, allowing for the efficient capture of target miRNAs for high sensitivity; however, this technique remains vulnerable to the impact of external interferences, resulting in the production of non-specific signals. This method allows for the detection of exosomal microRNA in its natural environment with attomolar sensitivity.

#### 4.5 Hairpin DNA-based sensors

Hairpin DNA sensors (HDS) were designed with a single-stranded DNA sequence that



features a stem structure created by base pairs and a loop containing unpaired nucleotides.<sup>184</sup> The stem-loop of HDS demonstrated notable recognizable properties of hybridization specificity and distinct selectivity towards target miRs. HDS is extensively utilized as biosensors for exo-miRs, predominantly employing HDS-based molecular beacons (MBs) and hybridization chain reaction (HCR)<sup>185,186</sup>.

MBs are hairpin DNA structures with fluorophores and a quencher specifically engineered to monitor exo-miRs.<sup>187</sup> Also, it is noteworthy that the choice of fluorophores for multiplexing biomarkers in an exosome is crucial because of its small size. Rhee et al. documented the *in vivo* single-step identification of particular miR-21 within intact exosomes using a nano-scale molecular beacon probe.<sup>130</sup> Permeabilization by streptolysin treatment further enhanced the delivery of MBs into exosomes, giving significantly increased signals from miR-21. Rhee et al., for the first time, further developed simultaneous and multiplexed detection exo-miRs (from breast cancer cell line, MCF-7) using MBs (**Figure 6E**).<sup>177</sup> MBs successfully hybridized with multiple miR-21, miR-375, and miR-27a in the cancer cell-derived exosomes even in the presence of high human serum concentration. The proposed method is beneficial to high-throughput analysis for disease diagnosis, prognosis, and response to treatment because it is a time-, labor-, and cost-saving technique. Zhang et al. developed accurate and rapid detection of exo-miR-21 by DNA zipper-mediated membrane fusion approach for cancer diagnosis.<sup>188</sup> In this detection platform, a lipid vesicle biosensor with MBs was synthesized and subsequently functionalized with zipper DNA nanostructure (ZDCs) on its surface. Additionally, complementary zipper DNA nanostructure (cZDCs) are incorporated on the exosome. Upon co-mixing ZDC and cZDC, the process activates membrane fusion between exosomes and vesicle probes, leading to the identification of exo-miR-21 by encapsulated MBs and subsequent fluorescence emission that can be detected within a 30-minute timeframe. Significantly, flow cytometry allows for the differentiation of miR-21-overexpressed tumor exosomes derived from cell culture medium or clinical patient serums from exosomes secreted by normal cells.

Additionally, HCR, known as hybridization chain reaction, proved to be yet another effective technique for isothermal amplification that operates based on the opening of functional hairpin DNA structures.<sup>189</sup> Once the specific target miRNA was identified, it resulted in the formation of lengthy double helix structures comprised of multiple repeated units through the process of



hybridization between two DNA hairpins.<sup>190</sup> Zou et al. utilized HCR-based platform to construct the sensitive imaging method for intracellular miRNA in living cells.<sup>191</sup> For the purpose of delivery and recruitment, exosomes were altered by incorporating three different cholesterol-modified hairpins (designated as H1, H2, and H3), with H1 specifically intended for anchoring the target miRNA, while H2 and H3 were equipped with photocleavable linkers (PC-linkers) to facilitate spatiotemporal hybridization chain reaction (HCR). By strategically releasing probes with concentrated quantities in specific locations to effectively initiate hybridization chain reaction (HCR) and subsequently gathering the resultant double-stranded DNA polymers rather than allowing them to disperse throughout the cytoplasm, the LCR-HCR technique has the potential to greatly enhance imaging contrast by confining all of the chemical substances involved on exosome carriers. The comparison of low- and high-contrast reverse immunosensor systems measured the microRNA-21 at a limit of detection as low as 3.3 pM (equivalent to 165 attomoles per 50 microliters) in laboratory conditions and exhibited a response in vivo that was four times greater than that achieved by conventional high-contrast reverse immunosensors. The LCR-HCR method offers a highly effective tool for the visualization of microRNAs in living cells and for the detection of cancer. Li and colleagues created a method that combines hybridization chain reaction (HCR) and DNAzyme technology to amplify signals without the need for enzymes. This method allows for the detection of specific exo-miRs in cancer cell culture media and serum samples from cancer patients through the self-assembly of polymer DNAzyme nanostructures triggered by the target.<sup>192</sup> He et al. synthesized hairpins that loaded DNAzyme subunits in 3' and 5' ends. When exo-miRs existed, the long DNA double helixes were produced, leading to various activated DNAzymes by two nearby units. After that, the DNAzymes prompted the cleavage of fluorescent reporter probes, resulting in the emission of fluorescence signals. Based on this mechanism, quantifying miRNAs assay were conducted in circulating exosomes. Choi et al. developed that present hydrogel-based HCR for multiplex signal amplification to detect urinary exo-miRs (hsa-miR-6090 and hsa-miR-3665) from human clinical samples.<sup>193</sup> They identified small amounts (~amol) of exo-miRs from 600  $\mu$ L of urine with up to ~35-fold amplification. Furthermore, they created ratiometric detection platform using a reference miRNA, which indicated this analysis platform have a great potential for differentiating prostate cancer patients from healthy controls. HDS-based molecular beacons (MBs) and HCR-based detection platform primarily relies on the target molecule initiating the



opening of a hairpin, leading to fluorescence signal enhancement or amplification reactions. The assay necessitates carefully designed DNA sequences that specifically target regions within hairpins to minimize non-specific binding. Currently, this method can detect at the level of femtomolar sensitivity.

The analytical performance of the various strategies for exo-miRNA detection is summarized in **Table 1**. Even though there is a recent emergence of exo-miRNA as potential biomarkers, their clinical applications continue to be constrained by their low concentration in small quantities of clinical samples. Consequently, the endeavor to create a diagnostic tool that is non-invasive and specific, and to profile markers of exosomal microRNAs from urine, presents a considerable and pressing challenge. Even though the identification of terminal stem-loop/hairpin DNA-based exo-miRs brings various benefits due to its straightforward structure and exceptional sensitivity, it is also confronted with obstacles linked to elevated background signals and potential external disruptions.

**Table 1.** Different DNA nanostructure-based biosensors for exo-miRs detection.

| DNA Nanomaterials | Methods                | Target miRNA             | Linear range                      | LOD                               | Diseases   | Refs |
|-------------------|------------------------|--------------------------|-----------------------------------|-----------------------------------|------------|------|
| TDNs              | Electrochemical sensor | miR-27a                  | 100 aM-1 nM                       | 11.7 aM                           | Tumors     | 150  |
|                   | Electrochemical sensor | miR-21                   | 100 aM-1 nM                       | 34.0 aM                           | Tumors     | 149  |
|                   | Fluorescent sensor     | miR-21, miR-141,         | --                                | 100.0 fM                          | Cancer     | 194  |
|                   | Fluorescent sensing    | miR-21, miR-27a, miR-375 | 5-140 nM;<br>2-100 nM,<br>7-70 nM | 462.0 pM<br>301.0 pM,<br>154.0 pM | cancer     | 170  |
| Branched DNA      | Electrochemical sensor | miR-21                   | 10-70 fM                          | 2.3fM                             | MCF-7 cell | 171  |
|                   | Fluorescent sensor     | miR-141                  | 100-900 pM                        | 57.6pM                            | Cancer     | 195  |
| Cubic DNA         | Fluorescent            | miR-21                   | 10–100 nM;                        | 3.4 nM,                           | U251 cells | 140  |



|               |                        |           |   |  |  |                                      |
|---------------|------------------------|-----------|---|--|--|--------------------------------------|
| Spherical DNA | sensor                 | miR-221   | 10-120 nM   | 2.3 nM                                       |  |                                      |
|               | Fluorescent sensor     | miR-21    | 100 nM  | pM-25  | 77.4 pM                                      | No 196                               |
|               | Fluorescent sensor     | miR-44438 | No  | No   | No   | Sleep disorder 197                   |
|               | Fluorescent sensor     | miR-1246  | 0-30 nM   | 0.68 nM                                      | 0.68 nM                                      | Breast Cancer 198                    |
|               | SERS                   | miR-10b   | No  | 0.21 fM                                      | 0.21 fM                                      | Pancreatic ductal adenocarcinoma 199 |
|               | Fluorescent sensor     | miR-21    | 10 <sup>3</sup> -10 <sup>8</sup> particles $\mu\text{L}^{-1}$ | 10 <sup>3</sup> particles $\mu\text{L}^{-1}$ | 10 <sup>3</sup> particles $\mu\text{L}^{-1}$ | Tumor 200                            |
| Hairpin DNA   | Electrochemical sensor | miR-181   | 10 nM   | fM-100                                       | 7.94 fM                                      | Cardiovascular disease 201           |

## 5 Importance and challenges

Exo-miRs are emerging as powerful early biomarkers, forging new directions in molecular diagnostics, prognosis, and personalized disease management, which represents a notable advancement, yielding quicker diagnostic results and greater accessibility for patients. In situ detection facilitates the simultaneous detection of exo-miRs, which is crucial for examining complex miRNA-involved regulatory networks. DNA nanomaterials have excellent cell permeability, biocompatibility, low toxicity, and resistance to enzyme degradation for their exceptional sensitivity, enabling the detection of low levels of exo-miR in confined spaces. Despite the many advancements in technologies for exo-miR detection, we must stress a few critical points about exo-miR diagnostics, establish ultrasensitive biosensing in site, reliable biomarkers, and multiplexed and sensitive assays.





The concentration of exo-miRs in exosomes is low, and these molecules' exclusive expression or downregulation content was not significant in specific human diseases. Therefore, it is essential that the detection method be both accurate and highly sensitive to target miRs effectively. The integration of DNA nanomaterials with both electrochemical and optical detection systems has now facilitated the detection of exo-miRs at unprecedentedly low attomolar levels, thereby representing a notable breakthrough in the field of clinical disease diagnostics. However, numerous factors influence miRNA expression levels in real complex samples, including age, gender, lifestyle, and an individual's multiple factors collaboratively influence the expression level of exo-miRs. The expression levels of target miRNAs were reportedly upregulated or downregulated in patients with various diseases. Considering these factors, certain miRNA and concentrations as standards for disease detection need to be first found. Further research is needed to explore the regulation of miRNA expression levels and its correlation with disease in relation to interfering factors. Identify the precise target exosomal microRNA molecule for reliable disease's diagnostics. As mentioned above, in specific diseases, multiplexed exo-miRs should be used as biomarkers simultaneously for accurate disease diagnostics. Therefore, it is important to create DNA nanomaterials that feature several target molecular junction sites, with each site corresponding to a unique signal to enable recognition of multiple targets. The above-mentioned challenges of DNA nanomaterial-based detection platforms still need to be further studied for exo-miRs as biomarkers of disease's diagnostics.

### Author contributions

Zhikun Zhang: Conceptualization (equal); writing – original draft (equal); writing – review and editing (equal). Md. Ahasan Ahamed: – original draft (equal); writing – review and editing (equal). Dayong Yang: Conceptualization (equal); funding acquisition (lead); supervision (lead); writing – original draft (equal); writing – review and editing (equal).

### Declaration of Competing Interest

All authors declare no conflict of interests.

### Acknowledgements

This work was supported by National Natural Science Foundation of China (Grant No.:



22225505).

## Reference

- 1 R. Shang, S. Lee, G. Senavirathne and E. C. Lai, *Nat Rev Genet*, 2023, **24**, 816–833.
- 2 F. Wahid, A. Shehzad, T. Khan and Y. Y. Kim, *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 2010, **1803**, 1231–1243.
- 3 Z. Sun, Y. Wu, F. Gao, H. Li, C. Wang, L. Du, L. Dong and Y. Jiang, *Acta Biomaterialia*, 2023, **155**, 80–98.
- 4 S. Shin, Y. H. Park, S.-H. Jung, S.-H. Jang, M. Y. Kim, J. Y. Lee and Y. Chung, *npj Genom. Med.*, 2021, **6**, 45.
- 5 M. Jang, G. Choi, Y. Y. Choi, J. E. Lee, J.-H. Jung, S. W. Oh, D. H. Han, H. Lee, J.-H. Park, J.-H. Cheong and P. Kim, *NPG Asia Mater*, 2019, **11**, 79.
- 6 L. Saadatpour, E. Fadaee, S. Fadaei, R. Nassiri Mansour, M. Mohammadi, S. M. Mousavi, M. Goodarzi, J. Verdi and H. Mirzaei, *Cancer Gene Ther*, 2016, **23**, 415–418.
- 7 N. Jafari, P. Llevenes and G. V. Denis, *Nat Rev Endocrinol*, 2022, **18**, 327–328.
- 8 Z. Zhang, Y. Zou, C. Song, K. Cao, K. Cai, S. Chen, Y. Wu, D. Geng, G. Sun, N. Zhang, X. Zhang, Y. Zhang, Y. Sun and Y. Zhang, *Journal of Advanced Research*, 2023, S2090123223004022.
- 9 Y. Yu, K. Hou, T. Ji, X. Wang, Y. Liu, Y. Zheng, J. Xu, Y. Hou and G. Chi, *Mol Cell Biochem*, 2021, **476**, 2111–2124.
- 10 N. Mahtal, O. Lenoir, C. Tinel, D. Anglicheau and P.-L. Tharaux, *Nat Rev Nephrol*, 2022, **18**, 643–662.
- 11 B. Mohan, S. Kumar, S. Kumar, K. Modi, D. Tyagi, D. Papukashvili, N. Rcheulishvili and A. J. L. Pombeiro, *Sens. Diagn.*, 2023, **2**, 78–89.
- 12 J. Tang, X. Jia, Q. Li, Z. Cui, A. Liang, B. Ke, D. Yang and C. Yao, *Proc. Natl. Acad. Sci. U.S.A.*, 2023, **120**, e2303822120.
- 13 N. P. Hessvik and A. Llorente, *Cell. Mol. Life Sci.*, 2018, **75**, 193–208.
- 14 M. A. U. Khalid, Md. A. Ahamed, M. Dong, A. Kshirsagar and W. Guan, *Biosensors and Bioelectronics*, 2025, **268**, 116895.
- 15 Y. Xue, K. Wang, Y. Jiang, Y. Dai, X. Liu, B. Pei, H. Li, H. Xu and G. Zhao, *Biosensors and Bioelectronics*, 2024, **247**, 115927.
- 16 H. Zhang, L. Cao, J. Brodsky, I. Gablech, F. Xu, Z. Li, M. Korabecna and P. Neuzil, *TrAC Trends in Analytical Chemistry*, 2024, **174**, 117676.
- 17 H. Wang, *Cells*, 2024, **13**, 1277.
- 18 M. A. Ahamed, A. J. Politza, T. Liu, M. A. U. Khalid, H. Zhang and W. Guan, *Nanotechnology*, 2025, **36**, 042001.
- 19 S. Tam, R. De Borja, M.-S. Tsao and J. D. McPherson, *Laboratory Investigation*, 2014, **94**, 350–358.
- 20 H.-S. Tan, T. Wang, H.-N. Sun, A. Liu and S.-S. Li, *TrAC Trends in Analytical Chemistry*, 2023, **167**, 117253.
- 21 W. Wu, X. Yu, J. Wu, T. Wu, Y. Fan, W. Chen, M. Zhao, H. Wu, X. Li and S. Ding, *Biosensors and Bioelectronics*, 2021, **175**, 112835.
- 22 J. Peng, B. Li, Z. Ma, Z. Qiu, H. Hu, Y. Jiang and D. Gao, *Talanta*, 2025, **281**,



126838.

23 Y. Xia, Z. Huang, T. Chen, L. Xu, G. Zhu, W. Chen, G. Chen, S. Wu, J. Lan, X. Lin and J. Chen, *Biosensors and Bioelectronics*, 2022, **209**, 114259.

24 T. Liu, A. J. Politza, Md. A. Ahamed, A. Kshirsagar, Y. Zhu and W. Guan, *Biosensors and Bioelectronics*, 2024, 116997.

25 X. Li, X. Li, D. Li, M. Zhao, H. Wu, B. Shen, P. Liu and S. Ding, *Biosensors and Bioelectronics*, 2020, **168**, 112554.

26 C. Seeman, *J. Theor. Biol.*, **99**, 237–247.

27 J. Chen and N. C. Seeman, *Nature*, 1991, **350**, 631–633.

28 N. C. Seeman and H. F. Sleiman, *Nat Rev Mater*, 2017, **3**, 17068.

29 J. O'Brien, H. Hayder, Y. Zayed and C. Peng, *Front. Endocrinol.*, 2018, **9**, 402.

30 B. Hrdlickova, R. C. De Almeida, Z. Borek and S. Withoff, *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, 2014, **1842**, 1910–1922.

31 J. Krol, I. Loedige and W. Filipowicz, *Nat Rev Genet*, 2010, **11**, 597–610.

32 M. A. Kumar, S. K. Baba, H. Q. Sadida, S. Al. Marzooqi, J. Jerobin, F. H. Altemani, N. Algehainy, M. A. Alanazi, A.-B. Abou-Samra, R. Kumar, A. S. Al-Shabeeb Akil, M. A. Macha, R. Mir and A. A. Bhat, *Sig Transduct Target Ther*, 2024, **9**, 27.

33 A. Gallo, M. Tandon, I. Alevizos and G. G. Illei, *PLoS ONE*, 2012, **7**, e30679.

34 A. Turchinovich, L. Weiz, A. Langheinz and B. Burwinkel, *Nucleic Acids Research*, 2011, **39**, 7223–7233.

35 J. Zhang, S. Li, L. Li, M. Li, C. Guo, J. Yao and S. Mi, *Genomics, Proteomics & Bioinformatics*, 2015, **13**, 17–24.

36 S. Mathivanan, H. Ji and R. J. Simpson, *Journal of Proteomics*, 2010, **73**, 1907–1920.

37 A. J. Politza, T. Liu, A. Kshirsagar, M. Dong, M. A. Ahamed and W. Guan, 2024, preprint, 4781228, DOI: 10.2139/ssrn.4781228.

38 A. J. Politza, T. Liu, A. Kshirsagar, M. Dong, Md. Ahasan Ahamed and W. Guan, 2024, preprint, DOI: 10.17504/protocols.io.kxygxyj4wl8j/v1.

39 S. Modani, D. Tomar, S. Tangirala, A. Sriram, N. K. Mehra, R. Kumar, D. K. Khatri and P. K. Singh, *Journal of Drug Targeting*, 2021, **29**, 925–940.

40 Q. Guo, H. Wang, Y. Yan, Y. Liu, C. Su, H. Chen, Y. Yan, R. Adhikari, Q. Wu and J. Zhang, *Front. Oncol.*, 2020, **10**, 472.

41 B. Yue, H. Yang, J. Wang, W. Ru, J. Wu, Y. Huang, X. Lan, C. Lei and H. Chen, *Cell Proliferation*, 2020, **53**, e12857.

42 Y. Chen, Y. Zhao, Y. Yin, X. Jia and L. Mao, *Bioengineered*, 2021, **12**, 8186–8201.

43 F. Fabbiano, J. Corsi, E. Gurrieri, C. Trevisan, M. Notarangelo and V. G. D'Agostino, *J of Extracellular Vesicle*, 2020, **10**, e12043.

44 B. Hannafon and W.-Q. Ding, *IJMS*, 2013, **14**, 14240–14269.

45 Y. Wang, T. Xiao, C. Zhao and G. Li, *IJMS*, 2023, **25**, 255.

46 B. Février and G. Raposo, *Current Opinion in Cell Biology*, 2004, **16**, 415–421.

47 P. Perrin, L. Janssen, H. Janssen, B. Van Den Broek, L. M. Voortman, D. Van Elstrand, I. Berlin and J. Neefjes, *Current Biology*, 2021, **31**, 3884–3893.e4.

48 X. Tang, M. Leng, W. Tang, Z. Cai, L. Yang, L. Wang, Y. Zhang and J. Guo, *Molecules*, 2024, **29**, 1199.

49 N. W. Lin, C. Liu, I. V. Yang, L. A. Maier, D. L. DeMeo, C. Wood, S. Ye, M. H.



Cruse, V. L. Smith, C. A. Vyhlidal, K. Kechris and S. Sharma, *Front. Genet.*, 2022, **13**, 762834.

50 P. Kanlikilicer, R. Bayraktar, M. Denizli, M. H. Rashed, C. Ivan, B. Aslan, R. Mitra, K. Karagoz, E. Bayraktar, X. Zhang, C. Rodriguez-Aguayo, A. A. El-Arabey, N. Kahraman, S. Baydogan, O. Ozkayar, M. L. Gatza, B. Ozpolat, G. A. Calin, A. K. Sood and G. Lopez-Berestein, *eBioMedicine*, 2018, **38**, 100–112.

51 G. K. Patel, M. A. Khan, A. Bhardwaj, S. K. Srivastava, H. Zubair, M. C. Patton, S. Singh, M. Khushman and A. P. Singh, *Br J Cancer*, 2017, **116**, 609–619.

52 C. Li, T. Zhou, J. Chen, R. Li, H. Chen, S. Luo, D. Chen, C. Cai and W. Li, *J Transl Med*, 2022, **20**, 6.

53 H. T. Nguyen, S. E. O. Kacimi, T. L. Nguyen, K. H. Suman, R. Lemus-Martin, H. Saleem and D. N. Do, *Biology*, 2021, **10**, 417.

54 S. U. Khan, M. U. Khan, M. Azhar Ud Din, I. M. Khan, M. I. Khan, S. Bungau and S. S. U. Hassan, *Front. Immunol.*, 2023, **14**, 1166487.

55 Z. Zhang, T. Xing, Y. Chen and J. Xiao, *Biomedicine & Pharmacotherapy*, 2018, **106**, 1135–1143.

56 K. Liu, X. Gao, B. Kang, Y. Liu, D. Wang and Y. Wang, *Front. Oncol.*, 2022, **12**, 836548.

57 J. C. Santos, N. D. S. Lima, L. O. Sarian, A. Matheu, M. L. Ribeiro and S. F. M. Derchain, *Sci Rep*, 2018, **8**, 829.

58 J. M. Cozar, I. Robles-Fernandez, A. Rodriguez-Martinez, I. Puche-Sanz, F. Vazquez-Alonso, J. A. Lorente, L. J. Martinez-Gonzalez and M. J. Alvarez-Cubero, *Mutation Research/Reviews in Mutation Research*, 2019, **781**, 165–174.

59 J. A. Deiliis, *Int J Obes*, 2016, **40**, 88–101.

60 J. Wang, B.-L. Yue, Y.-Z. Huang, X.-Y. Lan, W.-J. Liu and H. Chen, *IJMS*, 2022, **23**, 2461.

61 Z. Zhang, T. Liu, M. Dong, Md. A. Ahamed and W. Guan, *WIREs Nanomed Nanobiotechnol*, 2024, **16**, e1969.

62 N. Al Rawi, N. Elmabrouk, R. Abu Kou, S. Mkadmi, Z. Rizvi and Z. Hamdoon, *Archives of Oral Biology*, 2021, **125**, 105108.

63 M. Song, H. Bai, P. Zhang, X. Zhou and B. Ying, *Int J Oral Sci*, 2023, **15**, 2.

64 Z. Xie, G. Chen, X. Zhang, D. Li, J. Huang, C. Yang, P. Zhang, Y. Qin, Y. Duan, B. Gong and Z. Li, *PLoS ONE*, 2013, **8**, e57502.

65 J. Du and L. Zhang, *Oncology Letters*, 2017, **14**, 1387–1394.

66 F. Li, J. M. Yoshizawa, K.-M. Kim, J. Kanjanapangka, T. R. Grogan, X. Wang, D. E. Elashoff, S. Ishikawa, D. Chia, W. Liao, D. Akin, X. Yan, M.-S. Lee, R. Choi, S.-M. Kim, S.-Y. Kang, J.-M. Bae, T.-S. Sohn, J.-H. Lee, M.-G. Choi, B.-H. Min, J.-H. Lee, J. J. Kim, Y. Kim, S. Kim and D. T. W. Wong, *Clinical Chemistry*, 2018, **64**, 1513–1521.

67 J. Yang, R. Xu, C. Wang, J. Qiu, B. Ren and L. You, *Cancer Communications*, 2021, **41**, 1257–1274.

68 M. Koopaie, F. Abedinejad, S. Manifar, R. Mousavi, S. Kolahdooz and A. Shamshiri, *Gene Reports*, 2021, **25**, 101317.

69 T. P. N. Nguyen, M. Kumar, E. Fedele, G. Bonanno and T. Bonifacino, *IJMS*, 2022, **23**, 4718.

70 H. Li, L. Yu, M. Li, X. Chen, Q. Tian, Y. Jiang and N. Li, *Molec Gen & Gen*



*Med*, 2020, **8**, e1189.

71 I. Kawikova and P. W. Askenase, *Brain Research*, 2015, **1617**, 63–71.

72 X. Xia, Y. Wang, Y. Huang, H. Zhang, H. Lu and J. C. Zheng, *Progress in Neurobiology*, 2019, **183**, 101694.

73 Y. W. Chen, H. V. Lee and S. B. Abd Hamid, *Carbohydrate Polymers*, 2017, **178**, 57–68.

74 R. Shi, P.-Y. Wang, X.-Y. Li, J.-X. Chen, Y. Li, X.-Z. Zhang, C.-G. Zhang, T. Jiang, W.-B. Li, W. Ding and S.-J. Cheng, *Oncotarget*, 2015, **6**, 26971–26981.

75 E. V. Filatova, A. Kh. Alieva, M. I. Shadrina and P. A. Slominsky, *Biochemistry Moscow*, 2012, **77**, 813–819.

76 E. V. Filatova, A. Kh. Alieva, M. I. Shadrina and P. A. Slominsky, *Biochemistry Moscow*, 2012, **77**, 813–819.

77 L. Leggio, S. Vivarelli, F. L'Episcopo, C. Tirolo, S. Caniglia, N. Testa, B. Marchetti and N. Iraci, *Int. J. Mol. Sci.*, 2017, **18**, 2698.

78 Q. Shi, W. Kang, Z. Liu and X. Zhu, *Heliyon*, 2023, **9**, e20595.

79 I. Manna, S. De Benedittis, A. Quattrone, D. Maisano, E. Iaccino and A. Quattrone, *Pharmaceuticals*, 2020, **13**, 243.

80 X. Wu, T. Zheng and B. Zhang, *Neurosci. Bull.*, 2017, **33**, 331–338.

81 L. Yuan and J.-Y. Li, *ACS Chem. Neurosci.*, 2019, **10**, 964–972.

82 E. Paccosi and L. Proietti-De-Santis, *IJMS*, 2023, **24**, 9547.

83 I. Manna, A. Quattrone, S. De Benedittis, B. Vescio, E. Iaccino and A. Quattrone, *Parkinsonism & Related Disorders*, 2021, **93**, 77–84.

84 D. Chandran, S. Krishnan, M. Urulangodi and S. Gopala, *Neurol Sci*, 2024, **45**, 3625–3639.

85 J. R. Pinnell, M. Cui and K. Tieu, *Journal of Neurochemistry*, 2021, **157**, 413–428.

86 H. Zhang, X. Liu, J. Shi, X. Su, J. Xie, Q. Meng and H. Dong, *Front. Cell. Infect. Microbiol.*, 2024, **14**, 1418168.

87 R. E. Drury, D. O'Connor and A. J. Pollard, *Front. Immunol.*, 2017, **8**, 1182.

88 S. Zhou and J. Xu, *Exp Ther Med*, DOI:10.3892/etm.2019.7528.

89 S. D. Hicks, D. Zhu, R. Sullivan, N. Kannikeswaran, K. Meert, W. Chen, S. Suresh and U. Sethuraman, *IJMS*, 2023, **24**, 8175.

90 I. Saulle, M. Garziano, G. Cappelletti, F. Limanaqi, S. Strizzi, C. Vanetti, S. Lo Caputo, M. Polisenio, T. A. Santantonio, M. Clerici and M. Biasin, *IJMS*, 2023, **24**, 10992.

91 R. Mishra and A. C. Banerjea, *Front. Immunol.*, 2021, **12**, 656700.

92 J. Zhang, X. Li, J. Hu, P. Cao, Q. Yan, S. Zhang, W. Dang and J. Lu, *Virol J*, 2020, **17**, 51.

93 R. D. Estep, A. N. Govindan, M. Manoharan, H. Li, S. S. Fei, B. S. Park, M. K. Axthelm and S. W. Wong, *PLoS ONE*, 2020, **15**, e0228484.

94 M. A. Ahamed and W. Guan, *Biophysical Journal*, 2024, **123**, 145a.

95 J. T. Granados-Riveron and G. Aquino-Jarquin, *Cells*, 2021, **10**, 1655.

96 Md. A. Ahamed, M. A. U. Khalid, M. Dong, A. J. Politza, Z. Zhang, A. Kshirsagar, T. Liu and W. Guan, *Biosensors and Bioelectronics*, 2024, **246**, 115866.

97 R. Ojha, R. Nandani, R. K. Pandey, A. Mishra and V. K. Prajapati, *Journal Cellular Physiology*, 2019, **234**, 1030–1043.

98 V. Martini, F. D'Avanzo, P. M. Maggiora, F. M. Varughese, A. Sica and A.



- Gennari, *ecancer*, DOI:10.3332/ecancer.2020.1149.
- 99 R. Kaur and K. Kumar, *European Journal of Medicinal Chemistry*, 2021, **215**, 113220.
- 100 P. Zhou, M. Dong, J. Wang, F. Li, J. Zhang and J. Gu, *Exp Ther Med*, DOI:10.3892/etm.2018.6685.
- 101 A. Khanam, J. V. Chua and S. Kottlilil, *IJMS*, 2021, **22**, 5497.
- 102 Y. Yin, Y. Zhao, Q. Chen, Y. Chen and L. Mao, *Front. Microbiol.*, 2022, **13**, 1044832.
- 103 T.-L. Liao, Y.-M. Chen, S.-L. Hsieh, K.-T. Tang, D.-Y. Chen, Y.-Y. Yang, H.-J. Liu and S.-S. Yang, *mBio*, 2021, **12**, e02764-21.
- 104 S. Li, S. Li, S. Wu and L. Chen, *BioMed Research International*, 2019, **2019**, 1–9.
- 105 Y. Hu, F. Cui, S. Wang, C. Liu, S. Zhang, R. Wang, J. Song and Y. Zhang, *Front. Cell. Infect. Microbiol.*, 2023, **13**, 1217984.
- 106 J. S. Nahand, M. Mahjoubin-Tehran, M. Moghoofei, M. H. Pourhanifeh, H. R. Mirzaei, Z. Asemi, A. Khatami, F. Bokharaei-Salim, H. Mirzaei and M. R. Hamblin, *Epigenomics*, 2020, **12**, 353–370.
- 107 J. Li, Y. Zhang and B. Luo, *Cancers*, 2022, **14**, 3552.
- 108 M. Mohammadinasr, S. Montazersaheb, V. Hosseini, H. Kahroba, M. Talebi, O. Molavi, H. Ayromlou and M. S. Hejazi, *Cytokine*, 2024, **179**, 156624.
- 109 H. Dong, Q. Gao, X. Peng, Y. Sun, T. Han, B. Zhao, Y. Liu, C. Wang, X. Song, J. Wu and L. Yang, *Front. Vet. Sci.*, 2017, **4**, 186.
- 110 A. Bayer, E. Delorme-Axford, C. Sleighter, T. K. Frey, D. W. Trobaugh, W. B. Klimstra, L. A. Emert-Sedlak, T. E. Smithgall, P. R. Kinchington, S. Vadia, S. Seveau, J. P. Boyle, C. B. Coyne and Y. Sadovsky, *American Journal of Obstetrics and Gynecology*, 2015, **212**, 71.e1-71.e8.
- 111 M. V. Liberti and J. W. Locasale, *Trends in Biochemical Sciences*, 2016, **41**, 211–218.
- 112 R. Mirzaei, S. Babakhani, P. Ajorloo, R. H. Ahmadi, S. R. Hosseini-Fard, H. Keyvani, Y. Ahmadyousefi, A. Teimoori, F. Zamani, S. Karampoor and R. Yousefimashouf, *Mol Med*, 2021, **27**, 34.
- 113 D. De Gonzalo-Calvo, R. W. Van Der Meer, L. J. Rijzewijk, J. W. A. Smit, E. Revuelta-Lopez, L. Nasarre, J. C. Escola-Gil, H. J. Lamb and V. Llorente-Cortes, *Sci Rep*, 2017, **7**, 47.
- 114 D. Zheng, M. Huo, B. Li, W. Wang, H. Piao, Y. Wang, Z. Zhu, D. Li, T. Wang and K. Liu, *Front. Cell Dev. Biol.*, 2021, **8**, 616161.
- 115 N. Yu-Qing, L. Xiao, Z. Jun-Kun and L. You-Shuo, *Aging and disease*, 2020, **11**, 164.
- 116 M. Lu, S. Yuan, S. Li, L. Li, M. Liu and S. Wan, *J. of Cardiovasc. Trans. Res.*, 2019, **12**, 68–74.
- 117 J. Heo and H. Kang, *IJMS*, 2022, **23**, 1002.
- 118 F. Sessa, M. Salerno, M. Esposito, G. Cocimano and C. Pomara, *IJMS*, 2023, **24**, 5192.
- 119 H. Ling, Z. Guo, Y. Shi, L. Zhang and C. Song, *Front. Physiol.*, 2020, **11**, 654.
- 120 C. Iaconetti, S. Sorrentino, S. De Rosa and C. Indolfi, *Physiology*, 2016, **31**, 16–24.



- 121 L. Wang, J. Liu, B. Xu, Y. Liu and Z. Liu, *The Kaohsiung J of Med Scie*, 2018, **34**, 626–633.
- 122 F. Vacante, L. Denby, J. C. Sluimer and A. H. Baker, *Vascular Pharmacology*, 2019, **112**, 24–30.
- 123 H. Jia, L. Kang, Z. Ma, S. Lu, B. Huang, C. Wang, Y. Zou and Y. Sun, *J Cardiothorac Surg*, 2021, **16**, 230.
- 124 W. Pan, J. Liang, H. Tang, X. Fang, F. Wang, Y. Ding, H. Huang and H. Zhang, *The International Journal of Biochemistry & Cell Biology*, 2020, **118**, 105645.
- 125 H. Li, J. Zhan, C. Chen and D. Wang, *Medical Review*, 2022, **2**, 140–168.
- 126 R. Xue, W. Tan, Y. Wu, B. Dong, Z. Xie, P. Huang, J. He, Y. Dong and C. Liu, *Front. Cardiovasc. Med.*, 2020, **7**, 592412.
- 127 S. Gurunathan, M. H. Kang and J.-H. Kim, *Front. Immunol.*, 2021, **12**, 716407.
- 128 Y. Fu and H. Sun, *J Ovarian Res*, 2022, **15**, 66.
- 129 C. Li, Y.-Q. Ni, H. Xu, Q.-Y. Xiang, Y. Zhao, J.-K. Zhan, J.-Y. He, S. Li and Y.-S. Liu, *Sig Transduct Target Ther*, 2021, **6**, 383.
- 130 J. H. Lee, J. A. Kim, M. H. Kwon, J. Y. Kang and W. J. Rhee, *Biomaterials*, 2015, **54**, 116–125.
- 131 Y. Wu, Y. Zhang, X. Zhang, S. Luo, X. Yan, Y. Qiu, L. Zheng and L. Li, *Biosensors and Bioelectronics*, 2021, **177**, 112962.
- 132 A. Kögel, A. Keidel, M.-J. Loukeri, C. C. Kuhn, L. M. Langer, I. B. Schäfer and E. Conti, *Nature*, 2024, **635**, 237–242.
- 133 Y. Dong, C. Yao, Y. Zhu, L. Yang, D. Luo and D. Yang, *Chem. Rev.*, 2020, **120**, 9420–9481.
- 134 C. Yao, Y. Xu, P. Hu, J. Ou and D. Yang, *Acc. Mater. Res.*, 2022, **3**, 42–53.
- 135 W. Ma, Y. Zhan, Y. Zhang, C. Mao, X. Xie and Y. Lin, *Sig. Transduct. Target Ther.*, 2021, **6**, 351.
- 136 D.-X. Wang, J. Wang, Y.-X. Wang, Y.-C. Du, Y. Huang, A.-N. Tang, Y.-X. Cui and D.-M. Kong, *Chem. Sci.*, 2021, **12**, 7602–7622.
- 137 T. Zhang, T. Tian, R. Zhou, S. Li, W. Ma, Y. Zhang, N. Liu, S. Shi, Q. Li, X. Xie, Y. Ge, M. Liu, Q. Zhang, S. Lin, X. Cai and Y. Lin, *Nat Protoc*, 2020, **15**, 2728–2757.
- 138 R. P. Goodman, R. M. Berry and A. J. Turberfield, *Chem. Commun.*, 2004, 1372.
- 139 J. A. Adaskaveg and B. Blanco-Ulate, *Curr. Opin. Biotech.*, 2023, **79**, 102872.
- 140 D. Mao, M. Zheng, W. Li, Y. Xu, C. Wang, Q. Qian, S. Li, G. Chen, X. Zhu and X. Mi, *Biosens. Bioelectron.*, 2022, **204**, 114077.
- 141 H. Li, M. Han, X. Weng, Y. Zhang and J. Li, *ACS Nano*, 2021, **15**, 1710–1717.
- 142 N. Xie, S. Liu, X. Yang, X. He, J. Huang and K. Wang, *Analyst*, 2017, **142**, 3322–3332.
- 143 L. Yu, S. Yang, Z. Liu, X. Qiu, X. Tang, S. Zhao, H. Xu, M. Gao, J. Bao, L. Zhang, D. Luo, K. Chang and M. Chen, *Materials Today Bio*, 2022, **15**, 100276.
- 144 H. Pei, N. Lu, Y. Wen, S. Song, Y. Liu, H. Yan and C. Fan, *Advanced Materials*, 2010, **22**, 4754–4758.
- 145 H. Ding, J. Li, N. Chen, X. Hu, X. Yang, L. Guo, Q. Li, X. Zuo, L. Wang, Y. Ma and C. Fan, *ACS Cent. Sci.*, 2018, **4**, 1344–1351.
- 146 L. Shan, Y. Chen, X. Tan, S. Ge, L. Zhang, L. Li, J. Yu and L. Li, *Anal. Chem.*, 2023, **95**, 4760–4767.



- 147 L. Wang, Y. Ji, Y. Chen, S. Zheng, F. Wang and C. Li, *TrAC Trends in Analytical Chemistry*, 2024, **180**, 117962.
- 148 J. Wu, H. Liu, W. Chen, B. Ma and H. Ju, *Nat Rev Bioeng*, 2023, **1**, 346–360.
- 149 N. Liu, H. Lu, L. Liu, W. Ni, Q. Yao, G.-J. Zhang and F. Yang, *Anal. Chem.*, 2021, **93**, 5917–5923.
- 150 J. Zhang, J. Zhu, F. Guo, J. Jiang, M. Xie, L. Hao and J. Chao, *Chem. Commun.*, 2023, **59**, 6869–6872.
- 151 H. Chai, Y. Tang and P. Miao, *Anal. Chem.*, 2022, **94**, 9975–9980.
- 152 Md. A. Ahamed, G. Kim, Z. Li and S.-J. Kim, *Analytica Chimica Acta*, 2022, **1236**, 340587.
- 153 J. Gao, H. Zhang and Z. Wang, *Analyst*, 2020, **145**, 3535–3542.
- 154 S. Wang, J. Guang, Y. Gao, B. Fan, Y. Liang, J. Pan, L. Li, W. Meng and F. Hu, *Microchim. Acta*, 2024, **191**, 462.
- 155 X. Wang and N. C. Seeman, *J. Am. Chem. Soc.*, 2007, **129**, 8169–8176.
- 156 Y. Wang, J. E. Mueller, B. Kemper and N. C. Seeman, *Biochemistry*, 1991, **30**, 5667–5674.
- 157 N. R. Kallenbach, R.-I. Ma and N. C. Seeman, *Nature*, 1983, **305**, 829–831.
- 158 A. Idili, A. Bonini, C. Parolo, R. Alvarez-Diduk, F. Di Francesco and A. Merkoçi, *Adv. Funct. Materials*, 2022, **32**, 2201881.
- 159 Y. Li, Y. D. Tseng, S. Y. Kwon, L. d’Espaux, J. S. Bunch, P. L. McEuen and D. Luo, *Nature Mater*, 2004, **3**, 38–42.
- 160 S. H. Um, J. B. Lee, S. Y. Kwon, Y. Li and D. Luo, *Nat Protoc*, 2006, **1**, 995–1000.
- 161 S. W. Shin, B. S. Lee, K. Yang, L. Amornkitbamrung, M. S. Jang, B. M. Ku, S.-W. Cho, J. H. Lee, H. Bae, B.-K. Oh, M.-J. Ahn, Y. T. Lim and S. H. Um, *Sci Rep*, 2017, **7**, 13499.
- 162 L. Yang, C. Yao, F. Li, Y. Dong, Z. Zhang and D. Yang, *Small*, 2018, **14**, 1800185.
- 163 Z. Zhang, Y. Liu, P. Liu, L. Yang, X. Jiang, D. Luo and D. Yang, *Nanoscale*, 2017, **9**, 19367–19373.
- 164 M. R. Hartman, D. Yang, T. N. N. Tran, K. Lee, J. S. Kahn, P. Kiatwuthinon, K. G. Yancey, O. Trotsenko, S. Minko and D. Luo, *Angew Chem Int Ed*, 2013, **52**, 8699–8702.
- 165 F. Li, Y. Dong, Z. Zhang, M. Lv, Z. Wang, X. Ruan and D. Yang, *Biosens. Bioelectron.*, 2018, **117**, 562–566.
- 166 L. Cheng, Z. Zhang, D. Zuo, W. Zhu, J. Zhang, Q. Zeng, D. Yang, M. Li and Y. Zhao, *ACS Appl. Mater. Interfaces*, 2018, **10**, 34869–34877.
- 167 M. Lin, J. Wang, G. Zhou, J. Wang, N. Wu, J. Lu, J. Gao, X. Chen, J. Shi, X. Zuo and C. Fan, *Angew Chem Int Ed*, 2015, **54**, 2151–2155.
- 168 L. Mo, Y. Hong, M. Mo, D. Liang, R. Yuan, C. Yang and W. Lin, *Sensors and Actuators B: Chemical*, 2024, **401**, 134973.
- 169 J. Deng, J. Xu, M. Ouyang, Z. Zou, Y. Lei, J. Li, Z. Qing and R. Yang, *Chinese Chemical Letters*, 2022, **33**, 773–777.
- 170 H. Wang, D. He, K. Wan, X. Sheng, H. Cheng, J. Huang, X. Zhou, X. He and K. Wang, *Analyst*, 2020, **145**, 3289–3296.
- 171 L. Luo, L. Wang, L. Zeng, Y. Wang, Y. Weng, Y. Liao, T. Chen, Y. Xia, J. Zhang and J. Chen, *Talanta*, 2020, **207**, 120298.
- 172 X. Sun, Y. Chen, H. Li, W. Xing, M. Chen, J. Wang and L. Ye, *Chem. Commun.*, 2024, **60**, 4777–4780.





- 173 J. Chen, M. Xie, M. Shi, K. Yuan, Y. Wu, H.-M. Meng, L. Qu and Z. Li, *Anal. Chem.*, 2022, **94**, 2227–2235.
- 174 S. Zhao, S. Zhang, H. Hu, Y. Cheng, K. Zou, J. Song, J. Deng, L. Li, X. Zhang, G. Ke and J. Sun, *Angew Chem Int Ed*, 2023, **62**, e202303121.
- 175 W. Wang, C. Li, S. Luo and Z.-S. Wu, *Anal. Chem.*, 2024, **96**, 7091–7100.
- 176 J. Zhao, C. Liu, Y. Li, Y. Ma, J. Deng, L. Li and J. Sun, *J. Am. Chem. Soc.*, 2020, **142**, 4996–5001.
- 177 J. H. Lee, J. A. Kim, S. Jeong and W. J. Rhee, *Biosens. Bioelectron.*, 2016, **86**, 202–210.
- 178 H. Yang, J. Zhao, J. Dong, L. Wen, Z. Hu, C. He, F. Xu, D. Huo and C. Hou, *Chem. Eng. J.*, 2022, **438**, 135594.
- 179 C. A. Mirkin, R. L. Letsinger, R. C. Mucic and J. J. Storhoff, *Nature*, 1996, **382**, 607–609.
- 180 C. A. Mirkin and S. H. Petrosko, *ACS Nano*, 2023, **17**, 16291–16307.
- 181 L. Liu, H. Lu, R. Shi, X.-X. Peng, Q. Xiang, B. Wang, Q.-Q. Wan, Y. Sun, F. Yang and G.-J. Zhang, *Anal. Chem.*, 2019, **91**, 13198–13205.
- 182 Y. Lei, X. Fei, Y. Ding, J. Zhang, G. Zhang, L. Dong, J. Song, Y. Zhuo, W. Xue, P. Zhang and C. Yang, *Sci. Adv.*, 2023, **9**, eadi1556.
- 183 Q.-Q. Wu, Y.-Y. Wu, Y.-W. Liang, H.-Y. Yang, J. Xie, M.-M. Li, B.-P. Xie, J. Chen and W.-J. Duan, *Sens. Actuat. B-Chem.*, 2024, **418**, 136298.
- 184 R. Abolhasan, A. Mehdizadeh, M. R. Rashidi, L. Aghebati-Maleki and M. Yousefi, *Biosensors and Bioelectronics*, 2019, **129**, 164–174.
- 185 J. Huang, X. Yang, X. He, K. Wang, J. Liu, H. Shi, Q. Wang, Q. Guo and D. He, *TrAC Trends in Analytical Chemistry*, 2014, **53**, 11–20.
- 186 X. Liu, M. Zou, D. Li, R. Yuan and Y. Xiang, *Analytica Chimica Acta*, 2019, **1076**, 138–143.
- 187 S. Tyagi and F. R. Kramer, *Nat Biotechnol*, 1996, **14**, 303–308.
- 188 M. Xie, F. Wang, J. Yang, Y. Guo, F. Ding, X. Lu, Y. Huang, Y. Li, X. Zhu and C. Zhang, *Anal. Chem.*, 2022, **94**, 13043–13051.
- 189 S. Bi, S. Yue and S. Zhang, *Chem. Soc. Rev.*, 2017, **46**, 4281–4298.
- 190 R. M. Dirks and N. A. Pierce, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 15275–15278.
- 191 Y. Zhang, J. Chen, H. Yang, W. Yin, C. Li, Y. Xu, S.-Y. Liu, Z. Dai and X. Zou, *Anal. Chem.*, 2022, **94**, 9665–9673.
- 192 D. He, L. Hai, H. Wang, R. Wu and H.-W. Li, *Analyst*, 2018, **143**, 813–816.
- 193 J. Kim, J. S. Shim, B. H. Han, H. J. Kim, J. Park, I.-J. Cho, S. G. Kang, J. Y. Kang, K. W. Bong and N. Choi, *Biosensors and Bioelectronics*, 2021, **192**, 113504.
- 194 E. S. Lee, J. Woo, J. Shin, B. S. Cha, S. Kim and K. S. Park, *Biosens. Bioelectron.*, 2024, **250**, 116055.
- 195 Y. Li, X. Tang, R. Deng, L. Feng, S. Xie, M. Chen, J. Zheng and K. Chang, *ACS Omega*, 2024, **9**, 19723–19731.
- 196 Y. Xu, X. Li, C. Niu, H. Wu, Y. Yong, C. Qi, W. Gong, H. Bai, Y. Chen, S. Ding and P. Liao, *Biosens. Bioelectron.*, 2022, **212**, 114405.
- 197 Z. Yu, Y. Zheng, H. Cai, S. Li, G. Liu, W. Kou, C. Yang, S. Cao, L. Chen, X. Liu, Z. Wan, N. Zhang, X. Li, G. Cui, Y. Chang, Y. Huang, H. Lv and T. Feng, *Sci. Adv.*, 2024,



**10**, ead16442.

198 L.-Y. Zhai, M.-X. Li, W.-L. Pan, Y. Chen, M.-M. Li, J.-X. Pang, L. Zheng, J.-X. Chen and W.-J. Duan, *ACS Appl. Mater. Interfaces*, 2018, **10**, 39478–39486.

199 S. Jiang, Q. Li, C. Wang, Y. Pang, Z. Sun and R. Xiao, *ACS Sens.*, 2021, **6**, 852–862.

200 D. He, H. Wang, S.-L. Ho, H.-N. Chan, L. Hai, X. He, K. Wang and H.-W. Li, *Theranostics*, 2019, **9**, 4494–4507.

201 R.-Y. Zhang, S.-H. Luo, X.-M. Lin, X.-M. Hu, Y. Zhang, X.-H. Zhang, C.-M. Wu, L. Zheng and Q. Wang, *Anal. Chim. Acta*, 2021, **1157**, 338396.



[View Article Online](#)

DOI: 10.1039/D4SD00373J

No primary research results, software or code have been included and no new data were

generated or analysed as part of this review.

