



Cite this: *Environ. Sci.: Adv.*, 2024, 3, 1271

Stormwater alters the resistome of urban surface water, an impact that can be mitigated by green stormwater infrastructure†

Kassidy O'Malley,  Walter McDonald and Patrick McNamara *

Antibiotic resistance poses an escalating threat to global health, with environmental reservoirs being pivotal areas of concern, as well as opportunities for potential mitigation. Stormwater systems are an important type of environmental reservoir in the urban water cycle with a dearth of research related to impacts on antibiotic resistance. In particular, there has been limited research exploring the impact of diverse antibiotic resistance genes (ARGs) carried by stormwater from various land uses on surface water, nor has there been an examination of the role played by green stormwater infrastructure (GSI) in mitigating this impact. Therefore, this study sought to elucidate the variability of ARGs across diverse land uses and evaluate the efficacy of GSI in mitigating ARG dissemination. Five distinct stormwater samples—representing mixed, residential, urban, and GSI-treated effluents—were taken to assess variations in ARG resistomes based on land use types. The ARGs in stormwater collected from different land uses were found to be similar in composition and represent a similar level of diversity. A GSI system with a rock swale and bioretention cell connected in series, was also sampled to see how GSI impacted ARGs, and this GSI system did substantially alter the diversity of ARGs. Moreover, the bioretention cell was found to reduce ARG concentrations by 30%. This research also sought to assess the impact of all five stormwater samples on the resistome of surface water *via* lab-scale microcosm experiments. The urban and residential stormwater significantly ($p < 0.05$) altered the resistome of surface water, while the mixed-land use sample did not. This finding underscored stormwater's pivotal role in introducing distinct ARG resistome compositions into downstream waters, heightening the chances for development of antibiotic resistant bacteria. The effluent stormwater from the GSI system, however, had less of an impact on the resistome of surface water in the microcosm experiments in comparison to the influent (untreated) stormwater. In managing stormwater runoff through GSI systems, this study's findings highlight the potential of GSI designs and practices to limit the dissemination of diverse and abundant ARGs, safeguard public health, and contribute to sustainable stormwater management by minimizing the impact on downstream surface waters.

Received 10th April 2024
Accepted 10th July 2024

DOI: 10.1039/d4va00111g

rsc.li/esadvances

Environmental significance

The study addresses the pressing issue of antibiotic resistance gene (ARG) dissemination through stormwater runoff into surface water bodies. Understanding this phenomenon is crucial due to the potential risks posed by the emergence of antibiotic resistant bacteria in aquatic ecosystems. The key finding reveals that stormwater, especially from residential and urban areas, significantly impacts surface water resistomes, potentially leading to long-term consequences for environmental and public health. This underscores the urgency of implementing effective mitigation strategies, such as green stormwater infrastructure (GSI), to reduce ARG dissemination. Moreover, this work reveals that GSI systems, particularly bioretention cells, can reduce ARG concentrations and alter stormwater's resistome, ultimately lessening the immediate risk posed to aquatic ecosystems.

1. Introduction

Antibiotic resistance is a public health crisis that presents a growing concern in the environment, due to the interconnectedness of environmental and human health.¹ The environment is a reservoir for antibiotic resistance genes (ARGs) and can contribute to the development of resistance in environmental and pathogenic bacteria.^{2–4} Stormwater has recently been

Department of Civil, Construction, and Environmental Engineering, Marquette University, 1637 W Wisconsin Ave, Milwaukee, WI 53233, USA. E-mail: patrick.mcnamara@marquette.edu; Tel: +414 288-2188

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4va00111g>



identified as an area of concern for the environmental occurrence of antibiotic resistance,^{5–7} due to its capacity to transport ARGs from various sources, such as urban soil environments and impervious surfaces, into downstream environments.^{8–10} Moreover, stormwater contains concentrated and diverse ARGs,^{11,12} in addition to other resistance elements, specifically mobile genetic elements (MGEs), and selective pressures (*e.g.*, heavy metals) that can contribute to resistant bacteria propagation.^{8,13,14}

Stormwater is a concern for antibiotic resistance in the environment particularly for the role it plays in spreading ARGs across an urban environment.⁹ As stormwater runs off of urban surfaces, it can accumulate a unique composition of ARGs and subsequently deposit them into downstream urban surface waters.^{8,15} Urban surface waters, as such, can serve as a critical area in the public health crisis of antibiotic resistance, acting as dynamic environments where the convergence of human, environmental, and bacterial interactions heighten the risk and dissemination of antibiotic resistance.¹⁶ However, a notable gap in the literature is the downstream impact that stormwater has on the antibiotic resistance profile of receiving surface waters. This knowledge gap hinders a comprehensive understanding of the environmental implications of ARGs in stormwater and the subsequent risks they pose to aquatic ecosystems. In understanding the risks stormwater poses as it mixes with urban surface waters, specific management strategies can be proposed on how stormwater should be managed to mitigate antibiotic resistance. Green stormwater infrastructure (GSI) is a specific management system that can be utilized to mitigate the transport of pollutants *via* stormwater runoff in urban settings.¹⁷ These systems are designed to capture and filter stormwater, removing pollutants before transport to downstream waters.¹⁸ As it relates to ARGs, GSI systems have very scarcely been investigated for their ability to remove ARGs from stormwater runoff. In one field-scale biofilter, a 0.9 and 2.5-log reduction in the ARGs *sul1* and *ermB* was achieved, respectively.¹⁹ Consequently, it is possible that GSI systems play a role in shifting stormwater's resistome prior to mixing with urban surface waters.

Stormwater is a further concern for antibiotic resistance in the environment due to the concentration and diversity of elements that contribute to the development of resistance in microbial communities.^{5,15,20} ARGs, MGEs, and selective pressures in stormwater, however, are not homogenous and exhibit significant variations across geographical locations.²¹ These variations lead to distinctions and region-specific pressures on the microbial communities in stormwater, differentially influencing antibiotic resistance propagation.²² Stormwater has thus been found to vary in ARG concentrations across land uses.^{12,23} Specifically, stormwater from residential areas was found to harbor a higher concentration of ARGs compared to commercial areas.²³ Moreover, specific land uses have been found to result in the occurrence of distinct ARGs, such as the *bla*_{SHV-02} ARG which was only detected in samples collected from a campus setting.¹²

Research has not clarified the diversity of ARGs in stormwater across land uses. Diversity, which indicates the variety and composition of ARGs in an environment, is crucial to providing insights into the complexity and potential evolution of antibiotic resistance.^{24,25} By understanding these variations,

a deeper comprehension of environmental resistomes and the specific environmental factors shaping them can be gained. Such knowledge is essential for predicting resistance patterns and informing strategies to mitigate the risks associated with antibiotic resistance dissemination in various ecosystems.

The aim of this research was to elucidate how the dissemination of stormwater's resistome impacts the resistome of urban surface water. The impact of stormwater from different land uses was evaluated, as well as stormwater before and after treatment in a GSI system. With this knowledge, effective strategies can be developed for managing and mitigating the potential adverse effects of stormwater runoff on downstream surface water. Specifically in this study, microcosms were conducted in which surface water was combined with five different stormwater samples and monitored for five days. The stormwater samples included runoff collected at an outfall of a mixed, residential, and urban land use as well as stormwater from the effluent of a GSI rock swale and bioretention cell. The specific objectives of this research were to: (1) determine how the diversity of ARGs in stormwater varies across land uses, (2) investigate the impact of stormwater from varying land uses on the resistome of a receiving surface water, and (3) determine how a GSI bioretention cell changes the impact of stormwater on the resistome of urban surface water.

2. Methods

2.1. Sampling locations and events

One surface water and five stormwater locations were targeted for sample collection for the microcosm experiments (Fig. S1†). The surface water sample was taken downstream of all stormwater locations at the convergence of the Menomonee, Milwaukee, and Kinnikinic Rivers in Lake Michigan. The first sampling location was a stormwater outfall in Wauwatosa, WI, USA, that collects runoff from a roughly 1.10 km² drainage area of mixed land use. The second location was a stormwater outfall also in Wauwatosa, WI, USA that discharges to the Menominee River and collects stormwater from a 4.4 km² drainage area that is dominated by residential land use. The third sampling location was the downspout of a highway overpass that discharges into a GSI system and that collects urban runoff exclusively from a highway surface. The fourth and fifth samples were collected at sequential locations within the GSI system. The fourth location was the effluent of a rock swale and the fifth location was the effluent of a bioretention cell at the most downstream end of the GSI treatment system. Land use characteristics of each sampling location were defined using the National Land Cover Database and the City of Milwaukee's Land Use Citywide Policy Plan.²⁶ Surface water samples were collected from August 3rd through August 13th, 2023, after an antecedent dry weather period of 5 days.²⁷ Each of the five stormwater sites were then sampled from on August 14th, 2023.

2.2. Sample collection and microcosm processing

A microcosm, which combined surface water with five stormwater samples separately, was conducted to evaluate the impact of stormwater from various land use types on the downstream



surface water bacterial community and resistomes (Fig. S2†).^{28,29} Prior to the microcosm, surface water was collected for 10 days to characterize the baseline microbial community (designated as days –10 through –1). All surface water samples were grab samples collected at the same time each day and transported back to the lab and analyzed immediately. On day 0 a rain event occurred, and grab stormwater samples were collected at all five locations. Upon transport to the lab, the stormwater samples were analyzed immediately for their initial day 0 condition. Then the microcosms were assembled, in which the stormwater samples were mixed with an equal volume of surface water and placed on a shaker table (100 rpm) for 5 days at ambient temperatures. To determine the mixing ratio, streamflow data from USGS stream gages across Milwaukee County was evaluated during rainfall events. It was determined that during peak flows, the streamflow could more than quadruple during intense rainfall. Therefore, the decision to mix stormwater and surface water at a 1 : 1 ratio by volume, simulating a doubling in streamflow, was made to simplify the experimental setup and enhance reproducibility, as it provides a clear, standardized condition under which the response of surface water to stormwater introduction can be systematically monitored and analyzed. The surface water that was mixed with the stormwater was the sample collected for day –1. The day –1 surface water was kept in a refrigerator (5 °C) for 24 hours after collection and a sample was collected from that surface water (designated as surface water day 0) prior to mixing with stormwater to analyze any changes in the surface water community within the time that it was stored in the lab. Samples were collected from the mixed stormwater/surface water microcosms immediately after mixing, as well as on days 1, 3, and 5 thereafter. The duration of the experiment was set at five days based on prior research finding that ARG concentrations in surface water return to a pre-rain state five days after a rainfall event.²⁷ Therefore, samples were taken on days 1, 3, and 5 to track the evolving changes during this period. From each sample collected, water quality analyses were performed, DNA was extracted, and metagenomic sequencing was completed. Microcosm controls were also included in which surface water and each stormwater location were run by themselves to monitor any changes due to time without mixing of waters. All conditions, the mixed microcosms and controls, were run in triplicate ($n = 3$).

2.3. DNA extraction and ARG quantification

For each sample collected prior to and during the microcosm experiment, DNA was extracted by first filtering the sample through a 0.22 μm Merck Millipore Express Plus® membrane filter and then extracting the DNA from the filters *via* FastDNA Spin Kit (MP Biomedicals, Santa Ana, CA). The manufacturer's protocol was followed with the addition of three liquid nitrogen freeze–thaw cycles for cell lysis as described previously.^{29,30} Subsequently, qPCR was utilized to quantify three ARGs, *sul1*, *tetW*, and *ermF*, and the 16S rRNA gene which were selected based on their reported frequency and abundances in stormwater.⁵ The protocol followed for qPCR has been previously described.^{31–33}

2.4. Metagenomic sequencing and bioinformatics

Samples from this work were sequenced at the SeqCenter (previously Microbial Genome Sequencing Center, Pittsburgh, PA) on the Illumina NextSeq 2000 platform (151-bp paired end) at a sequencing depth of 650 Mbp. The raw metagenomic sequencing data have been deposited in the Sequence Read Archive under accession number PRJNA1026961. The sequenced reads were first quality filtered using Trimmomatic to remove adaptors and low-quality sequences.³⁴ The quality filtered reads were utilized to assign taxonomy using MetaPhlan.³⁵ Next, reads were *de novo* assembled into contigs using metaSPAdes.³⁶ After each sample had its own assembly created using metaSPAdes, these assemblies were individually annotated using resistance gene identifier (RGI) to predict ARGs.³⁷ To estimate abundances of annotated ARGs, quality-filtered reads were mapped FASTA sequences from RGI output files using Kallisto.³⁸ Following, relative gene abundance was calculated for each ARG as the mapped reads per kilo base of gene length per million total reads (RPKM).³⁹

2.5. Water quality analysis

Water quality analysis for each sample included pH, conductivity, dissolved organic carbon (DOC), metals (chromium, iron, copper, zinc, cadmium, nickel) and ions (sodium and magnesium), total phosphate, nitrate, and ammonia. In detail, pH and conductivity were measured with Thermo Scientific Orion probes (Thermo Fisher Scientific, Waltham, MA). The US EPA Method 415.3 was employed to measure dissolved organic carbon (DOC) using a TOC-VCSN analyzer (Shimadzu, Kyoto, JP). Metals and ions were quantified through inductively coupled plasma mass spectrometry (ICP-MS) as previously described.⁴⁰ Furthermore, total phosphate was determined using the Hach Phosphate Color Disc Test Kit (0–40 ppm PO_4), nitrate levels were assessed with the Hach Nitrate Color Disc Test Kit (0–10 ppm $\text{NO}_3\text{-N}$), and ammonia concentrations were quantified with the Hach TNTplus Vial Test (0.015–2.000 ppm $\text{NH}_3\text{-N}$).

2.6. Statistical analyses

All genes and water quality parameters were measured in technical triplicates from each microcosm, which were completed in experimental triplicates. Error between replicate values was calculated through the standard deviation of the mean, and statistically significant relationships across sampling locations was evaluated with one-way analysis of variance (ANOVA) with the *post hoc* Tukey's multiple comparisons test. Significant relationships were assessed at a p -value ≤ 0.05 and all analysis were completed using GraphPad Prism 7® (GraphPad Software, La Jolla, CA). Gene relative abundances were determined by dividing the gene's absolute concentration by the 16S rRNA gene absolute concentration.

Statistical analyses for metagenomic results were also performed in R (v.4.2.2). The package phyloseq (v.1.42.0) was used to calculate the alpha and beta diversities of the ARG. The metric utilized for alpha diversity was the Shannon diversity



index. Alpha diversity analysis was completed to measure the diversity in each sample individually, accounting for the number of different ARGs and their relative abundance. The metric utilized for beta diversity was the Bray–Curtis dissimilarity index. Beta diversity analysis was completed to comparatively analyze the number and relative abundance of ARGs in one sample *versus* another sample to determine the degree of similarity. Principle coordinate analysis (PCoA) was further used to ordinate and plot the beta diversity dissimilarity distances. To assess statistical significance between samples, an ANOVA test with the Tukey's post hoc multiple comparison test was applied for alpha diversity and the permutational multivariate analysis of variance (PERMANOVA) (999 permutations) test *adonis2* (vegan package v.2.6.4) was applied to the Bray–Curtis distance matrices. Both were assessed at a significance level of p -values ≤ 0.05 .

3. Results and discussion

3.1. Stormwater contains a more diverse and distinct ARG profile than surface water: grab sample comparison

Diversity analyses were completed on individual samples to define the initial condition of the stormwater and surface water collected in this work, prior to mixing. First, the number of different ARGs identified in the metagenomic datasets were quantified (Fig. 1A). Only the mixed land use stormwater sample had a greater number of uniquely identified ARGs in comparison to the surface water sample. The mixed land use was followed by the residential site in terms of the number of uniquely identified ARGs, and then the urban site had the least. Despite the lower number of uniquely identified ARGs in the

mixed and residential land use stormwater samples in comparison to the surface water sample, the shared proportion of ARGs between the surface water and all stormwater samples was relatively low (Fig. S3†). This finding carries implications for the potential unique composition that is created when stormwater is transported into downstream surface waters. Moreover, as the ARGs introduced by stormwater into surface water are new and unique, further risks arise for the development of new antibiotic resistant bacteria (ARB) and resistance mechanisms from the potential exchange and mutations of ARGs among the microbial communities in aquatic environments.⁴¹

Alpha diversity analysis indicates that stormwater harbors a greater diversity of ARGs in comparison to surface water (Fig. 1B). All stormwater samples exhibited similar diversity values, with the diversity index ranging from 3.19 to 3.45. Though the difference was minimal, the residential stormwater sample had the highest diversity, followed by the urban stormwater, and then the mixed land use stormwater. The higher diversity of ARGs in residential areas could have been the result of a multitude of sources, such as pet waste left on surfaces and lawns, as well as fecal matter contamination when transported through the storm sewer network from possible cross contamination with wastewater sources such as leakages in broken pipes or joints.^{23,42,43} Furthermore, the residential stormwater sample location had the largest drainage area, indicating that ARGs accumulated from a broad geographical region. In addition, the alpha diversity of the resistomes in the residential and urban land uses samples were greater than the diversity of the resistome in the surface water, even though the surface water had a higher number of unique ARGs (*e.g.*,




Fig. 1 (A) The number of uniquely identified ARGs identified in surface water and stormwater collected from a mixed, residential, and urban land use area and (B) Shannon alpha diversity indices.



richness). This contrasting trend suggests that the higher diversity in the stormwater samples from the residential and urban areas was driven primarily by a greater evenness of ARGs in the resistomes.

Beta diversity analysis – comparing the composition of ARGs identified in each sample – revealed statistically significant differences (PERMANOVA, $p < 0.05$) in the mixed, residential, and urban land use stormwater grab samples relative to surface water (Fig. 2). As indicated in Fig. 2, there are notably significant variations between the ARG profiles of the surface water samples collected over a period of 10 days and those of stormwater collected on Day 0. Conversely, when comparing the diversity of ARGs across the stormwater samples themselves, there is minimal difference, suggesting a relatively stable ARG composition in stormwater collected from this rainfall event. The findings of this study contradict prior research, such as the studies conducted by Ahmed *et al.*, 2018 and X. Zuo *et al.*, 2022, which identified varying concentrations of ARGs in stormwater based on different land uses. Moreover, ARG diversity has also been found to vary by land use in other environments, including soil and surface water environments.^{44–46} As such, it is likely that quantitative analysis of a limited set of ARGs may not be sufficient in drawing broad conclusions about the nature of ARGs in stormwater runoff, and also that stormwater might maintain a more uniform profile of ARGs in comparison to other environments. This conclusion is supported by previous research which found that land use had less of an impact on ARG diversity in surface water during the rainy season in comparison to the dry season. This was the result of stormwater's ability to dilute physicochemical parameters that are land use specific and can indirectly affect microbial activity and ARG diversity.⁴⁷ This work, however, only represents one stormwater event and a relatively small spatial scale. Therefore, future work should aim to assess these results over a larger temporal scale and

include additional land uses (*e.g.*, agriculture, industrial, and undisturbed). Importantly, the beta diversity analysis, when paired with the alpha diversity results, indicates that the ARGs in stormwater collected from different land uses are similar in composition and represent a similar level of diversity.

Interestingly, the diversity of ARGs in the surface water samples collected across 10 days varied significantly (Fig. S4†). A hierarchical clustering analysis was completed to determine which samples were most similar in terms of ARG diversity (Fig. S5†) and no trend was observed based on time. A statistical difference in ARG diversity was however observed between samples collected 1, 3, 4, 7, and 8 days prior to the rain event and samples collected 2, 5, 6, 9, and 10 days prior to the rain event (PERMANOVA, $p = 0.005$). Consequently, other factors, such as pollution sources (*e.g.*, fecal contamination, surface sediments, or biofilms) or upstream dynamics, may be leading to the substantial difference in ARG diversity.⁴⁸ To the authors' knowledge, no other study has sequenced a surface water's resistome daily during a dry period. Seasonal and monthly sampling has been conducted, revealing variations in ARG concentrations and diversity on account of differences in water quality and rainfall conditions.^{8,49–51} Further work will be needed to confirm what factors are influencing the diversity of ARGs in surface water over a daily period.

3.2. Stormwater collected from varying land uses have a distinct impact on the resistome of surface water: microcosm experiments

Despite the similarities in ARG composition in the original stormwater samples, the different stormwater had distinct impacts on surface water (Fig. 3). After collecting the stormwater, the samples were mixed with surface water to monitor the impact on the resistome. The mixed land use stormwater had an immediate impact on the diversity of ARGs in the




Fig. 2 PCoA results of the ARG beta diversity dissimilarity distances for the surface water samples collected for 10 days prior to a stormwater event (day –10 through day –1) and stormwater collected from three outfall locations with different land use classifications for the drainage area. PERMANOVA analysis indicates a statistically significant difference ($p = 0.014$) between the beta diversity of the surface water samples ($n = 10$) and the stormwater samples ($n = 3$).






Fig. 3 ARG diversity across the 5 day microcosm in which surface water was mixed with mixed (A), residential (B), and urban (C) land use stormwater. The diversity of ARGs in the grab surface water (day -1) and stormwater (day 0) samples are shown along with the surface water controls taken in the lab on days 0 and 5. Microcosm results indicating the diversity of ARGs in surface water immediately after, and 1, 3, and 5 days after being mixed with the respective stormwater are also shown.

surface water, as the sample collected immediately after mixing had a similar diversity of ARGs to the stormwater day 0 sample, that was statistically different (PERMANOVA, $p < 0.05$) from the surface water samples (grab and control). The significant change in the surface water's resistome diversity was not maintained throughout the 5 day experimental period as no statistical difference was observed in the ARG diversity in the surface water grab and controls samples from the mixed surface–stormwater samples (days 1, 3, and 5) (PERMANOVA, $p > 0.05$). Residential stormwater on the other hand had an immediate and steady impact on the composition of ARGs in surface water. Statistically, a significant difference (PERMANOVA, $p < 0.05$) in ARG diversity was found between the surface water samples (grab and controls) and the residential stormwater impacted surface water (immediate, day 1, 3, and 5). Finally, the urban land use did not have an immediate impact on the diversity of ARGs in surface water, $p > 0.05$ between surface water (grab and control) and the urban stormwater grab sample (day 0) and immediately mixed sample. However, a statistical difference (PERMANOVA, $p < 0.05$) was found between the surface water (grab and control) and the mixed surface–stormwater samples (days 1, 3, and 5). In addition, as can be observed in Fig. 3, the beta diversity of the surface water did vary throughout the 5 day experiment. All statistical analyses were thus performed with all surface water control samples (days 0, -1 , and 5) to incorporate this variability into the analyses.

ARG concentrations have previously been shown to increase in downstream receiving environments after rainfall events but return to a baseline condition shortly after;^{27,52,53} this is the first instance in which the diversity of ARGs in surface water was monitored for their response to stormwater from various land uses. The implications of these results are that stormwater from residential and urban areas could be having a lasting impact on the resistome of surface water, altering the profile of ARGs for some time after a stormwater event. The stormwater from the mixed-land use, while impacting the diversity of ARGs in surface water immediately, did not have an impact that lasted beyond 5 days. Further research will be needed to determine how these results apply to real-world environmental systems. This includes sampling and evaluating the duration of the

impact of residential and urban stormwater *in situ*, and assessing whether the resistome ultimately returns to a pre-rain state or if the community is permanently altered.

3.3. The role of GSI in shifting the concentration and diversity of ARGs in stormwater and in altering the impact of stormwater on surface water

Through the microcosm experiments conducted, we observed that stormwater can significantly alter the resistome of surface water in the short-term aftermath of rainfall events. Given the increasing implementation of GSI in urban settings for managing stormwater volume and quality, it is crucial to investigate its potential in mitigating such impacts.^{54,55} Thus, in this work, a bioretention cell was sampled to analyze the capability of GSI to remove ARGs and change the resistome. In addition, the stormwater effluent of GSI was similarly mixed with surface water to determine if the impact of stormwater on surface water can be lessened by passing the stormwater through GSI.

3.3.1. GSI has minimal removal capability for ARGs: grab sample comparison. The capability of a GSI system to remove ARGs was initially assessed by qPCR for the samples collected at the influent of the GSI system (*i.e.*, the urban stormwater samples), and the effluent of the treatment processes (sample locations 4 and 5 as defined in the methods). The bioretention cell was found to have an average percent removal rate of 30% for the ARGs *sul1*, *ermF*, and *tetW* (Fig. S6†). The GSI system sampled though had two treatment processes: a rock swale and a bioretention cell. The rock swale was found to increase ARG concentration by 149% and therefore the GSI displayed a net increase in ARG concentrations of 87%, even though the bioretention portion did achieve ARG removal. All ARGs quantified, *sul1*, *tetW*, and *ermF*, followed the same trend through the GSI, with no statistical differences observed in the removal rates ($p > 0.05$). In addition, the total bacterial load was quantified *via* enumerating the 16S rRNA gene. This gene displayed a net increase in concentration through both the rock swale and the bioretention cell, with an average ($n = 3$) increase through the system of 0.66-log. The purpose of the rock swale is to mitigate stormwater velocities and promote stormwater infiltration, not



remove pollutants.⁵⁶ Consequently, no change in ARG and bacterial load concentrations was expected. The increase in ARG concentrations suggests that there is a source of ARGs within the rock swale that is being mobilized as the stormwater passes through. The rock swale also had variable physicochemical parameter results (Table S1†). For instance, the concentration of sodium and magnesium increased along with pH, nitrate, TOC, and the metals cadmium, copper, and iron. Phosphorus, ammonium, and the metals chromium, zinc, and nickel decreased. Interestingly, the physiochemical changes – such as an increase in antibiotic resistance metal selecting agents, nutrients, and carbon – may be creating favorable conditions for the proliferation of ARB in the bioretention cell, therefore explaining the observed increases in ARG diversity. Following, physiochemical parameters were also analyzed through the bioretention system to monitor their removal as bioretention cells are typically designed to remove stormwater contaminants. In this work, pH, ammonium, TOC, zinc, and nickel were reduced through the bioretention cell (Table S1†). Though limited, previous research has indicated that ARGs can be removed in GSI through adsorption mechanisms. Specifically, in a field scale study, a 0.9 and 2.5-log reduction in the ARGs *sul1* and *ermB* was achieved, respectively, through biofilters.¹⁹ Removal of ARGs has also been enhanced through engineering design considerations. For instance, under a specific media composition, hydraulic loading rate, and submerged area depth a 5-log removal of ARGs was attained through bioretention cells.³⁷ The findings of this work support the hypothesis that bioretention cells can be utilized to remove ARG contaminants. This study, however, only captured one stormwater event at one bioretention site, and thus further storms and sites will need to be monitored to confirm the capabilities of the GSI system to remove ARGs.

3.3.2. GSI alters the resistome of influent stormwater: grab sample comparison. The alpha diversity of ARGs was also monitored through the GSI system and was found to vary through the bioretention treatment process (Fig. 4). The largest shift was in the number of uniquely identified ARGs; 196 ARGs were observed in the effluent of the bioretention that were not found in the influent (Fig. 4A). Among these, 51 carbapenem resistance genes, 33 macrolide resistance genes, and 21 tetracycline resistance genes were identified (Fig. S7 and S8†). Carbapenems are often considered as last resort antibiotics for treating severe bacterial infections, and their resistance poses a significant threat to public health, limiting treatment options for serious infections.⁵⁸ Macrolides and tetracyclines on the other hand are widely used antibiotics, and resistance to these drugs can compromise the effectiveness of common treatment regimens. These ARGs were only found in the effluent of the GSI system, indicating they likely originated from within the GSI treatment system and subsequently were mobilized as the stormwater passed through the GSI system. Furthermore, 82 ARGs were found in the influent sample, but not the effluent, supporting the qPCR results of some ARG removal through the GSI treatment system (Fig. S6 and S7†). Further analysis of the ARGs present in both the influent and effluent samples was completed and it was determined that in the influent *lpeB*, *vatB*,

CBP-1, *cdeA*, *lpsB*, *smeE*, and *taeA* (91 – 1719 RPKM) dominated in terms of relative abundance. In contrast, the most dominant ARGs in the GSI effluent were ACC-1d, GRD33-1, and *oqxB* (19 – 51 RPKM). Further, some ARGs exhibited significant shifts in their relative abundance; the ARGs that decreased most significantly in relative abundance through the system were *vanY*, *vatB*, *CBP-1*, *cdeA*, *lpsB*, *smeE*, and *taeA* (93.2–99.0%) while the ARGs that increased most significantly in relative abundance through the system were GRD33-1 and *oqxB* (406–1286%). In general, the reduction in ARG relative concentration from the influent to effluent was much more common and larger in magnitude compared to the ARGs that increased through the GSI. These findings provide further insights into the ability of GSI to reduce the presence of ARGs from stormwater. In analyzing ARG alpha Shannon diversity, an inverse trend was observed in which both factors were less in the effluent in comparison to the influent (Fig. 4B). Such a result suggests that, while a number of different ARGs are being mobilized into the effluent, their relative abundance is minimal, having little impact on the overall resistome diversity.

3.3.3. GSI limits the change in ARG diversity in downstream surface water resistomes by stormwater: microcosm experiments. When stormwater from the GSI system was mixed with surface water, different trends in ARG diversity were observed over time (Fig. 5). For instance, the influent of the GSI did not immediately impact the surface water resistome but over time, a significant change was observed from the surface water controls and the days 1, 3, and 5 mixed samples. The effluent of the rock swale on the other hand had an immediate and steady impact on the surface water resistome, with minimal change in ARG diversity from the immediate mixed sample to that on day 5. Moreover, similar to the influent stormwater, the diversity of the ARGs in the mixed samples were also statistically different (PERMANOVA, $p = 0.006$) from the surface water controls. The effluent of the bioretention though had a minimal impact on surface water, as there was no statistical difference ($p > 0.05$) in ARG beta diversity between the stormwater mixed with surface water and the surface water controls. This result suggests that GSI can not only alter the resistome of stormwater but does so in such a way that the impact to surface water is lessened.

Metagenomic ARG annotation has revealed that, within the bioretention cell, the fate of ARGs includes both removal (ARGs in influent and not in effluent) and release (ARGs in effluent and not in influent). Removal is likely due to soil adsorption, whereas release might stem from factors like surface competition and desorption, findings consistent with observations related to other bacterial contaminants in GSI such as *E. coli* and fecal indicator bacteria.^{59–62} It is also a possibility that these ARGs, which are subject to mobilization from the soil, may have been formed within the GSI, potentially due to the presence of metals or other selective pressures in the soil. This suggests that ARGs don't solely originate from stormwater inputs but could also be locally generated within the GSI environment. Crucially, when mixed with surface water, the effluent from the GSI system had less of a prolonged impact compared to the influent. Consequently, even though there is mobilization within the bioretention cell and a subsequent rise in the number of ARGs






Fig. 4 Change in ARG count (A) and Shannon alpha diversity (B) through a GSI system, at the influent, and effluent of a sequential rock swale and bioretention cell.




Fig. 5 ARG diversity across the 5 day microcosm in which surface water was mixed with influent (A), rock swale effluent (B), and bioretention cell effluent (C) stormwater. The diversity of ARGs in the grab surface water (day -1) and stormwater (day 0) samples are shown along with the surface water controls taken in the lab on days 0 and 5. Also shown are the microcosm results, which indicate the diversity of ARGs in surface water immediately after, and 1, 3, and 5 days after being mixed with the respective stormwater.

detected in the effluent stormwater, the ARGs are not causing substantial effects on the downstream surface waters. To explain this result, the relative abundance of the ARGs identified was calculated as RPKM (Fig. S9[†]). The relative abundance notably declined from 8460 RPKM in the influent to 5608 RPKM in the effluent. This decrease could potentially explain the limited change in ARG diversity in the surface water after its mixing with the effluent stormwater. Such findings underscore the potential of GSI systems in mitigating the proliferation of ARB and highlight their role in safeguarding public health. The implications of these results suggest that optimizing GSI designs and practices could further enhance their efficacy in ensuring water quality, thereby reinforcing their value in sustainable stormwater management.

4. Conclusion

In unveiling the dynamics of ARGs in stormwater, this work sought to specifically explore the impact of stormwater's resistome on the resistome of urban surface water. Initially, stormwater exhibited a richer and more diverse profile of ARGs than surface water, suggesting that the introduction of stormwater into downstream surface waters can result in a unique composition of ARGs, posing potential risks for the emergence of ARB. Furthermore, the ARG profile in stormwater varied in the number of unique ARGs based on land use types. However, the diversity of ARGs in the stormwater samples did not vary across the residential, mixed, and urban land uses, highlighting



that this geographical factor had minimal impact on shaping ARG diversity in stormwater runoff.

Stormwater originating from different land uses did lead to distinct impacts on the resistome of surface water. Interestingly, while stormwater from mixed land use areas had an immediate impact on the resistome of surface water, its influence did not last beyond its initial impact on day 0. In contrast, stormwater from residential and urban areas had a more lasting impact on the resistome of surface waters, suggesting potential long-term consequences for aquatic ecosystems. This observation only reflects one stormwater event however, and thus further research is necessary to ascertain the duration and extent of this impact, and whether these changes have a long-lasting effect on surface water's resistome.

Additionally, this study delved into the role of GSI in modulating the diversity and concentration of ARGs in stormwater runoff. Results indicated that, within the GSI system monitored, the rock swale process promoted the dissemination of ARGs while the bioretention cell reduced ARG concentrations, and through the entire system the resistome of stormwater was significantly altered. Importantly, the shifts in the resistome observed between the influent and the effluent of the GSI system led to a lessened impact on the resistome of surface water. This result suggests that bioretention cells can be effective in removing ARGs from the environment and the observed modifications in the resistome profiles indicate a crucial role for GSI in limiting the dissemination of diverse ARGs into aquatic ecosystems.

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This material is based upon work supported by the National Science Foundation Graduate Research Fellowship [grant number 2041851]. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. This work was also supported by the Lafferty Family Charitable Foundation. The assistance of graduate researcher Veronika Folvarska and undergraduate researchers Maya Adelgren, Laura Johnson, Maya Martinez-Bates, and Sasha Razin is also greatly appreciated.

References

1 United Nations Environment Programme, *Bracing for Superbugs: Strengthening Environmental Action in the One Health Response to Antimicrobial Resistance*, 2023.

- 2 R. L. Finley, P. Collignon, D. G. J. Larsson, S. A. McEwen, X.-Z. Li, W. H. Gaze, R. Reid-Smith, M. Timinouni, D. W. Graham and E. Topp, *Clin. Infect. Dis.*, 2013, **57**, 704–710.
- 3 G. D. Wright, *Curr. Opin. Microbiol.*, 2010, **13**, 589–594.
- 4 H. K. Allen, J. Donato, H. H. Wang, K. A. Cloud-Hansen, J. Davies and J. Handelsman, *Nat. Rev. Microbiol.*, 2010, **8**, 251–259.
- 5 K. O'Malley, W. McDonald and P. McNamara, *Environ. Sci.*, 2023, **9**, 2188–2212.
- 6 E. Garner, R. Benitez, E. von Wagoner, R. Sawyer, E. Schaberg, W. C. Hession, L. A. H. Krometis, B. D. Badgley and A. Pruden, *Water Res.*, 2017, **123**, 144–152.
- 7 S. Lee, M. Suits, D. Wituszynski, R. Winston, J. Martin and J. Lee, *Sci. Total Environ.*, 2020, **723**, 138033.
- 8 K. O'Malley, P. J. McNamara and W. M. McDonald, *Environ. Sci.: Adv.*, 2022, **1**, 380–390.
- 9 D. Baral, B. I. Dvorak, D. Admiraal, S. Jia, C. Zhang and X. Li, *Environ. Sci. Technol.*, 2018, **52**, 9033–9044.
- 10 S. Zhang, S. Pang, P. F. Wang, C. Wang, N. Han, B. Liu, B. Han, Y. Li and K. Anim-Larbi, *Environ. Sci. Pollut. Res.*, 2016, **23**, 9984–9992.
- 11 K. O'Malley, W. McDonald and P. McNamara, *J. Environ. Eng.*, 2022, **148**, 04022017.
- 12 X. Zuo, P. Suo, Y. Li and Q. Xu, *Environ. Pollut.*, 2022, **292**, 118470.
- 13 S. Saifur and C. M. Gardner, *Water Sci. Technol.*, 2021, **83**, 2863–2885.
- 14 L. Hou, J. Li, H. Wang, Q. Chen, J. Q. Su, M. Gad, W. Ahmed, C. P. Yu and A. Hu, *Environ. Int.*, 2022, **168**, 107457.
- 15 K. A. Hamilton, E. Garner, S. Joshi, W. Ahmed, N. Ashbolt, G. Medema and A. Pruden, *Curr. Opin. Environ. Sci. Health.*, 2020, **16**, 101–112.
- 16 S. Lee, M. Suits, D. Wituszynski, R. Winston, J. Martin and J. Lee, *Sci. Total Environ.*, 2020, **723**, 138033.
- 17 M. Deeb, P. M. Groffman, J. L. Joyner, G. Lozefski, A. Paltseva, B. Lin, K. Mania, D. L. Cao, J. McLaughlin, T. Muth, B. Prithiviraj, J. Kerwin and Z. Cheng, *Ecol. Eng.*, 2018, **125**, 68–75.
- 18 B. Bodus, K. O. Malley, G. Dieter, C. Gunawardana and W. McDonald, *Sci. Total Environ.*, 2024, **906**, 167195.
- 19 M. B. Rugh, S. B. Grant, W.-C. Hung, J. A. Jay, E. A. Parker, M. Feraud, D. Li, S. Avasarala, P. A. Holden, H. Liu, M. A. Rippey, L. C. Van De Werfhorst, T. Kefela, J. Peng, S. Shao, K. E. Graham, A. B. Boehm, S. Choi, S. K. Mohanty and Y. Cao, *Water Res.*, 2022, **219**, 118525.
- 20 W. Ahmed, K. Hamilton, S. Toze, S. Cook and D. Page, *Sci. Total Environ.*, 2019, **692**, 1304–1321.
- 21 H. J. Beck and G. F. Birch, *Water, Air, Soil Pollut.*, 2012, **223**, 1005–1015.
- 22 A. Selvakumar and M. Borst, *J. Water Health*, 2006, **4**, 109–124.
- 23 W. Ahmed, Q. Zhang, A. Lobos, J. Senkbeil, M. J. Sadowsky, V. J. Harwood, N. Saeidi, O. Marinoni and S. Ishii, *Environ. Int.*, 2018, **116**, 308–318.
- 24 J. Bengtsson-Palme, E. Kristiansson and D. G. J. Larsson, *FEMS Microbiol. Rev.*, 2018, **42**, 68–80.



- 25 S. Sengupta, M. K. Chattopadhyay and H. P. Grossart, *Front. Microbiol.*, 2013, **4**, 1–13.
- 26 D. of C. Development, *Land Use*, 2016.
- 27 N. L. R. Williams, N. Siboni, S. L. McLellan, J. Potts, P. Scanes, C. Johnson, M. James, V. McCann and J. R. Seymour, *Environ. Pollut.*, 2022, **307**, 119456.
- 28 L. K. Kimbell, E. Lou Lamartina, S. Kohls, Y. Wang, R. J. Newton and P. J. McNamara, *mSphere*, 2023, **8**(5), e00307.
- 29 K. R. Harrison, A. D. Kappell and P. J. McNamara, *Environ. Pollut.*, 2020, **257**, 113472.
- 30 K. O'Malley, P. McNamara and W. McDonald, *J. Water Health*, 2021, **19**, 885–894.
- 31 L. K. Kimbell, E. Lou Lamartina, A. D. Kappell, J. Huo, Y. Wang, R. J. Newton and P. J. McNamara, *Environ. Sci.: Water Res. Technol.*, 2021, **7**, 584.
- 32 A. D. Kappell, L. K. Kimbell, M. D. Seib, D. E. Carey, M. J. Choi, T. Kalayil, M. Fujimoto, D. H. Zitomer and P. J. McNamara, *Environ. Sci.: Water Res. Technol.*, 2018, **4**, 1783–1793.
- 33 L. K. Kimbell, A. D. Kappell and P. J. McNamara, *Environ. Sci.: Water Res. Technol.*, 2018, **4**, 1807–1818.
- 34 A. M. Bolger, M. Lohse and B. Usadel, *Bioinformatics*, 2014, **30**, 2114–2120.
- 35 A. Blanco-Míguez, F. Beghini, F. Cumbo, L. J. McIver, K. N. Thompson, M. Zolfo, P. Manghi, L. Dubois, K. D. Huang, A. M. Thomas, W. A. Nickols, G. Piccinno, E. Piperni, M. Punčochář, M. Valles-Colomer, A. Tett, F. Giordano, R. Davies, J. Wolf, S. E. Berry, T. D. Spector, E. A. Franzosa, E. Pasolli, F. Asnicar, C. Huttenhower and N. Segata, *Nat. Biotechnol.*, 2023, **41**, 1633–1644.
- 36 S. Nurk, D. Meleshko, A. Korobeynikov and P. A. Pevzner, *Genome Res.*, 2017, **27**, 824–834.
- 37 B. P. Alcock, A. R. Raphenya, T. T. Y. Lau, K. K. Tsang, M. Bouchard, A. Edalatmand, W. Huynh, A. L. V. Nguyen, A. A. Cheng, S. Liu, S. Y. Min, A. Miroshnichenko, H. K. Tran, R. E. Werfalli, J. A. Nasir, M. Oloni, D. J. Speicher, A. Florescu, B. Singh, M. Faltyn, A. Hernandez-Koutoucheva, A. N. Sharma, E. Bordeleau, A. C. Pawlowski, H. L. Zubyk, D. Dooley, E. Griffiths, F. Maguire, G. L. Winsor, R. G. Beiko, F. S. L. Brinkman, W. W. L. Hsiao, G. V. Domselaar and A. G. McArthur, *Nucleic Acids Res.*, 2020, **48**, D517–D525.
- 38 N. L. Bray, H. Pimentel, P. Melsted and L. Pachter, *Nat. Biotechnol.*, 2016, **34**, 525–527.
- 39 K. O'Malley, P. McNamara, C. W. Marshall, E. Lou LaMartina, T. Lam, N. Ali and W. McDonald, *J. Hazard. Mater.*, 2024, **469**, 133923.
- 40 E. K. Maher, K. N. O'Malley, M. E. Dollhopf, B. K. Mayer and P. J. McNamara, *Environ. Eng. Sci.*, 2020, **37**(2), 99–108.
- 41 H. Sanderson, C. Fricker, R. S. Brown, A. Majury and S. N. Liss, *Environ. Rev.*, 2016, **24**, 205–218.
- 42 S. L. McLellan, E. J. Hollis, M. M. Depas, M. Van Dyke, J. Harris and C. O. Scopel, *J. Great Lake. Res.*, 2007, **33**, 566–580.
- 43 A. K. Salmore, E. J. Hollis and S. L. McLellan, *J. Water Health*, 2006, **4**, 247–262.
- 44 J. Wu, S. Guo, H. Lin, K. Li, Z. Li, J. Wang, W. H. Gaze and J. Zou, *J. Environ. Manage.*, 2023, **344**, 118920.
- 45 P. A. Fernanda, S. Liu, T. Yuan, B. Ramalingam, J. Lu and R. Sekar, *Front. Microbiol.*, 2022, **13**, 1–17.
- 46 C. E. Sanderson, J. T. Fox, E. R. Dougherty, A. D. S. Cameron and K. A. Alexander, *Front. Microbiol.*, 2018, **9**, 1–13.
- 47 L. Zhang, L. Ji, X. Liu, X. Zhu, K. Ning and Z. Wang, *Water Res.*, 2022, **215**, 118279.
- 48 G. Reichert, S. Hilgert, J. Alexander, J. C. Rodrigues de Azevedo, T. Morck, S. Fuchs and T. Schwartz, *Sci. Total Environ.*, 2021, **768**, 144526.
- 49 H. Jiang, R. Zhou, M. Zhang, Z. Cheng, J. Li, G. Zhang, B. Chen, S. Zou and Y. Yang, *Ecotoxicol. Environ. Saf.*, 2018, **161**, 64–69.
- 50 I. Herrig, S. Fleischmann, J. Regnery, J. Wesp, G. Reifferscheid and W. Manz, *PLoS One*, 2020, **15**(4), e0232289.
- 51 C. Dang, Y. Xia, M. Zheng, T. Liu, W. Liu, Q. Chen and J. Ni, *Environ. Int.*, 2020, **136**, 105449.
- 52 R. L. Carney, M. Labbate, N. Siboni, K. A. Tagg, S. M. Mitrovic and J. R. Seymour, *Water Res.*, 2019, **167**, 115081.
- 53 A. Di Cesare, E. M. Eckert, M. Rogora and G. Corno, *Environ. Pollut.*, 2017, **226**, 473–478.
- 54 A. R. McFarland, L. Larsen, K. Yeshitela, A. N. Engida and N. G. Love, *Environ. Sci.: Water Res. Technol.*, 2019, **5**, 643–659.
- 55 M. Keeley, A. Koburger, D. P. Dolowitz, D. Medearis, D. Nickel and W. Shuster, *Environ. Manage.*, 2013, **51**, 1093–1108.
- 56 S. A. Ekka, H. Rujner, G. Leonhardt, G. T. Blecken, M. Viklander and W. F. Hunt, *J. Environ. Manage.*, 2021, **279**, 111756.
- 57 X. Zuo, Q. Xu, Y. Li and K. Zhang, *J. Hazard. Mater.*, 2022, **429**, 128336.
- 58 L. Poirel, J. D. Pitout and P. Nordmann, *Future Microbiol.*, 2007, **2**, 501–512.
- 59 S. K. Mohanty, A. A. Torkelson, H. Dodd, K. L. Nelson and A. B. Boehm, *Environ. Sci. Technol.*, 2013, **47**, 10791–10798.
- 60 B. P. Kranner, A. R. M. Nabiul Afrooz, N. J. M. Fitzgerald and A. B. Boehm, *PLoS One*, 2019, **14**, 1–23.
- 61 S. K. Mohanty, K. B. Cantrell, K. L. Nelson and A. B. Boehm, *Water Res.*, 2014, **61**, 288–296.
- 62 P. Shen, A. Deletic, C. Urich, G. I. Chandrasena and D. T. McCarthy, *Sci. Total Environ.*, 2018, **630**, 992–1002.

