

# ChemComm

Chemical Communications

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: R. H. Kern, P. L. Schmiedel, H. Schubert and L. Wesemann, *Chem. Commun.*, 2025, DOI: 10.1039/D5CC00639B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## COMMUNICATION

## Heterocycles in reactions with boradigermaallyl

Ralf H. Kern, Paul L. Schmiedel, Hartmut Schubert and Lars Wesemann\*

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Reactions of boradigermaallyl, which can also be regarded as a chloroborylene stabilized by two germylene donors, with thiophene, furan, pyridazine and 2,2'-bipyridine are presented. Insertion of the boron atom into the heterocycles is observed and the resulting heterocycles continue to react with the bis(germylene) molecule.

The chemistry of low valent boron compounds is in the case of free borylenes dominated by matrix experiments and high temperature gas-phase synthesis.<sup>1-8</sup> Preparative studies of borylene chemistry are concerned with trapping experiments with particular interest currently focused on Lewis base adducts of borylenes (examples for Lewis bases: CAAC,<sup>9</sup> CO,<sup>10</sup> isonitriles,<sup>10</sup> oxazol-2-ylidene,<sup>11</sup> silylenes<sup>12, 13</sup>).<sup>9-11, 14-20</sup> Furthermore, borylene transition metal complexes have also been studied extensively and display a unique reactivity.<sup>21-24</sup> We are interested in low valent tetrylene chemistry and have prepared intramolecular tetrylene phosphine Lewis pairs and bis(tetrylenes) of germanium and tin.<sup>25-28</sup> Stabilization of low valent boron in these chelating vicinities lead to phosphine stabilized germa- and stannaborenes and boradigermaallyl **1** (Scheme 1).<sup>28-30</sup> Chloro substituted boradigermaallyl, which can also be regarded as chloroborylene stabilized by two germylene donors, is a highly reactive molecule and shows at room temperature an insertion reaction with benzene. In this publication we present the reactions of boradigermaallyl with the organic heterocycles thiophene, furan, pyridazine and 2,2'-bipyridine.

Boron insertion into thiophenes was observed by photoinitiated transformation of a N,C-chelated boron compound bearing at boron a benzothienyl substituent in 2-position **A** to yield a 1,2-

benzothiaborinine **B** derivative (Chart 1).<sup>31</sup> 2H-1,2-Thiaborinine **D** has been synthesized by Ashe, III and co-workers using a carbenoid ring expansion of 2,5-dihydro-1,2-thiaborole **C**.<sup>32</sup> Sulphur insertion into boroles was employed by Martin et al. for pentaphenylborole **E** in reaction with sulphur at 65 °C.<sup>33, 34</sup> In the case of 9-borafluorene **G** derivatives Finze, Braunschweig and co-workers incorporated sulphur under UV light irradiation.<sup>35</sup> Low valent boron chemistry has not yet been employed for insertion reactions of thiophene. However, Crimmin et al. studied the reaction of a low valent aluminium(I) complex  $[\{HC(CMeNDipp)_2\}Al]$  (Dipp = 2,6-di-*isopropylphenyl*) with thiophenes.<sup>36, 37</sup> At 60 °C formation of a mixture of ring-expansion products, insertion of one and two aluminium fragments, and desulphurisation was observed.<sup>36</sup>




Chart 1. Syntheses of 1,2-thiaborinine derivatives and splitting of NN-double bonds by electrophilic boron compounds.


Boradigermaallyl **1** was treated with 1.2 equiv. of thiophene in pentane and shows an immediate reaction, which is documented by a spontaneous colour change from turquoise to almost colorless (Scheme 1). The solution was stirred for another four days at 40 °C to ensure complete conversion. The reaction product **2** was isolated at -38 °C as a colorless solid in a yield of 70%. The molecular structure (Figure 1, left top)

Institut für Anorganische Chemie, Eberhard Karls Universität Tübingen, Auf der Morgenstelle 18, 72076 Tübingen, Germany.

Electronic supplementary information (ESI) available. CCDC 2420519-2420522. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/x0xx00000x



features the insertion of the boron atom into a C-S bond of the thiophene ring. The Cl-Ge-B unit could be the product of an oxidative addition of the B-Cl moiety at a germylene. Finally, the other germylene unit exhibits a connectivity to the six membered SBC<sub>4</sub> ring, which could be the result of a (1+4) cycloaddition between germylene and diene moiety of the thiaborinine cycle. The <sup>11</sup>B NMR signal of **2** was observed as a very broad signal at 83 ppm and can be compared with the signal for the 1,2-thiaborinine **F** at 50.8 ppm.<sup>33</sup> Because of the missing electronic delocalization of a boron neighboring double bond like in **F** the <sup>11</sup>B NMR signal of **2** was observed at much higher frequency. The B-S bond length of 1.778(3) Å can be compared with B-S distances observed in **H** [1.775(7)–1.796(7) Å], **F** [1.7934(17) Å], and Mes<sub>2</sub>BMe [1.787(6) Å] which exhibit partial B-S double bond character.<sup>33, 35, 38</sup> The found B-Ge1 distance of 2.060(3) Å is similar to a single bond between these elements.<sup>29</sup> The homologous furan was also treated with boradigermaallyl (Scheme 1). In the product compound **3** a BCl unit is inserted into a C-O bond of the furan ring and furthermore a germylene exhibits an oxidative addition with the C-O moiety of the oxaborinine to give a seven membered oxagermaborin ring compound. Finally, a (1+4) cycloaddition of the diene unit with another germylene was observed (Figure 1). 1,2-Oxaborinines have been synthesized by ring closure treating 2-methoxy-3,3'-bithiophene with BCl<sub>3</sub>, the carbenoid ring-expansion route starting with 1,2-oxaborol and oxygen insertion into boroles.<sup>39–41</sup> To date oxaborinines have been synthesized via insertion of a CO unit of aldehydes, ketones or ketenes into boroles and boriranes.<sup>34, 42–45</sup> In **3** a B-O bond length of 1.332(2) Å was found which is slightly shorter than B-O distances observed in oxaborinines [B-O 1.384(6), 1.380(2) Å].<sup>40, 41</sup> The Ge-O bond in **3** [1.8218(12) Å] is comparable with a Ge-O single bond.<sup>46</sup>



Scheme 1. Boradigermaallyl **1** in reactions with heterocycles thiophene, furan, pyridazine, and 2,2'-bipyridine.

Reactions of 1,2-diazenes such as pyridazine and benzo[*c*]cinnoline with Lewis acidic diborane(4) compounds, *o*-phenylbisborate and 9,10-dihydro-9,10-diboraanthracene lead to the formation of cyclic BN adducts.<sup>47–51</sup> Yamashita, Lin and co-workers explored the reactivity of tetra(*o*-tolyl)diborane(4) (Chart 1, **I**) and reported adduct formation with pyridazine and a N=N-splitting reaction towards phthalazine to give 1,2-bis(borylimidoyl)benzene (**J**).<sup>51</sup> Splitting of a N=N-bond and insertion of a boron atom was presented by Martin and co-worker treating pentaphenylborole **E** with azobenzene derivatives under photolytic conditions to give a BN<sub>2</sub>C<sub>4</sub> seven membered boracycle **K**.<sup>52</sup> In the case of the benzo[*c*]cinnoline reaction with pentaphenylborole insertion of a PhB unit into the N-N bond was observed to give a bicyclic molecule with a 1,3,2-diazaborepin moiety.<sup>52</sup> Boradigermaallyl was treated with three equivalents of pyridazine in cyclohexane at room temperature. The color of the solution changed after addition of pyridazine from turquoise to orange-red in a period of 15 minutes. To complete the conversion the solution has been stirred for another three days to yield a microcrystalline yellow-orange solid of **4** (Scheme 1). Crystals suitable for X-ray diffraction were obtained from toluene (Figure 1).




Figure 1. ORTEPs of the molecular structures of **2**, **3**, **4** and **5**. Ellipsoids set at 50% probability. Selected interatomic distances (Å) and angles (°). Trip units and hydrogen atoms of the substituents have been omitted. **2**: Ge1-B 2.060(3), B-S 1.778(3), B-C1 1.572(3), Ge2-C1 2.015(2), Ge2-C4 2.028(2), Ge1-Cl 2.1896(6), S-C4 1.857(2), C4-C3 1.498(3), C3-C2 1.338(3), C1-C2 1.512(3), S-B-Ge1 123.8(1), S-B-C1 114.0(2), C1-B-Ge 121.1(2); **3**: B-O1 1.332(2), B-Cl 1.7986(19), B-C1 1.567(2), C1-C2 1.505(2), C2-C3 1.339(3), C3-C4 1.509(2), C4-Ge1 1.9780(16), C4-Ge2 2.0251(16), C1-Ge2 1.9994(16), Ge1-O 1.8218(12), C1-B-Cl 118.0(1), O1-B-Cl 116.1(1), O1-B-C1 125.8(2); **4**: B-N1 1.424(3), B-N2 1.434(3), B-N3 1.443(3), N3-N4 1.368(2), N1-Ge1 1.8527(17), N2-Ge2 1.8715(18), Ge1-Cl 2.2436(6), Ge2-C5 1.991(2), N1-C1 1.427(3), C1-C2 1.329(3), C2-C3 1.457(3), C3-C4 1.331(3), N2-C4 1.421(3), N3-C5 1.488(3), C5-C6 1.492(3), C6-C7 1.328(3), C7-C8 1.447(3), N4-C8 1.293(3), N1-B-N2 128.0(2), N1-B-N3 117.26(19), N2-B-N3 114.5(2), C5-N3-N4 123.6(2), B-N3-



N4 113.7(2), B-N3-C5 119.3(2), B-N1-C1 118.7(2), B-N1-Ge 118.6(1), Ge1-N1-C1 122.5(1), C4-N2-Ge2 118.6(1), B-N2-Ge2 108.2(1), C4-N2-B 119.4(2); **5**: B-N1 1.402(3), N1-C5 1.437(3), C4-C5 1.496(3), C3-C4 1.520(3), C2-C3 1.330(3), C1-C2 1.521(3), C1-B 1.558(3), N1-Ge1 1.8455(16), Ge1-N2 1.8721(18), N2-C6 1.429(3), C5-C6 1.361(3), C6-C7 1.448(3), C7-C8 1.349(3), C8-C9 1.444(4), C9-C10 1.343(4), N2-C10 1.373(3), Ge2-C1 1.9902(19), Ge2-C4 2.0064(19), B-Cl 1.804(2), N1-Ge1-N2 88.5(1), Cl-B-N1 116.2(2), Cl-B-C1 120.9(2), C1-B-N1 122.9(2).

Due to strong broadening, an  $^{11}\text{B}$  NMR signal for **4** could not be observed in the  $^{11}\text{B}$  NMR spectrum. The NMR signals for the CH unit of C5 (Figure 1) were observed at  $^1\text{H}$  NMR 4.23–4.27 and  $^{13}\text{C}\{^1\text{H}\}$  NMR 47.3 ppm indicating an increase of the coordination number. The boron atom in **4** is trigonal planar coordinated by three nitrogen atoms (N1 - N3), showing a slight pyramidalization ( $\Sigma$ angles N1: 359.8, N2: 346.2, N3: 356.6°), and the BN-distances [1.424(3), 1.434(3), 1.443(3) Å] are comparable with  $\text{BN}_3$  motifs, which exhibit delocalization of the electron pairs at nitrogen into the empty p-orbital at the boron atom.<sup>53–56</sup> The found Ge-N bond lengths [1.8527(17), 1.8715(18) Å] are similar to single bonds between these elements.<sup>57</sup> The structure of the 1,3,2-diazaboropin moiety in **4** can be compared with the diazaboropin structures published by Martin et al.<sup>52</sup> Interestingly, Martin did not observe an NN bond cleavage in the reaction of unsubstituted pyridazine with pentaphenylborol, even at elevated temperatures. In a presumed mechanistic sequence for the formation of **4** (see SI) an adduct between the pyridazine nitrogen atoms and the germanium atoms could be the first step. Due to Ge-N adduct formation the borylene electron pair should be localized to a higher degree at the boron atom resulting in an increase of reactivity at boron and formation of a diazaboriridine.<sup>58</sup> The chloride atom at the boron atom is substituted by a second equivalent of pyridazine followed by chloride germanium bond formation. The N-N bond opens, and the other germanium atom reacts as a nucleophile with the carbon atom of the coordinated, neighbored pyridazine to form a Ge-C bond.

2,2'-Bipyridine shows a redox reaction with low valent boron compounds like borylene transition metal complexes and abstracts an RB unit under formation of BN bonds.<sup>59</sup> To abstract the BCl unit from boradigermaallyl **1** the reaction with 2,2'-bipyridine was investigated. After 2 days stirring at room temperature and filtration red to purple-coloured crystals were obtained from hexane (Scheme 1). In contrast to the borylene abstraction from metal borylene complexes we observed in reaction of the allyl-compound **1** with 2,2'-bipyridine an insertion of the BCl unit into a C-N bond and a hitherto unknown redox reaction of the bipyridine with a diarylgermylene moiety. Interestingly, the bis(germylene), boradigermaallyl **1** without BCl unit, does not react with 2,2'-bipy. The bipyridine is reduced to the dianion and exhibits two Ge-N bonds [1.8455(16), 1.8721(18) Å] with distances lying in the range of anionic amide coordination at germanium.<sup>57, 60</sup> Longer Ge-N bond lengths were observed for bipyridine coordination at  $\text{GeCl}_2$  [2.0738(16), 2.0637(15) Å].<sup>61</sup> The bond length between the two carbon

atoms C5-C6, connecting the two pyridine units, of 1.361(3) Å in **5** is shorter than the corresponding C-C bond in [bipyGeCl<sub>2</sub>] [1.474(3) Å] and comparable with the reduced bipy in DurB(bipy) [1.387(3) Å] (Dur = 2,3,5,6-tetramethylphenyl).<sup>59, 61</sup> The B-N interatomic distance of 1.402(3) Å in **5** is comparable with compounds exhibiting BN moieties conjugated with unsaturated organic units.<sup>62</sup> The  $^1\text{H}$  NMR signals of the six membered  $\text{C}_5\text{N}$  cycle were found in the range of 4.29–5.59 ppm, and are thus at considerably lower frequencies than those observed for the  $^1\text{H}$  NMR signals of [bipyGeCl<sub>2</sub>] (7.51–8.84 ppm).<sup>61</sup> The signal in the  $^{11}\text{B}$  NMR for compound **5** was found at 40.9 ppm and is similar to the resonance observed for PhBCl[N(H)Dipp] at 40 ppm.<sup>63</sup>

In conclusion, as a highly reactive compound boradigermaallyl shows insertion reactions with thiophene, furan, pyridazine and 2,2'-bipyridine. The low valent germylene units participate in the reactions, leading to the formation of products of an oxidative addition of a BCl and B-O unit, as well as a (1+4) cycloaddition with an organic diene. Pyridazine adds a boron atom between the two nitrogen atoms to yield a seven membered ring compound. 2,2'-Bipyridine does not abstract a BCl unit into the NN-pocket. Instead bipy inserts a BCl unit into a CN-bond and furthermore shows a previously unknown redox reaction with a diarylgermylene to give a Ge(IV)diamide and a reduced bipy derivative. With this exciting reactivity study of boradigermaallyl towards heterocycles we are looking forward to experiments with further groups of compounds.

L. W. thanks the German research foundation (DFG) for a grant (WE 1876/16-1).

## Author contributions

R. H. K. experiments, writing of manuscript and ESI, P. L. S. synthesis of **2**, H. S. crystallographic investigations, L. W. supervision, funding acquisition, manuscript writing and review.

## Data availability

Full experimental details are provided as part of the ESI.

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- B. Xu, H. Beckers, H. Ye, Y. Lu, J. Cheng, X. Wang and S. Riedel, *Angew. Chem., Int. Ed.*, 2021, **60**, 17205–17210.
- H. F. Bettinger, *J. Am. Chem. Soc.*, 2006, **128**, 2534–2535.
- P. L. Timms, *Acc. Chem. Res.*, 1973, **6**, 118–123.
- R. W. Kirk and P. L. Timms, *J. Am. Chem. Soc.*, 1969, **91**, 6315–6318.
- P. L. Timms, *J. Am. Chem. Soc.*, 1968, **90**, 4585–4589.
- P. L. Timms, *J. Am. Chem. Soc.*, 1967, **89**, 1629–1632.
- J. A. J. Pardoe, N. C. Norman and P. L. Timms, *Polyhedron*, 2002, **21**, 543–548.



8. C. R. Brazier, *J. Mol. Spectrosc.*, 1996, **177**, 90-105.
9. R. Kinjo, B. Donnadiou, M. A. Celik, G. Frenking and G. Bertrand, *Science*, 2011, **333**, 610-613.
10. H. Braunschweig, R. D. Dewhurst, F. Hupp, M. Nutz, K. Radacki, C. W. Tate, A. Vargas and Q. Ye, *Nature*, 2015, **522**, 327-330.
11. L. Kong, Y. Li, R. Ganguly, D. Vidovic and R. Kinjo, *Angew. Chem., Int. Ed.*, 2014, **53**, 9280-9283.
12. H. Wang, L. Wu, Z. Lin and Z. Xie, *Angew. Chem., Int. Ed.*, 2018, **57**, 8708-8713.
13. H. Wang, L. Wu, Z. Lin and Z. Xie, *J. Am. Chem. Soc.*, 2017, **139**, 13680-13683.
14. B. Pachaly and R. West, *Angew. Chem., Int. Ed.*, 1984, **23**, 454-455.
15. M. Ito, N. Tokitoh, T. Kawashima and R. Okazaki, *Tetrahedron Lett.*, 1999, **40**, 5557-5560.
16. A. Meller, U. Seebold, W. Maringgele, M. Noltemeyer and G. M. Sheldrick, *J. Am. Chem. Soc.*, 1989, **111**, 8299-8300.
17. P. Bissinger, H. Braunschweig, K. Kraft and T. Kupfer, *Angew. Chem., Int. Ed.*, 2011, **50**, 4704-4707.
18. W. J. Grigsby and P. P. Power, *J. Am. Chem. Soc.*, 1996, **118**, 7981-7988.
19. F. Dahchek, D. Martin, D. W. Stephan and G. Bertrand, *Angew. Chem. Int. Ed.*, 2014, **53**, 13159-13163.
20. A. D. Ledet and T. W. Hudnall, *Dalton Trans.*, 2016, **45**, 9820-9826.
21. H. Braunschweig, C. Kollann and D. Rais, *Angew. Chem., Int. Ed.*, 2006, **45**, 5254-5274.
22. H. Braunschweig, R. D. Dewhurst and A. Schneider, *Chem. Rev.*, 2010, **110**, 3924-3957.
23. H. Braunschweig, R. D. Dewhurst and V. H. Gessner, *Chem. Soc. Rev.*, 2013, **42**, 3197-3208.
24. J. T. Goettel and H. Braunschweig, *Coord. Chem. Rev.*, 2019, **380**, 184-200.
25. J. Schneider, K. M. Krebs, S. Freitag, K. Eichele, H. Schubert and L. Wesemann, *Chem.-Eur. J.*, 2016, **22**, 9812-9826.
26. J. Henning and L. Wesemann, *Angew. Chem., Int. Ed.*, 2012, **51**, 12869-12873.
27. J. Henoch, A. Auch, F. Diab, K. Eichele, H. Schubert, P. Sirsch, T. Block, R. Pöttgen and L. Wesemann, *Inorg. Chem.*, 2018, **57**, 4135-4145.
28. R. H. Kern, M. Schneider, K. Eichele, H. Schubert, H. F. Bettinger and L. Wesemann, *Angew. Chem., Int. Ed.*, 2023, **62**, e202301593.
29. D. Raiser, C. P. Sindlinger, H. Schubert and L. Wesemann, *Angew. Chem., Int. Ed.*, 2020, **59**, 3151-3155.
30. M. Zweigart, K. Eichele, H. Schubert, C. P. Sindlinger and L. Wesemann, *J. Am. Chem. Soc.*, 2023, **145**, 12452-12458.
31. S. K. Mellerup, C. Li, J. Radtke, X. Wang, Q.-S. Li and S. Wang, *Angew. Chem., Int. Ed.*, 2018, **57**, 9634-9639.
32. A. D. Rohr, M. M. Banaszak Holl, J. W. Kampf and A. J. Ashe, III, *Organometallics*, 2011, **30**, 3698-3700.
33. S. Yruegas and C. D. Martin, *Chem.-Eur. J.*, 2016, **22**, 18358-18361.
34. X. Su, J. J. Baker and C. D. Martin, *Chem. Sci.*, 2020, **11**, 126-131.
35. T. Bischof, N. Wieprecht, S. Fuchs, L. Endres, I. Krummenacher, M. Michel, C. Mihm, H. Braunschweig and M. Finze, *Inorg. Chem.*, 2023, **62**, 21329-21335.
36. J. S. McMullen, A. J. P. White and M. R. Crimmin, *Chem. Commun.*, 2023, **59**, 14681-14684.
37. L. Zhang, S. Kaukver, J. McMullen, A. J. P. White and M. R. Crimmin, *Organometallics*, 2023, **42**, 1711-1716. VIEW ARTICLE ONLINE  
CC00639B
38. M. T. Ashby and N. A. Sheshtawy, *Organometallics*, 1994, **13**, 236-243.
39. L. Menduti, C. Baldoli, S. Manetto, M. Bolte, H.-W. Lerner, G. Longhi, C. Villani, E. Licandro and M. Wagner, *Angew. Chem., Int. Ed.*, 2023, **62**, e202215468.
40. J. Chen, Z. Bajko, J. W. Kampf and A. J. Ashe, *Organometallics*, 2007, **26**, 1563-1564.
41. S. Yruegas, D. C. Patterson and C. D. Martin, *Chem. Commun.*, 2016, **52**, 6658-6661.
42. K. Huang and C. D. Martin, *Inorg. Chem.*, 2015, **54**, 1869-1875.
43. K. R. Bluer, L. E. Laperriere, A. Pujol, S. Yruegas, V. A. K. Adiraju and C. D. Martin, *Organometallics*, 2018, **37**, 2917-2927.
44. T. Bischof, L. Beßler, I. Krummenacher, L. Erhard, H. Braunschweig and M. Finze, *Chem.-Eur. J.*, 2023, **29**, e202300210.
45. Y. Wei, J. Wang, W. Yang, Z. Lin and Q. Ye, *Chem.-Eur. J.*, 2023, **29**, e202203265.
46. P. Kitschke, A. A. Auer, A. Seifert, T. Ruffer, H. Lang and M. Mehring, *Inorg. Chim. Acta*, 2014, **409**, 472-478.
47. A. Lorbach, M. Bolte, H.-W. Lerner and M. Wagner, *Chem. Commun.*, 2010, **46**, 3592-3594.
48. Z. Lu, H. Quanz, O. Burghaus, J. Hofmann, C. Logemann, S. Beeck, P. R. Schreiner and H. A. Wegner, *J. Am. Chem. Soc.*, 2017, **139**, 18488-18491.
49. J. Ruhl, N. Oberhof, A. Dreuw and H. A. Wegner, *Angew. Chem., Int. Ed.*, 2023, **62**, e202300785.
50. Y. Katsuma, H. Asakawa and M. Yamashita, *Chem. Sci.*, 2018, **9**, 1301-1310.
51. Y. Katsuma, L. Wu, Z. Lin, S. Akiyama and M. Yamashita, *Angew. Chem., Int. Ed.*, 2019, **58**, 317-321.
52. V. A. K. Adiraju and C. D. Martin, *Chem.-Eur. J.*, 2017, **23**, 11437-11444.
53. G. J. Bullen, *Dalton Trans.*, 1981, 511-514.
54. K. Huang, J. L. Dutton and C. D. Martin, *Chem.-Eur. J.*, 2017, **23**, 10532-10535.
55. M. A. Rodriguez and T. T. Borek, *Acta Cryst., Sec. E.*, 2006, **62**, o3341-o3343.
56. K. Lee, C. M. Donahue and S. R. Daly, *Dalton Trans.*, 2017, **46**, 9394-9406.
57. M. J. Evans, F. M. Burke, P. M. Chapple and J. R. Fulton, *Inorg. Chem.*, 2021, **60**, 8293-8303.
58. R. Boese and U. Klingebiel, *J. Organomet. Chem.*, 1986, **306**, 295-302.
59. S. Liu, M.-A. Légaré, J. Seufert, D. Prieschl, A. Rempel, L. Englert, T. Dellermann, V. Paprocki, A. Stoy and H. Braunschweig, *Inorg. Chem.*, 2020, **59**, 10866-10873.
60. M. F. Lappert, M. J. Slade, J. L. Atwood and M. J. Zaworotko, *Chem. Commun.*, 1980, 621-622.
61. F. Cheng, J. M. Dyke, F. Ferrante, A. L. Hector, W. Levason, G. Reid, M. Webster and W. Zhang, *Dalton Trans.*, 2010, **39**, 847-856.
62. M. Franceschini, M. Crosta, R. R. Ferreira, D. Poletto, N. Demitri, J. P. Zobel, L. González and D. Bonifazi, *J. Am. Chem. Soc.*, 2022, **144**, 21470-21484.
63. T. Chivers, C. Fedorchuk and M. Parvez, *Inorg. Chem.*, 2004, **43**, 2643-2653.



Full experimental details are provided as part of the ESI.

View Article Online  
DOI: 10.1039/D5CC00639B

Open Access Article. Published on 27 February 2025. Downloaded on 2/27/2025 10:23:51 PM.  
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

