ChemComm



View Article Online

CORRECTION

Check for updates

Cite this: DOI: 10.1039/d5cc90059j

Correction: Macrocyclic peptides as inhibitors of WDR5–lncRNA interactions

Jen-Yao Chang, Cora Neugebauer, Stefan Schmeing, Gulshan Amrahova and Peter 't Hart*

DOI: 10.1039/d5cc90059j

Correction for 'Macrocyclic peptides as inhibitors of WDR5–lncRNA interactions' by Jen-Yao Chang et al., Chem. Commun., 2023, **59**, 10656–10659, https://doi.org/10.1039/D3CC03221C.

rsc.li/chemcomm

The authors regret that the K_D values given for entries 14 and 15, in column 3 of Table 1, were incorrectly given in the original article. The authors would like to clarify the correct K_D values for entries 14 and 15 are 0.53 \pm 0.06 μ M and 5.2 \pm 0.3 μ M respectively. These errors do not affect the conclusions of the work and the updated version of Table 1 is shown below.

The authors also regret the omission of an acknowledgements section from the article at the time of publication. The acknowledgements section for the article is:

We would like to acknowledge Christiane Heitbrink (TU Dortmund) for measuring HRMS data. We acknowledge the European Synchrotron Radiation Facility for provision of synchrotron radiation facilities and we would like to thank P. Carpentier for assistance in using beamline ID30A-3. We also thank the Ministry of Education (Taiwan) for funding Jen-Yao Chang with a Government Scholarship for Overseas Study. Open Access funding provided by the Max Planck Society.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

Chemical Genomics Centre of the Max Planck, Society Max Planck Institute of Molecular Physiology, Otto-Hahn-Strasse 11, 44227 Dortmund, Germany. E-mail: peter.t-hart@mpi-dortmund.mpg.de

 Table 1
 Sequences of WBM site binding peptides and their affinity for WDR5 as determined by fluorescence polarization. Residues used for side-chain macrocyclization are highlighted in bold. See Fig. S1 (ESI) for full structural details of the peptides

_

No.	Sequence	<i>К</i> ₀ (µМ)
	371 381	
1	$FITC-PEG-EDEEVDVTSVD-NH_2$	2.05 ± 0.42
2	FITC-PEG-EDEEVDVTSV-NH2	7.11 ± 2.80
3	FITC-PEG-EDEEVDVTS-NH2	20.92 ± 8.72
4	FITC-PEG-EDEEVDVT-NH2	29.91 ± 15.13
5	Ac-EDEEVDVTSVD-PEG-FITC	2.42 ± 0.14
6	Ac-DEEVDVTSVD-PEG-FITC	6.74 ± 1.04
7	Ac-EEVDVTSVD-PEG-FITC	23.45 ± 6.89
8	Ac-EVDVTSVD-PEG-FITC	>37.5
9	Ac-VDVTSVD-PEG-FITC	>37.5
10	FITC-020c-DEEVDVTSV-NH2	10.39 ± 2.69
11	FITC-020c-DEEV D VT Dap V-NH ₂	21.86 ± 5.29
12	FITC-020c-DEEVDVTDabV-NH2	1.18 ± 0.07
13	FITC-020c-DEEV D VT Orn V-NH ₂	2.67 ± 0.10
	258 268	
14	FITC-020c-DEEEIDVVSVE-NH2	0.53 ± 0.06
	407 417	
15	FITC-020c-FSDDLDVVGDG-NH2	9.87 ± 4.78
16	FITC-020c-DEEEIDVVDabV-NH2	0.33 ± 0.02
17	FITC-020c-DEEEIDVVOrnV-NH2	1.96 ± 0.18
18	FITC-020c-DEEEIDVVDabVE-NH2	0.11 ± 0.02
19	FITC-020c-DEEEIDVVOrnVE-NH2	0.56 ± 0.04
20	FITC-020c-DEEEIDVTDabVE-NH2	0.23 ± 0.01
21	FITC-020c-DEEEIDIVDabVE-NH2	0.18 ± 0.01
22	FITC-020c-DEEEIDITDabVE-NH2	0.43 ± 0.02
23	FITC-020c-DEEEIEVVDapVE-NH2	>37.5
24	FITC-020c-DEEEI Dab VV D VE-NH ₂	>37.5
25	FITC-020c-DEEEIDapVVEVE-NH2	>37.5

8